TITLE:

Strichartz Estimates for Schrödinger Equations on Scattering Manifolds

AUTHOR(S):

Mizutani, Haruya

CITATION:

Mizutani, Haruya. Strichartz Estimates for Schrödinger Equations on Scattering Manifolds. Communications in Partial Differential Equations 2012, 37(2): 169-224

ISSUE DATE:

2012-01

URL:

<http://hdl.handle.net/2433/169704>

RIGHT:

© Taylor & Francis Group, LLC; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

STRICHARTZ ESTIMATES FOR SCHRÖDINGER EQUATIONS ON SCATTERING MANIFOLDS

HARUYA MIZUTANI

ABSTRACT. The present paper is concerned with Schrödinger equations on non-compact Riemannian manifolds with asymptotically conic ends. It is shown that, for any admissible pair (including the endpoint), local in time Strichartz estimates outside a large compact set centered at origin hold. Moreover, we prove global in space Strichartz estimates under the nontrapping condition on the metric.

1. INTRODUCTION

Let us recall that the *Strichartz estimates* for the free Schrödinger equation on the Euclidean space $\mathbb{R}^d$ ([21, 10, 23, 14]) state that

$$\|e^{i\frac{t}{2}\Delta}u_0\|_{L^p([-T,T];L^q(\mathbb{R}^d))} \leq C_T\|u_0\|_{L^2(\mathbb{R}^d)}, \quad u_0 \in L^2(\mathbb{R}^d),$$

where the pair (p, q) satisfies the following *admissible* condition:

$$(1.1) \quad \frac{2}{p} + \frac{d}{q} = \frac{d}{2}, \quad p, q \geq 2, \quad (d, p, q) \neq (2, 2, \infty).$$

It is well known that these estimates are fundamental in studying low regularity well-posedness of the Cauchy problem for nonlinear Schrödinger equations. It is a natural question if the same estimates hold for Schrödinger equations on manifolds. Though the global in time estimates (*i.e.*, $\sup_T C_T < \infty$) do not hold in general, the local in time estimates have been proved by many authors under several geometric conditions. The purpose of the present paper is to prove sharp Strichartz estimates on scattering manifolds for any admissible pair (including the endpoint $(2, \frac{2d}{d-2})$), where the scattering manifolds are non-compact manifolds which have asymptotically conic ends. The paper is also concerned with studying a relationship between Strichartz estimates and microlocal properties of the solution. More precisely, we show that (local in time) Strichartz estimates follow from the dispersive estimates for spatially and frequency localized solutions which we call *microlocal dispersive estimates*.

We consider the following model. We mean by the scattering manifold a non-compact manifold with an asymptotically conic structure. Let M be a non-compact Riemannian manifold of dimension $d \geq 2$ such that M can be decomposed as

$$M = M_c \cup M_\infty,$$

where $M_c \Subset M$ is relatively compact, and there exists a $(d-1)$ -dimensional closed Riemannian manifold ∂M such that M_∞ is diffeomorphic to $(0, \infty) \times \partial M$. Let

$$\iota : M_\infty \ni z \mapsto (r(z), \theta(z)) \in (0, \infty) \times \partial M$$

be an identification mapping which is called a boundary decomposition. Suppose that $M_c \cap M_\infty \subset (0, 1) \times \partial M$ under this identification. Throughout the paper we fix a boundary decomposition ι and do not write it explicitly, and denote local coordinates on $(0, \infty) \times \partial M$ by (r, θ) .

2000 *Mathematics Subject Classification.* Primary 35A40; Secondary 35B45.

Key words and phrases. Strichartz estimate; Microlocal dispersive estimate; Scattering manifold; Schrödinger equation.

The author was partly supported by JSPS Research Fellowships for Young Scientists 08J06914(2008–2010); GCOE ‘Fostering top leaders in mathematics’, Kyoto University.

We next recall the definition of the scattering metric on M . Let g be a Riemannian metric on M such that, for sufficiently large $R_M > 0$, g takes the form

$$(1.2) \quad g = dr^2 + r^2(h_{jk} + a_{jk})d\theta^j d\theta^k \quad \text{for } (r, \theta) \in (R_M, \infty) \times \partial M,$$

where (h_{jk}) is the Riemannian metric on ∂M and (a_{jk}) is a smooth and real-valued tensor. Here we used the Einstein summation convention. We may assume that $R_M = 1$ without loss of generality, and define the *scattering region* by $\widetilde{M}_\infty := (1, \infty) \times \partial M$. We also assume throughout the paper that there exists $\mu > 0$ such that for any $(l, \alpha) \in \mathbb{Z}_+^d$,

$$(1.3) \quad |\partial_r^l \partial_\theta^\alpha a_{jk}(r, \theta)| \leq C_{l\alpha} r^{-\mu-l}, \quad (r, \theta) \in \widetilde{M}_\infty.$$

Such a g is said to be a long-range *scattering metric* (in normal form).

Let Δ_g be the Laplace-Beltrami operator associated to g on $L^2(M)$:

$$\Delta_g = \frac{1}{G(z)} \partial_{z_i} G(z) g^{lm}(z) \partial_{z_l}, \quad (g^{lm}(z)) = (g_{lm}(z))^{-1}, \quad G(z) = \sqrt{\det g_{lm}(z)},$$

and set $P = -\frac{1}{2}\Delta_g$, where $L^p(M) := L^p(M, G(z)dz)$, $1 \leq p \leq \infty$. In the setting we consider the Schrödinger equation:

$$(1.4) \quad \begin{cases} i\partial_t u(t) = Pu(t), & t \in \mathbb{R}, \\ u(0) = u_0 \in L^2(M). \end{cases}$$

Since P is essentially self-adjoint on $C_0^\infty(M)$ under the above condition, (1.4) has a unique solution $u(t) = e^{-itP}u_0$ by the Stone theorem. The main result is the following.

Theorem 1.1 (Strichartz estimates near infinity). *There exist a large compact subset $M_c \subset K \subset M$ and $\chi_K \in C_0^\infty(M)$ with $\chi_K \equiv 1$ on K such that*

$$(1.5) \quad \|(1 - \chi_K)e^{-itP}u_0\|_{L^p([0,1];L^q(M))} \leq C\|u_0\|_{L^2(M)}, \quad u_0 \in C_0^\infty(M),$$

provided that (p, q) satisfies the admissible condition (1.1).

Let $p(z, \xi)$ be the principal symbol of P . We say that M is nontrapping if for any $(z_0, \xi^0) \in T^*M$ with $\xi^0 \neq 0$, the geodesic flow $(z(t, z_0, \xi^0), \xi(t, z_0, \xi^0))$ generated by H_p satisfies

$$|z(t, z_0, \xi^0)| \rightarrow +\infty \text{ as } t \rightarrow \pm\infty,$$

where $H_p = \sum_{j=1}^d \left(\frac{\partial p}{\partial \xi_j} \frac{\partial}{\partial z^j} - \frac{\partial p}{\partial z^j} \frac{\partial}{\partial \xi_j} \right)$ is the Hamilton vector field associated to $p(z, \xi)$. If M is nontrapping, then using local smoothing effects which follow from resolvent estimates proved by [7] (see also [8]), we obtain local in time Strichartz estimates (without loss of derivatives) for $\chi_K e^{-itP}u_0$ (see Appendix A for more details). Combining with Theorem 1.1, we have the following:

Theorem 1.2 (Global in space estimates). *Suppose that M is a nontrapping scattering manifold. Then, for any (p, q) satisfying (1.1), we have*

$$(1.6) \quad \|e^{-itP}u_0\|_{L^p([0,1];L^q(M))} \leq C\|u_0\|_{L^2(M)}, \quad u_0 \in C_0^\infty(M).$$

Remark 1.3. (i) Let $T > 0$. Since e^{-itP} is unitary on $L^2(M)$, the time interval $[0, 1]$ in (1.5) and (1.6) can be replaced by $[-T, T]$ provided that we replace the constant $C > 0$ by some $C_T > 0$ depending on T .

(ii) (Metrics in not normal form) Let g be a more general Riemannian metric than in normal form such that g takes the form

$$g = (1 + a^1)dr^2 + ra_j^2(dr d\theta^j + d\theta^j dr) + r^2(h_{jk} + a_{jk}^3)d\theta^j d\theta^k$$

on $(1, \infty) \times \partial M$, where $a^N(r, \theta)$ are smooth and real-valued tensors satisfying (1.3) with some $0 < \mu_N < 1$. Then, there exists a change of coordinates $(r', \theta') = (R(r, \theta), \Theta(r, \theta))$ such that, for some $0 < \mu < \mu_N$,

$$R(r, \theta) = r + O(r^{1-\mu}), \quad \Theta(r, \theta) = \theta + O(r^{-\mu}) \text{ as } r \rightarrow +\infty,$$

and g can be brought to a long-range metric in normal form as above. Hence the statements of Theorem 1.1 and Theorem 1.2 also hold for such a metric in not normal form. We refer to [11, Section 10.6] for more details.

(iii) (Potential perturbations) Let V be a smooth and real-valued potential on M of short-range type *i.e.*,

$$|\partial_r^l \partial_\theta^\alpha V(r, \theta)| \leq C_{k\alpha} r^{-1-\nu-l} \quad \text{for } (r, \theta) \in (1, \infty) \times \partial M,$$

with some $\nu > 0$. Our proof still works well if we replace P with $P + V$. Hence the statements of Theorem 1.1 and Theorem 1.2 are still hold for $P + V$.

The local in time Strichartz estimates on manifolds recently have been studied by many authors. Staffilani-Tataru [20], Robbiano-Zuily [17] and Bouclet-Tzvetkov [4] studied the case of Schrödinger equations on the Euclidean space with the asymptotically flat metric under several settings. In [6], Burq-Gérard-Tzvetkov proved Strichartz estimates with a loss of derivative $1/p$ on any compact manifolds without boundaries. They also proved that the loss $1/p$ is optimal in the case of $M = \mathbb{S}^d$. Hassell-Tao-Wunsch [11] considered the case of nontrapping scattering manifolds except the endpoint estimate. Our result thus is regarded as a generalization of their result to the critical exponent case, however the method of the proof is considerably different. More recently, Bouclet [1, 2, 3] studied the case of an asymptotically hyperbolic manifold which is a non-compact Riemannian manifold with the metric of the form $dr^2 + e^{2r}(h_{jk} + a_{jk})d\theta^j d\theta^k$ in the scattering region, and he proved Strichartz estimates localized near infinity without the nontrapping condition. The present paper is motivated by his works. Global in time Strichartz estimates has been studied by [5, 22, 16] in the case of Euclidean space with an asymptotically flat metric.

On the other hand, dispersive estimates for Schrödinger equations with potentials on the flat Euclidean space $(\mathbb{R}^d, \delta_{jk})$ also have been studied by many authors. In particular, it was shown by Fujiwara [9] that if $V(x)$ is smooth, real-valued and increases at most quadratically at infinity, namely $|\partial_x^\alpha V(x)| \leq C_\alpha$ if $|\alpha| \geq 2$, then the fundamental solution $E(t, x, y)$ of the propagator e^{-itP} for $P = -\frac{1}{2}\Delta + V(x)$ satisfies the dispersive estimate $|E(t, x, y)| \leq C|t|^{-d/2}$ on $\mathbb{R}^d$ provided that $t \neq 0$ is small enough. Local in time Strichartz estimates are immediate consequences of this estimate and the TT^* -argument due to [14]. Long time dispersive estimates for $e^{-itP}P_{ac}(P)$, which implies global in time Strichartz estimates, also have been proved by many authors (*e.g.*, Journé-Soffer-Sogge [13], Yajima [24, 25]) under suitable conditions of potentials and assumptions for the zero energy, where $P_{ac}(P)$ is the projection onto the absolutely continuous spectrum of P . For more references on dispersive estimates for Schrödinger equations with potentials, we refer to Schlag's survey [19].

The rest of the paper is devoted to the proof of Theorems 1.1 and 1.2. Though the proof is based on Bouclet's argument in [3], the behavior of classical trajectories at infinity $r \rightarrow +\infty$ is different from the case of an asymptotically hyperbolic manifold and the class of the phase function of the parametrix becomes even worse. We thus cannot apply straightforwardly his method to the case of a scattering manifold. To overcome this difficulty, we introduce a localization in the r -variable by using the dyadic decomposition. The proof of Theorem 1.1 is then reduced to that of microlocal dispersive estimates.

The paper is organized as follows. In Section 2 we fix notations and the pseudodifferential setup, and collect results on the functional calculus recently proved by Bouclet [1, 2]. Section 3 discusses a localization of both space and energy, and we show that Theorem 1.1 follows from microlocal dispersive estimates. We study some properties of the geodesic flow in Section 4. In Section 5 we construct the semiclassical Isozaki-Kitada parametrix and prove microlocal dispersive estimates on the strongly outgoing and incoming regions (cf. Definition 2.5). By using an Egorov type theorem, we prove microlocal smoothing properties of the propagator in Section 6 which imply microlocal dispersive estimates on intermediate regions. In Section 7, we construct the semiclassical WKB parametrix and prove short time microlocal dispersive estimates on the

outgoing and incoming regions. We complete the proof of Theorem 1.1 in Section 8. We give the sketch of the proof of Theorem 1.2 in Appendix A.

Throughout the paper we use the following notations: We denote the set of multi-indices by $\mathbb{Z}_+^d$. For Banach spaces X and Y , $\mathcal{L}(X, Y)$ denotes the Banach space of bounded operators from X to Y , and we write $\mathcal{L}(X) := \mathcal{L}(X, X)$. For $a \in \mathbb{R}$, we use the notation $a_+ = \max(a, 0)$. $\langle r \rangle$ stands for $\sqrt{1 + |r|^2}$. For $A, B \geq 0$, $A \lesssim B$ means that there exists some universal constant $C > 0$ such that $A \leq CB$.

Acknowledgments: The author would like to thank Jean-Marc Bouclet for helpful discussions and comments regarding the construction of the parametrix, in particular the spatial localization. He also would like to thank his advisor Shu Nakamura for much of comments and suggestions. He also thanks the anonymous referee for careful reading the manuscript and for giving useful comments.

2. PRELIMINARIES

In this section we set up some standard notations on scattering manifolds. Notice that a boundary decomposition is always fixed. The first step is to choose a suitable atlas and a partition of unity on the scattering region $\widetilde{M}_\infty$. Let

$$\{\kappa : \widetilde{M}_\infty \supset (1, \infty) \times V_\kappa \rightarrow (1, \infty) \times U_\kappa \subset \mathbb{R}^d\}_\kappa$$

be a finite atlas on $\widetilde{M}_\infty$ such that $\kappa = \text{Id} \otimes \kappa_b$, where $\{\kappa_b : \partial M \ni V_\kappa \rightarrow U_\kappa \subset \mathbb{R}^{d-1}\}_\kappa$ is a finite atlas on ∂M . We denote the associated pull-back and push-forward by κ^* and $\kappa_* = (\kappa^{-1})^*$, respectively. We also denote the induced chart diffeomorphism $T^*((1, \infty) \times V_\kappa) \rightarrow T^*((1, \infty) \times U_\kappa) \cong (1, \infty) \times U_\kappa \times \mathbb{R}^d$ by the same symbol κ_* if there is no confusion. Let $\{\psi_{\kappa_b}\}_\kappa \subset C_0^\infty(V_\kappa)$ be a partition of unity subordinate to $\{V_\kappa\}$ and $\psi \in C^\infty(\mathbb{R})$ a cut-off function such that $\text{supp } \psi \subset (2, \infty)$ and $\psi \equiv 1$ for $r \geq 3$. We set $\psi_\kappa := \psi \times \psi_{\kappa_b}$. Then, $\{\psi_\kappa\}_\kappa$ is a partition of unity subordinate to $\{(1, \infty) \times V_\kappa\}_\kappa$. Let $\tilde{\psi}_\kappa \in C^\infty(V_\kappa)$ be a cut-off function such that $\tilde{\psi}_\kappa$ takes the form $\tilde{\psi}_\kappa = \tilde{\psi} \times \tilde{\psi}_{\kappa_b}$, where $\tilde{\psi} \in C^\infty(\mathbb{R})$, $\tilde{\psi}_{\kappa_b} \in C_0^\infty(V_\kappa)$, $\text{supp } \tilde{\psi} \subset (3/2, \infty)$, $\tilde{\psi} \equiv 1$ on $(2, \infty)$ and $\tilde{\psi}_{\kappa_b} \equiv 1$ close to $\text{supp } \psi_{\kappa_b}$. Define smooth functions Ψ_κ and $\tilde{\Psi}_\kappa$ on $(1, \infty) \times U_\kappa$ by $\Psi_\kappa = \kappa_* \psi_\kappa$ and $\tilde{\Psi}_\kappa = \kappa_* \tilde{\psi}_\kappa$, respectively. Then, $\{\Psi_\kappa\}_\kappa$ is a partition of unity subordinate to $\{(1, \infty) \times U_\kappa\}_\kappa$ and $\tilde{\Psi}_\kappa$ satisfies $\tilde{\Psi}_\kappa = 1$ on $\text{supp } \Psi_\kappa$.

2.1. Manifolds with asymptotically cylindrical ends. Noting that $r(z)$ can be extended to a positive smooth function on M and is bounded on M_c from above and below, we define a new density

$$\widehat{G}(z)dz := r(z)^{1-d}G(z)dz, \quad z \in M,$$

and set $L^p(\widehat{M}) := L^p(M, \widehat{G}(z)dz)$ for $1 \leq p < \infty$, $L^\infty(\widehat{M}) := L^\infty(M)$. (1.3) implies that

$$|\partial_r^l \partial_\theta^\alpha (\widehat{G}(r, \theta) - (\det h_{jk}(\theta))^{1/2})| \leq C_{l\alpha} r^{-\mu-l} \quad \text{for } r > 1,$$

and that $\widehat{G}(r, \theta)drd\theta$ is comparable with $drd\theta$ for $r > 1$. This fact implies that, for every function u supported in $(1, \infty) \times V$ with $V \Subset V_\kappa$, $\|u\|_{L^p(\widehat{M})}$ is equivalent to $\|\kappa_* u\|_{L^p(\mathbb{R}^d)}$ for all $1 \leq p \leq \infty$.

We next define an operator $\widehat{P}$ on $L^2(\widehat{M})$ by $\widehat{P} = r(z)^{\frac{d-1}{2}} P r(z)^{-\frac{d-1}{2}}$. It is easy to see that $\widehat{P}$ is unitarily equivalent to P under the unitary map

$$L^2(\widehat{M}) \ni u \mapsto r(z)^{-\frac{d-1}{2}} u \in L^2(M),$$

and $\widehat{P}$ is essentially self-adjoint on $C_0^\infty(M)$. We denote the unique self-adjoint extension by the same symbol $\widehat{P}$. By definition, $(M, \widehat{G})$ can be regarded as a manifold with an asymptotically cylindrical end. For the most part of the paper we shall work with $\widehat{P}$ instead of P . By the formula (1.2) of the scattering metric g , $\widehat{P}_\kappa := \kappa_* \widehat{P} \kappa^*$ takes the form

$$\widehat{P}_\kappa = -\frac{1}{2} \widehat{G}^{-1}(\partial_r, \partial_\theta/r) \widehat{G} \begin{pmatrix} 1 & 0 \\ 0 & h+a \end{pmatrix} \begin{pmatrix} \partial_r \\ \partial_\theta/r \end{pmatrix} + W, \quad (r, \theta) \in (1, \infty) \times U_\kappa,$$

where $h = (h^{jk}) := (h_{jk})^{-1}$ and $a = (a^{jk})$ are smooth and real-valued tensors, and W is a smooth and real-valued potential such that

$$(2.1) \quad |\partial_r^l \partial_\theta^\alpha a^{jk}(r, \theta)| \leq C_{k\alpha} r^{-\mu-l}, \quad |\partial_r^l \partial_\theta^\alpha W(r, \theta)| \leq C_{k\alpha} r^{-1-\mu-l} \quad \text{for } r > 1.$$

Denoting by (ρ, ω) the dual coordinate to the vector field $(\partial_r, \partial_\theta)$, the principal symbol of $\widehat{P}_\kappa$ is written in the form

$$(2.2) \quad p_\kappa(r, \theta, \rho, \omega) = \frac{1}{2}\rho^2 + \frac{1}{2r^2}(h^{jk}(\theta) + a^{jk}(r, \theta))\omega_j\omega_k \quad \text{on } T^*((1, \infty) \times U_\kappa).$$

Moreover, the full symbol of $\widehat{P}_\kappa$ takes the form $p_\kappa + p_{\kappa 1} + p_{\kappa 2}$, where $p_{\kappa j}$ can be written in the form

$$p_{\kappa j}(r, \theta, \rho, \omega) = \sum_{k+|\beta|=2-j} b_{\kappa j}^{k\beta}(r, \theta) \rho^k (\omega/r)^\beta,$$

with some smooth functions $b_{\kappa j}^{k\beta}(r, \theta)$ on $(1, \infty) \times U_\kappa$ satisfying

$$(2.3) \quad |\partial_r^l \partial_\theta^\alpha b_{\kappa j}^{k\beta}(r, \theta)| \leq C_{l\alpha k\beta} r^{-1-\mu-l}.$$

2.2. Pseudodifferential calculus on scattering manifolds. In this subsection we define pseudodifferential operators and study their properties. Moreover we collect known results on the functional calculus on scattering manifolds which were proved in [1] in a more general setting. We begin with the definition of our symbol class.

Definition 2.1. Let X be an open subset of $T^*((1, \infty) \times \mathbb{R}^{d-1}) = (1, \infty)_r \times \mathbb{R}_\theta^{d-1} \times \mathbb{R}_\rho \times \mathbb{R}_\omega^{d-1}$ such that $\pi_\theta(X)$ is relatively compact, and that

$$(2.4) \quad |\omega| \lesssim r \quad \text{in } X,$$

where π_θ is the projection onto the θ -space. We define the symbol class $S_{\text{sc}}(X)$ as the set of all $a \in C^\infty(\mathbb{R}^{2d})$ such that $\text{supp } a \subset X$, and that

$$(2.5) \quad |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta a(r, \theta, \rho, \omega)| \leq C_{j\alpha k\beta} r^{-j-|\beta|} \quad \text{on } X.$$

Example 2.2. Let $\tilde{U}_\kappa \Subset U_\kappa$ be an open subset with $\tilde{\Psi}_\kappa = 1$ on $(2, \infty) \times \tilde{U}_\kappa$, p_κ the principal symbol of P_κ and $\varphi \in C_0^\infty((0, \infty))$. For $R \geq 2$ and an open interval $J \Subset (0, \infty)$ with $\text{supp } \varphi \Subset J$, we set $X_\kappa = (R, \infty) \times \tilde{U}_\kappa \times \mathbb{R}^d \cap p_\kappa^{-1}(J)$. Since $|\rho|^2 + |\omega/r|^2 \leq Cp_\kappa(r, \theta, \rho, \omega) \leq C \sup J$, X_κ satisfies (2.4). Moreover, by (2.1) and (2.2), we have

$$|\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta p_\kappa(r, \theta, \rho, \omega)| \leq C_{j\alpha k\beta} r^{-j-|\beta|} (1 + |\rho|^2 + |\omega/r|^2).$$

Therefore, $\varphi \circ p_\kappa \in S_{\text{sc}}(X_\kappa)$. $\varphi \circ p_\kappa$ is the principal symbol of the semiclassical pseudodifferential approximation of $\varphi(h^2 \widehat{P})$ in the coordinated neighborhood $(1, \infty) \times U_\kappa$ (cf. Lemma 2.4).

Suppose that $X_\kappa \subset (1, \infty) \times U_\kappa \times \mathbb{R}^d$ satisfies (2.4) and $\tilde{\Psi}_\kappa = 1$ near $\pi(X_\kappa)$, where $\pi : T^*\mathbb{R}^d \rightarrow \mathbb{R}^d$ is the projection onto the base space. For all $h \in (0, 1]$ and $a \in S_{\text{sc}}(X_\kappa)$, we define the semiclassical pseudodifferential operator (h -PDO for short) by

$$\text{Op}_\kappa(a)u := \kappa^* \left(a(r, \theta, hD_r, hD_\theta) \kappa_* (\tilde{\Psi}_\kappa u) \right) : C_0^\infty(M) \rightarrow C^\infty(M),$$

where $a(r, \theta, hD_r, hD_\theta)$ is the standard h -PDO which has the kernel

$$(2\pi h)^{-d} \int_{\mathbb{R}^d} e^{\frac{i}{h}[(r-r')\rho + (\theta-\theta')\omega]} a(r, \theta, \rho, \omega) d\rho d\omega.$$

Note that since $\pi(\text{supp } a) \subset (2, \infty) \times \tilde{U}_\kappa$, for any $f_\kappa \in C^\infty(M)$ with $\kappa_* f_\kappa \equiv 1$ on $\pi(\text{supp } a)$, we see that $\text{Op}_\kappa(a)u = f_\kappa \text{Op}_\kappa(a)u$. $\text{Op}_\kappa(a)$ is thus well-defined on M . Moreover, the Calderón-Vaillancourt theorem shows that $\text{Op}_\kappa(a)$ extends to a bounded operator on $L^2(\widehat{M})$ and satisfies

$$(2.6) \quad \|\text{Op}_\kappa(a)\|_{\mathcal{L}(L^2(\widehat{M}))} \leq C_d \sum_{|\gamma| \leq M_d} \|\partial_{(r, \theta, \rho, \omega)}^\gamma a\|_{L^\infty(\mathbb{R}^{2d})} < \infty,$$

uniformly with respect to $h \in (0, 1]$, where $C_d, M_d \geq 0$ depend only on d .

We next describe basic symbolic calculus for $S_{\text{sc}}(X_\kappa)$. We first note that (2.4) and (2.5) are invariant under coordinate transformations since any chart diffeomorphism κ takes the form $\kappa = \text{Id} \otimes \kappa_b$. Let $a \in S_{\text{sc}}(X_\kappa)$ and $b \in S_{\text{sc}}(X_{\kappa'})$. The above fact allow us to define the composition $\text{Op}_\kappa(a) \circ \text{Op}_{\kappa'}(b)$, though $\text{Op}_{\kappa'}(b)$ is not a *properly supported* h -PDO (see Definition 2.3) in general. A standard symbolic calculus implies that the symbol of composition has the following semiclassical expansion:

$$\sum_{l+|\alpha|=0}^N \frac{h^{l+|\alpha|}}{l!|\alpha|!} \partial_\rho^l \partial_\omega^\alpha a(r, \theta, \rho, \omega) D_r^l D_\theta^\alpha \nu_{b*}(\tilde{\Psi}_{\kappa'} b)(r, \theta, \rho, \omega) + h^{N+1} r_N(r, \theta, \rho, \omega),$$

where $r_N \in S_{\text{sc}}(X_\kappa)$ and ν_{b*} is the induced chart diffeomorphism with respect to $\nu_b = \kappa'_b \circ \kappa_b^{-1}$. Note that if $V_\kappa \cap V_{\kappa'} = \emptyset$, then $\text{Op}_\kappa(a_\kappa) \circ \text{Op}_{\kappa'}(b_{\kappa'}) = 0$.

Following [1], we also define the *properly supported* h -PDO.

Definition 2.3. Let $\Psi \in C_0^\infty(\mathbb{R}^d)$ be a cut-off function such that $\text{supp } \Psi \subset \{|z| < \delta\}$ and $\Psi \equiv 1$ on $\{|z| < \delta/2\}$ with some small $\delta > 0$. For $a \in S_{\text{sc}}(X_\kappa)$, we then define the properly supported h -PDO

$$\text{Op}_\kappa^{pr}(a) : C_0^\infty(M) \rightarrow C_0^\infty(M)$$

as the operator so that $\kappa_* \text{Op}_\kappa^{pr}(a) \kappa^*$ has the kernel

$$(2\pi h)^{-d} \int e^{\frac{i}{h}[(r-r')\rho + (\theta-\theta')\omega]} a(r, \theta, \rho, \omega) \Psi(r-r', \theta-\theta') d\rho d\omega.$$

When δ is small, $\Psi \equiv 1$ sufficiently near $\pi(X_\kappa)$ since $\text{supp } a \subset X_\kappa$. We hence removed the factor $\tilde{\Psi}_\kappa$ of the amplitude. We also note that $\text{Op}_h^{pr}(a)$ is uniquely determined on $L^2(\widehat{M})$ up to $O(h^\infty)$. More precisely, we have for any $N \geq 0$,

$$(2.7) \quad \|\text{Op}_\kappa(a) - \text{Op}_\kappa^{pr}(a)\|_{\mathcal{L}(L^2(\widehat{M}))} \leq C_N h^N,$$

uniformly with respect to $h \in (0, 1]$ since $\text{supp}(1 - \Psi)$ away from the diagonal. We here describe a simple property of the properly supported h -PDO. Choose arbitrarily $\chi_0, \chi_1 \in C_0^\infty((2, \infty))$ so that $\chi_1 \equiv 1$ on $\{r \mid \text{dist}(\text{supp } \chi_0, r) < 2\delta\}$. We then have

$$\chi_0(r) \Psi(r-r', \theta-\theta') = \chi_0(r) \Psi(r-r', \theta-\theta') \chi_1(r').$$

In particular,

$$\chi_0(r) \text{Op}_\kappa^{pr}(a) = \chi_0(r) \text{Op}_\kappa^{pr}(a) \chi_1(r'), \quad \text{Op}_\kappa^{pr}(a) \chi_0(r') = \chi_1(r) \text{Op}_\kappa^{pr}(a) \chi_0(r').$$

This property plays an important role in the spatial localization.

Fix $\varphi \in C_0^\infty((0, \infty))$ and a relatively compact open interval $J \Subset (0, \infty)$ so that $\text{supp } \varphi \Subset J$. Let $\chi \in C_0^\infty(M)$ be a smooth cut-off function such that $\chi(z) = 1$ for $z \in M_c \cup \iota^{-1}((0, R_0) \times \partial M)$, $\chi(z) = 0$ for $z \in \iota^{-1}((R_0 + 1, \infty) \times \partial M)$ with some $R_0 > 1$. By using above h -PDO's, we have two kinds of the semiclassical approximations of $(1 - \chi)\varphi(h^2 \hat{P})$.

Lemma 2.4 ([1]). *Let $\delta > 0$ be small enough. Then, for each κ and all $N \geq 0$, there exist semiclassical symbols*

$$a_{\kappa, h} = \sum_{j=0}^N h^j a_{\kappa, j} \quad \text{with} \quad a_{\kappa, j} \in S_{\text{sc}}((R_0, \infty) \times \tilde{U}_\kappa \times \mathbb{R}^d \cap p_\kappa^{-1}(J)),$$

such that

$$(1 - \chi)\varphi(h^2 \hat{P}) = \sum_{\kappa} \text{Op}_\kappa(a_{\kappa, h}) + h^{N+1} R_N(h) = \sum_{\kappa} \text{Op}_\kappa^{pr}(a_{\kappa, h}) + h^{N+1} R_N^{pr}(h)$$

on $L^2(\widehat{M})$. Moreover, there exists $C_N > 0$ such that the followings hold true uniformly with respect to $h \in (0, 1]$:

(i) ($L^2(\widehat{M})$ -boundedness)

$$(2.8) \quad \|\text{Op}_\kappa(a_{\kappa, h})\|_{\mathcal{L}(L^2(\widehat{M}))} + \|R_N(h)\|_{\mathcal{L}(L^2(\widehat{M}))} \leq C_N;$$

(ii) (*semiclassical Sobolev embedding*) for $2 \leq q \leq \infty$,

$$(2.9) \quad \|r^{-\frac{d-1}{2}} \text{Op}_\kappa^{pr}(a_{\kappa,h})\|_{\mathcal{L}(L^2(\widehat{M}), L^q(M))} \leq C_N h^{-d(1/2-1/q)},$$

$$(2.10) \quad \|r^{-\frac{d-1}{2}} R_N^{pr}(h)\|_{\mathcal{L}(L^2(\widehat{M}), L^q(M))} \leq C_N h^{-d(1/2-1/q)};$$

(iii) (*weighted $L^q(\widehat{M})$ -boundedness*) for $1 \leq q \leq \infty$ and all $s \in \mathbb{R}$,

$$(2.11) \quad \|r^{-s} \text{Op}_\kappa^{pr}(a_{\kappa,h}) r^s\|_{\mathcal{L}(L^q(\widehat{M}))} \leq C_N.$$

Proof. The proof was essentially given by [1]. We hence only check that $a_{\kappa,j} \in S_{\text{sc}}((R_0, \infty) \times \tilde{U}_\kappa \times \mathbb{R}^d \cap p_\kappa^{-1}(J))$. $a_{\kappa,0}$ is explicitly given by $a_{\kappa,0} := \kappa_*(1 - \chi) \cdot \varphi \circ p_\kappa$. Moreover, for each j , $a_{\kappa,j}$ is of the form

$$\sum_{k \leq N_j} d_{jk} \cdot (\partial^k \varphi) \circ p_\kappa \quad \text{for some } 0 < N_j < \infty.$$

For each k , d_{jk} is a polynomial of degree $2k - j \geq 0$ with respect to $(\rho, \omega/r)$, and its coefficients are linear combinations of products of derivatives of $\tilde{\Psi}_\kappa$, $\kappa_*(1 - \chi)$ and the full symbol of P_κ . Therefore, $a_{\kappa,j}$ takes the form

$$a_{\kappa,j}(r, \theta, \rho, \omega) = b_{\kappa,j}(r, \theta, \rho, \omega/r),$$

where $b_{\kappa,j}$ is compactly supported with respect to ρ and ω , and satisfies

$$|\partial_r^l \partial_\theta^\alpha \partial_\rho^m \partial_\omega^\beta b_{\kappa,j}(r, \theta, \rho, \omega)| \leq C_{l\alpha k \beta R_0} r^{-l}.$$

We hence obtain $a_{\kappa,j} \in S_{\text{sc}}((R_0, \infty) \times \tilde{U}_\kappa \times \mathbb{R}^d \cap p_\kappa^{-1}(J))$. $\square$

2.3. Outgoing and incoming regions. In this subsection we recall the definition of the outgoing and incoming regions and study some basic properties of these regions needed later. Let $R \geq 1$, $\tilde{U}_\kappa \Subset U_\kappa$ an open subset, $J \Subset (0, \infty)$ an open interval and $\sigma \in (-1, 1)$.

Definition 2.5. (i) We set

$$\Gamma^\pm(R, \tilde{U}_\kappa, J, \sigma) = \{(r, \theta, \rho, \omega) \in \mathbb{R}^{2d} \mid r > R, \theta \in \tilde{U}_\kappa, p_\kappa \in J, \pm\rho > -\sigma\sqrt{2p_\kappa}\},$$

where $p_\kappa = p_\kappa(r, \theta, \rho, \omega)$. $\Gamma^+(R, \tilde{U}_\kappa, J, \sigma)$ (resp. $\Gamma^-(R, \tilde{U}_\kappa, J, \sigma)$) is said to be the outgoing (resp. incoming) region.

(ii) Let $\tilde{U}_{\kappa, \sqrt{\varepsilon}}$ and J_{ε^2} be an $\sqrt{\varepsilon}$ -neighborhood of $\tilde{U}_\kappa$ and an ε^2 -neighborhood of J , respectively:

$$\tilde{U}_{\kappa, \sqrt{\varepsilon}} := \{\theta \in \mathbb{R}^{d-1} \mid \text{dist}(\tilde{U}_\kappa, \theta) < \sqrt{\varepsilon}\}, \quad J_{\varepsilon^2} := \{\rho \pm \varepsilon^2 \in (0, \infty) \mid \rho \in J\}.$$

For sufficiently small $\varepsilon > 0$ such that $\tilde{U}_{\kappa, \sqrt{\varepsilon}} \Subset U_\kappa$, we define the strongly outgoing and incoming regions as follows:

$$\Gamma_s^\pm(R, \tilde{U}_\kappa, J, \varepsilon) := \Gamma^\pm(R, \tilde{U}_{\kappa, \sqrt{\varepsilon}}, J_{\varepsilon^2}, -\sqrt{1 - \varepsilon^2}).$$

(iii) For sufficiently small $\varepsilon, \delta > 0$, and for any $L > 0$ and $\sigma_l \in (-1, 1/2]$, $l = 0, 2, \dots, L$, satisfying

$$(2.12) \quad \begin{cases} (-\sqrt{1 - \varepsilon^2/4}, 1/2) = \bigcup_{l=1}^{L-1} (\sigma_{l-1}, \sigma_{l+1}), \\ -\sqrt{1 - \varepsilon^2/4} = \sigma_0 < \sigma_1 < \dots < \sigma_L = 1/2, \quad |\sigma_{l+1} - \sigma_{l-1}| \leq \delta, \end{cases}$$

the intermediate outgoing region and incoming region are defined by

$$\begin{aligned} & \Gamma_i^\pm(R, \tilde{U}_\kappa, J, \varepsilon, \delta, l) \\ & := \Gamma^\pm(R, \tilde{U}_{\kappa, \sqrt{\varepsilon}}, J_{\varepsilon^2}, 1/2) \cap \{-\sigma_{l+1}\sqrt{2p_\kappa} < \pm\rho < -\sigma_{l-1}\sqrt{2p_\kappa}\}. \end{aligned}$$

We describe basic properties of these regions. $\Gamma^\pm(R, \tilde{U}_\kappa, J, \sigma)$ are monotonously decreasing with respect to R , and increasing with respect to $\tilde{U}_\kappa$, J and σ . By definition, we have

$$(R, \infty) \times \tilde{U}_\kappa \times \mathbb{R}^d \cap p_\kappa^{-1}(J) \subset \bigcup_{\pm} \Gamma^\pm(R, \tilde{U}_\kappa, J, 1/2).$$

We also obtain

$$\Gamma^\pm(R, \tilde{U}_\kappa, J, 1/2) \subset \Gamma_s^\pm(R, \tilde{U}_\kappa, J, \varepsilon) \cup \bigcup_{l=1}^{L-1} \Gamma_i^\pm(R, \tilde{U}_\kappa, J, \varepsilon, \delta, l),$$

respectively. Moreover for sufficiently large $R_0, C > 0$, all $0 < \varepsilon < 1/2$ and $R \geq R_0$,

$$|\omega/r| \leq C\varepsilon \quad \text{on} \quad \Gamma_s^\pm(R, \tilde{U}_\kappa, J, \varepsilon).$$

Indeed, since $\rho^2 \geq (1 - \varepsilon^2) \left(\rho^2 + \frac{1}{r^2} (h^{jk} + a^{jk}) \omega_j \omega_k \right)$ on $\Gamma_s^\pm(R, \tilde{U}_\kappa, J, \varepsilon)$, taking $R_0 > 0$ large enough so that $(h^{jk} + a^{jk})_{j,k} \geq C_0^{-1} \text{Id}$ for some fixed $C_0 > 0$, we obtain $|\omega/r|^2 \leq C_0 \rho^2 \varepsilon^2 / (1 - \varepsilon^2)$.

We also define spatial localized regions as follows.

Definition 2.6. Let $\varepsilon, \delta, L, \sigma_l$ be as above. For $R_2 > R_1 > 1$, we define the spatial localized outgoing and incoming regions $\Omega^\pm(R_1, R_2, \tilde{U}_\kappa, J, \sigma)$ by

$$\Omega^\pm(R_1, R_2, \tilde{U}_\kappa, J, \sigma) := \Gamma^\pm(R_1, \tilde{U}_\kappa, J, \sigma) \cap \{R_1 < r < 4R_2\}.$$

We shall use the notation $\Omega^\pm(R, \tilde{U}_\kappa, J, \sigma) = \Omega^\pm(R, R, \tilde{U}_\kappa, J, \sigma)$. We also define the corresponding spatial localized strongly outgoing (incoming) and intermediate regions

$$\Omega_s^\pm(R_1, R_2, \tilde{U}_\kappa, J, \varepsilon), \quad \Omega_s^\pm(R, \tilde{U}_\kappa, J, \varepsilon), \quad \Omega_i^\pm(R_1, R_2, \tilde{U}_\kappa, J, \varepsilon, \delta, l),$$

and $\Omega_i^\pm(R, \tilde{U}_\kappa, J, \varepsilon, \delta, l)$ in the same manner, respectively.

Remark 2.7. Since the principal symbol of P is invariant under coordinate transformations, these regions define invariant subsets in $T^*\widetilde{M}_\infty$ except the choice of the boundary decomposition. Moreover we will prove in Section 4 that these regions are also invariant under the geodesic flow generated by p . This property will be used to prove microlocal smoothing properties of the propagator (see Section 6).

3. REDUCTION TO MICROLOCAL DISPERSIVE ESTIMATES

In this section we shall show that (1.5) follows from microlocal dispersive estimates. We first recall the frequency localization using the Littlewood-Paley decomposition. The following theorem was proved by Bouclet [2] for a large class of non-compact manifolds with ends (including scattering manifolds).

Proposition 3.1 ([2]). *Let $\psi \in C_0^\infty((0, \infty))$ be a smooth cut-off function such that*

$$\text{supp } \psi \subset [1/4, 4], \quad 0 \leq \psi \leq 1, \quad \sum_{j=0}^{\infty} \psi(2^{-2j}\lambda) = 1 \quad \text{for } \lambda \in [1, \infty).$$

Then, for all $\chi \in C_0^\infty(M)$ with $\text{supp}(1 - \chi) \subset M_\infty$ and $2 \leq q < \infty$ with $0 \leq d(1/2 - 1/q) \leq 1$, there exists $C > 0$ such that

$$\|(1 - \chi)u\|_{L^q(M)} \leq C\|u\|_{L^2(M)} + C \left(\sum_{j=0}^{\infty} \|(1 - \chi)\psi(2^{-2j}P)u\|_{L^q(M)}^2 \right)^{\frac{1}{2}}.$$

Using this proposition, we see that (1.5) follows from semiclassical Strichartz estimates. More precisely, it suffices to prove that for $\chi \in C_0^\infty(M)$ as above and every $\varphi \in C_0^\infty((0, \infty))$,

$$(3.1) \quad \|(1 - \chi)\varphi(h^2P)e^{-itP}u_0\|_{L^p([0,1];L^q(M))} \leq C\|u_0\|_{L^2(M)}$$

uniformly with respect to $h \in (0, 1]$, where (p, q) satisfies the admissible condition (1.1). This reduction is a standard and can be proved by using the L^2 -functional calculus and the almost orthogonality of $\psi(2^{-2j}P)$. By the definition of $\widehat{P}$, (3.1) is equivalent to

$$(3.2) \quad \|r^{-\frac{d-1}{2}}(1-\chi)\varphi(h^2\widehat{P})e^{-it\widehat{P}}u_0\|_{L^p([0,1];L^q(M))} \leq C\|u_0\|_{L^2(\widehat{M})}, \quad h \in (0, 1].$$

For $R > 2$, we take a cut-off $\chi \in C_0^\infty(M)$ so that $\chi(z) = 1$ for $r(z) \leq R$, $\chi(z) = 0$ for $r(z) \geq R+1$. Let $J \Subset (0, \infty)$ be an open interval so that $\text{supp } \varphi \Subset J$ and $N \geq d/2$ an integer. By Lemma 2.4, we then can find a semiclassical symbol $a_{\kappa,h} \in S_{\text{sc}}((R, \infty) \times \tilde{U}_\kappa \times \mathbb{R}^d \cap p_\kappa^{-1}(J))$ such that $(1-\chi)\varphi(h^2\widehat{P})$ is well approximated by $A_h := \sum_\kappa \text{Op}_\kappa^{pr}(a_{\kappa,h})$. Moreover, the following holds.

Proposition 3.2. *To prove (3.2), it suffices to show that for sufficiently large $R > 1$ and all $a \in S_{\text{sc}}((R, \infty) \times \tilde{U}_\kappa \times \mathbb{R}^d \cap p_\kappa^{-1}(J))$,*

$$(3.3) \quad \|r^{-\frac{d-1}{2}}A_h \text{Op}_\kappa(a)e^{-it\widehat{P}}u_0\|_{L^p([0,1];L^q(M))} \leq C\|u_0\|_{L^2(\widehat{M})}, \quad u_0 \in C_0^\infty(M),$$

uniformly with respect to $h \in (0, 1]$, where (p, q) satisfies (1.1).

This proposition is a direct consequence of Lemma 2.4, and the proof is completely same as [3, Proposition 2.18].

We next describe the spatial localization and give the main step of the proof of Theorem 1.1. The following theorem is the main result of this section. We here fix $\tilde{U}_\kappa$ and J , and hence do not write explicitly. We also use a notation such as $\Gamma^\pm(R, \tilde{U}_\kappa, J, \sigma) = \Gamma^\pm(R, \sigma)$ for short.

Theorem 3.3 (Microlocal dispersive estimates). *There exist $R_0 \geq 0$ and $\varepsilon_0 > 0$ such that the following hold for all $R_2 \geq R_1 \geq R_0$, $0 < \varepsilon < \varepsilon_0$ and $h \in (0, 1]$.*

(i) *There exists $t_0 > 0$, independent of R_2 , such that for all symbols*

$$a^\pm \in S_{\text{sc}}(\Gamma^\pm(R_1, 1/2)), \quad b^\pm \in S_{\text{sc}}(\Omega^\pm(R_2, 1/2)),$$

and $0 < \pm t \leq \min(R_2 t_0, h^{-1})$, we have

$$(3.4) \quad \|r^{-\frac{d-1}{2}}A_h \text{Op}_\kappa(a^\pm)e^{-ith\widehat{P}}\text{Op}_\kappa(b^\pm)^*A_h^*r^{-\frac{d-1}{2}}\|_{\mathcal{L}(L^1(\widehat{M}), L^\infty(\widehat{M}))} \leq C_0|th|^{-\frac{d}{2}}.$$

(ii) *For all symbols*

$$a_s^\pm \in S_{\text{sc}}(\Gamma_s^\pm(R_1, \varepsilon)), \quad b_s^\pm \in S_{\text{sc}}(\Omega_s^\pm(R_2, \varepsilon)),$$

and $0 < \pm t \leq h^{-1}$, we have

$$(3.5) \quad \|r^{-\frac{d-1}{2}}A_h \text{Op}_\kappa(a_s^\pm)e^{-ith\widehat{P}}\text{Op}_\kappa(b_s^\pm)^*A_h^*r^{-\frac{d-1}{2}}\|_{\mathcal{L}(L^1(\widehat{M}), L^\infty(\widehat{M}))} \leq C_1|th|^{-\frac{d}{2}}.$$

(iii) *For all $t_1 > 0$, we can find $\delta_{\varepsilon, t_1} > 0$ and $L_{\varepsilon, t_1} > 0$ such that for all $\sigma_l \in (-1, 1/2]$ satisfying (2.12), all symbols*

$$a_l^\pm \in S_{\text{sc}}(\Gamma_i^\pm(R_1, \varepsilon, \delta_{\varepsilon, t_0}, l)), \quad b_l^\pm \in S_{\text{sc}}(\Omega_i^\pm(R_2, \varepsilon, \delta_{\varepsilon, t_0}, l)),$$

and $R_2 t_1 \leq \pm t \leq h^{-1}$, we have

$$(3.6) \quad \|r^{-\frac{d-1}{2}}A_h \text{Op}_\kappa(a_l^\pm)e^{-ith\widehat{P}}\text{Op}_\kappa(b_l^\pm)^*A_h^*r^{-\frac{d-1}{2}}\|_{\mathcal{L}(L^1(\widehat{M}), L^\infty(\widehat{M}))} \leq C_2|th|^{-\frac{d}{2}}.$$

Moreover $C_0, C_1, C_2 > 0$ may be taken uniformly with respect to h, t and R_2 .

We give the proof of Theorem 3.3 in Section 8. Before proving Theorem 1.1, we prepare the following lemma.

Lemma 3.4. *Let $R, \delta, L > 0$, $0 < \varepsilon < 1$ and $(\sigma_l)_{1 \leq l \leq L} \subset (-1, 1/2]$ so that (2.12) is satisfied. Then for any $a \in S_{\text{sc}}((R, \infty) \times \tilde{U}_\kappa \times \mathbb{R}^d \cap p_\kappa^{-1}(J))$, there exist symbols $a^\pm \in S_{\text{sc}}(\Gamma^\pm(R, 1/2))$, $a_s^\pm \in S_{\text{sc}}(\Gamma_s^\pm(R, \varepsilon))$ and $a_l^\pm \in S_{\text{sc}}(\Gamma_i^\pm(R, \varepsilon, \delta, l))$, $l = 1, 2, \dots, L-1$, such that*

$$a = a^+ + a^- = a_s^+ + a_s^- + \sum_{l=1}^{L-1} (a_l^+ + a_l^-).$$

Proof. Define $v(r, \theta, \rho, \omega) := \rho / \sqrt{2p_\kappa(r, \theta, \rho, \omega)}$. v is then smooth on $\Gamma^\pm(R, 1/2)$, and satisfies

$$|\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta v(r, \theta, \rho, \omega)| \leq C_{j\alpha k\beta} r^{-j-|\beta|} \quad \text{on } \Gamma^\pm(R, 1/2),$$

since p_κ is of the form (2.2). Take a cut-off function $\chi \in C^\infty(\mathbb{R})$ such that

$$\chi(t) + \chi(-t) = 1, \quad \text{supp } \chi \subset (-1/2, \infty),$$

and define $a^\pm = \chi(\pm v) \cdot a$. Then $a^\pm \in S_{\text{sc}}(\Gamma^\pm(R, 1/2))$ and $a = a^+ + a^-$. Next, choose $\chi_s \in C^\infty(\mathbb{R})$ and $\chi_l \in C_0^\infty(\mathbb{R})$, $l = 1, 2, \dots, L-1$, so that

$$\text{supp } \chi_s \subset (\sqrt{1-\varepsilon^2}, \infty), \quad \text{supp } \chi_l \subset (-\sigma_{l+1}, -\sigma_{l-1}), \quad \text{supp } \chi_L \subset (-1, -\sigma_{L-1}),$$

and that $\chi_s + \sum_{l=1}^L \chi_l = 1$. We define $a_s^\pm := \chi_s(\pm v) \cdot a^\pm$ and $a_l^\pm := \chi_l(\pm v) \cdot a^\pm$, respectively. It is easy to see that they satisfy the assertion. $\square$

Proof of Theorem 1.1. Choose $R_0, \varepsilon_0, t_0 > 0$ so that (i) and (ii) in Theorem 3.3 hold for all $R_2 \geq R_1 \geq R_0$ and $0 < \varepsilon \leq \varepsilon_0$. Next, fix $\varepsilon, \delta_{\varepsilon, t_0}, L_{\varepsilon, t_0} > 0$ so that (iii) holds with $t_1 = t_0$. By virtue of Proposition 3.2, it suffices to show (3.3) for any $a \in S_{\text{sc}}((R_1, \infty) \times \tilde{U}_\kappa \times \mathbb{R}^d \cap p_\kappa^{-1}(J))$. Using Lemma 3.4 with $\delta = \delta_{\varepsilon, t_0}$ and $L = L_{\varepsilon, t_0}$, we split $\text{Op}_\kappa(a)$ as follows:

$$\text{Op}_\kappa(a) = \text{Op}_\kappa(a_s^+) + \text{Op}_\kappa(a_s^-) + \sum_{l=1}^{L_{\varepsilon, t_0}-1} (\text{Op}_\kappa(a_l^+) + \text{Op}_\kappa(a_l^-)),$$

where $a_s^\pm \in S_{\text{sc}}(\Gamma_s^\pm(R_1, \varepsilon))$ and $a_l^\pm \in S_{\text{sc}}(\Gamma_l^\pm(R_1, \varepsilon, \delta_{\varepsilon, t_0}, l))$. By virtue of the TT^* -argument [14], it suffices to show the $\mathcal{L}(L^2(\widehat{M}), L^2(M))$ -boundedness and $L^1(M) - L^\infty(M)$ estimates for corresponding operators uniformly with respect to $h \in (0, 1]$.

By (2.6), (2.9) and the fact that $e^{-it\hat{P}}$ is unitary on $L^2(\widehat{M})$, we obtain

$$\begin{aligned} \|r^{-\frac{d-1}{2}} A_h \text{Op}_\kappa(a_s^\pm) e^{-it\hat{P}} u_0\|_{L^2(M)} &\leq C \|u_0\|_{L^2(\widehat{M})}, \\ \|r^{-\frac{d-1}{2}} A_h \text{Op}_\kappa(a_l^\pm) e^{-it\hat{P}} u_0\|_{L^2(M)} &\leq C \|u_0\|_{L^2(\widehat{M})}, \end{aligned}$$

uniformly with respect to $h \in (0, 1]$, $t \in \mathbb{R}$ and $l = 1, 2, \dots, L_{\varepsilon, t_0} - 1$. After the time rescaling $t \rightarrow th$, we set

$$U_s^\pm(t) := r^{-\frac{d-1}{2}} A_h \text{Op}_\kappa(a_s^\pm) e^{-ith\hat{P}}, \quad U_l^\pm(t) := r^{-\frac{d-1}{2}} A_h \text{Op}_\kappa(a_l^\pm) e^{-ith\hat{P}}.$$

It remains to show that $U_s^\pm(t)U_s^\pm(s)^*$ and $U_l^\pm(t)U_l^\pm(s)^*$ satisfy dispersive estimates for $0 < |t-s| \leq h^{-1}$. We here use a trick by [4, Lemma 4.3]. We denote by $K^\pm(t-s, r, \theta, r', \theta')$ the kernel of $U_s^\pm(t)U_s^\pm(s)^*$, respectively. Since $U_s^\pm(t)U_s^\pm(s)^* = (U_s^\pm(s)U_s^\pm(t)^*)^*$, we see that

$$K^\pm(t-s, r, \theta, r', \theta') = \overline{K^\pm(s-t, r', \theta', r, \theta)}.$$

A same property holds for the kernel of $U_l^\pm(t)U_l^\pm(s)^*$. We hence can restrict the sign of $t-s$ so that $0 \leq \pm(t-s) \leq h^{-1}$, and it is enough to prove the following:

$$\begin{aligned} (3.7) \quad &\|U_s^\pm(t)U_s^\pm(s)^* u_0\|_{L^\infty(M)} + \|U_l^\pm(t)U_l^\pm(s)^* u_0\|_{L^\infty(M)} \\ &\leq C |(t-s)h|^{-\frac{d}{2}} \|u_0\|_{L^1(M)}, \end{aligned}$$

uniformly with respect to $h \in (0, 1]$ and $0 < \pm(t-s) \leq h^{-1}$, $l = 1, 2, \dots, L_{\varepsilon, t_0} - 1$, respectively. Combining with the facts $L^1(M) = r^{-(d-1)} L^1(\widehat{M})$, $L^\infty(M) = L^\infty(\widehat{M})$ and

$$(r^{-\frac{d-1}{2}})^* = r^{\frac{d-1}{2}} : L^2(M) \rightarrow L^2(\widehat{M}),$$

(3.7) follows from

$$\begin{aligned} (3.8) \quad &\|U_s^\pm(t)U_s^\pm(s)^* u_0\|_{L^\infty(\widehat{M})} + \|U_l^\pm(t)U_l^\pm(s)^* u_0\|_{L^\infty(\widehat{M})} \\ &\leq C |(t-s)h|^{-\frac{d}{2}} \|u_0\|_{L^1(\widehat{M})}, \end{aligned}$$

where $V_s^\pm(s)$ and $V_l^\pm(s)$ are given by

$$V_s^\pm(s) := e^{ish\hat{P}} \text{Op}_\kappa(a_s^\pm)^* A_h^* r^{-\frac{d-1}{2}}, \quad V_l^\pm(s) := e^{ish\hat{P}} \text{Op}_\kappa(a_l^\pm)^* A_h^* r^{-\frac{d-1}{2}}.$$

We now introduce a spatial localization. Let $\chi \in C_0^\infty((0, \infty))$ be a smooth cut-off function such that

$$\text{supp } \chi \subset [1, 4], \quad 0 \leq \chi \leq 1, \quad \sum_{j=0}^{\infty} \chi(2^{-j}r) = 1 \quad \text{for } r \in [2, \infty).$$

Choose $\chi' \in C_0^\infty((0, \infty))$ so that $\text{supp } \chi' \subset [1/2, 8]$, $\chi' \equiv 1$ on $\text{supp } \chi$, and set $\chi_j(r) := \chi(2^{-j}r)$, $\chi'_j(r) := \chi'(2^{-j}r)$. Since Ψ is supported in a small ball centered at origin (see Definition 2.3), we obtain that

$$\chi_j(r)\Psi(r-r', \theta-\theta') = \chi_j(r)\Psi(r-r', \theta-\theta')\chi'_j(r'), \quad j \geq j_0,$$

with some large $j_0 \geq 0$. In particular, we have

$$\text{Op}_\kappa(\chi_j a)^* A_h^* = \text{Op}_\kappa(\chi_j a)^* A_h^* \chi'_j \quad \text{for } j \geq j_0.$$

If we choose j_0 as $2^{j_0} > R_0$ and set $R_1 = 2^{j_0}$, then

$$a_s^\pm = \sum_{j \geq j_0} \chi_j a_s^\pm, \quad a_l^\pm = \sum_{j \geq j_0} \chi_j a_l^\pm.$$

Since $\chi_j a_s^\pm \in S_{\text{sc}}(\Omega_s^\pm(2^j, \varepsilon))$ and $\chi_j a_l^\pm \in S_{\text{sc}}(\Omega_l^\pm(2^j, \varepsilon, \delta_{\varepsilon, t_0}, l))$, applying Theorem 3.3 with $R_2 = 2^j$ to $U_s^\pm(t)V_s^\pm(s)$ and $U_l^\pm(t)V_l^\pm(s)$, we have

$$\begin{aligned} & \|U_s^\pm(t)V_s^\pm(s)u_0\|_{L^\infty(\widehat{M})} + \|U_l^\pm(t)V_l^\pm(s)u_0\|_{L^\infty(\widehat{M})} \\ & \leq C|(t-s)h|^{-\frac{d}{2}} \sum_{j \geq j_0} \|\chi'_j u_0\|_{L^1(\widehat{M})} \\ & \leq 4C|(t-s)h|^{-\frac{d}{2}} \|u_0\|_{L^1(\widehat{M})}, \end{aligned}$$

uniformly with respect to $h \in (0, 1]$ and $0 < \pm(t-s) \leq h^{-1}$. We hence obtain (3.8) and conclude the proof of Theorem 1.1. $\square$

Remark 3.5. Since $b^\pm, b_s^\pm$ and $b_l^\pm$ are compactly supported with respect to both the space and the frequency, the above argument tells us that Strichartz estimates follows from microlocal dispersive estimates.

4. CLASSICAL TRAJECTORIES

In this section we study the behavior of the geodesic flow which we denote by

$$\exp tH_p : T^*M \rightarrow T^*M.$$

Recall that the principal symbol p_κ of $\widehat{P}_\kappa$ is of the form

$$p_\kappa(r, \theta, \rho, \omega) = \frac{1}{2}\rho^2 + \frac{1}{2r^2}(h^{jk}(\theta) + a^{jk}(r, \theta))\omega_j\omega_k, \quad (r, \theta, \rho, \omega) \in (1, \infty) \times U_\kappa \times \mathbb{R}^d,$$

where $a^{jk}(r, \theta)$ satisfies (2.1). We put

$$(r(t), \theta(t), \rho(t), \omega(t)) = \exp tH_{p_\kappa}(r, \theta, \rho, \omega),$$

which solves the following Hamilton equation:

$$(4.1) \quad \begin{cases} \dot{r}(t) = \rho(t), \\ \dot{\theta}^j(t) = \frac{1}{r(t)^2} h^{jk}(\theta(t)) \omega_k(t) + \frac{1}{r(t)^2} a^{jk}(r(t), \theta(t)) \omega_k, \\ \dot{\rho}(t) = \frac{1}{r(t)^3} h^{jk}(\theta(t)) \omega_j(t) \omega_k(t) + \frac{1}{r(t)^3} a^{jk}(r(t), \theta(t)) \omega_j(t) \omega_k(t) \\ \quad - \frac{1}{2r^2} \frac{\partial a^{jk}}{\partial r}(r(t), \theta(t)) \omega_j(t) \omega_k(t), \\ \dot{\omega}_j(t) = -\frac{1}{2r(t)^2} \frac{\partial h^{kl}}{\partial \theta^j}(\theta(t)) \omega_k(t) \omega_l(t) - \frac{1}{2r(t)^2} \frac{\partial a^{kl}}{\partial \theta^j}(r(t), \theta(t)) \omega_k(t) \omega_l(t). \end{cases}$$

We first prepare an a priori estimate for $\exp tH_{p_\kappa}$.

Lemma 4.1. *Let $J \in (0, \infty)$ and $-1 < \sigma < 1$. Then, there exist $R_0 > 0$ such that for all $R \geq R_0$,*

$$(4.2) \quad C^{-1}(r + |t|) \leq r(t) \leq C(r + |t|), \quad |\rho(t)| + |\omega(t)/r| \leq C,$$

uniformly with respect to $(r, \theta, \rho, \omega) \in \Gamma^\pm(R, \tilde{U}_\kappa, J, \sigma)$ and $\pm t \geq 0$, where the constant C may be taken uniformly with respect to R and t .

Using Lemma 4.1, we obtain the behavior of the geodesic flow near infinity.

Proposition 4.2. *Let $R_0 > 0$ be as in Lemma 4.1 and $R \geq R_0$. Then the following estimates hold for all $(r, \theta, \rho, \omega) \in \Gamma^\pm(R, \tilde{U}_\kappa, J, \sigma)$, $\pm t \geq 0$ and $(j, \alpha, k, \beta) \in \mathbb{Z}_+^{2d}$ as long as the trajectory belongs to the same coordinate neighborhood $(1, \infty) \times U_\kappa$:*

$$(4.3) \quad \begin{cases} |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (r(t) - r \mp t\rho)| \leq Cr^{-j-|\beta|} |\omega/r|^{(2-|\beta|)+} \langle r/t \rangle^{-1} |t|, \\ |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\theta(t) - \theta)| \leq Cr^{-j-|\beta|} |\omega/r|^{(1-|\beta|)+} \langle r/t \rangle^{-1}, \\ |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\rho(t) - \rho)| \leq Cr^{-j-|\beta|} |\omega/r|^{(2-|\beta|)+} \langle r/t \rangle^{-1}, \\ |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\omega(t) - \omega)| \leq Cr^{1-j-|\beta|} |\omega/r|^{(2-|\beta|)+} \langle r/t \rangle^{-1}, \end{cases}$$

$$(4.4) \quad \begin{cases} |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (r(t) - r)| + |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\omega(t) - \omega)| \leq Cr^{-j-|\beta|} |t|, \\ |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\theta(t) - \theta)| + |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\rho(t) - \rho)| \leq Cr^{-1-j-|\beta|} |t|, \end{cases}$$

Moreover, for all $(r, \theta, \rho, \omega) \in \Gamma^\pm(R, \tilde{U}_\kappa, J, \sigma)$ and $\pm t \geq 0$, we have

$$(4.5) \quad |\sqrt{2E_0} \mp \rho(t)| \leq C|\omega/r|^2 \langle t/r \rangle^{-1},$$

where $E_0 := p_\kappa(r, \theta, \rho, \omega)$ is the initial energy.

Proof of Lemma 4.1. We prove the lemma for $t \geq 0$, and the proof for $t \leq 0$ is analogous. We first note that the energy conservation law, namely

$$E_0 = p_\kappa(r(t), \theta(t), \rho(t), \omega(t)), \quad t \in \mathbb{R},$$

implies $|\rho(t)| + |\omega(t)/r(t)| \leq C_0$ as long as $r(t)$ large enough, where $C_0 > 0$ depends only on $\sqrt{E_0}$. In particular, we can find $R_1, C_1 > 0$ so large that

$$\dot{r}(0) = O(1), \quad \dot{\rho}(0) = \frac{1}{r^3} h^{jk}(\theta) \omega_j \omega_k + O(r^{-1-\mu}),$$

for $r > R_1$, and hence

$$\left. \frac{d^2}{dt^2} r(t)^2 \right|_{t=0} = 2(\dot{r}(t)\rho(t) + r(t)\dot{\rho}(t))|_{t=0} \geq 4E_0 - C_1 r^{-\mu}.$$

Since $\sigma \in (-1, 1)$, we can choose $0 < \delta < 1$ and $C_2 > 0$ so that

$$r^2 - 2\sigma r \sqrt{2E_0} t + 2(1 - \delta)^2 E_0 t^2 \geq \frac{1}{C_2^2} (r + t \sqrt{2E_0})^2, \quad r \geq 0, \quad t \geq 0.$$

We now suppose that there exist $R > 0$ and $t_0 > 0$ such that

$$(4.6) \quad 4E_0 - C_1 r(t)^{-\mu} \geq 4(1 - \delta)^2 E_0$$

holds true for $r > R$ and $0 \leq t \leq t_0$. We then have

$$\begin{aligned} r(t)^2 &\geq r^2 + 2r\rho t + 2(1 - \delta)^2 E_0 t^2 \\ &\geq r^2 - 2\sigma r \sqrt{2E_0} t + 2(1 - \delta)^2 E_0 t^2 \\ &\geq \frac{1}{C_2^2} (r + t \sqrt{2E_0})^2, \end{aligned}$$

for $0 \leq t \leq t_0$ and $(r, \theta, \rho, \omega) \in \Gamma^+(R, \tilde{U}_\kappa, J, \sigma)$. Moreover, if we put $q_0(t) = h^{jk}(\theta(t))\omega_j(t)\omega_k(t)$, then a direct computation yields

$$\begin{aligned} \dot{q}_0(t) &= \dot{\theta}^l \frac{\partial h^{jk}}{\partial \theta^l} \omega_j \omega_k + 2h^{jk} \dot{\omega}_j \omega_k \\ &= \frac{1}{r^2} (h^{lm} + a^{lm}) \omega_m \frac{\partial h^{jk}}{\partial \theta^l} \omega_j \omega_k - h^{jk} \omega_k \frac{1}{r^2} \left(\frac{\partial h^{lm}}{\partial \theta^j} + \frac{\partial a^{lm}}{\partial \theta^j} \omega_l \omega_m \right) \\ &= O(r(t)^{-1-\mu} q_0(t)). \end{aligned}$$

Integrating with respect to $s \in [0, t]$, we have

$$q_0(t) \leq q_0(0) + C \int_0^t (r + |s| \sqrt{2E_0})^{-1-\mu} q_0(s) ds, \quad 0 \leq t \leq t_0.$$

By Gronwall's inequality and the ellipticity of q_0 , we obtain

$$(4.7) \quad |\omega(t)| \leq C_3 \sqrt{q_0} \leq C_3 r \sqrt{2E_0} \quad \text{for } 0 \leq t \leq t_0.$$

Applying (4.7) to (4.1), we have

$$(4.8) \quad \dot{r}(t) = \rho(t), \quad \dot{\theta}(t) = O(r(t)^{-2} r), \quad \dot{\rho}(t) = O(r(t)^{-3} r^2), \quad \dot{\omega}(t) = O(r(t)^{-2} r^2).$$

In particular, we see that $|\rho(t)| \leq C_4 \sqrt{2E_0}$ for $0 \leq t \leq t_0$ with some large $C_4 > 0$. Therefore, it is enough to check that (4.6) holds with $t_0 = \infty$. Define

$$S := \{t \geq 0 \mid (4.6) \text{ holds for all } s \in [0, t]\}.$$

For sufficiently large $r > R_1$, the above argument shows that $0 \in S$ and $S \neq \emptyset$. Set $t_0 = \sup S$. The above argument then implies

$$r(t) \geq C_2^{-1} (r + t \sqrt{2E_0}) \quad \text{for } r > R_2, \quad 0 \leq t \leq t_0,$$

with some $R_2 > R_1$ large enough. Taking $R_3 > R_2$ so that

$$4\delta E_0 - \delta^2 E_0 \geq C_1 C_2^\mu r^{-\mu} \quad \text{for } r > R_3,$$

we have

$$\begin{aligned} 4E_0 - C_1 r(t)^{-\mu} &\geq 4(1 - \frac{\delta}{2})^2 E_0 + 4\delta E_0 - \delta^2 E_0 - C_1 C_2^\mu (r + t \sqrt{2E_0})^{-\mu} \\ &\geq 4(1 - \frac{\delta}{2})^2 E_0 \end{aligned}$$

for $r > R_3$ and $0 \leq t \leq t_0$. Therefore, $t_0 + \varepsilon \in S$ for some $\varepsilon > 0$ which implies $t_0 = \infty$ by the definition of t_0 . The estimate of $r(t)$ from above is obvious. $\square$

Proof of Proposition 4.2. Let $t \geq 0$. The proof for $t < 0$ is similar. Take $R_0 > 0$ as in Lemma 4.1. (4.3) with $j + |\alpha| + k + |\beta| = 0$ is a direct consequence of Lemma 4.1 and (4.8) since

$$\int_0^t (r + |s|)^{-1-a} ds \leq C r^{-a} \langle r/t \rangle^{-1} \quad \text{for any } a > 0.$$

We next consider the derivatives. Put $z(t) := r(t) - t\rho(t)$. It is easy to see that

$$W(t) := (z(t), \theta(t), \rho(t), \omega(t)) = \exp(-tH_{\frac{1}{2}\rho^2}) \circ \exp tH_{p_\kappa}(r, \theta, \rho, \omega)$$

solves the following Hamilton equations:

$$\dot{z} = \frac{\partial K}{\partial \rho}, \quad \dot{\theta} = \frac{\partial K}{\partial \omega}, \quad \dot{\rho} = -\frac{\partial K}{\partial z}, \quad \dot{\omega} = -\frac{\partial K}{\partial \theta},$$

with a time dependent Hamiltonian:

$$K(t, z, \theta, \rho, \omega) := \frac{1}{2}\rho^2 - p_\kappa(z + t\rho, \theta, \rho, \omega) = -\frac{h^{jk}(\theta) + a_3^{jk}(z + t\rho, \theta)}{2(z + t\rho)^2}\omega_j\omega_k.$$

Lemma 4.1 shows that $K(t, W(t))$ satisfies

$$(4.9) \quad |(\partial_z^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta K)(t, W(t))| \leq Cr(t)^{-2-j}|\omega|^{(2-|\beta|)_+} \quad \text{if } |\beta| \leq 2,$$

and $(\partial_z^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta K)(t, W(t)) = 0$ if $|\beta| \geq 3$. Let $\gamma = (j, \alpha, k, \beta)$, $|\gamma| = 1$, and denote $\partial^\gamma = \partial_z^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta$ for short. By differentiating the Hamilton equation with respect to ∂^γ , we have

$$(4.10) \quad \partial_t \partial^\gamma (W(t) - W(0)) = A(t) \partial^\gamma (W(t) - W(0)) + A(t) \partial^\gamma W(0),$$

where $A(t) = (A_j(t))_{1 \leq j \leq 4} := dH_K(W(t))$ satisfies

$$(4.11) \quad \begin{aligned} |A_1(t)| &\leq Cr(t)^{-2}|\omega|^2, \quad |A_2(t)| \leq Cr(t)^{-2}|\omega|, \\ |A_3(t)| &\leq Cr(t)^{-3}|\omega|^2, \quad |A_4(t)| \leq Cr(t)^{-2}|\omega|^2. \end{aligned}$$

If we put

$$\begin{aligned} f(t) &= (f_1(t), f_2(t), f_3(t), f_4(t)) \\ &:= \left(\frac{\partial^\gamma(z(t) - r)}{r}, \partial^\gamma(\theta(t) - \theta), \partial^\gamma(\rho(t) - \rho), \frac{\partial^\gamma(\omega(t) - \omega)}{r} \right), \end{aligned}$$

then, by using (4.2), (4.9), (4.10) and (4.11), we have

$$(4.12) \quad |f_1(t)| \leq C \int_0^t \left(r^{-1}r(s)^{-2-j}|\omega|^{(2-|\beta|)_+} + r(s)^{-2}r|f(s)| \right) ds.$$

We similarly obtain

$$(4.13) \quad |f_2(t)| \leq C \int_0^t \left(r(s)^{-2-j}|\omega|^{(1-|\beta|)_+} + r(s)^{-2}r|f(s)| \right) ds,$$

$$(4.14) \quad |f_3(t)| \leq C \int_0^t \left(r^{-1}r(s)^{-2-j}|\omega|^{(2-|\beta|)_+} + r(s)^{-2}r|f(s)| \right) ds,$$

$$(4.15) \quad |f_4(t)| \leq C \int_0^t \left(r^{-1}r(s)^{-2-j}|\omega|^{(2-|\beta|)_+} + r(s)^{-2}r|f(s)| \right) ds.$$

Gronwall's inequality then implies

$$(4.16) \quad |f(t)| \leq C \int_0^t r(s)^{-2-j}|\omega|^{(1-|\beta|)_+} ds \lesssim r^{-1-j}|\omega|^{(1-|\beta|)_+} \langle r/t \rangle^{-1},$$

and we obtain the estimate for $\partial^\gamma(\theta(t) - \theta)$. For the proof on other variables, we set

$$g(t) := (r^{-1}\partial^\gamma(z(t) - r), \partial^\gamma(\rho(t) - \rho), r^{-1}\partial^\gamma(\omega(t) - \omega)).$$

Combining the second estimate of (4.3) with (4.9), (4.12), (4.14) and (4.15), we have

$$|g(t)| \lesssim \int_0^t \left(r^{-1-j}r(s)^{-2}|\omega|^{(2-|\beta|)_+} + r(s)^{-2}r|g(s)| \right) ds.$$

Again Gronwall's inequality implies

$$|g(t)| \leq Cr^{-2-j}|\omega|^{(2-|\beta|)_+} \langle r/t \rangle^{-1},$$

and we obtain the estimates for $\partial^\gamma \rho(t)$, $\partial^\gamma \omega(t)$. Moreover the first estimate of (4.3) follows from

$$\partial^\gamma(\dot{r}(t) - \rho) = \partial^\gamma(\rho(t) - \rho) = O(r^{-2-j}|\omega|^{(2-|\beta|)_+} \langle r/t \rangle^{-1}),$$

since $\langle r/t \rangle^{-1}$ is monotone increasing with respect to t . Next, let l be a non-negative integer and suppose that (4.13) holds for any γ with $|\gamma| \leq l$. Let $\gamma = (j, \alpha, k, \beta)$, $|\gamma| = l + 1$. A direct computation yields

$$\partial_t \partial^\gamma (W(t) - W(0)) = A(t) \partial^\gamma (W(t) - W(0)) + A(t) \partial^\gamma W(0) + B(t),$$

where $B(t)$ is a linear combination of products of

$$(\partial_z^{\alpha_1} \partial_\theta^{\alpha'} \partial_\rho^{\alpha_{d+1}} \partial_\omega^{\alpha''} dH_K)(W(t)),$$

$$\partial^{\gamma_1^1} z(t) \cdots \partial^{\gamma_{\alpha_1}^1} z(t) \times \cdots \times \partial^{\gamma_1^{2d}} \omega_{d-1}(t) \cdots \partial^{\gamma_{\alpha_{2d}}^{2d}} \omega_{d-1}(t),$$

with $\alpha, \gamma_1^1, \gamma_2^1, \dots, \gamma_{\alpha_{2d}}^{2d} \in \mathbb{Z}_+^{2d}$ such that

$$\alpha = (\alpha_1, \dots, \alpha_{2d}) = (\alpha_1, \alpha', \alpha_{d+1}, \alpha'') \in \mathbb{Z}_+^{1+(d-1)+1+(d-1)}, \quad 1 \leq |\alpha| \leq |\gamma|,$$

$$\gamma_1^1 + \gamma_2^1 + \cdots + \gamma_{\alpha_{2d}}^{2d} = \gamma, \quad 1 \leq \gamma_l^k \leq |\gamma| - 1.$$

The induction hypothesis implies that each entry of $B(t) = (B_j(t))$ satisfies

$$\begin{aligned} & |B_1(t)| + |B_4(t)| \\ & \leq C \sum_{\substack{1 \leq |\alpha| \leq |\gamma|, \\ |\alpha''| \leq 2}} r(t)^{-2-\alpha_1} r^{2-|\alpha''|} |\nu|^{2-|\alpha''|} r^{\alpha_1+|\alpha''|} r^{-j-|\beta|} |\nu|^{(2|\alpha|-|\alpha'|+|\alpha''|-|\beta|)_+} \\ & \leq Cr(t)^{-2} r^{-j-|\beta|} |\nu|^{(2-|\beta|)_+}, \end{aligned}$$

and we similarly obtain

$$|B_2(t)| \leq Cr(t)^{-2} r^{-j-|\beta|} |\nu|^{(1-|\beta|)_+}, \quad |B_3(t)| \leq Cr(t)^{-3} r^{-j-|\beta|} |\nu|^{(2-|\beta|)_+},$$

where $\nu = \omega/r$. By a similar argument as that in the case for $|\gamma| = 1$, we obtain the assertion. The proof of (4.4) is more simpler than (4.3), and we hence omit it.

Finally, we prove (4.5). Note that if $R > 0$ is large enough, then

$$\dot{\rho}(t) = r(t)^{-3} \left(h^{jk}(\theta(t)) + O(r(t)^{-\mu}) \right) \omega_j(t) \omega_k(t) \geq 0, \quad t \in \mathbb{R}.$$

Therefore, integrating $\dot{\rho}(t)$ with respect to t , we have (4.5). $\square$

Proposition 4.2 implies that the trajectory belongs to a fixed coordinate neighborhood as long as either $(r, \theta, \rho, \omega) \in \Gamma^\pm(R, \tilde{U}_\kappa, J, \sigma)$ and $0 \leq \pm t \leq rt_0$ or $(r, \theta, \rho, \omega) \in \Gamma_s^\pm(R, \tilde{U}_\kappa, J, \varepsilon_0)$ and $\pm t \geq 0$, respectively, provided $t_0, \varepsilon_0 > 0$ are small enough. It also follows from Proposition 4.2 that the outgoing and incoming regions are invariant under the geodesic flow (except the choice of the boundary decomposition). More precisely we have the following.

Corollary 4.3. *Let $R_0 > 0$ be as in Lemma 4.1. Fix $U_\kappa, \tilde{U}_\kappa, U_{\kappa'}, \tilde{U}_{\kappa'}$ and $\kappa' : V_{\kappa'} \rightarrow U_{\kappa'}$ so that $\tilde{U}_\kappa \Subset U_\kappa$ and $\tilde{U}_{\kappa'} \Subset U_{\kappa'}$. Then there exists $C > 0$ such that, for all $R \geq R_0$, $T_0 > 0$ and $(r, \theta, \rho, \omega) \in \Gamma^+(R, \tilde{U}_\kappa, J, \sigma)$, we have*

$$(4.17) \quad \kappa'_* \exp T_0 H_{p_\kappa} \kappa_*^{-1}(r, \theta, \rho, \omega) \in \Gamma^+((R + T_0)/C, \tilde{U}_{\kappa'}, J, \sigma).$$

In particular we can find $t_0 > 0$ such that, for all $(r, \theta, \rho, \omega) \in \Gamma^+(R, \tilde{U}_\kappa, J, \sigma)$ and $0 \leq t \leq rt_0$,

$$(4.18) \quad \exp t H_{p_\kappa}(r, \theta, \rho, \omega) \in \Gamma^+((R + t)/C, \tilde{U}_{\kappa, \sqrt{t_0}}, J, \sigma),$$

where $\tilde{U}_{\kappa, \sqrt{t_0}}$ is the $\sqrt{t_0}$ -neighborhood of $\tilde{U}_\kappa$. Moreover there exists a small constant $\varepsilon_0 > 0$ such that if $0 < \varepsilon \leq \varepsilon_0$, $(r, \theta, \rho, \omega) \in \Gamma_s^+(R, \tilde{U}_\kappa, J, \varepsilon)$ and $t \geq 0$, then

$$(4.19) \quad \exp t H_{p_\kappa}(r, \theta, \rho, \omega) \in \Gamma_s^+(R/C, \tilde{U}_\kappa, J, \varepsilon).$$

When $t < 0$, analogous results hold in the incoming region.

Proof. We first prove (4.18) and (4.19). By Lemma 4.1, we can find $R > 0$ large enough such that

$$\dot{\rho}(t) = \frac{1}{r(t)^3} \left(h^{jk}(\theta(t)) - O(r(t)^{-\mu}) \right) \omega_j(t) \omega_k(t) \geq 0, \quad r > R, \quad t \geq 0,$$

and hence $\rho(t) \geq \rho$ for $t \geq 0$. Therefore, (4.18) follows from (4.2), (4.4) and the energy conservation $p_\kappa(r(t), \theta(t), \rho(t), \omega(t)) = p_\kappa(r, \theta, \rho, \omega)$. Next, let $(r, \theta, \rho, \omega) \in \Gamma_s^+(R, \tilde{U}_\kappa, J, \varepsilon)$. Since $\langle r/t \rangle^{-1} \leq 1$ and

$$|\omega/r| \leq C\varepsilon \quad \text{on} \quad \Gamma_s^\pm(R, \tilde{U}_\kappa, J, \varepsilon),$$

(4.19) follows from (4.3). To prove (4.17), divide the time interval $[0, T_0]$ as

$$[0, T_0] \subset [0, Rt_0] \cup [Rt_0, 2Rt_0] \cup \cdots \cup [T_0 - Rt_0, T_0].$$

In each interval $[jRt_0, (j+1)Rt_0]$, the flow is contained some fixed coordinate neighborhood. Since the outgoing region is invariant under coordinate transformations, applying (4.18) on each chart, we have the assertion. $\square$

5. THE ISOZAKI-KITADA PARAMETRIX

In this section we construct the Isozaki-Kitada parametrix of $e^{-ithP} \text{Op}_\kappa(a_s^\pm)$, where the symbols a_s^+ and a_s^- are supported in the strongly outgoing incoming regions, respectively. Though the method of construction is similar to the case of an asymptotically hyperbolic manifold, the class of the phase function of the parametrix becomes even worse (see Remark 5.2). We thus give the full details of the proof.

By Corollary 4.3, we can always work on one fixed coordinate chart (U_κ, κ) , and hence drop the subscript κ if there is no confusion. Fix an open set $\tilde{U}_\kappa \Subset U_\kappa$ with $\tilde{\Psi}_\kappa = 1$ on $(2, \infty) \times \tilde{U}_\kappa$ and an open interval $J \Subset (0, \infty)$ arbitrarily. We denote $\Gamma^\pm(R, \varepsilon) = \Gamma_s^\pm(R, \tilde{U}_\kappa, J, \varepsilon)$ for short. For a large parameter $\lambda \geq 1$, we also denote $\Gamma_s^\pm(\lambda) = \Gamma_s^\pm(R/\lambda, \tilde{U}_\kappa, J, \lambda\varepsilon)$. Notice that $\Gamma_s^\pm(\lambda)$ and $\Omega_s^\pm(\lambda)$ is increasing with respect to λ : $\Gamma_s^\pm(R, \varepsilon) \subset \Gamma_s^\pm(\lambda_1) \subset \Gamma_s^\pm(\lambda_2)$, $1 \leq \lambda_1 < \lambda_2$.

5.1. Fourier integral operators for the Isozaki-Kitada parametrix. We here study Fourier integral operators (FIO's for short) on $\mathbb{R}^d$ which will be used to construct the Isozaki-Kitada parametrix. The first step is to construct the corresponding phase function.

Theorem 5.1. *There exist $R_0, \lambda_0 > 0$ large enough and $\varepsilon_0 > 0$ small enough such that for all $R, \varepsilon, \lambda > 0$ satisfying $\lambda \geq \lambda_0$, $R \geq \lambda R_0$ and $0 < \varepsilon \leq \varepsilon_0/\lambda$, we can find smooth and real-valued functions $S^\pm \in C^\infty(\mathbb{R}^{2d}, \mathbb{R})$ satisfying the Eikonal equation:*

$$p(r, \theta, \partial_r S^\pm(r, \theta, \rho, \omega), \partial_\theta S^\pm(r, \theta, \rho, \omega)) = \frac{1}{2} \rho^2, \quad (r, \theta, \rho, \omega) \in \Gamma_s^\pm(R, \varepsilon).$$

If we put $\varphi^\pm(r, \theta, \rho, \omega) := S^\pm(r, \theta, \rho, \omega) - r\rho - \theta \cdot \omega$, then $\varphi^\pm$ satisfy $\text{supp } \varphi^\pm \subset \Gamma_s^\pm(\lambda)$, and

$$(5.1) \quad |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \varphi^\pm(r, \theta, \rho, \omega)| \leq C r^{1-j-|\beta|} |\omega/r|^{(2-|\beta|)_+} \quad \text{on} \quad \Gamma_s^\pm(\lambda).$$

Furthermore, we can write

$$(5.2) \quad \varphi^\pm(r, \theta, \rho, \omega) = \frac{1}{2r\rho} q_0(\theta, \omega) + R^\pm(r, \theta, \rho, \omega) \quad \text{on} \quad \Gamma_s^\pm(R, \varepsilon),$$

where $q_0(\theta, \omega) := h^{jk}(\theta) \omega_j \omega_k$ and $R^\pm(r, \theta, \rho, \omega)$ satisfy

$$|\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta R^\pm(r, \theta, \rho, \omega)| \leq C r^{1-j-|\beta|} (|\omega/r|^{(3-|\beta|)_+} + r^{-\mu} |\omega/r|^{(2-|\beta|)_+}).$$

Here the constant $C > 0$ can be taken uniformly with respect to R, ε and λ .

Remark 5.2. We remark that $\varphi^\pm$ and its derivatives with respect to (θ, ρ) are not bounded with respect to r even for the perfectly conic ($a_{jk} \equiv 0$) case. This condition is even worse than that of the asymptotically flat case or asymptotically hyperbolic case. Indeed, we see that $\partial_x^\alpha \partial_\xi^\beta (S^\pm(x, \xi) - x \cdot \xi) = O(\langle x \rangle^{1-\varepsilon-|\alpha|})$ with some $\varepsilon > 0$ in the asymptotically flat case [4], and $\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (S^\pm(r, \theta, \rho, \omega) - r\rho - \theta \cdot \omega) = O(e^{-r|\beta|})$ in the asymptotically hyperbolic case [3]. We

refer to [4, 3] for more details. We also refer to the original paper by Isozaki-Kitada [12] which was concerned with a long-range potential scattering theory on $\mathbb{R}^d$.

We also note that (5.1) implies

$$(5.3) \quad |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \varphi^\pm(r, \theta, \rho, \omega)| \leq C \langle r \rangle^{1-j-|\beta|} \varepsilon_0^{(2-|\beta|)_+} \quad \text{on } \mathbb{R}^{2d},$$

since $|\omega/r| \leq C\lambda\varepsilon \leq C\varepsilon_0$ on $\text{supp } \varphi^\pm (\subset \Gamma_s^\pm(\lambda))$. For sufficiently large $R_0 > 0$ and sufficiently small $\varepsilon_0 > 0$, we hence have

$$(5.4) \quad 1/2 < |\det {}^t \partial_{\rho, \omega} \partial_{r, \theta} S^\pm(r, \theta, \rho, \omega)| < 3/2 \quad \text{on } \mathbb{R}^{2d},$$

though $|{}^t \partial_{\rho, \omega} \partial_{r, \theta} S^\pm - \text{Id}|$ is not bounded with respect to r in general. This estimate is crucial to obtain L^2 -boundedness of FIO's.

To prove Theorem 5.1, we prepare several lemmas.

Lemma 5.3. *There exist $R_0 > 0$ large enough and $\varepsilon_0 > 0$ small enough such that, for all $R, \varepsilon > 0$, $\lambda_0 \geq 1$ satisfying $R \geq \lambda_0 R_0$, $\varepsilon \leq \varepsilon_0/\lambda_0$ and all $\pm t \geq 0$, the maps*

$$f^\pm(t) : (r, \theta, \rho, \omega) \mapsto (r, \theta, \rho(t, r, \theta, \rho, \omega), \omega(t, r, \theta, \rho, \omega))$$

are diffeomorphisms from $\Gamma_s^\pm(\lambda_0)$ onto its range, respectively. Moreover, for sufficiently large $\lambda_0 > 0$, we have

$$(5.5) \quad \Gamma_s^\pm(R, \varepsilon) \subset f^\pm(t) (\Gamma_s^\pm(\lambda_0)), \quad \pm t \geq 0.$$

Proof. We prove the lemma for $t \geq 0$ only, and the proof for $t \leq 0$ is similar. Let $F : (r, \theta, \rho, \omega) \mapsto (r, \theta, \rho, \omega/r)$ be a global diffeomorphism from $(0, \infty) \times \mathbb{R}^{d-1}$ onto itself, and we define for $(r, \theta, \rho, \nu) \in F\Gamma_s^+(\lambda_0)$,

$$\tilde{f}^+(t)(r, \theta, \rho, \nu) = (r, \theta, \tilde{\rho}(t), \tilde{\omega}(t)) := (F \circ f^+(t) \circ F^{-1})(r, \theta, \rho, \nu),$$

where $\nu = \omega/r$. By (4.3), we can choose $R_0, \varepsilon_0 > 0$ and $C_0 > 0$ such that

$$(5.6) \quad \begin{aligned} |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\nu^\beta (\tilde{\rho}(t) - \rho)| &\leq C_0 r^{-j} |\nu|^{(2-|\beta|)_+} \leq C_0 R_0^{-j} \varepsilon_0^{(2-|\beta|)_+}, \\ |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\nu^\beta (\tilde{\omega}(t) - \nu)| &\leq C_0 r^{-j} |\nu|^{(2-|\beta|)_+} \leq C_0 R_0^{-j} \varepsilon_0^{(2-|\beta|)_+}, \end{aligned}$$

and hence

$$(5.7) \quad |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\nu^\beta (\partial \tilde{f}^+(t) - \text{Id})| \leq C \varepsilon_0 < 1/2,$$

uniformly with respect to $(r, \theta, \rho, \nu) \in F\Gamma_s^+(\lambda_0)$, where $\partial \tilde{f}^+(t)$ is the differential at (r, θ, ρ, ν) . Choose $\chi^+ \in C^\infty(\mathbb{R}^{2d})$ so that $0 \leq \chi^+ \leq 1$, $\chi^+ \equiv 1$ on $F\Gamma_s^+(\lambda_0)$, $\text{supp } \chi^+ \subset F\Gamma_s^+(2\lambda_0)$, and that

$$(5.8) \quad |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\nu^\beta \chi^+(r, \theta, \rho, \nu)| \leq C_1 \langle r \rangle^{-j} \quad \text{on } \mathbb{R}^{2d},$$

and define $\tilde{f}_\chi^+(t)(r, \theta, \rho, \nu) = (r, \theta, \tilde{\rho}_\chi(t), \tilde{\omega}_\chi(t))$ by

$$\tilde{f}_\chi^+(t)(r, \theta, \rho, \nu) := (r, \theta, (1 - \chi^+) \rho + \chi^+ \tilde{\rho}(t), (1 - \chi^+) \nu + \chi^+ \tilde{\omega}(t)).$$

Since $\tilde{f}_\chi^+(t) = \tilde{f}^+(t)$ on $F\Gamma_s^+(\lambda_0)$, $\tilde{f}_\chi^+(t) = \text{Id}$ outside $F\Gamma_s^+(2\lambda_0)$, we have

$$|\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\nu^\beta \tilde{f}_\chi^+(t)| \leq C, \quad (r, \theta, \rho, \omega) \in \mathbb{R}^{2d}, \quad j + |\alpha| + k + |\beta| \geq 1.$$

Moreover, (5.6) and (5.8) imply

$$(5.9) \quad \begin{aligned} |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\nu^\beta (\tilde{\rho}_\chi(t) - \rho)| &\leq C_0 C_1 \langle r \rangle^{-j} |\nu|^{(2-|\beta|)_+} \leq C_2 R_0^{-j} \varepsilon_0^{(2-|\beta|)_+}, \\ |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\nu^\beta (\tilde{\omega}_\chi(t) - \nu)| &\leq C_0 C_1 \langle r \rangle^{-j} |\nu|^{(2-|\beta|)_+} \leq C_2 R_0^{-j} \varepsilon_0^{(2-|\beta|)_+} \end{aligned}$$

on $\mathbb{R}^{2d}$ with some $C_2 > 4^{j+2} C_0 C_1$. $\tilde{f}_\chi^+(t)$ hence satisfies the same estimate as (5.7) on $\mathbb{R}^{2d}$ provided $R_0 > 0$ large enough and $\varepsilon_0 > 0$ small enough. By the Hadamard global inverse mapping theorem, we see that $\tilde{f}_\chi^+(t)$ is a diffeomorphism from $\mathbb{R}^{2d}$ onto its range. Since $f^+(t) = F^{-1} \circ \tilde{f}^+(t) \circ F$, $f^+(t)$ is a diffeomorphism from $\Gamma_s^+(\lambda_0)$ onto its range.

Next, we define $f_\chi^+(t) := F^{-1} \circ \tilde{f}_\chi^+ \circ F$, and shall prove

$$(5.10) \quad \Gamma_s^+(R, \varepsilon) \subset f_\chi^+(t) \left(\Gamma_s^+(\lambda_0) \right), \quad t \geq 0,$$

for sufficiently large $\lambda_0 > 0$. Since $f_\chi^+(t)$ is bijective, it suffices to show that

$$\mathbb{R}^{2d} \setminus \Gamma_s^+(R, \varepsilon) \supset f_\chi^+(t) \left(\mathbb{R}^{2d} \setminus \Gamma_s^+(\lambda_0) \right), \quad t \geq 0.$$

Suppose that $Z := (r, \theta, \rho, \omega) \in \mathbb{R}^{2d} \setminus \Gamma_s^+(\lambda_0)$. If $Z \in \mathbb{R}^{2d} \setminus \Gamma_s^+(2\lambda_0)$, then

$$f_\chi^+(t)(Z) = Z \in \mathbb{R}^{2d} \setminus \Gamma_s^+(2\lambda_0) \subset \mathbb{R}^{2d} \setminus \Gamma_s^+(R, \varepsilon).$$

If $Z \in \Gamma_s^+(2\lambda_0) \setminus \Gamma_s^+(\lambda_0)$, then we have

$$p(Z) \in J_{4\lambda_0^2\varepsilon^2} \setminus J_{\lambda_0^2\varepsilon^2}, \quad \sqrt{1 - 4\lambda_1^2\varepsilon^2} \leq \frac{\rho}{\sqrt{2p(Z)}} \leq \sqrt{1 - \lambda_0^2\varepsilon^2}.$$

Since $|p(f_\chi^+(t)(Z)) - p(Z)| \leq C|\omega/r|^2 \leq C\varepsilon_0^2$, Proposition 4.2 and the above argument imply that if ε_0 is small enough and λ_0 is large enough, then we obtain

$$p(f_\chi^+(t)(Z)) \notin J_{\varepsilon^2}, \quad \frac{\rho_\chi(t, Z)}{\sqrt{2p(f_\chi^+(t)(Z))}} < \sqrt{1 - \varepsilon^2} \quad \text{for } t \geq 0,$$

which implies (5.10). Here $\rho_\chi(t, Z) = (1 - \chi^+)\rho + \chi^+\rho(t, Z)$. Since $f_\chi^+(t) = f^+(t)$ on $\Gamma_s^+(\lambda_0)$, (5.5) follows from (5.10). $\square$

Let $\pm t \geq 0$ and $\Gamma_s^\pm(R, \varepsilon) \ni (r, \theta, \rho, \omega) \mapsto (r, \theta, \hat{\rho}^\pm(t), \hat{\omega}^\pm(t)) \in \Gamma_s^\pm(\lambda_0)$ the inverse mappings of $f^\pm(t)$, respectively.

Lemma 5.4. For $0 \leq \pm s \leq \pm t$ and $(r, \theta, \rho, \omega) \in \Gamma_s^\pm(R, \varepsilon)$, we define

$$(r_t^\pm(s), \theta_t^\pm(s), \rho_t^\pm(s), \omega_t^\pm(s)) := (r, \theta, \rho, \omega)(s, r, \theta, (\hat{\rho}^\pm, \hat{\omega}^\pm)(t, r, \theta, \rho, \omega)).$$

We then have, for all $(j, \alpha, k, \beta) \in \mathbb{Z}_+^{2d}$,

$$\begin{aligned} |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\hat{\rho}^\pm(t) - \rho)| &\leq C r^{-j-|\beta|} |\omega/r|^{(2-|\beta|)_+}, \\ |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\hat{\omega}^\pm(t) - \omega)| &\leq C r^{1-j-|\beta|} |\omega/r|^{(2-|\beta|)_+}, \end{aligned}$$

uniformly with respect to $(r, \theta, \rho, \omega) \in \Gamma_s^\pm(R, \varepsilon)$ and $\pm t \geq 0$.

Proof. Since $(r, \theta, \hat{\rho}^\pm(t), \hat{\omega}^\pm(t)) \in \Gamma_s^\pm(\lambda_0)$, we have

$$\begin{aligned} |\hat{\rho}^\pm(t) - \rho| + r^{-1} |\hat{\omega}^\pm(t) - \omega| &= |\rho_t(0) - \rho_t^\pm(t)| + r^{-1} |\omega_t(0) - \omega_t^\pm(t)| \\ (5.11) \quad &\leq C \sup_{\Gamma_s^\pm(\lambda_0)} (|\rho(t) - \rho| + r^{-1} |\omega(t) - \omega|) \\ &\leq C |\omega/r|^2, \end{aligned}$$

where C is independent of R_0, ε_0 and λ_0 . We next consider the derivatives. Let $\gamma = (j, \alpha, k, \beta)$, $|\gamma| = 1$. Applying $\partial^\gamma = \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta$ to the equality

$$(\rho, \omega) = (\rho, \omega)(t, r, \theta, \hat{\rho}^\pm(t), \hat{\omega}^\pm(t)),$$

we have

$$(5.12) \quad A(Z^\pm(t)) \begin{pmatrix} \partial^\gamma (\hat{\rho}^\pm(t) - \rho) \\ r^{-1} \partial^\gamma (\hat{\omega}^\pm(t) - \omega) \end{pmatrix} = \begin{pmatrix} \partial^\gamma (\rho - \rho(t)) \\ r^{-1} \partial^\gamma (\omega - \omega(t)) \end{pmatrix} \Big|_{Z^\pm(t)},$$

where $Z^\pm(t) = (r, \theta, \hat{\rho}^\pm(t), \hat{\omega}^\pm(t))$ and

$$A(Z^\pm(t)) = \begin{pmatrix} (\partial_\rho \rho)(t, Z^\pm(t)) & r(\partial_\omega \rho)(t, Z^\pm(t)) \\ r^{-1}(\partial_\rho \omega)(t, Z^\pm(t)) & (\partial_\omega \omega)(t, Z^\pm(t)) \end{pmatrix}.$$

(4.3) and (5.11) show that $A(Z^\pm(t))$ are invertible, $A(Z^\pm(t))$ and $A(Z^\pm(t))^{-1}$ are bounded on $\Gamma_s^\pm(R, \varepsilon)$ and the right hand side of (5.12) is bounded by

$$r^{-j-|\beta|} |\omega/r|^{(2-|\beta|)_+}.$$

The proof for higher derivatives is obtained by a simple induction with respect to $|\gamma|$, and we omit the details. $\square$

The following easily follows from Lemma 4.1, Proposition 4.2 and Lemma 5.4.

Corollary 5.5. *For all $\pm t \geq 0$, $(r, \theta, \rho, \omega) \in \Gamma_s^\pm(R, \varepsilon)$, we have*

$$r_t^\pm(t) \geq C^{-1}(r + |t|).$$

Moreover, for all $(j, \alpha, k, \beta) \in \mathbb{Z}_+^{2d}$,

$$\begin{aligned} |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (r_t^\pm(t) - r \mp t\rho)| &\leq Cr^{1-j-|\beta|} |\omega/r|^{(2-|\beta|)_+}, \\ |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\theta_t^\pm(t) - \theta)| &\leq Cr^{-j-|\beta|} |\omega/r|^{(1-|\beta|)_+}. \end{aligned}$$

Proof of Theorem 5.1. We give the proof for the case $t \geq 0$, and the proof for the case $t \leq 0$ is analogous. Define $\Lambda^+(t) \in C^\infty(\Gamma_s^+(R, \varepsilon))$ for $t \geq 0$ by

$$(5.13) \quad \Lambda^+(t, r, \theta, \rho, \omega) = r_t^+(t)\rho + \theta_t^+(t) \cdot \omega + \int_0^t L(r_t^+(s), \theta_t^+(s), \rho_t^+(s), \omega_t^+(s)) ds,$$

where $L(r, \theta, \rho, \omega) = \partial_\rho p(r, \theta, \rho, \omega)\rho + \partial_\omega p(r, \theta, \rho, \omega) \cdot \omega - p(r, \theta, \rho, \omega)$ is the Lagrangian associated with p . Note that the smoothness of $\Lambda^+(t)$ follows from the smoothness of $(r_t^+, \theta_t^+, \rho_t^+, \omega_t^+)$. By the standard Hamilton-Jacobi theory, $\Lambda^+(t)$ solves

$$\begin{cases} \partial_t \Lambda^+(t) = p(r, \theta, \partial_r \Lambda^+(t), \partial_\theta \Lambda^+(t)), \\ \Lambda^+(0) = r\rho + \theta \cdot \omega, \\ \partial_{r, \theta, \rho, \omega} \Lambda^+(t) = (\hat{\rho}^+(t), \hat{\omega}^+(t), r_t^+(t), \theta_t^+(t)). \end{cases}$$

Put $F^+(t) = \Lambda^+(t) - \frac{1}{2}t\rho^2$. The energy conservation, namely

$$p(r, \theta, \hat{\rho}^+(t), \hat{\omega}^+(t)) = p(r_t^+(t), \theta_t^+(t), \rho, \omega),$$

implies

$$\partial_t F^+(t) = \frac{1}{2r_t^+(t)^2} \left(h^{jk}(\theta_t^+(t)) + a^{jk}(r_t^+(t), \theta_t^+(t)) \right) \omega_j \omega_k^0.$$

By Lemma 5.4 and Corollary 5.5, we have

$$|\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \partial_t F^+(t)| \leq C(r + |t|)^{-2-j} r^{2-|\beta|} |\omega/r|^{(2-|\beta|)_+},$$

and hence

$$\left| \int_0^\infty \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \partial_t F^+(t) dt \right| \leq Cr^{1-j-|\beta|} |\omega/r|^{(2-|\beta|)_+}$$

for all $t \geq 0$, $(r, \theta, \rho, \omega) \in \Gamma_s^+(R, \varepsilon)$ and $(j, \alpha, k, \beta) \in \mathbb{Z}_+^{2d}$. If we put $q_0(\theta, \omega) = h^{jk}(\theta) \omega_j \omega_k$, then the mean value theorem and Corollary 5.5 imply

$$\begin{aligned} &\left| \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \left(\frac{1}{2r_t^+(t)^2} q_0(\theta_t^+(t), \omega) - \frac{1}{2(r+t\rho)^2} q_0(\theta, \omega) \right) \right| \\ &\leq C(r + |t|)^{-2} r^{2-j-|\beta|} |\omega/r|^{(3-|\beta|)_+}. \end{aligned}$$

Therefore, $\partial_t F^+(t)$ can be written in the form

$$\frac{1}{2(r+t\rho)^2} q_0(\theta, \omega) + \tilde{R}^+(t)$$

with $\tilde{R}^+(t) \in C^\infty(\Gamma^+(R, \varepsilon))$ satisfying

$$\begin{aligned} &|\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \tilde{R}^+(t, r, \theta, \rho, \omega)| \\ &\leq C(r + |t|)^{-2} r^{2-j-|\beta|} \left(|\omega/r|^{(3-|\beta|)_+} + r^{-\mu} |\omega/r|^{(2-|\beta|)_+} \right). \end{aligned}$$

Define $\tilde{\varphi}^+, \tilde{S}^+$ on $\Gamma_s^+(R, \varepsilon)$ by

$$\begin{aligned}\tilde{\varphi}^+(r, \theta, \rho, \omega) &:= \int_0^\infty \partial_t F^+(t, r, \theta, \rho, \omega) dt, \\ \tilde{S}^+(r, \theta, \rho, \omega) &:= r\rho + \theta \cdot \omega + \tilde{\varphi}^+(r, \theta, \rho, \omega).\end{aligned}$$

The above argument shows that $\tilde{\varphi}^+, \tilde{S}^+$ are smooth and $\tilde{\varphi}^+$ satisfies (5.1) and (5.2) on $\Gamma_s^+(R, \varepsilon)$. Moreover, we have

$$\partial_{r,\theta} \tilde{S}^+ = \lim_{t \rightarrow +\infty} \partial_{r,\theta} \Lambda^+(t) \quad \text{on } \Gamma_s^+(R, \varepsilon).$$

By using the energy conservation and Corollary 5.5, we see that $\tilde{S}^+$ satisfies the Eikonal equation on $\Gamma_s^+(R, \varepsilon)$:

$$\begin{aligned}p(r, \theta, \partial_r \tilde{S}^+, \partial_\theta \tilde{S}^+) &= \lim_{t \rightarrow +\infty} p(r, \theta, \partial_r \Lambda^+(t), \partial_\theta \Lambda^+(t)) \\ &= \lim_{t \rightarrow +\infty} p(r_t^+, \theta_t^+, \rho, \omega) \\ &= \rho^2/2.\end{aligned}$$

Choosing a smooth cut-off function χ^+ on $\mathbb{R}^{2d}$ so that $0 \leq \chi^+ \leq 1$, $\chi^+ \equiv 1$ on $\Gamma_s^+(R, \varepsilon)$, $\text{supp } \chi^+ \subset \Gamma_s^+(\lambda)$, $\lambda \geq \lambda_0$ and that

$$|\partial_r^j \partial_\theta^\alpha \partial_\rho^k \chi(r, \theta, \rho, \omega)| \leq C \langle r \rangle^{-j-|\alpha|} \quad \text{on } \mathbb{R}^{2d},$$

we define $\varphi^+, S^+ \in C^\infty(\mathbb{R}^{2d})$ by $\varphi^+ = \chi^+ \tilde{\varphi}^+$, $S^+ := r\rho + \theta \cdot \omega + \varphi^+$. Clearly, φ^+ and S^+ satisfy the statements of Theorem 5.1. $\square$

Definition 5.6. For $a^\pm \in S_{\text{sc}}(\Gamma^\pm(R, \varepsilon))$ and $h \in (0, 1]$, we define the FIO's for the Isozaki-Kitada parametrix by

$$I_{\text{IK}}^\pm(a^\pm)u(r, \theta) = (2\pi h)^{-d} \int e^{\frac{i}{h}(S^\pm(r, \theta, \rho, \omega) - r' \rho - \theta' \cdot \omega)} a^\pm(r, \theta, \rho, \omega) u(r', \theta') dr' d\theta' d\rho d\omega.$$

The following theorem shows that $I_{\text{IK}}^\pm(a^\pm)$ are bounded on $L^2(\mathbb{R}^d)$ uniformly with respect to $R > 0$ and $h \in (0, 1]$.

Proposition 5.7. Let $R_0, \varepsilon_0, \lambda_0 > 0$ be as in Theorem 5.1, $\lambda \geq \lambda_0$, $R \geq \lambda^4 R_0$ and $0 < \varepsilon \leq \lambda^{-4} \varepsilon_0$. Then, for all $N \geq 0$ and $a^\pm, b^\pm \in S_{\text{sc}}(\Gamma^\pm(\lambda^3))$, there exist symbols $c_j^\pm \in S_{\text{sc}}(\Gamma^\pm(\lambda^4))$, $j = 0, 1, \dots, N$, such that if we put $c_h^\pm = \sum_{j=0}^N h^j c_j^\pm$, then

$$\|I_{\text{IK}}^\pm(a^\pm)I_{\text{IK}}^\pm(b^\pm)^* - c_h^\pm(r, \theta, hD_r, hD_\theta)\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \leq C_N h^{N+1}, \quad h \in (0, 1].$$

In particular, we have

$$\|I_{\text{IK}}^\pm(a^\pm)\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \leq C, \quad h \in (0, 1].$$

Here $C_N, C > 0$ may be taken uniformly with respect to R and h .

The following shows that any elliptic FIO (in the semiclassical sense) has a microlocal approximate inverse.

Proposition 5.8. Let $R_0, \varepsilon_0, \lambda_0 > 0$ be as in Theorem 5.1, $\lambda \geq \lambda_0$, $R \geq \lambda^4 R_0$, $0 < \varepsilon \leq \lambda^{-4} \varepsilon_0$ and $N \geq 0$ a non-negative integer. Choose arbitrarily sequences of symbols $(a_j^\pm)_{0 \leq j \leq N} \subset S_{\text{sc}}(\Gamma^\pm(\lambda^3))$ satisfying

$$a_0^\pm > C_0^{-1} \quad \text{on } \Omega_s^\pm(\lambda)$$

with some fixed $C_0 > 0$, respectively. Then for all $c^\pm \in S_{\text{sc}}(\Gamma^\pm(R, \varepsilon))$, there exist sequences of symbols $(b_j^\pm)_{0 \leq j \leq N} \subset S_{\text{sc}}(\Gamma^\pm(\lambda))$ such that

$$\|I_{\text{IK}}^\pm(a^\pm)I_{\text{IK}}^\pm(b^\pm)^* - c^\pm(r, \theta, hD_r, hD_\theta)\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \leq C_N h^{N+1}, \quad h \in (0, 1],$$

where $a_h^\pm = \sum_{j=0}^N h^j a_j^\pm$ and $b_h^\pm = \sum_{j=0}^N h^j b_j^\pm$. Moreover $C_N > 0$ can be taken uniformly with respect to R and h .

To prove the above two propositions, we need the following lemma.

Lemma 5.9. Define $(\rho_1^\pm, \omega_1^\pm)(r, \theta, \rho, \omega, r', \theta') : \mathbb{R}^{3d} \rightarrow \mathbb{R}^d$ by

$$(5.14) \quad (\rho_1^\pm, \omega_1^\pm) = \int_0^1 (\partial_{r,\theta} S^\pm)(r' + \sigma(r - r'), \theta' + \sigma(\theta - \theta'), \rho, \omega) d\sigma.$$

Then the followings hold for all $\lambda \geq \lambda_0$, $R \geq \lambda R_0$ and $0 < \varepsilon \leq \varepsilon_0/\lambda$.

(i) If $(r, \theta, \rho, \omega) \in \Gamma_s^\pm(R, \varepsilon)$, then

$$(5.15) \quad (r, \theta, \rho_1^\pm, \omega_1^\pm)|_{r'=r, \theta'=\theta} \in \Gamma_s^\pm(\lambda).$$

Conversely, if $(r, \theta, \rho_1^\pm, \omega_1^\pm)|_{r'=r, \theta'=\theta} \in \Gamma_s^\pm(R, \varepsilon)$, then

$$(5.16) \quad (r, \theta, \rho, \omega) \in \Gamma_s^\pm(\lambda).$$

(ii) We set $\tilde{\Gamma}_s^\pm(\lambda) := \{(r, \theta, \rho, \omega, r', \theta') \in \mathbb{R}^{3d}; (r, \theta, \rho, \omega), (r', \theta', \rho, \omega) \in \Gamma^\pm(\lambda)\}$ and $\tilde{\Gamma}_s^\pm(R, \varepsilon) := \tilde{\Gamma}_s^\pm(1)$. Then, the map $(r, \theta, \rho, \omega, r', \theta') \mapsto (r, \theta, \rho_1^\pm, \omega_1^\pm, r', \theta')$ is a diffeomorphism from $\tilde{\Gamma}_s^\pm(\lambda)$ onto its range, and satisfies

$$\pi_{\rho, \omega}(\tilde{\Gamma}_s^\pm(R, \varepsilon)) \subset (\rho_1^\pm, \omega_1^\pm)(\pi_{\rho, \omega}(\tilde{\Gamma}_s^\pm(\lambda))),$$

respectively. Denoting by $(\rho_2^\pm, \omega_2^\pm)$ the corresponding inverse, the same properties as in (i) hold with $(\rho_1^\pm, \omega_1^\pm)$ replaced by $(\rho_2^\pm, \omega_2^\pm)$, respectively.

(iii) If $\lambda, \lambda' \geq \lambda_0$, $R \geq \max(\lambda, \lambda')R_0$ and $0 < \varepsilon \leq \min(\lambda^{-1}, \lambda'^{-1})\varepsilon_0$, then we have

$$(5.17) \quad |\partial^\gamma(\rho_2^\pm - \rho)| + \min(r, r')^{-1} |\partial^\gamma(\omega_2^\pm - \omega)| \\ \leq C \min(r, r')^{-j-j'-|\beta|} (\max(\lambda, \lambda')\varepsilon)^{(2-|\beta|)_+},$$

for all $(r, \theta, \rho_2^\pm, \omega_2^\pm) \in \Gamma_s^\pm(\lambda)$ and $(r', \theta', \rho_2^\pm, \omega_2^\pm) \in \Gamma_s^\pm(\lambda')$, where we use the notation $\partial^\gamma := \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \partial_{r'}^{j'} \partial_{\theta'}^{\alpha'}$ for $\gamma = (j, \alpha, k, \beta, j', \alpha') \in \mathbb{Z}_+^{3d}$.

Proof. We only consider the outgoing case. Remark that (5.14) is equivalent to

$$(\rho_1^+, \omega_1^+) = (\rho, \omega) + \int_0^1 (\partial_{r,\theta} \varphi^+)(r' + \sigma(r - r'), \theta' + \sigma(\theta - \theta'), \rho, \omega) d\sigma.$$

Suppose that $(r, \theta, \rho, \omega) \in \Gamma_s^+(R, \varepsilon)$. Since

$$(\rho_1^+ - \rho, \omega_1^+ - \omega)|_{r'=r, \theta'=\theta} = (\partial_{r,\theta} \varphi^+)(r, \theta, \rho, \omega),$$

(5.1) implies $(|\rho_1^+ - \rho| + r^{-1}|\omega_1^+ - \omega|)|_{r'=r, \theta'=\theta} \leq C\varepsilon^2$, and

$$|p_1 - p| \leq Cp\varepsilon^2, \quad \rho_1^+|_{r'=r, \theta'=\theta} \geq \sqrt{(1 - C\varepsilon^2)2p_1},$$

with some $C > 0$ which is independent of R, ε and λ_0 , where we denote $p = p(r, \theta, \rho, \omega)$ and $p_1 = p(r, \theta, \rho_1^+, \omega_1^+)|_{r'=r, \theta'=\theta}$. Choosing $\lambda > 0$ large enough such that $\lambda^2 > C$, we have $(r, \theta, \rho_1^+, \omega_1^+)|_{r'=r, \theta'=\theta} \in \Gamma^+(\lambda)$. Next, consider the mapping

$$f^+ : (r, \theta, \rho, \omega) \mapsto (r, \theta, \rho_1^+, \omega_1^+)|_{r'=r, \theta'=\theta}.$$

By (5.1), we have

$$(5.18) \quad \left| \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\rho_1^+|_{r'=r, \theta'=\theta} - \rho) \right| \leq C_{j\alpha k\beta} r^{-j-|\beta|} |\omega/r|^{(2-|\beta|)_+}, \\ \left| \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\omega_1^+|_{r'=r, \theta'=\theta} - \omega) \right| \leq C_{j\alpha k\beta} r^{1-j-|\beta|} |\omega/r|^{(2-|\beta|)_+}.$$

By same argument as that in the proof of Lemma 5.3, we obtain that f^+ is injective and $\Gamma_s^+(R, \varepsilon) \subset f^+(\Gamma_s^+(\lambda))$ provided that $\lambda > 0$ is large enough. We note that this λ can be taken uniformly with respect to R and ε . This fact implies (5.16).

We next prove (ii). We write $(r_\sigma, \theta_\sigma) = (r' + \sigma(r - r'), \theta' + \sigma(\theta - \theta'))$ for short. Since $|\omega| \lesssim \min(r, r')\lambda\varepsilon$, (5.1) implies, for $\gamma = (j, \alpha, k, \beta, j', \alpha') \in \mathbb{Z}_+^{3d}$,

$$\begin{aligned} |(\partial^\gamma \varphi^+)(r_\sigma, \theta_\sigma, \rho, \omega)| &\leq C_\gamma r_\sigma^{1-j-|\beta|} |\omega/r_\sigma|^{(2-|\beta|)_+} \\ &\leq C_\gamma \min(r^{1-j-|\beta|}, (r')^{1-j-|\beta|}) (\lambda\varepsilon)^{(2-|\beta|)_+}, \end{aligned}$$

and hence

$$(5.19) \quad |\partial^\gamma(\rho_1^+ - \rho)| + \frac{|\partial^\gamma(\omega_1^+ - \omega)|}{\min(r, r')} \leq C_\gamma R_0^{-j-|\beta|} (\lambda\varepsilon)^{(2-|\beta|)_+} \leq C_\gamma R_0^{-j-|\beta|} (\lambda\varepsilon\varepsilon_0)^{(1-|\beta|)_+}.$$

In particular, $(r, \theta, \rho_1^+, \omega_1^+, r', \theta') \in \tilde{\Gamma}_s^+(\lambda^2)$ if $\lambda > 0$ is large enough. By a same argument as that in the proof of Lemma 5.3, we obtain the first assertion of (ii). Applying (5.16) and (5.15) with $(\rho, \omega) = (\rho_2^+, \omega_2^+)|_{r'=r, \theta'=\theta}$, we obtain

$$\begin{aligned} (r, \theta, \rho_2^+, \omega_2^+)|_{r'=r, \theta'=\theta} &\in \Gamma_s^+(\lambda) \quad \text{if} \quad (r, \theta, \rho, \omega) \in \Gamma_s^+(R, \varepsilon), \\ (r, \theta, \rho, \omega) &\in \Gamma_s^+(\lambda) \quad \text{if} \quad (r, \theta, \rho_2^+, \omega_2^+)|_{r'=r, \theta'=\theta} \in \Gamma_s^+(R, \varepsilon). \end{aligned}$$

Finally we shall prove (iii). Since $(r, \theta, \rho_2^+, \omega_2^+)$ satisfy

$$(5.20) \quad (\rho_2^+, \omega_2^+) = (\rho, \omega) - \int_0^1 (\partial_{r, \theta} \varphi^+)(r' + \sigma(r - r'), \theta' + \sigma(\theta - \theta'), \rho_2^+, \omega_2^+) d\sigma,$$

(5.1) implies

$$(\rho_2^+ - \rho) + \min(r, r')^{-1} |\omega_2^+ - \omega| \leq C \max(\lambda, \lambda')^2 \varepsilon^2$$

for $(r, \theta, \rho_2^+, \omega_2^+) \in \Gamma_s^+(\lambda)$ and $(r', \theta', \rho_2^+, \omega_2^+) \in \Gamma_s^+(\lambda')$. For the derivatives, differentiating (5.20) with respect to $\partial_r^j \partial_\theta^\alpha \partial_{r'}^{j'} \partial_{\theta'}^{\alpha'} \partial_\rho^k \partial_\omega^\beta$ and using (5.1), we obtain (5.17) by an induction with respect to $j + |\alpha| + k + |\beta| + j' + |\alpha'|$. $\square$

Proof of Proposition 5.7. We only prove the outgoing case. Note that since $R \geq \lambda^4 R_0$ and $\varepsilon \leq \lambda^{-4} \varepsilon_0$, (ρ_2^+, ω_2^+) is well-defined for $r, r' > \lambda^{-3} R$. The Schwartz kernel of $I_{\text{IK}}^+(a^+) I_{\text{IK}}^+(b^+)^*$ can be written in the form

$$(2\pi h)^{-d} \int e^{\frac{i}{h}(S^+(r, \theta, \rho, \omega) - S^+(r', \theta', \rho, \omega))} a^+(r, \theta, \rho, \omega) \overline{b^+(r', \theta', \rho, \omega)} d\rho d\omega.$$

Since $(r, \theta, \rho, \omega, r', \theta') \in \tilde{\Gamma}_s^+(\lambda^4)$, Lemma 5.9 implies that (ρ_2^+, ω_2^+) is well-defined and

$$(5.21) \quad 1/2 < |\det \partial_{\rho, \omega}(\rho_2^+, \omega_2^+)| < 3/2.$$

for all $(r, \theta, \rho, \omega, r', \theta')$ with $(r, \theta, \rho_2^+, \omega_2^+, r', \theta') \in \tilde{\Gamma}_s^+(\lambda^4)$. We thus can make the change of variables $(\rho, \omega) \mapsto (\rho_2^+, \omega_2^+)(r, \theta, \rho, \omega, r', \theta')$, and the above integral can be brought to the form

$$(2\pi h)^{-d} \int e^{\frac{i}{h}[(r-r')\rho - (\theta-\theta')\cdot\omega]} A^+(r, \theta, \rho, \omega, r', \theta') d\rho d\omega,$$

where $A^+ = a^+(r, \theta, \rho_2^+, \omega_2^+) \overline{b^+(r', \theta', \rho_2^+, \omega_2^+)} |\det \partial_{\rho, \omega}(\rho_2^+, \omega_2^+)|$. By using (5.17), (5.21) and the support properties of a^+ and b^+ , $\partial_r^j \partial_\theta^\alpha \partial_{r'}^{j'} \partial_{\theta'}^{\alpha'} \partial_\rho^k \partial_\omega^\beta A^+$ are uniformly bounded on $\mathbb{R}^{3d}$ for all $(j, \alpha, k, \beta, j', \alpha') \in \mathbb{Z}_+^{3d}$, and the Calderón-Vaillancourt theorem implies

$$\|I_{\text{IK}}^+(a^+) I_{\text{IK}}^+(b^+)^*\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \leq C_d \sum_{|\gamma| \leq M_d} \|\partial^\gamma A^+\|_{L^\infty(\mathbb{R}^{3d})} \leq C,$$

uniformly with respect to $R > 0$ and $h \in (0, 1]$, with some $C_d, M_d > 0$ depending only on d . In particular $I_{\text{IK}}^+(a^+)_{\mathcal{L}(L^2(\mathbb{R}^d))}$ is bounded on $L^2(\mathbb{R}^d)$ uniformly with respect to $R > 0$ and $h \in (0, 1]$. Furthermore, by the standard symbolic calculus (e.g., see the textbook [15]), the simplified symbol of the above operator has the following asymptotic expansion

$$\sum_{l+|\alpha|=j} \frac{h^j}{l! \alpha!} \left(\partial_\rho^l \partial_\omega^\alpha D_{r'}^l D_{\theta'}^\alpha A^+ \right) (r, \theta, \rho, \omega, r, \theta).$$

By Lemma 5.9 (ii) and (iii), we see that $(\partial_\rho^l \partial_\omega^l D_{r'}^l D_{\theta'}^\alpha A^+)(r, \theta, \rho, \omega, r, \theta)$ is supported in $\Gamma_s^+(\lambda^4)$ and belongs to $S_{\text{sc}}(\Gamma^+(\lambda^4))$. $\square$

Proof of Proposition 5.8. Let $c^+ \in S_{\text{sc}}(\Gamma^+(R, \varepsilon))$. By Proposition 5.7, it suffices to show that there exist $b_j^+ \in S_{\text{sc}}(\Gamma^+(\lambda))$, $j = 0, 1, \dots, N$, such that $c_0^+ = c^+$ and $c_j^+ \equiv 0$ for $j = 1, 2, \dots, N$. We set $J_k = |\det \partial_{\rho, \omega}(\rho_k^+, \omega_k^+)|$. b_j^+ can be defined inductively as follows. We first note that, by the construction, b_h^+ should satisfy

$$\begin{aligned} c^+(r, \theta, \rho, \omega) \\ = \sum_{j=1}^N \sum_{l+|\alpha|=j} \frac{h^j}{l! \alpha!} \left(\partial_\rho^l \partial_\omega^\alpha D_{r'}^l D_{\theta'}^\alpha a_h^+(r, \theta, \rho_2^+, \omega_2^+) \overline{b_h^+(r', \theta', \rho_2^+, \omega_2^+) J_2} \right) \Big|_{r'=r, \theta'=\theta}. \end{aligned}$$

Define b_0^+ by

$$\overline{b_0^+(r, \theta, \rho, \omega)} := (a_0^+(r, \theta, \rho, \omega))^{-1} c^+(r, \theta, \rho_1^+, \omega_1^+) J_1 \Big|_{r'=r, \theta'=\theta}.$$

Since $\text{supp } c^+ \subset \Gamma_s^+(R, \varepsilon)$, Lemma 5.9 (i) implies that $c^+(r, \theta, \rho_1^+, \omega_1^+) \Big|_{r'=r, \theta'=\theta}$ is supported in $\Gamma_s^+(\lambda)$. Since a_0^+ is elliptic on $\Gamma_s^+(\lambda)$, b_0^+ is well-defined and supported in $\Gamma_s^+(\lambda)$. Moreover, (5.18) implies that $b_0^+ \in S_{\text{sc}}(\Gamma^+(\lambda))$. Next, let $j \geq 1$ and assume that $b_k^+ \in S_{\text{sc}}(\Gamma^+(\lambda))$ for all $k < j$. We then define b_j^+ by

$$(5.22) \quad a_0^+(r, \theta, \rho_2^+, \omega_2^+) \overline{b_j^+(r', \theta', \rho_2^+, \omega_2^+) J_2} \Big|_{r'=r, \theta'=\theta} = r_j^+(r, \theta, \rho, \omega),$$

where r_j^+ takes the form

$$\sum_{k=0}^j \sum_{\substack{l+|\alpha|=k, \\ k_0+k_1=j-k, \\ k_1 \leq j-1}} \frac{1}{l! \alpha!} \left(\partial_\rho^l \partial_\omega^\alpha D_{r'}^l D_{\theta'}^\alpha a_{k_0}^+(r, \theta, \rho_2^+, \omega_2^+) \overline{b_{k_1}^+(r', \theta', \rho_2^+, \omega_2^+) J_2} \right) \Big|_{r'=r, \theta'=\theta}.$$

Substituting $(\rho, \omega) = (\rho_1^+, \omega_1^+)$ for (5.22) and dividing by a_0^+ , we have

$$\overline{b_j^+(r, \theta, \rho, \omega)} = (a_0^+(r', \theta', \rho, \omega))^{-1} r_j^+(r, \theta, \rho_1^+, \omega_1^+) J_1 \Big|_{r'=r, \theta'=\theta}.$$

By induction hypothesis, we conclude that $b_j^+ \in S_{\text{sc}}(\Gamma^+(\lambda))$. The proof for the incoming case is similar. $\square$

5.2. Construction of the parametrix. By using the FIO defined in the previous subsection, we construct the semiclassical Isozaki-Kitada parametrix.

Theorem 5.10. *For any $N \geq 0$, there exist $R_{\text{IK}}, \lambda_{\text{IK}} > 0$ large enough and $\varepsilon_{\text{IK}} > 0$ small enough such that if $R \geq R_{\text{IK}}$, $0 < \varepsilon \leq \varepsilon_{\text{IK}}$ and $\lambda \geq \lambda_{\text{IK}}$, then we can find*

$$b_h^\pm = \sum_{j=0}^N h^j b_j^\pm \quad \text{with} \quad b_j^\pm \in S_{\text{sc}}(\Gamma^\pm(\lambda^3)), \quad j = 0, 1, \dots, N,$$

such that, for any $a_s^\pm \in S_{\text{sc}}(\Gamma^\pm(R, \varepsilon))$, there exist

$$c_h^\pm = \sum_{j=0}^N h^j c_j^\pm \quad \text{with} \quad c_j^\pm \in S_{\text{sc}}(\Gamma^\pm(\lambda)), \quad j = 0, 1, \dots, N,$$

such that, for all $T > 0$, $h \in (0, 1]$ and $0 \leq \pm t \leq Th^{-1}$,

$$(5.23) \quad \|e^{-ith\hat{P}} \text{Op}_\kappa(a_s^\pm) - \kappa^* I_{\text{IK}}^\pm(b_h^\pm) e^{-ith\frac{1}{2}D_r^2} I_{\text{IK}}^\pm(c_h^\pm)^* \kappa_*\|_{\mathcal{L}(L^2(\widehat{M}))} \leq C_{N,T} h^{N-1},$$

where $C_{N,T}$ may be taken uniformly with respect to h , t and R .

Remark 5.11. Since $\pi(\text{supp } b_h^\pm), \pi(\text{supp } c_h^\pm) \subset (2, \infty) \times \tilde{U}_\kappa$, we see that

$$I_{\text{IK}}^\pm(b_h^\pm) e^{-ith\frac{1}{2}D_r^2} I_{\text{IK}}^\pm(c_h^\pm)^* = \tilde{\Psi}_\kappa I_{\text{IK}}^\pm(b_h^\pm) e^{-ith\frac{1}{2}D_r^2} I_{\text{IK}}^\pm(c_h^\pm)^* \tilde{\Psi}_\kappa,$$

and $\kappa^* I_{\text{IK}}^\pm(b_h^\pm) e^{-ith\frac{1}{2}D_r^2} I_{\text{IK}}^\pm(c_h^\pm)^* \kappa_*$ is well-defined on M .

We shall prove Theorem 5.10 for the outgoing case, and the proof for the incoming case is completely analogous. Set $B_+ := I_{\text{IK}}^+(b_h^+)$, $C_+ := I_{\text{IK}}^+(c_h^+)$. By the Duhamel formula, we have

$$\begin{aligned} & e^{-ith\hat{P}} \kappa^* B_+ C_+^* \kappa_* \\ &= \kappa^* B_+ e^{-ith\frac{1}{2}D_r^2} C_+^* \kappa_* - \frac{i}{h} \int_0^t e^{-i(t-s)h\hat{P}} \kappa^* (h^2 \hat{P}_\kappa B_+ - B_+ h^2 \frac{1}{2} D_r^2) e^{-ish\frac{1}{2}D_r^2} C_+^* ds. \end{aligned}$$

To prove (5.23), it suffices to show that

$$(5.24) \quad \|(h^2 \hat{P}_\kappa B_+ - B_+ h^2 \frac{1}{2} D_r^2) e^{-ish\frac{1}{2}D_r^2} C_+^*\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \lesssim h^{N+1},$$

$$(5.25) \quad \|a_s^+(r, \theta, hD_r, hD_\theta) - B_+ C_+^*\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \lesssim h^N,$$

uniformly with respect to $h \in (0, 1]$, $0 \leq s \leq Th^{-1}$ and $R > 0$. To prove above two estimates, we prepare several lemmas.

Let $p + p_1 + p_2$ be the full symbol of $\hat{P}_\kappa$:

$$\hat{P}_\kappa = p(r, \theta, D_r, D_\theta) + p_1(r, \theta, D_r, D_\theta) + p_2(r, \theta).$$

Choosing $R_0 > 0$ and $\varepsilon_0 > 0$ so that S^+ is well-defined and solves the Hamilton-Jacobi equation on $\Gamma_s^+(R_0, \varepsilon_0)$, we define smooth tensors X^+ and Y^+ by

$$X^+ := \partial_{\rho, \omega} p(r, \theta, \partial_r S^+, \partial_\theta S^+), \quad Y^+ := (p + p_1)(r, \theta, \partial_r, \partial_\theta) S^+,$$

and define symbols d_j^+ , $j = 1, 2, \dots, N+2$, by

$$h^2 \hat{P}_\kappa B_+ - B_+ h^2 \frac{1}{2} D_r^2 = \sum_{j=1}^{N+2} h^j I_{\text{IK}}^+(d_j^+).$$

Then d_j^+ should satisfy

$$(5.26) \quad \begin{cases} id_1^+ = X^+ \cdot \partial_{r, \theta} b_0^+ + Y^+ b_0^+, \\ id_j^+ = X^+ \cdot \partial_{r, \theta} b_{j-1}^+ + Y^+ b_{j-1}^+ + i\hat{P}_\kappa b_{j-2}^+, \quad j = 2, \dots, N+2, \end{cases}$$

where $b_{N+1}^+ \equiv 0$. To construct b_j^+ , we solve transport equations.

Lemma 5.12. *There exist $R_1 \geq R_0$, $\lambda_1 > 1$ large enough and $\varepsilon_1 \leq \varepsilon_0$ small enough such that, for all $N \geq 0$, $R \geq \lambda_1^{N+5} R_1$ and $0 < \varepsilon \leq \lambda_1^{-N-5} \varepsilon_1$, we can find $b_j^+ \in S_{\text{sc}}(\Gamma_s^+(\lambda_1^3))$, $j = 0, 1, \dots, N$ such that b_0^+ is elliptic on $\Gamma_s^+(\lambda_1)$, and that b_j^+ solve transport equations on $\Gamma_s^+(\lambda_1^2)$:*

$$(5.27) \quad \begin{cases} X^+ \cdot \partial_{r, \theta} b_0^+ + Y^+ b_0^+ = 0, \\ X^+ \cdot \partial_{r, \theta} b_j^+ + Y^+ b_j^+ + i\hat{P}_\kappa b_{j-1}^+ = 0, \quad j = 1, \dots, N. \end{cases}$$

For $(r, \theta, \rho, \omega) \in \Gamma_s^+(\lambda_1^{N+4})$, we consider the flow $(r^+(t), \theta^+(t))$ generated by X^+ , i.e.,

$$(r^+(t), \theta^+(t)) = (r^+, \theta^+)(t, r, \theta, \rho, \omega)$$

is the solution to

$$\begin{cases} (\dot{r}^+(t), \dot{\theta}^+(t)) = X^+(r^+(t), \theta^+(t), \rho, \omega), \\ (r^+(0), \theta^+(0)) = (r, \theta). \end{cases}$$

Then $(r^+(t), \theta^+(t))$ is defined on $[0, \infty) \times \Gamma_s^+(\lambda_1^{N+4})$, and satisfies the following:

Lemma 5.13. *For all $t \geq 0$, $(r, \theta, \rho, \omega) \in \Gamma_s^+(\lambda_1^{N+4})$ and $(j, \alpha, k, \beta) \in \mathbb{Z}_+^{2d}$,*

$$(5.28) \quad \begin{aligned} & (r^+(t), \theta^+(t), \rho, \omega) \in \Gamma_s^+(\lambda_1^{N+5}), \\ & |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (r^+(t) - r - t\rho)| \leq C_{j\alpha k\beta} r^{1-j-|\beta|} |\omega/r|^{(2-|\beta|)_+}, \end{aligned}$$

$$(5.29) \quad |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\theta^+(t) - \theta)| \leq C_{j\alpha k\beta} r^{-j-|\beta|} |\omega/r|^{(1-|\beta|)_+},$$

Proof. Let $(r, \theta, \rho, \omega) \in \Gamma_s^+(\lambda_1^{N+4})$. Since $X^+ = (\partial_r S^+, r^{-2}(h^{jk} + a^{jk})\partial_{\theta^k} S^+)$, it follows from (5.1) that

$$|\dot{r}^+(t) - \rho| \leq C_0 |\omega/r^+(t)|^2, \quad |\dot{\theta}^+(t)| \leq C_0 r^+(t)^{-1} |\omega/r^+(t)|$$

with some $C_0 > 0$. In particular, we have

$$(5.30) \quad |\dot{r}^+(0) - \rho| \leq C_0 |\omega/r|^2 \leq C_1 \varepsilon_1, \quad |\dot{\theta}^+(0)| \leq C_0 r^{-1} |\omega/r| \leq C_1 r^{-1} \varepsilon_1$$

with some $C_1 > 0$. We set $C_2 = \inf J$ and

$$F := \{t \geq 0 \mid r^+(s) \geq r + s\rho/2, |\theta^+(s) - \theta| \leq 4C_0 C_2^{-1} |\omega/r| \text{ for } 0 \leq s \leq t\}.$$

By (5.30), it is easy to see that $0 \in F \neq \emptyset$. Let $t_0 = \sup F$. We then have

$$|\dot{r}^+(t) - \rho| \leq C_0 (r + s\rho/2)^{-2} |\omega|^2 \leq C_0 C_1 \varepsilon_1^2, \quad |\dot{\theta}^+(t)| \leq C_0 (r + s\rho/2)^{-2} |\omega|,$$

for $0 \leq t \leq t_0$, and hence

$$|r^+(t) - r - t\rho| \leq C_0 C_1 \varepsilon_1^2 t, \quad |\theta^+(t) - \theta| \leq 2C_0 C_2^{-1} |\omega/r|,$$

for $0 \leq t \leq t_0$. Choosing $\varepsilon_1 > 0$ such that $\rho - C_0 C_1 \varepsilon_1^2 > \rho/2$ and $\delta > 0$ small enough, we see that $t_0 + \delta \in F$ which implies $t_0 = \infty$. Therefore, $(r^+(t), \theta^+(t))$ is well-defined on $[0, \infty)$, and satisfies (5.28) and (5.29) with $(j, \alpha, k, \beta) = 0$ for all $t \geq 0$. In particular, $(r^+(t), \theta^+(t), \rho, \omega) \in \Gamma_s^+(\lambda_1^{N+5})$ for $t \geq 0$, provided that $\lambda_1 > 0$ large enough. The proof for higher derivatives is obtained by (5.1) and an induction with respect to $j + |\alpha| + k + |\beta|$. $\square$

Proof of Lemma 5.12. Define a smooth real-valued function Z^+ by

$$Z^+(t) := Y^+(r^+(t), \theta^+(t), \rho, \omega) \in C^\infty([0, \infty) \times \Gamma_s^+(\lambda_1^{N+4})).$$

By (2.3), (5.1) and Lemma (5.13), we have $Z^+(t) \in L_t^1([0, \infty))$ and

$$(5.31) \quad \left| \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \int_0^\infty Z^+(t, r, \theta, \rho, \omega) dt \right| \leq C_{j\alpha k\beta} r^{-j-|\beta|} \quad \text{on } \Gamma_s^+(\lambda_1^{N+4}).$$

We now define smooth functions $\tilde{b}_j$ on $\Gamma_s^+(\lambda_1^{N+4-j})$ by

$$\begin{cases} \tilde{b}_0(r, \theta, \rho, \omega) = e^{\int_0^\infty Z^+(t) dt}, \\ \tilde{b}_j(r, \theta, \rho, \omega) = \int_0^\infty (i\hat{P}_\kappa \tilde{b}_{j-1})(r^+(t), \theta^+(t), \rho, \omega) e^{\int_0^t Z^+(s) ds} dt, \quad j = 1, 2, \dots, N. \end{cases}$$

By a standard Hamilton-Jacobi theory, $\tilde{b}_j$ solve (5.27). Moreover, by (2.3), Lemma 5.13 and (5.31), we have

$$|\partial_r^l \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \tilde{b}_j(r, \theta, \rho, \omega)| \leq C_{l\alpha k\beta} r^{-l-|\beta|} \quad \text{on } \Gamma_s^+(\lambda_1^{N+4-j}).$$

Take $\chi^+ \in C_0^\infty(\mathbb{R}^{2d})$ so that $0 \leq \chi^+ \leq 1$, $\chi^+ \equiv 1$ on $\Gamma_s^+(\lambda_1^2)$, $\text{supp } \chi^+ \subset \Gamma_s^+(\lambda_1^3)$, and that

$$|\partial_r^l \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \chi^+(r, \theta, \rho, \omega)| \leq C_{l\alpha k\beta} r^{-l-|\beta|} \quad \text{on } \Gamma_s^+(\lambda_1^3),$$

and define $b_j^+ := \chi^+ \tilde{b}_j$. By the construction, b_j^+ solve (5.27) on $\Gamma_s^+(\lambda_1^2)$, b_0^+ is elliptic on $\Gamma_s^+(\lambda_1)$ and $b_j^+ \in S_{\text{sc}}(\Gamma_s^+(\lambda_1^3))$. $\square$

Proof of Theorem 5.10. Let $R \geq \lambda_1^{N+5} R_1$ and $0 < \varepsilon \leq \lambda_1^{-N-5} \varepsilon_1$. We first prove (5.25). By Proposition 5.8, there exists a symbol $c_h^+ = \sum_{j=0}^N h^j c_j^+ \in S_{\text{sc}}(\Gamma_s^+(\lambda_1))$ such that

$$\|a_s^+(r, \theta, hD_r, hD_\theta) - B_+ C_+^*\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \leq C_N h^{N+1}, \quad h \in (0, 1],$$

where C_N may be taken uniformly with respect to h and R .

We next prove (5.24). Let d_j^+ , $j = 1, 2, \dots, N+2$, be defined by (5.26). Then $d_j^+ \in S_{\text{sc}}(\Gamma_s^+(\lambda_1^3))$ and $d_j^+ \equiv 0$ on $\Gamma_s^+(\lambda_1^2)$. For each j and k , $I_{\text{IK}}^+(d_j^+) e^{-ish \frac{1}{2} D_r^2} I_{\text{IK}}^+(c_k^+)^*$ has the distribution kernel

$$I(s, h) = (2\pi h)^{-d} \int e^{\frac{i}{h}((r-r')\rho_1^+ + (\theta-\theta') \cdot \omega_1^+ - \frac{1}{2} s \rho^2)} d_j^+(r, \theta, \rho, \omega) \overline{c_k^+(r', \theta', \rho, \omega)} dp d\omega,$$

where $(\rho_1^+, \omega_1^+)(r, \theta, \rho, \omega, r', \theta')$ is defined in Lemma 5.9. Since $d_j^+ \equiv 0$ on $\Gamma_s^+(\lambda_1^2)$, one of the followings hold at least:

$$(5.32) \quad \lambda_1^{-3}R < r \leq \lambda_1^{-2}R,$$

$$(5.33) \quad \theta \in \tilde{U}_{\kappa, \sqrt{\lambda_1^3 \varepsilon}} \setminus \tilde{U}_{\kappa, \sqrt{\lambda_1^2 \varepsilon}},$$

$$(5.34) \quad p(r, \theta, \rho, \omega) \in J_{\lambda_1^6 \varepsilon^2} \setminus J_{\lambda_1^2 \varepsilon^2},$$

$$(5.35) \quad \lambda_1^2 \varepsilon \lesssim |\omega/r| \lesssim \lambda_1^3 \varepsilon.$$

We first assume (5.32) or (5.35). Then, it is easy to see that

$$r - r' < -R/(2\lambda_1)$$

if $R > 0$ is large enough. Therefore, by (5.19), we have

$$(r - r')\partial_\rho \rho_1^+ + (\theta - \theta')\partial_\rho \omega_1^+ - s\rho \leq -R/(4\lambda_1) - s\rho \leq -R_0/2 - s\rho \leq -\langle s \rangle,$$

and we obtain (5.24) by a standard integration by parts. We next assume (5.34). Since $p(r, \theta, \rho, \omega) - \rho^2/2 = O(|\omega/r|^2) = O(\varepsilon_0^2)$, for sufficiently small $\varepsilon_0 > 0$, the amplitude vanishes identically in this case, and we have (5.24). Finally, we assume $\theta \in \tilde{U}_{\kappa, \sqrt{\lambda_1^3 \varepsilon}} \setminus \tilde{U}_{\kappa, \sqrt{\lambda_1^2 \varepsilon}}$. Since $\theta' \in \tilde{U}_{\kappa, \sqrt{\lambda_1 \varepsilon}}$ and

$$|\partial_\omega \rho_1^+| \lesssim r^{-1}|\omega/r| \lesssim r^{-1}\lambda_1^3 \varepsilon \lesssim r^{-1}\sqrt{\lambda_1 \varepsilon_1}, \quad |\partial_\omega(\omega_1^+ - \omega)| \lesssim |\omega/r| \lesssim \lambda_1^3 \varepsilon \lesssim \varepsilon_1,$$

we have

$$(5.36) \quad |\theta - \theta'| \gtrsim \lambda_1 \sqrt{\varepsilon}.$$

Therefore,

$$\begin{aligned} & |\partial_\omega ((r - r')\rho_1^+ + (\theta - \theta') \cdot \omega_1^+ - s\rho^2/2)| \\ & \geq |\theta - \theta'| (1 - C\varepsilon_1) - C|r - r'|r^{-1}\sqrt{\lambda_1 \varepsilon_1} \\ & \gtrsim (\lambda_1 - C\varepsilon_1 - \sqrt{\lambda_1 \varepsilon_1})\sqrt{\varepsilon} \\ & > \sqrt{\varepsilon}, \end{aligned}$$

for sufficiently large $\lambda_1 > 0$ and small $\varepsilon_1 > 0$. We now fix the constants $R_{\text{IK}}, \varepsilon_{\text{IK}}, \lambda_{\text{IK}}$ so that $\lambda_{\text{IK}} = \lambda_1$, $R_{\text{IK}} = \lambda_{\text{IK}}^{N+5}R_1$, $\varepsilon_{\text{IK}} = \lambda_{\text{IK}}^{-N-5}\varepsilon_1$. Put

$$L(r, \theta, \rho, \omega, r', \theta') = \frac{h((r - r')\partial_\omega \rho_1^+ + (\theta - \theta') \cdot \partial_\omega \omega_1^+)}{i|(r - r')\partial_\omega \rho_1^+ + (\theta - \theta') \cdot \partial_\omega \omega_1^+|^2} \cdot \partial_\omega,$$

and integrate by parts $I(s, h)$ with respect to L . For any $n \geq 0$, $I(s, h)$ then reads

$$I(s, h) = h^n (2\pi h)^{-d} \int e^{\frac{i}{h}((r-r')\rho_1^+ + (\theta-\theta') \cdot \omega_1^+ - \frac{1}{2}s\rho^2)} G^+(r, \theta, \rho, \omega, r', \theta') d\rho d\omega,$$

where $G^+ = (L^*)^n(d_j^+(r, \theta, \rho, \omega)\overline{c_k^+(r', \theta', \rho, \omega)})$. Using the change of the variables $(\rho, \omega) \mapsto (\rho_2^+, \omega_2^+)$, we have

$$I(s, h) = h^n (2\pi h)^{-d} \int e^{\frac{i}{h}\Phi^+(s, r, \theta, \rho, \omega, r', \theta')} G^+(r, \theta, \rho_2^+, \omega_2^+, r', \theta') d\rho d\omega,$$

where $\Phi^+(s, r, \theta, \rho, \omega, r', \theta') := (r - r')\rho + (\theta - \theta') \cdot \omega - \frac{1}{2}s\rho_2^+(r, \theta, \rho, \omega, r', \theta')^2$, and (ρ_2^+, ω_2^+) is the inverse of (ρ_1^+, ω_1^+) . By Lemma 5.9 and (5.36), $\rho_2^+(r, \theta, \rho, \omega, r', \theta')$ and $G^+(r, \theta, \rho_2^+, \omega_2^+, r', \theta')$ are smooth and uniformly bounded functions on $\mathbb{R}^{3d}$. Applying the Calderón-Vaillancourt theorem, we hence have

$$\|I_{\text{IK}}^+(d_j^+)e^{-ish\frac{1}{2}D_r^2}I_{\text{IK}}^+(c_k^+)^*\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \leq C_{N,d,\varepsilon}h^{n-n_d}\langle s \rangle^{n_d} \leq C_{N,d,\varepsilon,T}h^{n-2n_d}$$

for all $n \geq 0$, $h \in (0, 1]$ and $0 \leq s \leq Th^{-1}$, where $n_d > 0$ depends only on d , and $C_{N,d,\varepsilon,T}$ is independent of h and R . Choosing $n > 0$ with $n - 2n_d > N + 1$, we complete the proof $\square$

5.3. Dispersive estimates. We here prove dispersive estimates for the Isozaki-Kitada parametrix.

Let $R_{\text{IK}}, \varepsilon_{\text{IK}}, \lambda_{\text{IK}} > 0$ be as in Theorem 5.10. In this subsection we use the notation $\partial^\gamma := \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \partial_{r'}^{j'} \partial_{\theta'}^{\alpha'}$ for $\gamma = (j, \alpha, k, \beta, j', \alpha') \in \mathbb{Z}_+^{3d}$.

Theorem 5.14. *For sufficiently large $R > R_{\text{IK}}$, small $0 < \varepsilon < \varepsilon_{\text{IK}}$, all $b^\pm \in S_{\text{sc}}(\Gamma^\pm(\lambda_{\text{IK}}^3))$ and $c^\pm \in S_{\text{sc}}(\Gamma^\pm(\lambda_{\text{IK}}))$, we can write*

$$I_{\text{IK}}^\pm(b^\pm) e^{-ith \frac{1}{2} D_r^2} I_{\text{IK}}^\pm(c^\pm)^* = U_{\text{IK}}^\pm(t, h) + R_{\text{IK}}^\pm(t, h),$$

where $U_{\text{IK}}^\pm(t, h)$ satisfy dispersive estimates

$$\|r^{-\frac{d-1}{2}} U_{\text{IK}}^\pm(t, h) r^{-\frac{d-1}{2}}\|_{\mathcal{L}(L^1(\mathbb{R}^d), L^\infty(\mathbb{R}^d))} \leq C |th|^{-d/2}, \quad 0 < \pm t \leq h^{-1}, \quad h \in (0, 1],$$

and $R_{\text{IK}}^\pm(t, h)$ are rapidly decaying with respect to h : for any $N \geq 0$,

$$\|R_{\text{IK}}^\pm(t, h)\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \leq C_N h^N, \quad 0 < \pm t \leq h^{-1}, \quad h \in (0, 1].$$

Moreover $C, C_N > 0$ can taken uniformly with respect to h, t and R .

We prove the theorem for the case $t \geq 0$, and the proof for the case $t \leq 0$ is similar. The distribution kernel of $I_{\text{IK}}^+(b^+) e^{-ith \frac{1}{2} D_r^2} I_{\text{IK}}^+(c^+)^*$ takes the form

$$(5.37) \quad I_{A^+}(t, h) = (2\pi h)^{-d} \int e^{\frac{i}{h} \Phi^+(t, r, \theta, \rho, \omega, r', \theta')} A^+(r, \theta, \rho, \omega, r', \theta') d\rho d\omega,$$

where

$$\Phi^+(t, r, \theta, \rho, \omega, r', \theta') := (r - r')\rho + (\theta - \theta') \cdot \omega - \frac{1}{2} t (\rho_2^+)^2,$$

$$A^+(r, \theta, \rho, \omega, r', \theta') := b^+(r, \theta, \rho_2^+, \omega_2^+) \overline{b^+(r', \theta', \rho_2^+, \omega_2^+)} |\det \partial_{\rho, \omega}(\rho_2^+, \omega_2^+)|,$$

and $(\rho_2^+, \omega_2^+) = (\rho_2^+, \omega_2^+)(r, \theta, \rho, \omega, r', \theta')$ is given by Lemma 5.9. Note that since $(r, \theta, \rho_2^+, \omega_2^+) \in \Gamma_s^+(\lambda_{\text{IK}}^3)$ and $(r', \theta', \rho_2^+, \omega_2^+) \in \Gamma_s^+(\lambda_{\text{IK}})$, (5.17) implies

$$C^{-1} \leq \rho \leq C, \quad |\omega / \sqrt{rr'}| \leq C\varepsilon_1 \quad \text{on} \quad \text{supp } A^+,$$

and $\partial^\gamma A^+$ and $\partial^\gamma \rho_2^+$ are uniformly bounded on $\mathbb{R}^{3d}$.

We first remove a smoothing term from $I_{A^+}(t, h)$. Let $\chi_\rho \in C_0^\infty(\mathbb{R})$, $\chi_\omega \in C_0^\infty(\mathbb{R}^{d-1})$ be smooth cut-off functions such that

$$\text{supp } \chi_\rho \subset (-1, 1), \quad \chi_\rho \equiv 1 \text{ on } (-1/2, 1/2),$$

$$\text{supp } \chi_\omega \subset \{|\theta| \leq 1\}, \quad \chi_\omega \equiv 1 \text{ on } \{|\theta| \leq 1/2\},$$

and define $A_\delta^+ := \chi_\rho (\partial_\rho \Phi^+) \chi_\omega (\partial_\omega \Phi^+ / \delta) A^+$, where $\delta > 0$ is a small parameter. We denote the operator having the Schwartz kernel $I_{A_\delta^+}(t, h)$ by $U_{A_\delta^+}(t, h)$.

Lemma 5.15. *For all $N > 0$ and $\delta > 0$, we have*

$$\|I_{\text{IK}}^+(b^+) e^{-ith \frac{1}{2} D_r^2} I_{\text{IK}}^+(c^+)^* - U_{A_\delta^+}(t, h)\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \leq C_N h^N,$$

uniformly with respect to $h \in (0, 1]$, $0 \leq \pm t \leq h^{-1}$ and $R > 0$.

Proof. We split $A^+ - A_\delta^+ = (1 - \chi_\rho) \chi_\omega A^+ - (1 - \chi_\omega) A^+ =: A_1^+ + A_2^+$, and denote by $I_{A_j^+}$ the oscillatory integral of the form (5.37) with the phase Φ^+ and the amplitude A_j^+ , respectively. We set $L_1 := (h/i \partial_\rho \Phi^+) \partial_\rho$, $L_2 := (h/i |\partial_\omega \Phi^+|^2) (\partial_\omega \Phi^+) \cdot \partial_\omega$. Since

$$|\partial^\gamma (1/\partial_\rho \Phi^+)| \lesssim 1 \quad \text{on} \quad \text{supp } A_1^+, \quad |\partial^\gamma (\partial_\omega \Phi^+ / |\partial_\omega \Phi^+|^2)| \lesssim 1/\delta \quad \text{on} \quad \text{supp } A_2^+,$$

n -times integration by parts $I_{A_j^+}$ with L_j , respectively, implies that $I_{A_j^+}$ reads

$$I_{A_j^+} = (2\pi h)^{-d} \int e^{\frac{i}{h} ((r-r')\rho + (\theta-\theta') \cdot \omega)} \tilde{A}_j^+(r, \theta, \rho, \omega, r', \theta') d\rho d\omega,$$

where $\tilde{A}_j^+ = e^{-\frac{i}{h} \frac{t}{2} (\rho_2^+)^2} (L_j^*)^n A_j$ satisfies

$$|\partial^\gamma \tilde{A}_j^+(r, \theta, \rho, \omega, r', \theta')| \leq C_n h^{n-|\gamma|} \langle t \rangle^{|\gamma|} \leq C_n h^{n-2|\gamma|} \quad \text{on } \mathbb{R}^{3d},$$

for all $\gamma \in \mathbb{Z}_+^{3d}$, $h \in (0, 1]$, and $0 \leq \pm t \leq h^{-1}$. By the Calderón-Vaillancourt theorem, there exists an integer $N_d > 0$ depending only on d such that

$$\|I_{\text{IK}}^+(b^+) e^{-ith \frac{1}{2} D_r^2} I_{\text{IK}}^+(c^+)^* - U_{A_\delta^+}(t, h)\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \leq C_n h^{n-2N_d}.$$

Choosing $n \geq 0$ with $n - 2N_d \geq N$, we obtain the assertion. $\square$

To prove dispersive estimates for $U_{A_\delta^+}(t, h)$, we next study the phase function more precisely.

Lemma 5.16. *On $\text{supp } A^+$, $\frac{1}{2} \rho_2^+(r, \theta, \rho, \omega, r', \theta')^2$ takes the form*

$$\frac{1}{2} \rho_2^+(r, \theta, \rho, \omega, r', \theta')^2 = \frac{1}{2} \rho^2 + \frac{1}{2rr'} q_0(\theta, \omega) + Q^+(r, \theta, \rho, \omega, r', \theta'),$$

where $q_0(\theta, \omega) = h^{jk}(\theta) \omega_j \omega_k$. Moreover we can write

$$Q^+ = Q_1^+ + (\theta - \theta') \cdot Q_2^+$$

such that, for all $(r, \theta, \rho, \omega, r', \theta') \in \text{supp } A^+$ and $\gamma = (j, \alpha, k, \beta, j', \alpha') \in \mathbb{Z}_+^{3d}$,

$$(5.38) \quad \begin{cases} |\partial^\gamma Q_1^+(r, \theta, \rho, \omega, r', \theta')| \leq C R_1^{-j-j'-|\beta|} \left(\varepsilon_1^{(3-|\beta|)_+} + R_1^{-\mu} \varepsilon_1^{(2-|\beta|)_+} \right), \\ |\partial^\gamma Q_2^+(r, \theta, \rho, \omega, r', \theta')| \leq C R_1^{-j-j'-|\beta|} \varepsilon_1^{(2-|\beta|)_+}, \end{cases}$$

where R_1, ε_1 are given by Lemma 5.12.

Proof. We start from the formula $\rho_2^+ = \rho - \int_0^1 (\partial_r \varphi^+)(r_s, \theta_s, \rho_2^+, \omega_2^+) ds$, where $(r_s, \theta_s) = (r', \theta') + s(r - r', \theta - \theta')$. By using the mean value theorem, we have

$$(\partial_r \varphi^+)(r_s, \theta_s, \rho_2^+, \omega_2^+) = (\partial_r \varphi^+)(r_s, \theta, \rho_2^+, \omega_2^+) - (\theta - \theta') \cdot F^+(s),$$

where $F^+(s)$ is defined by

$$F^+(s) := (1-s) \int_0^1 (\partial_\theta \partial_r \varphi^+)(r_s, \theta_{s\sigma}, \rho_2^+, \omega_2^+) d\sigma, \quad \theta_{s\sigma} = \theta + \sigma(\theta_s - \theta).$$

By (5.1) and (5.17), we obtain

$$\sup_{s \in [0,1]} |\partial^\gamma F^+(s)| \lesssim R_1^{-j-j'-|\beta|} \varepsilon_1^{(2-|\beta|)_+}.$$

Since $R_{\text{IK}} \geq \lambda_{\text{IK}}^4 R_1$ and $\varepsilon_{\text{IK}} \leq \lambda_{\text{IK}}^{-4} \varepsilon_1$, by the mean value theorem, we can write

$$(\partial_r \varphi^+)(r_s, \theta, \rho_2^+, \omega_2^+) = -\frac{1}{2r_s^2 \rho} q_0(\theta, \omega) - G^+(s),$$

where, by (5.2) and (5.17), $G^+(s)$ satisfies

$$\sup_{s \in [0,1]} |\partial^\gamma G^+(s)| \lesssim R_1^{-j-j'-|\beta|} \left(\varepsilon_1^{(3-|\beta|)_+} + R_1^{-\mu} \varepsilon_1^{(2-|\beta|)_+} \right).$$

Set $\tilde{F}^+ = \int_0^1 F^+(s) ds$ and $\tilde{G}^+ = \int_0^1 G^+(s) ds$. Since $\int_0^1 r_s^{-2} ds = 1/\sqrt{rr'}$, we have

$$(\rho_2^+)^2 = \left(\rho + \frac{1}{2rr'\rho} q_0(\theta, \omega) + (\theta - \theta') \cdot \tilde{F}^+ + \tilde{G}^+ \right)^2.$$

If we set

$$Q_1^+ = 4\rho \tilde{G}^+ + 2 \left(\frac{1}{2rr'\rho} q_0(\theta, \omega) + (\theta - \theta') \cdot \tilde{F}^+ + \tilde{G}^+ \right)^2, \quad Q_2^+ = 4\rho \tilde{F}^+,$$

then $Q^+ = Q_1^+ + Q_2^+$ satisfy (5.38). $\square$

Using the change of the variable $\omega \mapsto \sqrt{rr'}\nu$, $I_{A_\delta^+}(t, h)$ reads

$$I_{A_\delta^+}(t, h) = \frac{(rr')^{\frac{d-1}{2}}}{(2\pi h)^d} \int e^{\frac{i}{h}\Phi^+(t, r, \theta, \rho, \sqrt{rr'}\nu, r', \theta')} A_\delta^+(r, \theta, \rho, \sqrt{rr'}\nu, r', \theta') d\rho d\nu,$$

where $\tilde{A}_\delta^+(r, \theta, \rho, \nu, r', \theta') := A_\delta^+(r, \theta, \rho, \sqrt{rr'}\nu, r', \theta')$ is bounded on $\mathbb{R}^{3d}$ and compactly supported with respect to (ρ, ν) . Moreover, $\pi_{\rho, \nu} \circ \text{supp } \tilde{A}_\delta^+$ is bounded uniformly with respect to $R > 0$, where $\pi_{\rho, \nu} : \mathbb{R}^{3d} \rightarrow \mathbb{R}^d$ is a canonical projection onto the (ρ, ν) -space.

Proposition 5.17. *There exists $\delta > 0$ small enough such that, for all $h \in (0, 1]$ and $0 < t \leq h^{-1}$, we have*

$$\|r^{-\frac{d-1}{2}} U_{A_\delta^+}(t, h) r^{-\frac{d-1}{2}}\|_{\mathcal{L}(L^1(\mathbb{R}^d), L^\infty(\mathbb{R}^d))} \leq C|th|^{-d/2},$$

where C may be taken uniformly with respect to R, h and t .

Proof. For $0 < t \leq h$, it follows from

$$r^{-\frac{d-1}{2}} |I_{A_\delta^+}(t, h)| r^{-\frac{d-1}{2}} \leq Ch^{-d} \leq C|th|^{-d/2}.$$

Suppose that $h \leq t \leq h^{-1}$ and assume $2h \leq h^{-1}$ without loss of generality. Define

$$\begin{aligned} \tilde{\Phi}^+(t, r, \theta, \rho, \nu, r', \theta') &:= \frac{1}{t} \Phi^+(t, r, \theta, \rho, \sqrt{rr'}\nu, r', \theta') \\ &= \frac{r-r'}{t} \rho + \frac{\theta-\theta'}{t} \cdot \sqrt{rr'}\nu - \frac{1}{2} \rho_2^+(r, \theta, \rho, \sqrt{rr'}\nu, r', \theta')^2. \end{aligned}$$

By Lemma 5.16, we obtain

$$\partial_\rho \tilde{\Phi}^+(t) = \frac{r-r'}{t} - \rho + O(\varepsilon_1^2), \quad \partial_\nu \tilde{\Phi}^+(t) = \frac{\sqrt{rr'}(\theta-\theta')}{t} + O(\varepsilon_1)$$

on the support of $\tilde{A}_\delta^+$. Thus, if we restrict the support of $\tilde{A}_\delta^+$ to one of the regions

$$\left| \frac{r-r'}{t} - \rho \right| \geq \varepsilon_1 \quad \text{or} \quad \left| \frac{\sqrt{rr'}(\theta-\theta')}{t} \right| \geq 1,$$

then we have $|\partial_\rho \tilde{\Phi}(t)| + |\partial_\nu \tilde{\Phi}(t)| \geq \varepsilon_1$ with sufficiently small $\varepsilon_1 > 0$. Take $\chi_1 \in C_0^\infty(\mathbb{R})$ and $\chi_2 \in C_0^\infty(\mathbb{R}^{d-1})$ satisfying $\chi_1 \equiv 1$ on $(-1/2, 1/2)$, $\text{supp } \chi_1 \subset (-1, 1)$, $\chi_2 \equiv 1$ on $\{|\theta| \leq 1/2\}$ and $\text{supp } \chi_2 \subset \{|\theta| \leq 1\}$, and put

$$\tilde{\chi}_1(t, r, \theta, \rho, r', \theta') = \chi_1 \left(\frac{r-r'}{\varepsilon_1 t} - \frac{\rho}{\varepsilon_1} \right) \chi_2 \left(\frac{\sqrt{rr'}(\theta-\theta')}{t} \right).$$

Integrating by parts, we then obtain the non stationary estimates

$$\left| (2\pi h)^{-d} \int e^{\frac{i}{h}\tilde{\Phi}^+(t)} (1 - \tilde{\chi}_1) \tilde{A}_\delta^+ d\rho d\nu \right| \leq C_{\varepsilon_1, \lambda_{\text{IK}}} h^{-d} |t/h|^{-n} \leq C_{\varepsilon_1, \lambda_{\text{IK}}} |th|^{-d/2},$$

for all $h \in (0, 1]$, $h \leq t \leq 2h^{-1}$, $\delta > 0$ and $n \geq d/2$, where $C_{\varepsilon_1, \lambda_{\text{IK}}}$ may be taken uniformly in R . Since $C^{-1} \leq \rho \leq C$ for some $C > 0$, if $\varepsilon_1 > 0$ is sufficiently small, then

$$C_0^{-1}|t| \leq r-r' \leq C_0|t| \quad \text{with some } C_0 > 0,$$

on the support of $\tilde{\chi}_1 \tilde{A}_\delta^+$, and this estimate implies $|t/\sqrt{rr'}| \lesssim 1$. We now fix δ with $0 < \delta \leq \varepsilon_1$. Since $\partial_\nu \tilde{\Phi}^+(t) = (\sqrt{rr'}/t) \partial_\omega \Phi_+(t)$, we have

$$(5.39) \quad |\theta - \theta'| \leq C|t/\sqrt{rr'}| (|\partial_\nu \tilde{\Phi}^+(t)| + \varepsilon_1) \leq C_1 \varepsilon_1 \quad \text{with some } C_1 > 0,$$

on the support of $\tilde{\chi}_1 \tilde{A}_\delta^+$. We fix $\lambda'_{\text{IK}} > \lambda_{\text{IK}}$ with $2\lambda_{\text{IK}} > \lambda'_{\text{IK}}$, and choose $\chi_3, \chi_4 \in C_0^\infty(\mathbb{R}^{2d})$ so that $\chi_3 \equiv 1$ on $\Omega^+(\lambda_{\text{IK}}^3)$, $\text{supp } \chi_3 \subset \Omega^+(\lambda_{\text{IK}}^3)$, $\chi_4 \equiv 1$ on $\Omega^+(\lambda_{\text{IK}})$, $\text{supp } \chi_4 \subset \Omega^+(\lambda'_{\text{IK}})$ and

$$(5.40) \quad |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \chi_j(r, \theta, \rho, \omega)| \leq C_{j\alpha k \beta \lambda_1} \langle r \rangle^{-j-|\beta|} \quad \text{on } \mathbb{R}^{3d}, \quad j = 3, 4.$$

We also choose $\chi_5 \in C_0^\infty(\mathbb{R}^{d-1})$ with $\chi_5 \equiv 1$ on $\{|\theta| \leq 1/2\}$ and $\text{supp } \chi_5 \subset \{|\theta| \leq 1\}$. We now define

$$\begin{aligned} \Phi_0^+(t, r, \theta, \rho, \nu, r', \theta') \\ := \frac{r - r'}{t} \rho + \frac{\theta - \theta'}{t} \cdot \sqrt{rr'} \nu - \frac{1}{2} \rho^2 - \frac{1}{2} q_0(\theta, \nu) - \tilde{\chi}_3 \tilde{\chi}_4 \tilde{\chi}_5 Q^+(r, \theta, \rho, \sqrt{rr'} \nu, r', \theta'), \end{aligned}$$

where $\tilde{\chi}_3$, $\tilde{\chi}_4$ and $\tilde{\chi}_5$ are defined by

$$\begin{aligned} \tilde{\chi}_3(r, \theta, \rho, \nu, r', \theta') &:= \chi_3(r, \theta, (\rho_2^+, \omega_2^+)(r, \theta, \rho, \sqrt{rr'} \nu, r', \theta')) \\ \tilde{\chi}_4(r, \theta, \rho, \nu, r', \theta') &:= \chi_3(r', \theta', (\rho_2^+, \omega_2^+)(r, \theta, \rho, \sqrt{rr'} \nu, r', \theta')) \\ \tilde{\chi}_5(\theta, \theta') &:= \chi_5\left(\frac{\theta - \theta'}{2C_1 \varepsilon_1}\right). \end{aligned}$$

It is easy to see that Φ_0^+ is smooth on $[h, \infty) \times \mathbb{R}^{3d}$ and $\Phi_0^+ \equiv \tilde{\Phi}^+$ on $\text{supp } \tilde{\chi}_1 \tilde{A}_\delta^+$. Moreover by Lemma 5.16, (5.39) and (5.40), we have

$$|\partial_\rho^k \partial_\nu^\beta \Phi_0^+(t, r, \theta, \rho, \nu, r', \theta')| \leq C_{k\beta}$$

for all $(r, \theta, \rho, \nu, r', \theta') \in \mathbb{R}^{3d}$, $h \leq t \leq h^{-1}$ and $|k + \beta| \geq 2$, where $C_{k\beta} > 0$ may be taken uniformly in h , t and R . We also obtain

$$\partial_{\rho, \nu}^2 \Phi_0^+ = - \begin{pmatrix} 1 & 0 \\ 0 & h^{jk} \end{pmatrix} + O(\varepsilon_1 + R_1^{-\mu}) \quad \text{on } [h, h^{-1}] \times \mathbb{R}^{3d}.$$

Since $|\det(h^{jk})| \gtrsim 1$, if R_1 is large enough and ε_1 is small enough,

$$|\det \partial_{\rho, \nu}^2 \Phi_0^+| \gtrsim 1$$

on $[h, h^{-1}] \times \mathbb{R}^{3d}$ uniformly with respect to R, h and t . Therefore the mapping

$$(\rho, \nu) \mapsto \partial_{\rho, \nu} \Phi_0^+(t, r, \theta, \rho, \nu, r', \theta')$$

is a diffeomorphism from $\mathbb{R}^d$ to $\mathbb{R}^d$, and Φ_0^+ has a unique non-degenerate critical point $(\rho_c, \nu_c) = (\rho_c, \nu_c)(t, r, r', \theta, \theta')$. Moreover, $\partial_\rho^k \partial_\nu^\beta \Phi_0^+(t, r, \theta, \rho_c, \nu_c, r', \theta')$ are bounded uniformly with respect to R and t if $|k + \beta| \geq 2$. We hence can apply the stationary phase theorem and obtain

$$\begin{aligned} \left| (2\pi h)^{-d} \int e^{\frac{i}{h} t \tilde{\Phi}^+(t)} \tilde{\chi}_1 \tilde{A}_\delta^+ d\rho d\nu \right| &= \left| (2\pi h)^{-d} \int e^{\frac{i}{h} t \Phi_0^+(t)} \tilde{\chi}_1 \tilde{A}_\delta^+ d\rho d\nu \right| \\ &\leq C'_{\varepsilon_1, \lambda_{\text{IK}}} h^{-d} |t/h|^{-d/2} \\ &\leq C'_{\varepsilon_1, \lambda_{\text{IK}}} |th|^{-d/2}, \end{aligned}$$

for $h \in (0, 1]$ and $h \leq t \leq h^{-1}$, where $C'_{\varepsilon_1, \lambda_{\text{IK}}} > 0$ does not depend on $R > 0$. We complete the proof $\square$

Proof of Theorem 5.14. We set

$$U_{\text{IK}}^+(t, h) = U_{A_\delta^+}(t, h), \quad R_{\text{IK}}^+(t, h) = I_{\text{IK}}^+(b^+) e^{-ith \frac{1}{2} D_r^2} I_{\text{IK}}^+(c^+)^* - U_{A_\delta^+}(t, h).$$

Clearly, they satisfy the assertion. When $-h^{-1} \leq t < 0$, the proof is analogous. $\square$

6. MICROLOCAL SMOOTHING PROPERTIES

Fix arbitrarily a coordinate chart $\kappa_0 : V_{\kappa_0} \rightarrow U_{\kappa_0}$. Let $\tilde{U}_{\kappa_0} \Subset U_{\kappa_0}$, $J \Subset (0, \infty)$ and $-1 < \sigma < 1$ be an in Definition 2.5. In this section we prove the following:

Theorem 6.1. *Fix arbitrarily $t_1 > 0$ and let $\varepsilon > 0$ be small enough. Then there exist $\delta_{\varepsilon, t_1} > 0$ and $L_{\varepsilon, t_1} > 0$ such that for all $(\sigma_l)_{0 \leq l \leq L_{\varepsilon, t_1}} \subset (-1, 1/2]$ satisfying (2.12), sufficiently large $R_0 > 0$, all $R_2 \geq R_1 \geq R_0$, all symbols*

$$a_l^\pm \in S_{\text{sc}}(\Gamma_i^\pm(R_1, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l)) \quad b_l^\pm \in S_{\text{sc}}(\Omega_i^\pm(R_2, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l)),$$

and $N \geq 0$, we have

$$(6.1) \quad \|\mathrm{Op}_{\kappa_0, h}(a_l^\pm) e^{-ith\hat{P}} \mathrm{Op}_{\kappa_0, h}(b_l^\pm)\|_{\mathcal{L}(L^2(\widehat{M}))} \leq C_{N, l} h^N,$$

uniformly with respect to $h \in (0, 1]$, $R_2 t_1 \leq \pm t \leq h^{-1}$ and R_2 .

We prove Theorem 6.1 for the case $t \geq 0$, and the proof for case $t \leq 0$ is analogous. We need the following Egorov theorem.

Theorem 6.2 (The Egorov theorem). *For any $T > 0$, $N \geq 0$ and symbol $b^+ \in S_{\mathrm{sc}}(\Omega^+(R_2, \tilde{U}_{\kappa_0}, J, \sigma))$, there exist symbols*

$$b_{\kappa, h}^+(t) = \sum_{j=0}^N h^j b_{\kappa, j}^+(t) \quad \text{with} \quad b_{\kappa, j}^+(t) \in S_{\mathrm{sc}}(\kappa_* \exp tH_p(\kappa_0^{-1} \mathrm{supp} b^+)),$$

and a constant $C_{N, T} > 0$, independent of R_2 , such that

$$(6.2) \quad \left\| e^{-ith\hat{P}} \mathrm{Op}_{\kappa_0}(b^+) e^{ith\hat{P}} - \sum_{\kappa} \mathrm{Op}_{\kappa}(b_{\kappa, h}^+(t)) \right\|_{\mathcal{L}(L^2(\widehat{M}))} \leq C_{N, T} h^{N+1} |t|$$

uniformly with respect to $0 \leq t \leq R_2 T$ and $h \in (0, 1]$. Moreover, $b_{\kappa, j}^+(t)$ is uniformly bounded in $S_{\mathrm{sc}}(\kappa_* \exp tH_p(\kappa_0^{-1} \mathrm{supp} b^+))$ with respect to $0 \leq t \leq R_2 T$.

Proof. This theorem is basically well known, and we hence give the sketch of the proof. By (4.19), we can choose $t_1 > 0$, independent of R_2 , such that the geodesic is contained one fixed coordinate neighborhood if $0 \leq t \leq R_2 t_1$. Define the map $\tilde{\varphi}(t) = (\tilde{r}(t), \tilde{\theta}(t), \tilde{\rho}(t), \tilde{\omega}(t))$ by

$$(\tilde{r}(t), \tilde{\theta}(t), \tilde{\rho}(t), \tilde{\omega}(t)) = F_{R_2} \circ \exp tH_{p_{\kappa_0}} \circ F_{R_2}^{-1}(r, \theta, \rho, \omega),$$

where $F_{R_2}(r, \theta, \rho, \omega) = (r/R_2, \theta, \rho, \omega/R_2)$. By (4.4), we have for all $(x, \theta, \rho, \nu) \in F_{R_2} \Omega^+(R_2, \tilde{U}_{\kappa_0}, J, \sigma)$, $0 \leq t \leq R_2 t_1$ and $\gamma = (j, \alpha, k, \beta)$,

$$|\partial^\gamma(\tilde{r}(t) - x)| + |\partial^\gamma(\tilde{\theta}(t) - \theta)| + |\partial^\gamma(\tilde{\rho}(t) - \rho)| + |\partial^\gamma(\tilde{\omega}(t) - \nu)| \leq C_\gamma R_2^{-j-|\beta|} t_1,$$

and hence

$$|\partial^\gamma(\partial\tilde{\varphi}(t) - \mathrm{Id})| \leq C t_1 < 1/2,$$

where $x = r/R_2$, $\nu = \omega/R_2$ and $\partial^\gamma = \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta$. For all $0 \leq t \leq R_2 t_1$ and $(r, \theta, \rho, \omega) \in \tilde{\varphi}(t, F_{R_2} \Omega^+(R_2, \tilde{U}_{\kappa_0}, J, \sigma))$, $\tilde{\varphi}(t)$ thus has the inverse

$$\tilde{\varphi}(t)^{-1} = (\tilde{r}(t)^{-1}, \tilde{\theta}(t)^{-1}, \tilde{\rho}(t)^{-1}, \tilde{\omega}(t)^{-1}),$$

and $\tilde{\varphi}(t)^{-1}$ satisfies

$$\begin{aligned} |\partial^\gamma(\tilde{r}(t)^{-1} - x)| + |\partial^\gamma(\tilde{\theta}(t)^{-1} - \theta)| &\leq C_\gamma R_2^{-j-|\beta|} t_1, \\ |\partial^\gamma(\tilde{\rho}(t)^{-1} - \rho)| + |\partial^\gamma(\tilde{\omega}(t)^{-1} - \nu)| &\leq C_\gamma R_2^{-j-|\beta|} t_1. \end{aligned}$$

After the rescaling $(x, \nu) \mapsto (R_2 x, R_2 \nu)$, we see that

$$(r(t)^{-1}, \theta(t)^{-1}, \rho(t)^{-1}, \omega(t)^{-1}) = (\exp tH_{p_{\kappa_0}})^{-1}(r, \theta, \rho, \omega)$$

exists for all $(r, \theta, \rho, \omega) \in \exp tH_{p_{\kappa_0}}(\Omega^+(R_2, \tilde{U}_{\kappa_0}, J, \sigma))$ and $0 \leq t \leq R_2 t_1$, and satisfies

$$(6.3) \quad \begin{aligned} |\partial^\gamma(r(t)^{-1} - r)| + |\partial^\gamma(\omega(t)^{-1} - \omega)| &\leq C_{j\alpha k\beta} R_2^{1-j-|\beta|} t_1, \\ |\partial^\gamma(\theta(t)^{-1} - \theta)| + |\partial^\gamma(\rho(t)^{-1} - \rho)| &\leq C_{j\alpha k\beta} R_2^{-j-|\beta|} t_1. \end{aligned}$$

We now define $b_j^+(t)$ inductively as follows. Put

$$b_0^+(t, r, \theta, \rho, \omega) = b^+ \circ (\exp tH_{p_{\kappa_0}})^{-1}(r, \theta, \rho, \omega) \quad \text{on} \quad \exp tH_{p_{\kappa_0}}(\mathrm{supp} b^+),$$

and $b_0^+(t, r, \theta, \rho, \omega) = 0$ outside $\exp tH_{p_{\kappa_0}}(\text{supp } b^+)$. By (6.3), $(b_0^+(t))_{0 \leq t \leq R_2 t_1}$ is bounded in $S_{\text{sc}}(\exp tH_{p_{\kappa_0}}(\text{supp } b^+))$. It is well known that $b_0^+(t)$ solves the first transport equation

$$\frac{\partial b_0^+}{\partial t} + \{p_{\kappa_0}, b_0^+\} = 0, \quad b_0^+(0) = b^+,$$

where $\{\cdot, \cdot\}$ is the Poisson bracket. Moreover, a standard semiclassical symbolic calculus yields that

$$\frac{\partial}{\partial t} \text{Op}_{\kappa_0, h}(b_0^+(t)) + \frac{i}{h} [h^2 \widehat{P}_{\kappa_0}, \text{Op}_{\kappa_0, h}(b_0^+(t))] = h \text{Op}_{\kappa_0, h}(r_0^+(t)),$$

where $r_0^+(t)$ is supported in $\text{supp } b_0^+(t)$ modulo $O(h^\infty)$ on $L^2(\widehat{M})$, i.e., for all $n \geq 0$, there exists a symbol $\tilde{r}_0^+(t) \in S_{\text{sc}}(\text{supp } b_0^+(t))$ such that

$$\|\text{Op}_{\kappa_0, h}(r_0^+(t)) - \text{Op}_{\kappa_0, h}(\tilde{r}_0^+(t))\|_{\mathcal{L}(L^2(\widehat{M}))} \leq C_n h^n, \quad h \in (0, 1],$$

for $0 \leq t \leq R_2 t_1$. Next, put

$$b_1^+(t, r, \theta, \rho, \omega) = \int_0^t \tilde{r}_0^+(s, \exp sH_{p_{\kappa_0}} \circ (\exp tH_{p_{\kappa_0}})^{-1}(r, \theta, \rho, \omega)) ds$$

on $\exp tH_{p_{\kappa_0}}(\text{supp } b^+)$, and $b_1^+(t, r, \theta, \rho, \omega) = 0$ otherwise. Again (6.3) implies $(b_1^+(t))_{0 \leq t \leq R_2 t_1}$ is bounded in $S_{\text{sc}}(\exp tH_{p_{\kappa_0}}(\text{supp } b^+))$. $b_1^+(t)$ is a solution to the second transport equation

$$\frac{\partial b_1^+}{\partial t} + \{p_{\kappa_0}, b_1^+\} = \tilde{r}_0^+, \quad b_1^+(0) = 0,$$

which implies

$$\frac{\partial}{\partial t} \text{Op}_{\kappa_0, h}(b_0^+(t) + hb_1^+(t)) + \frac{i}{h} [h^2 \widehat{P}_{\kappa_0}, \text{Op}_{\kappa_0, h}(b_0^+(t) + hb_1^+(t))] = h^2 \text{Op}_{\kappa_0, h}(r_1^+(t)),$$

where $r_1^+(t)$ is supported in $\text{supp } b_0^+(t)$ modulo $O(h^\infty)$ on $L^2(\widehat{M})$. Iterating this procedure and putting $b_h^+(t) = \sum_{j=0}^N h^j b_j^+(t)$, we have

$$\frac{\partial}{\partial t} \text{Op}_{\kappa_0, h}(b_h^+(t)) + \frac{i}{h} [h^2 \widehat{P}_{\kappa_0}, \text{Op}_{\kappa_0, h}(b_h^+(t))] = O(h^{N+1}),$$

and $\text{Op}_{\kappa_0, h}(b_h^+(0)) = \text{Op}_{\kappa_0, h}(b^+)$. Integrating the above equation with respect to $t \in [0, R_2 t_1]$, we obtain the assertion for $t \in [0, R_2 t_1]$. For general $T > 0$, we divide the geodesics into a finite number of small curves, as well as the proof of Corollary 4.3, so that each curve is contained some fixed coordinate neighborhood. Applying the above argument on each chart, we have the assertion by a partition of unity argument. $\square$

The following tells us that the support of $e^{-ith\widehat{P}} \text{Op}_{\kappa_0}(b_l^+)$ is essentially away from the support of $\text{Op}_{\kappa_0}(a_l^+)$ if $t = O(r)$, which is crucial to prove Theorem 6.1.

Proposition 6.3. *There exists $c_0 > 0$ that for all $0 < \varepsilon < 1/2$ and $t_1 > 0$, if we choose $\delta_{\varepsilon, t_1} < c_0 \varepsilon^2 \langle 1/t_1 \rangle^{-1}$ and $L_{\varepsilon, t_1} = \delta_{\varepsilon, t_1}^{-1}$, then for all $(\sigma_l)_{0 \leq l \leq L_{\varepsilon, t_1}} \subset (-1, 1/2]$ satisfying (2.12) with $\delta = \delta_{\varepsilon, t_1}$, $L = L_{\varepsilon, t_1}$ and all $t \geq R_2 t_1$,*

$$(6.4) \quad \exp tH_{p_{\kappa_0}} \kappa_{0*}^{-1} \Omega_i^+(R_2, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l) \cap \kappa_{0*}^{-1} \Gamma_i^+(R_1, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l) = \emptyset.$$

Proof. It suffices to show that (6.4) holds with $\Omega_i^+(R_2, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l)$ replaced by

$$\Omega_i^+(R_1, R_2, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l),$$

since $\Omega_i^+(R_2, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l) \subset \Omega_i^+(R_1, R_2, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l)$. We also note that, by Corollary 4.3, $\Omega_i^+(R_1, R_2, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l)$ is invariant with respect to ρ under the geodesic flow and coordinate transformations. Therefore, by the definition of the intermediate regions and the energy conservation law, it suffices to check that

$$\frac{\rho(t)}{\sqrt{2E_0}} > \frac{\rho}{\sqrt{2E_0}} + 2\delta_{\varepsilon, t_1}, \quad t \geq R_2 t_1,$$

where $E_0 = p_{\kappa_0}(r, \theta, \rho, \omega)$ and $(r, \theta, \rho, \omega)$ belongs to

$$\{R_1 < r < 4R_2, \theta \in \tilde{U}_{\kappa_0}, E_0 \in J, \rho/\sqrt{2E_0} \in [-1/2, \sqrt{1-\varepsilon^2/4}]\}.$$

Note that all $\Omega_i^+(R_1, R_2, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l)$ are contained in the above region. By (4.1) and (4.3), for sufficiently large $r > R_1$ and all $t \geq 0$,

$$\dot{\rho}(t) \geq \frac{1}{r(t)^3} (h^{jk}(\theta(t)) - O(r(t)^{-\mu})) \omega_j(t) \omega_k(t) \gtrsim (r + |t|)^{-3} r^2 \varepsilon^2 \sqrt{2E_0}.$$

Since

$$\int_0^{R_2 t_1} (r + |t|)^{-3} r^2 dt \gtrsim \frac{R_2 t_1}{r + R_2 t_1} \geq \frac{t_1}{4 + t_1},$$

we can find $c_0 > 0$ small enough such that

$$\frac{\rho(R_2 t_1)}{\sqrt{2E_0}} \geq \frac{\rho}{\sqrt{2E_0}} + 2c_0 \varepsilon^2 \frac{t_1}{1 + t_1} > \frac{\rho}{\sqrt{2E_0}} + 2\delta_{\varepsilon, t_1},$$

which implies the assertion since $\dot{\rho}(t)$ is non-negative. $\square$

Proof of Theorem 6.1. Since (4.5) implies

$$|\sqrt{2E_0} - \rho(t)| \lesssim R_2/(R_2 + |t|), \quad t \geq 0,$$

it follows from (4.17) and (4.18) that there exists a constant $C_0 > 0$ such that for any $0 < \varepsilon_0 < 1/2$, we can find $T_{\varepsilon_0} > 0$ such that

$$(6.5) \quad \begin{aligned} & \exp R_2 T_{\varepsilon_0} H_p(\kappa_{0*}^{-1} \Omega_i^+(R_2, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l)) \\ & \subset \bigcup_{\kappa} \kappa_*^{-1} \Omega_s^+(R_2 T_{\varepsilon_0}/C_0, C_0 R_2 T_{\varepsilon_0}, \tilde{U}_{\kappa}, J, \varepsilon_0), \end{aligned}$$

where $\tilde{U}_{\kappa} \subseteq U_{\kappa}$. Note that C_0 and T_{ε_0} may be taken uniformly with respect to R_2 . We fix such a T_{ε_0} with $T_{\varepsilon_0} \geq C_0$. By Theorem 6.2, we obtain that for all $N \geq 0$,

$$\text{Op}_{\kappa_0}(a_l^+) e^{-ith\hat{P}} \text{Op}_{\kappa_0}(b_l^+) = \text{Op}_{\kappa_0}(a_l^+) \sum_{\kappa} \text{Op}_{\kappa}(b_{\kappa, h}^+(t)) e^{-ith\hat{P}} + O(h^{N+1}|t|)$$

on $L^2(\widehat{M})$ uniformly with respect to R_2 and $0 \leq t \leq R_2 T_{\varepsilon_0}$, where

$$b_{\kappa, h}^+(t) \in S_{\text{sc}}(\kappa_* \exp t H_p(\kappa_{0*}^{-1} \Omega_i^+(R_2, \tilde{U}_{\kappa_0}, J, \varepsilon, \delta_{\varepsilon, t_1}, l))).$$

Suppose that $R_2 t_1 \leq t \leq R_2 T_{\varepsilon_0}$. Since the support of a_l^+ does not intersect with the support of $(\kappa \circ \kappa_0^{-1})_* b_{\kappa, h}^+(t)$ for any κ with $V_{\kappa_0} \cap V_{\kappa} \neq \emptyset$ by Proposition 6.3, The semiclassical symbolic calculus (see subsection 2.2) implies that the above operator is bounded on $L^2(\widehat{M})$ with the norm dominated by $C_N h^{N+1} \langle t \rangle$, where the constant $C_N > 0$ may be taken uniformly with respect to R_2, t and h . If $R_2 T_{\varepsilon_0} \geq h^{-1}$, then we obtain (6.1). We thus assume $R_2 T_{\varepsilon_0} \leq t \leq h^{-1}$. By (6.5) and Theorem 6.2, there exist symbols $b_{\kappa, h}^+(R_2 T_{\varepsilon_0}) \in S_{\text{sc}}(\Omega_s^+(R_2 T_{\varepsilon_0}/C_0, C_0 R_2 T_{\varepsilon_0}, \tilde{U}_{\kappa}, J, \varepsilon_0))$ such that

$$\begin{aligned} e^{-ith\hat{P}} \text{Op}_{\kappa_0}(b_l^+) &= e^{-i(t-R_2 T_{\varepsilon_0})h\hat{P}} e^{-iR_2 T_{\varepsilon_0} h\hat{P}} \text{Op}_{\kappa_0}(b_l^+) \\ &= e^{-i(t-R_2 T_{\varepsilon_0})h\hat{P}} \sum_{\kappa} \text{Op}_{\kappa}(b_{\kappa, h}^+(R_2 T_{\varepsilon_0})) e^{-iR_2 T_{\varepsilon_0} h\hat{P}} + O(h^N) \end{aligned}$$

on $L^2(\widehat{M})$. Put $B = \sum_{\kappa} \text{Op}_{\kappa}(b_{\kappa, h}^+(R_2 T_{\varepsilon_0}))$ and divide B as follows

$$B = \varphi_{\kappa_0} B \tilde{\varphi}_{\kappa_0} + (1 - \varphi_{\kappa_0}) B \tilde{\varphi}_{\kappa_0} + B(1 - \tilde{\varphi}_{\kappa_0}),$$

where $\varphi_{\kappa_0}, \tilde{\varphi}_{\kappa_0} \in C^\infty(\widehat{M}_\infty)$ such that

$$\text{supp } \tilde{\varphi}_{\kappa_0} \subset \text{supp } \varphi_{\kappa_0} \subset (2, \infty) \times V_{\kappa_0}, \quad \varphi_{\kappa_0} \equiv 1 \text{ close to } \text{supp } \tilde{\varphi}_{\kappa_0}.$$

Since $\text{supp}(1 - \varphi_{\kappa_0}) \cap \text{supp } \tilde{\varphi}_{\kappa_0} = \emptyset$, the second term is $O(h^\infty)$ on $L^2(\widehat{M})$. The third term is also $O(h^\infty)$ on $L^2(\widehat{M})$ since $\text{supp } b_{\kappa, h}^+(R_2 T_{\varepsilon_0}) \cap \text{supp}(1 - \tilde{\varphi}_{\kappa_0}) = \emptyset$, which follows from the facts

$$r > R_2 > 2 \quad \text{on } \text{supp } b_{\kappa, h}^+(R_2 T_{\varepsilon_0}), \quad 0 < r < 2 \quad \text{on } \text{supp}(1 - \tilde{\varphi}_{\kappa_0}).$$

By invariance properties of the strongly outgoing region and h -PDO under coordinates transformations, we can write

$$\varphi_{\kappa_0} B \tilde{\varphi}_{\kappa_0} = \text{Op}_{\kappa_0}(\tilde{b}_{\kappa_0, h}^+(R_2 T_{\varepsilon_0})) + O(h^\infty)$$

on $L^2(\widehat{M})$ with some $\tilde{b}_{\kappa_0, h}^+(R_2 T_{\varepsilon_0}) \in S_{\text{sc}}(\Omega_s^+(R_2 T_{\varepsilon_0}/C_0, C_0 R_2 T_{\varepsilon_0}, \tilde{U}'_{\kappa_0}, J, \varepsilon_0))$, where $\tilde{U}'_{\kappa_0} \subseteq U_{\kappa_0}$. Consider a splitting of the interval $[R_2 T_{\varepsilon_0}/C_0, C_0 R_2 T_{\varepsilon_0}]$:

$$R_2 T_{\varepsilon_0}/C_0 = \tilde{R}_0 < \tilde{R}_1 < \tilde{R}_2 < \cdots < \tilde{R}_k, \quad \tilde{R}_j = 2^j \tilde{R}_0, \quad \tilde{R}_{k-1} < C_0 R_2 T_{\varepsilon_0} < \tilde{R}_k.$$

Clearly $2^k \leq 2C_0^2$. Using a method of the partition of unity, we split $\tilde{b}_{\kappa_0, h}^+(R_2 T_{\varepsilon_0})$ as $\tilde{b}_{\kappa_0, h}^+(R_2 T_{\varepsilon_0}) = \sum_{j=0}^{k-2} \tilde{b}_{\kappa_0, h, j}^+$ with $\tilde{b}_{\kappa_0, h, j}^+ \in S_{\text{sc}}(\Omega_s^+(\tilde{R}_j, \tilde{U}'_{\kappa_0}, J, \varepsilon_0))$. By Theorem 5.10, we can construct the Isozaki-Kitada parametrix of $e^{-i(t-R_2 T_{\varepsilon_0})h\hat{P}} \text{Op}_{\kappa_0}(\tilde{b}_{\kappa_0, h, j}^+)$ for sufficiently large $R_2 \geq R_{\text{IK}}, \lambda \geq \lambda_{\text{IK}}$ and small $0 < \varepsilon_0 \leq \varepsilon_{\text{IK}}$, and we obtain

$$\begin{aligned} & e^{-ith\hat{P}} \text{Op}_{\kappa_0}(b_l^+) \\ &= \sum_{j=0}^{k-2} \kappa_0^* I_{\text{IK}}^+(c_{h, j}^+) e^{-i(t-R_2 T_{\varepsilon_0})h\frac{1}{2}D_r^2} I_{\text{IK}}^+(d_{h, j}^+)^* \kappa_0 e^{-iR_2 T_{\varepsilon_0}h\hat{P}} + Q(t, h, N, R_2), \end{aligned}$$

where, for each j , $I_{\text{IK}}^+(c_{h, j}^+)$ and $I_{\text{IK}}^+(d_{h, j}^+)$ are FIO's defined in Definition 5.6 with some phase function $S_{\kappa_0, j}^+ \in C^\infty(\mathbb{R}^d; \mathbb{R})$, which satisfies the statement of Theorem 5.1 with $R = \tilde{R}_j$, and some amplitudes

$$\begin{aligned} c_{h, j}^+ &\in S_{\text{sc}}(\Omega_s^+(\lambda^{-3}\tilde{R}_j, \lambda^3\tilde{R}_j, \tilde{U}'_{\kappa_0}, J, \lambda^3\varepsilon_0)), \\ d_{h, j}^+ &\in S_{\text{sc}}(\Omega_s^+(\lambda^{-1}\tilde{R}_j, \lambda\tilde{R}_j, \tilde{U}'_{\kappa_0}, J, \lambda\varepsilon_0)). \end{aligned}$$

The remainder term $Q(t, h, N, R_2)$ is uniformly bounded on $L^2(\widehat{M})$ with the norm of order h^N with respect to R_2 and $R_2 T_{\varepsilon_0} \leq t \leq h^{-1}$. By the composition rule of FIO's with PDO's, $a_l^+(r, \theta, hD_r, hD_\theta) I_{\text{IK}}^+(c_{h, j}^+)$ is also FIO's (up to the smoothing term $O(h^\infty)$ on $L^2(\widehat{M})$) with the phase $S_{\kappa_0, j}^+$ and the amplitude supported in

$$X = \{(r, \theta, \rho, \omega) \mid a_l^+(r, \theta, \partial_r S_{\kappa_0, j}^+, \partial_\theta S_{\kappa_0, j}^+) c_{h, j}^+(r, \theta, \rho, \omega) \neq 0\}.$$

By the support property of a_l^+ , we see that

$$\partial_r S_{\kappa_0, j}^+ \leq \sqrt{2(1 - \varepsilon^2/4)p_{\kappa_0}(r, \theta, \partial_r S_{\kappa_0, j}^+, \partial_\theta S_{\kappa_0, j}^+)}.$$

Since $(r, \theta, \rho, \omega) \in \text{supp } c_{h, j}^+ \subset \Omega_s^+(\lambda^{-3}\tilde{R}_j, \lambda^3\tilde{R}_j, \tilde{U}'_{\kappa_0}, J, \lambda^3\varepsilon_0)$, by the above estimate and (5.1), we have

$$\rho/\sqrt{2p_{\kappa_0}(r, \theta, \rho, \omega)} \leq \sqrt{1 - \varepsilon^2/4} - C|\lambda^3\varepsilon_0|^2 \leq \sqrt{1 - \varepsilon^2/8}.$$

On the other hand, choosing $\varepsilon_0 > 0$ small enough so that $\lambda^3\varepsilon_0 < \varepsilon^2/8$ (note that $\lambda > \lambda_{\text{IK}}$ is fixed), the support property of $c_{h, j}^+$ implies

$$\rho/\sqrt{2p_{\kappa_0}(r, \theta, \rho, \omega)} \geq \sqrt{1 - \lambda^3\varepsilon_0} > \sqrt{1 - \varepsilon^2/8}.$$

The above two inequalities show that $X = \emptyset$ and hence $a_l^+(r, \theta, hD_r, hD_\theta) I_{\text{IK}}^+(c_{h, j}^+)$ is $O(h^\infty)$ in $L^2(\widehat{M})$. Since $\kappa_0^* e^{-i(t-R_2 T_{\varepsilon_0})h\frac{1}{2}D_r^2} I_{\text{IK}}^+(d_{h, j}^+)^* \kappa_0 e^{-iR_2 T_{\varepsilon_0}h\hat{P}}$ is uniformly bounded on $L^2(\widehat{M})$ with respect to R_2, h and t , we obtain the assertion and conclude the proof. $\square$

7. THE WKB PARAMETRIX

In the previous section we proved that $\text{Op}_{\kappa_0, h}(a_l^\pm) e^{-ith\hat{P}} \text{Op}_{\kappa_0, h}(b_l^\pm)$ are rapidly decaying with respect to $h \in (0, 1]$ if $Rt_1 \leq \pm t \leq h^{-1}$ with any $t_1 > 0$ and large $R > 0$. Therefore, it remains to control the above operators for $0 \leq \pm t \leq Rt_0$ with sufficiently small t_0 . This section discuss construction of the WKB parametrix of propagator $e^{-it\hat{P}} \text{Op}_\kappa(a^\pm)$ for $0 \leq \pm t \leq Rt_0$, where a^+ (resp. a^-) is supported in an outgoing (resp. incoming) region. By (4.18), we can always work on a fixed coordinate neighborhood U_κ and hence do not write the subscript κ explicitly. Let

$\tilde{U}_\kappa \Subset U_\kappa$ be as in Example 2.2 and fix open subsets $U \Subset U_0 \Subset U_1 \Subset U_2 \Subset \tilde{U}_\kappa$, open intervals $J \Subset J_0 \Subset J_1 \Subset J_2 \Subset (0, \infty)$ and constants $-1 < \sigma < \sigma_0 < \sigma_1 < \sigma_2 < 1$ arbitrarily.

7.1. Fourier integral operators for the WKB parametrix. We here study time dependent FIO's which will be used to construct the WKB parametrix. We first construct the phase function.

Theorem 7.1. *We can choose $t_0 > 0$ small enough such that, for sufficiently large $R_2 > 0$ and all $R_1 > R_2$, there exist smooth and real-valued functions*

$$\Psi^+ \in C^\infty((0, R_1 t_0) \times \mathbb{R}^{2d}), \quad \Psi^- \in C^\infty((-R_1 t_0, 0) \times \mathbb{R}^{2d}),$$

satisfying the following Hamilton Jacobi equation on $\Gamma^\pm(R_1, U_1, J_1, \sigma_1)$:

$$(7.1) \quad \begin{cases} \partial_t \Psi^\pm + p(r, \theta, \partial_r \Psi^\pm, \partial_\theta \Psi^\pm) = 0, & 0 \leq \pm t \leq R_1 t_0, \\ \Psi^\pm|_{t=0} = r\rho + \theta \cdot \omega, \end{cases}$$

such that we have the followings all $0 \leq \pm t \leq R_1 t_0$:

$$\text{supp}(\Psi^\pm(t, r, \theta, \rho, \omega) - r\rho - \theta \cdot \omega) \subset \Gamma^\pm(R_2, U_2, J_2, \sigma_2),$$

$$(7.2) \quad \left| \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\Psi^\pm(t, r, \theta, \rho, \omega) - r\rho - \theta \cdot \omega) \right| \leq C_{j\alpha k\beta} \langle r \rangle^{-j-|\beta|} |t| \quad \text{on } \mathbb{R}^{2d}.$$

Moreover, for all $(r, \theta, \rho, \omega) \in \Gamma^\pm(R_1, U_1, J_1, \sigma_1)$, we have

$$(7.3) \quad \left| \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\Psi^\pm(t, r, \theta, \rho, \omega) - r\rho - \theta \cdot \omega + tp(r, \theta, \rho, \omega)) \right| \leq C_{j\alpha k\beta} r^{-j-|\beta|} |t| t_0.$$

Proof. This theorem can be proved similarly to Theorem 5.1. We only prove the theorem for the case $t \geq 0$, and the proof for the case $t \leq 0$ is similar. Let $R_1 > R'_1 > R_1/2$, $U_1 \Subset U'_1 \Subset \tilde{U}_\kappa$, $J_1 \Subset J'_1 \Subset (0, \infty)$ and $\sigma_1 < \sigma'_1 < 1$. Let $F_{R_1} : (r, \theta, \rho, \omega) \mapsto (r/R_1, \theta, \rho, \omega)$, and define $g^+(t)$ and $\tilde{g}^+(t)$ by

$$\begin{aligned} g^+(t) &:= (r(t, r, \theta, \rho, \omega), \theta(t, r, \theta, \rho, \omega), \rho, \omega), \\ \tilde{g}^+(t) &= (\tilde{r}(t, x, \theta, \rho, \omega), \tilde{\theta}(t, x, \theta, \rho, \omega), \rho, \omega) \\ &:= F_{R_1} \circ g^+(t) \circ F_{R_1}^{-1}(x, \theta, \rho, \omega), \end{aligned}$$

where $(r(t, r, \theta, \rho, \omega), \theta(t, r, \theta, \rho, \omega))$ is the Hamilton flow generated by p , and $x = r/R_1$. By Proposition 4.2, we have

$$\begin{aligned} |\partial_x^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\tilde{r}(t) - x)| &\leq C_{j\alpha k\beta} R_1^{-1} |t| \leq C_{j\alpha k\beta} t_0, \\ |\partial_x^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\tilde{\theta}(t) - \theta)| &\leq C_{j\alpha k\beta} R_1^{-1} |t| \leq C_{j\alpha k\beta} t_0, \end{aligned}$$

and

$$|\partial \tilde{g}^+(t) - \text{Id}| \leq C t_0 < 1/2,$$

for all $(x, \theta, \rho, \omega) \in F_{R_1} \Gamma^+(R'_1, U'_1, J'_1, \sigma'_1)$ and $0 \leq t \leq R_1 t_0$ as long as $t_0 > 0$ is small enough. Applying a same argument as that in the proof of Lemma 5.3 to $\tilde{g}^+(t)$, we see that $g^+(t)$ is diffeomorphic from $\Gamma^+(R'_1, U'_1, J'_1, \sigma'_1)$ onto its range for all $0 \leq t \leq R_1 t_0$, and satisfies

$$\Gamma^+(R_1, U_1, J_1, \sigma_1) \subset g^\pm(t)(\Gamma^+(R'_1, U'_1, J'_1, \sigma'_1)), \quad 0 \leq \pm t \leq R_1 t_0.$$

Let $\Gamma^+(R_1, U_1, J_1, \sigma_1) \ni (r, \theta, \rho, \omega) \mapsto (\hat{r}^+(t), \hat{\theta}^+(t), \rho, \omega)$ be the inverse of $g^+(t)$, and put $(r^+(t, s), \theta^+(t, s), \rho^+(t, s), \omega^+(t, s)) := (r, \theta, \rho, \omega)(s, \hat{r}^+(t), \hat{\theta}^+(t), \rho, \omega)$ for $0 \leq s \leq t \leq R_1 t_0$. Then, by a same argument as that in the proof of Lemma 5.4 and Lemma 5.5, we have

$$(7.4) \quad \begin{cases} |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\hat{r}^+(t) - r)| + |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\omega^+(t, t) - \omega)| \leq C_{j\alpha k\beta} r^{-j-|\beta|} |t|, \\ |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\hat{\theta}^+(t) - \theta)| + |\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta (\rho^+(t, t) - \rho)| \leq C_{j\alpha k\beta} r^{-j-|\beta|} t_0, \end{cases}$$

Define $\tilde{\Psi}^+ \in C^\infty((0, R_1 t_0) \times \Gamma^+(R_1, U_1, J_1, \sigma_1))$ by

$$\tilde{\Psi}^+(t) := r\rho + \theta \cdot \omega + \int_0^t L(r^+(t, s), \theta^+(t, s), \rho^+(t, s), \omega^+(t, s)) ds,$$

where $L = (\rho \partial_\rho p + \omega \cdot \partial_\omega p - p)(r, \theta, \rho, \omega)$. By a standard Hamilton-Jacobi theory, it is easy to see that $\tilde{\Psi}^+(t)$ solves (7.1) and satisfies

$$\partial \tilde{\Psi}(t) = (\rho^+(t, t), \omega^+(t, t), \hat{r}^+(t), \hat{\theta}^+(t)).$$

By (7.4) and the energy conservation law

$$p(r, \theta, \partial_r \tilde{\Psi}^+(t), \partial_\theta \tilde{\Psi}^+(t)) = p(\hat{r}^+(t), \hat{\theta}^+(t), \rho, \omega),$$

we have

$$\left| \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \left(p(r, \theta, \partial_r \tilde{\Psi}^+(t), \partial_\theta \tilde{\Psi}^+(t)) - p(r, \theta, \rho, \omega) \right) \right| \leq C_{j\alpha k\beta} r^{-j-|\beta|} t_0.$$

Therefore,

$$\left| \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \left(\tilde{\Psi}^+(t) - r\rho - \theta \cdot \omega + tp(r, \theta, \rho, \omega) \right) \right| \leq C_{j\alpha k\beta} r^{-j-|\beta|} |t| t_0.$$

Choose $\chi_+ \in C^\infty(\mathbb{R}^{2d})$ so that

$$0 \leq \chi_+ \leq 1, \quad \chi_+ \equiv 1 \quad \text{on} \quad \Gamma^+(R_1, U_1, J_1, \sigma_1), \quad \text{supp } \chi_+ \subset \Gamma^+(R_2, U_2, J_2, \sigma_2),$$

and that

$$\left| \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \chi_+(r, \theta, \rho, \omega) \right| \leq C_{j\alpha k\beta} \langle r \rangle^{-j-|\beta|} \quad \text{on} \quad \mathbb{R}^{2d}.$$

We now define $\Psi^+(t) := r\rho + \theta \cdot \omega + \chi_+(\tilde{\Psi}^+(t) - r\rho + \theta \cdot \omega)$. Clearly, $\Psi(t)^+$ satisfies the statement of theorem 7.1. $\square$

Suppose $(a^\pm(t))_{0 \leq \pm t \leq R_1 t_0}$ are bounded in $S_{\text{sc}}(\Gamma^\pm(R_1, U_1, J_1, \sigma_1))$, respectively. We define the FIO's for the WKB parametrix $I_{\text{WKB}}^\pm(a^\pm(t)) : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}(\mathbb{R}^d)$ by

$$\begin{aligned} & I_{\text{WKB}}^\pm(a^\pm(t))u(r, \theta) \\ &:= \frac{1}{(2\pi h)^d} \int e^{\frac{i}{h}(\Psi^\pm(t, r, \theta, \rho, \omega) - r'\rho - \theta' \cdot \omega)} a^\pm(t, r, \theta, \rho, \omega) u(r', \theta') dr' d\theta' d\rho d\omega. \end{aligned}$$

Proposition 7.2. $I_{\text{WKB}}^\pm(a^\pm(t))$ are bounded on $L^2(\mathbb{R}^d)$ uniformly with respect to $0 \leq \pm t \leq R_1 t_0$:

$$\|I_{\text{WKB}}^\pm(a^\pm(t))u\|_{L^2(\mathbb{R}^d)} \leq C\|u\|_{L^2(\mathbb{R}^d)}.$$

Proof. For $(r, \theta, \rho, \omega, r', \theta') \in \mathbb{R}^{3d}$ with $r, r' > R_1$ and $0 \leq t \leq R_1 t_0$, define the map (ρ_+^1, ω_+^1) by

$$(\rho_+^1, \omega_+^1)(t, r, \theta, \rho, \omega, r', \theta') = \int_0^1 (\partial_{r, \theta} \Psi^+)(t, r' + s(r - r'), \theta' + s(\theta - \theta'), \rho, \omega) ds.$$

By (7.2), we have

$$\sup_{0 \leq t \leq R_1 t_0} (|\partial^\gamma(\rho_+^1 - \rho)| + R_1^{-1} |\partial^\gamma(\omega_+^1 - \omega)|) \leq C_\gamma R_1^{-j-j'-|\beta|} t_0,$$

where $\partial^\gamma = \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \partial_{r'}^{j'} \partial_{\theta'}^{\alpha'}$ with $\gamma = (j, \alpha, k, \beta, j', \alpha')$. By this estimate and a similar argument as in the proof of Lemma 5.9, we obtain that, for all $(r, \theta, r', \theta') \in \mathbb{R}^{2d}$ with $r, r' > R_1$ and $0 \leq t \leq R_1 t_0$, the map $(\rho, \omega) \mapsto (\rho_+^1, \omega_+^1)(t, r, \theta, \rho, \omega, r', \theta')$ is a diffeomorphism from $\mathbb{R}^d$ onto itself provided that $t_0 > 0$ is small enough. We also see that the corresponding inverse $(\rho, \omega) \mapsto (\rho_+^2, \omega_+^2)(t, r, \theta, \rho, \omega, r', \theta')$ satisfies

$$(7.5) \quad \sup_{0 \leq t \leq R_1 t_0} (|\partial^\gamma(\rho_+^2 - \rho)| + R_1^{-1} |\partial^\gamma(\omega_+^2 - \omega)|) \leq C_\gamma R_1^{-j-j'-|\beta|} t_0.$$

Using the change of variable $(\rho, \omega) \mapsto (\rho_+^2, \omega_+^2)$, $I_{\text{WKB}}^+(a^+(t)) I_{\text{WKB}}^+(a^+(t))^*$ can be regarded as a h -PDO with the amplitude

$$a^+(t, r, \theta, \rho_+^2, \omega_+^2) \overline{a^+(t, r', \theta', \rho_+^2, \omega_+^2)} |\det \partial_{\rho, \omega}(\rho_+^2, \omega_+^2)|.$$

By (7.5), this amplitude and its all derivatives with respect to ∂^γ are bounded on $\mathbb{R}^{3d}$ uniformly with respect to $0 \leq t \leq R_1 t_0$. The assertion then follows from the Calderón-Vaillancourt theorem and the L^2 -functional calculus. When $-R_1 t_0 \leq t \leq 0$, the proof is similar. $\square$

7.2. Construction of the parametrix. The main result in this section is the following.

Theorem 7.3. *There exists $R_{\text{WKB}} > 0$ large enough and $t_{\text{WKB}} > 0$ small enough such that, for all $R > R_0 > R_1 > R_{\text{WKB}}$, $a^\pm \in S_{\text{sc}}(\Gamma^\pm(R, U, J, \sigma))$ and $N \geq 0$, we can find*

$$b_h^\pm(t) = \sum_{j=0}^N h^j b_j^\pm(t)$$

with $(b_j^\pm(t))_{0 \leq \pm t \leq R_1 t_{\text{WKB}}}$ bounded in $S_{\text{sc}}(\Gamma^\pm(R_0, U_0, J_0, \sigma_0))$ such that, for all $h \in (0, 1]$ and $0 \leq \pm t \leq R_1 t_{\text{WKB}}$,

$$\|e^{-ith\hat{P}} \text{Op}_\kappa(a^\pm) - \kappa^* I_{\text{WKB}}^\pm(b_h^\pm(t)) \kappa_*\|_{\mathcal{L}(L^2(\widehat{M}))} \leq C_N h^{N+1} |t|$$

where $C_N > 0$ can be taken uniformly with respect to h , t and R_1 .

Remark 7.4. The essential point of Theorem 7.3 is to construct the parametrix on the time interval $0 \leq |t| \leq R_1 t_{\text{WKB}}$ which allow us to choose the constant $\delta > 0$ in Theorem 3.3 independently with respect to R_1 . When $|t| > 0$ is small and independent of R_1 , such a parametrix construction is basically well known (see [18] for the case of elliptic operators on the Euclidean space, and [3] for the case of the Laplace-Beltrami operator on an asymptotically hyperbolic manifold).

We prove the theorem for the case when $t \geq 0$. Put $B_+(t) = I_{\text{WKB}}^+(b_h^+(t))$. By the Duhamel formula, we have

$$e^{-ith\hat{P}} \kappa^* B_+(0) \kappa_* = \kappa^* B_+(t) \kappa_* + \frac{i}{h} \int_0^t e^{-i(t-s)h\hat{P}} \kappa^* (hD_s + h^2 \hat{P}_\kappa) B_+(s) \kappa_* ds,$$

where $D_s = i^{-1} \partial_s$. Since the off-diagonal decay of the h -PDO implies

$$\|\text{Op}_\kappa(a^+) - \kappa^* a^+(r, \theta, hD_r, hD_\theta) \kappa_*\|_{\mathcal{L}(L^2(\widehat{M}))} \leq C_n h^n,$$

for all $n \geq 0$, it suffices to show that there exists $b_h^+(t)$ such that $b_h^+(0) = a^+$ and

$$(7.6) \quad \|(hD_s + h^2 \hat{P}_\kappa) B_+(s)\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \leq C_N h^{N+2}$$

for $h \in (0, 1]$ and $0 \leq s \leq R_1 t_{\text{WKB}}$.

Define smooth tensors $x^+(t)$ and $y^+(t)$ by

$$x^+(t) := \partial_{\rho, \omega} p_\kappa(r, \theta, \partial_r \Psi^+(t), \partial_\theta \Psi^+(t)), \quad y^+(t) := (p + p_1)(r, \theta, \partial_r, \partial_\theta) \Psi^+(t).$$

Then, $b_h^+(t)$ satisfying (7.6) can be constructed by solving the following transport equations.

Lemma 7.5. *For sufficiently small $t_0 > 0$, there exist*

$$(b_j^+(t))_{0 \leq t \leq R_1 t_0} \text{ bounded in } S_{\text{sc}}(\Gamma^\pm(R_0, U_0, J_0, \sigma_0)), \quad j = 0, 1, \dots, N,$$

such that $b_j^+(t)$ solve the transport equations:

$$(7.7) \quad \begin{cases} \partial_t b_0^+(t) + x^+(t) \cdot \partial_{r, \theta} b_0^+(t) + y^+(t) b_0^+(t) = 0, \\ \partial_t b_j^+(t) + x^+(t) \cdot \partial_{r, \theta} b_j^+(t) + y^+(t) b_j^+(t) + i \hat{P}_\kappa b_{j-1}^+(t) = 0, \quad j \geq 1, \end{cases}$$

with the initial condition $b_0^+(0) = a^+$, $b_j^+(0) = 0$ for $j = 1, 2, \dots, N$.

Proof. We mimic Bouclet's argument [3, Lemma 6.4]. Choose $R'_0, R''_0 > 0$, $U'_0, U''_0 \in \mathbb{R}^{d-1}$ and $J'_0, J''_0 \in (0, \infty)$ so that

$$\begin{aligned} R_0 &> R'_0 > R''_0 > R_1, & U_0 &\Subset U'_0 \Subset U''_0 \Subset U_1, \\ J_0 &\Subset J'_0 \Subset J''_0 \Subset J_1, & \sigma_0 &< \sigma'_0 < \sigma''_0 < \sigma_1. \end{aligned}$$

For $0 \leq s, t \leq R_1 t_0$ and $(r, \theta, \rho, \omega) \in \Gamma^+(R_0'', U_0'', J_0'', \sigma_0'')$, we consider the flow generated by $x^+(t)$, that is the solution to the ODE:

$$\begin{cases} (\partial_t r^+(t, s), \partial_t \theta^+(t, s)) = x^+(t, r^+(t, s), \theta^+(t, s), \rho, \omega), \\ (r^+(s, s), \theta^+(s, s)) = (r, \theta). \end{cases}$$

Since $x^+(t) = (\partial_r \Psi^+(t), r^{-2}(h^{jk} + a^{jk})\partial_{\theta^k} \Psi^+(t))$, by using (7.2), we have

$$|\partial_t r^+(t, s)| \leq C(1 + r^+(t, s)^{-1}|t|), \quad |\partial_t \theta^+(t, s)| \leq C r^+(t, s)^{-2}|t|.$$

In particular,

$$|\partial_t r^+(s, s)| \leq C, \quad |\partial_t \theta^+(s, s)| \leq C R_0''^{-1} t_0.$$

Therefore, a same argument as that in the proof of Lemma 5.13 implies that there exists $t_0 > 0$, independent of R_0 , such that $(r^+(t, s), \theta^+(t, s))$ is well-defined on $\Gamma^+(R_0'', U_0'', J_0'', \sigma_0'')$ for all $0 \leq s, t \leq R_1 t_0$, and that

$$|r^+(t, s) - r| \leq C|t|, \quad |\theta^+(t, s) - \theta| \leq t_0 \quad \text{on} \quad \Gamma^+(R_0'', U_0'', J_0'', \sigma_0'').$$

In particular, $(r^+(t, s), \theta^+(t, s), \rho, \omega) \in \Gamma^+(R_1, U_1, J_1, \sigma_1)$ for all $0 \leq s, t \leq R_1 t_0$. Moreover, differentiating the integral equation

$$(r^+(t, s), \theta^+(t, s)) = (r, \theta) + \int_s^t x^+(u, r^+(u, s), \theta^+(u, s), \rho, \omega) du$$

with respect to $\partial^\gamma = \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta$, by using (7.2) and an induction on $|\gamma|$, we have

$$(7.8) \quad |\partial^\gamma (r^+(t, s) - r)| \leq C_\gamma r^{-j-\beta} |t|, \quad |\partial^\gamma (\theta^+(t, s) - \theta)| \leq C_\gamma r^{-j-\beta} t_0,$$

for all $0 \leq s, t \leq R_1 t_0$ and $(r, \theta, \rho, \omega) \in \Gamma^+(R_0'', U_0'', J_0'', \sigma_0'')$. We put

$$\Phi_+(t, s) := (r^+(t, s), \theta^+(t, s), \rho, \omega),$$

and define $b_j^+(t)$ by

$$\begin{cases} b_0^+(t) := a^+(\Phi_+(0, t)) e^{\int_0^t y^+(s, \Phi_+(s, t)) ds}, \\ b_j^+(t) := - \int_0^t (i \hat{P}_\kappa b_{j-1}^+)(s, \Phi_+(s, t)) e^{\int_s^t y^+(u, \Phi_+(u, t)) du} ds, \quad j = 1, \dots, N. \end{cases}$$

We remark that if we choose t_0 small enough, then (7.8) implies

$$\begin{aligned} \Phi_+(t, s)(\Gamma^+(R, U, J, \sigma)) &\subset \Gamma^+(R_0, U_0, J_0, \sigma_0), \\ \Phi_+(t, s)(\Gamma^+(R_0, U_0, J_0, \sigma_0)) &\subset \Gamma^+(R'_0, U'_0, J'_0, \sigma'_0), \\ \Phi_+(t, s)(\Gamma^+(R'_0, U'_0, J'_0, \sigma'_0)) &\subset \Gamma^+(R_0'', U_0'', J_0'', \sigma_0'') \subset \Gamma^+(R_1, U_1, J_1, \sigma_1), \end{aligned}$$

for all $0 \leq s, t \leq R_1 t_0$. $\Phi_+(s, t)$ is thus well-defined on $\Gamma^+(R'_0, U'_0, J'_0, \sigma'_0)$, smooth on $\Gamma^+(R_0, U_0, J_0, \sigma_0)$, and satisfies $\Phi_+(s, t) = \Phi_+(t, s)^{-1}$. Moreover, $\text{supp } b_j^+(t) \subset \Gamma^+(R_0, U_0, J_0, \sigma_0)$ since $\text{supp } a^+ \subset \Gamma^+(R, U, J, \sigma)$. If we extend $b_j^+(t)$ on $\mathbb{R}^{2d}$ so that $b_j^+(t) = 0$ outside $\Gamma^+(R_0, U_0, J_0, \sigma_0)$, then $b_j^+(t)$ are still smooth with respect to $(r, \theta, \rho, \omega)$. Furthermore, by (2.3), (7.2) and (7.8), we see that $(b_j^+(t))_{0 \leq t \leq R_1 t_0}$ is bounded in $S_{\text{sc}}(\Gamma^\pm(R_0, U_0, J_0, \sigma_0))$. Finally, a standard Hamilton-Jacobi theory shows that $b_j^+(t)$ solve (7.7) for $0 \leq t \leq R_1 t_0$. $\square$

Proof of Theorem 7.1. By the construction, $B_+(0) = a^+(r, \theta, hD_r, hD_\theta)$. Ψ^+ solves (7.1) on $\Gamma^+(R_1, U_1, J_1, \sigma_1)$ which contains $\Gamma^+(R_0, U_0, J_0, \sigma_0)$, and b_j^+ satisfy (7.7) on the latter region. Therefore, a direct computation yields

$$(hD_s + h^2 \hat{P}_\kappa) B_+(s) = h^{N+2} i \hat{P}_\kappa b_N^+(s).$$

Since $(\hat{P}_\kappa b_N^+(s))_{0 \leq s \leq R_1 t_0}$ is bounded in $S_{\text{sc}}(\Gamma^\pm(R_0, U_0, J_0, \sigma_0))$, (7.1) implies that $I_{\text{WKB}}^+(i \hat{P}_\kappa b_j^+(s))$ is bounded on $L^2(\mathbb{R}^d)$ uniformly with respect to $h \in (0, 1]$ and $0 \leq s \leq R_1 t_0$. We hence proved (7.6). $\square$

7.3. Dispersive estimates. We here prove dispersive estimates for the WKB parametrix. Let $R_0 > R_1 > R_{\text{WKB}}$ and $t_{\text{WKB}} > 0$ be as in Theorem 7.3.

Theorem 7.6. *For any $(b^\pm(t))_{0 \leq \pm t \leq R_1 t_{\text{WKB}}}$ bounded in $S_{\text{sc}}(\Omega_s^\pm(R_0, U_0, J_0, \sigma_0))$, we can write*

$$I_{\text{WKB}}^\pm(b^\pm(t)) = U_{\text{WKB}}^\pm(t, h) + R_{\text{WKB}}^\pm(t, h),$$

where $U^\pm(t, h)$ satisfy

$$(7.9) \quad \|r^{-\frac{d-1}{2}} U_{\text{WKB}}^\pm(t, h) r^{-\frac{d-1}{2}}\|_{\mathcal{L}(L^1(\mathbb{R}^d), L^\infty(\mathbb{R}^d))} \leq C|th|^{-d/2},$$

for all $0 < \pm t \leq \min(R_1 t_{\text{WKB}}, h^{-1})$, $h \in (0, 1]$. Moreover the remainder terms $R_{\text{WKB}}^\pm(t, h)$ are rapidly decaying with respect to h : for any $N \geq 0$,

$$(7.10) \quad \|R_{\text{WKB}}^\pm(t, h)\|_{\mathcal{L}(L^2(\mathbb{R}^d))} \leq C_N h^N, \quad 0 \leq \pm t \leq \min(R_1 t_{\text{WKB}}, h^{-1}), \quad h \in (0, 1].$$

Here constants $C, C_N > 0$ may be taken uniformly with respect to h and R_1 .

Proof. Since the proof is similar as that of Theorem 5.14, we omit details and give the sketch of the proof. We consider the outgoing case only. The distribution kernel of $I_{\text{WKB}}^+(b^+(t))$ is written in the form

$$I_{b^+}(t, h) = (2\pi h)^{-d} \int e^{\frac{i}{h}\psi^+(t, r, \theta, \rho, \omega, r', \theta')} b^+(t, r, \theta, \rho, \omega) d\rho d\omega,$$

where $\psi^+(t, r, \theta, \rho, \omega, r', \theta') = \Psi^+(t, r, \theta, \rho, \omega) - r'\rho - \theta' \cdot \omega$. Let $\chi_\rho \in C_0^\infty(\mathbb{R})$, $\chi_\omega \in C_0^\infty(\mathbb{R}^{d-1})$ be as in the proof of Theorem 5.14, and set

$$B^+(t) = \chi_\rho(\partial_\rho \psi^+) \chi_\omega(\partial_\omega \psi^+) b^+(t).$$

We then have $|\partial_\rho \psi^+| < 1$ and $|\partial_\omega \psi^+| < 1$ on $\text{supp } B^+$. We denote the operator having the kernel $I_{B^+}(t, h)$ by $U_{B^+}(t, h)$. By a same proof as that in Lemma 5.15, we obtain that $R_{\text{WKB}}^+(t, h) = I_{\text{WKB}}^+(b^+(t)) - U_{B^+}(t, h)$ satisfies (7.10).

We next prove (7.9) for $U_{\text{WKB}}^+(t, h) := U_{B^+}(t, h)$. Assume $R_1 > 4$ without loss of generality. We first note that $r' \geq R_1/2$ on $\text{supp } B^+$; otherwise, there exists $C_0 > 0$, independent of R_0 , such that

$$|\partial_\rho \psi^+| \geq r - r' - C_0 R_1 t_{\text{WKB}} \geq (1/2 - C_0 t_{\text{WKB}}) R_0 > 1$$

if $0 < t_{\text{WKB}} < (4C_0)^{-1}$, since $r > R_0 > R_1$ on $\text{supp } b^+(t)$ and $\partial_\rho \psi = r - r' + O(|t|)$ by (7.2). This contradicts the support property of B^+ . We thus have

$$r'/4 \leq r' - |\partial_\rho \psi^+| - C_0 R_1 t_{\text{WKB}} \leq r \leq r' + |\partial_\rho \psi^+| + C_0 R_1 t_{\text{WKB}} \leq 4r'$$

on the support of B^+ . It follows from (7.3) that ψ^+ may be written in the form

$$\psi^+(t, r, \theta, \rho, \nu, r', \theta') = (r - r')\rho + (\theta - \theta') \cdot \omega - tp(r, \theta, \rho, \omega) + Q^+(t, r, \theta, \rho, \omega)$$

on $\text{supp } B^+$, where the remainder Q^+ satisfies

$$\left| \partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta Q^+(t, r, \theta, \rho, \omega) \right| \leq C r^{-j-|\beta|} |t| t_{\text{WKB}} \quad \text{on } \text{supp } B^+.$$

Let $\chi \in C^\infty(\mathbb{R}^{2d})$ be such that $0 \leq \chi \leq 1$, $\text{supp } \chi \subset \Gamma^+(R_1, U_1, J_1, \sigma_1)$, $\chi \equiv 1$ on $\Gamma^+(R_0, U_0, J_0, \sigma_0)$, and that $\partial_r^j \partial_\theta^\alpha \partial_\rho^k \partial_\omega^\beta \chi = O(\langle r \rangle^{-j-|\beta|})$. We then define

$$\begin{aligned} \psi_0^+(t, r, \theta, \rho, \omega, r', \theta') \\ := (r - r')\rho + (\theta - \theta') \cdot \omega - tp(r, \theta, \rho, \omega) + \chi(r, \theta, \rho, \omega) Q^+(t, r, \theta, \rho, \omega). \end{aligned}$$

Since $\psi_0^+ \equiv \psi^+$ on $[0, R_1 t_{\text{WKB}}] \times \text{supp } B^+$, using the change of the variable $\omega \mapsto r\nu$, $I_{B^+}(t, h)$ reads

$$I_{B^+}(t, h) = \frac{r^{d-1}}{(2\pi h)^d} \int e^{\frac{i}{h}\psi_0^+(t, r, \theta, \rho, r\nu, r', \theta')} \tilde{B}^+(r, \theta, \rho, \nu, r', \theta') d\rho d\nu,$$

where $\tilde{B}^+(r, \theta, \rho, \nu, r', \theta') := B^+(r, \theta, \rho, r\nu, r', \theta')$ is compactly supported with respect to (ρ, ν) , and all derivatives of $\tilde{B}^+$ are bounded on $\mathbb{R}^{3d}$. Since (7.9) is obvious for $0 < t < h$ (note that $r'/4 < r < 4r'$), we may assume that $h < t \leq R_1 t_{\text{WKB}}$. Set $\tilde{\psi}_0^+(t, r, \theta, \rho, \nu, r', \theta') :=$

$t^{-1}\psi_0^+(t, r, \theta, \rho, r\nu, r', \theta')$. Choose $\chi_1 \in C_0^\infty(\mathbb{R})$ so that $\chi_1 \equiv 1$ on $(-1/2, 1/2)$ and $\text{supp } \chi_1 \subset (-1, 1)$, and set

$$\tilde{B}_1 = \chi_1 \left(\frac{r - r'}{C_1 t} \right) \tilde{B}^+, \quad \tilde{B}_2 = \tilde{B}^+ - \tilde{B}_1,$$

with some large $C_1 > 0$. Since $\partial_\rho \tilde{\psi}_0^+ = (r - r')/t - \rho - O(t_{\text{WKB}})$, if $C_1 > 0$ is large enough, then we have $|\partial_\rho \tilde{\psi}_0^+| \gtrsim 1$ on $\text{supp } \tilde{B}_2$. An integration by parts then implies that

$$\left| \frac{1}{(2\pi h)^d} \int e^{\frac{i}{h} t \tilde{\psi}_0^+(t, r, \theta, \rho, \nu, r', \theta')} \tilde{B}_2^+(r, \theta, \rho, \nu, r', \theta') d\rho d\nu \right| \leq C_{t_{\text{WKB}}} |th|^{-d/2}$$

for $h \leq t \leq R_1 t_{\text{WKB}}$, where $C_{t_{\text{WKB}}}$ is independent of R_1 . Since

$$\partial_{\rho, \nu}^2 p(r, \theta, \rho, r\nu) = \begin{pmatrix} 1 & 0 \\ 0 & h^{jk}(\theta) + a^{jk}(r, \theta) \end{pmatrix}$$

is bounded from above and below on $\mathbb{R}^d$, we obtain

$$|\partial_\rho^k \partial_\nu^\beta \tilde{\psi}_0^+(t, r, \theta, \rho, \nu, r', \theta')| \leq C \quad \text{for} \quad |k + \beta| \geq 2,$$

uniformly with respect to $(r, \theta, \rho, \omega, r', \theta') \in \mathbb{R}^{3d}$ and $0 < t \leq R_1 t_{\text{WKB}}$. Moreover, if $t_{\text{WKB}} > 0$, which can be taken uniformly with respect to R_1 , is small enough, then we have

$$|\det \partial_{\rho, \nu}^2 \tilde{\psi}_0^+(t, r, \theta, \rho, \nu, r', \theta')| \gtrsim 1, \quad (r, \theta, \rho, \omega, r', \theta') \in \mathbb{R}^{3d}, \quad 0 < t \leq R_1 t_{\text{WKB}}.$$

By a same argument as that in Section 5, we can apply the stationary phase theorem, and have the assertion since $r'/4 < r < 4r'$ on the support of $\tilde{B}^+$. $\square$

8. PROOF OF THEOREM 3.3

We here prove Theorem 3.3. We only consider the case $t \geq 0$, and the proof for the case $t \leq 0$ is similar. Recall that $A_h = \sum_\kappa \text{Op}_\kappa^{pr}(a_{\kappa, h})$ is a sum of properly supported h -PDO's. By (2.9), (2.11) and their adjoint estimates with respect to $\widehat{G}dz$, we have

$$(8.1) \quad \begin{cases} \|r^{-\frac{d-1}{2}} A_h\|_{\mathcal{L}(L^2(\widehat{M}), L^\infty(\widehat{M}))} + \|A_h^* r^{-\frac{d-1}{2}}\|_{\mathcal{L}(L^1(\widehat{M}), L^2(\widehat{M}))} \leq Ch^{-\frac{d}{2}}, \\ \|r^{-\frac{d-1}{2}} A_h r^{\frac{d-1}{2}}\|_{\mathcal{L}(L^\infty(\widehat{M}))} + \|r^{\frac{d-1}{2}} A_h^* r^{-\frac{d-1}{2}}\|_{\mathcal{L}(L^1(\widehat{M}))} \leq C, \end{cases}$$

uniformly with respect to $h \in (0, 1]$.

Choose $R_0 > 0$ and $\varepsilon_0 > 0$ so that Theorems 5.10, 5.14, 6.1, 7.3 and 7.6 hold for all $R \geq R_0$ and $0 < \varepsilon \leq \varepsilon_0$, and let $N \geq d$.

We first prove (3.5). A standard symbolic calculus implies that $\text{Op}_\kappa(b_s^+)^*$ can be replaced by $\text{Op}_\kappa(\tilde{b}_s^+)$ modulo a smoothing term $O(h^{N+1})$ on $L^2(\widehat{M})$, where $\tilde{b}_s^+ \in S_{\text{sc}}(\Omega_s^+(R_2, \tilde{U}_\kappa, J, \varepsilon))$. Moreover, $\text{Op}_\kappa(a_s^+) - \text{Op}_\kappa^{pr}(a_s^+)$ has the $\mathcal{L}(L^2(\widehat{M}))$ -norm of order h^{N+1} by (2.7). Therefore, Theorems 5.10, 5.14 with $R = R_2$ and (2.11) with $(q, s) = (\infty, (d-1)/2)$ imply that $\text{Op}_\kappa(a_s^+)e^{-ith\widehat{P}} \text{Op}_\kappa(b_s^+)^*$ can be brought to the form

$$U_s^+(t, h, N) + h^{N+1} R_s^+(t, h, N),$$

where $U_s^+(t, h, N)$ and $R_s^+(t, h, N)$ satisfy

$$\begin{aligned} \|r^{-\frac{d-1}{2}} U_s^+(t, h, N) r^{-\frac{d-1}{2}}\|_{\mathcal{L}(L^1(\widehat{M}), L^\infty(\widehat{M}))} &\leq C_N |th|^{-\frac{d}{2}}, \\ \|R_s^+(t, h, N)\|_{\mathcal{L}(L^2(\widehat{M}))} &\leq C_N, \end{aligned}$$

uniformly with respect to $h \in (0, 1]$ and $0 < \pm t \leq h^{-1}$. By using (8.1), we obtain (3.5) since $1 \leq |th|^{-\frac{d}{2}}$ for $0 < t \leq h^{-1}$.

Next, let $R_2 \geq R_0$ and $R_1 > 0$ so that $2R_1 > R_2 > R_1$. Then, there exists $t_{\text{WKB}} > 0$, independent of R_2 , such that Theorems 7.3 and 7.6 with $R = R_2$ hold for $0 < t \leq R_2 t_{\text{WKB}}/2$. By a same argument as above, we obtain (3.4) with $t_0 = t_{\text{WKB}}/2$.

For any $t_1 > 0$, a same argument as above and Theorem 6.1 imply

$$\|\text{Op}_\kappa(a_l^+)e^{-ith\widehat{P}} \text{Op}_\kappa(b_l^+)^*\|_{\mathcal{L}(L^2(\widehat{M}))} \leq C_N h^{N+1},$$

for $h \in (0, 1]$ and $R_2 t_1 \leq t \leq h^{-1}$. Combining this estimate with (8.1), we obtain (3.6). We complete the proof. $\square$

APPENDIX A. PROOF OF THEOREM 1.2

We here give the sketch of the proof of Theorem 1.2. It is sufficient to prove that, for any $K \Subset M$, $\chi_K \in C_0^\infty(M)$ with $\chi_K \equiv 1$ on K and admissible pair (p, q) , the following estimate holds under the nontrapping condition:

$$(A.1) \quad \|\chi_K e^{-itP} u_0\|_{L^p([0,1]; L^q(M))} \leq C \|u_0\|_{L^2(M)}.$$

We mimic the Bouclet-Tzvetkov argument [4, Section 5 and 6].

Proposition A.1. *Let $\varphi \in C_0^\infty((0, \infty))$. Then there exist $t_0 > 0$ and $C > 0$ such that for all $h \in (0, 1]$ and admissible pair (p, q) ,*

$$(A.2) \quad \|\chi_K \varphi(h^2 P) e^{-itP} u_0\|_{L^p([0,t_0 h]; L^q(M))} \leq C \|u_0\|_{L^2(M)}.$$

Proof. Let $\{\kappa : V_\kappa \rightarrow U_\kappa\}_\kappa$ be a finite atlas on K and $\{\psi_\kappa\}_\kappa \subset C_0^\infty(M)$ be a partition of unity subordinate to $\{V_\kappa\}$. Let $\tilde{\psi}_\kappa \in C_0^\infty(M)$ so that $\tilde{\psi}_\kappa \equiv 1$ on a neighborhood of $\text{supp } \psi$. For a symbol $a \in C_0^\infty(U_\kappa \times \mathbb{R}^d)$, we define $\text{Op}_\kappa(a)u = \kappa^*(a(z, hD_z) \kappa_*(\tilde{\psi}_\kappa u))$. Since a has a compact support with respect z and ξ , the Schur lemma implies that for $1 \leq q \leq r \leq \infty$ and $h \in (0, 1]$,

$$(A.3) \quad \|a(z, hD_z)\|_{\mathcal{L}(L^q(\mathbb{R}^d), L^r(\mathbb{R}^d))} \leq Ch^{-d(1/q-1/r)}.$$

By a same argument as that in [6, Section 2] which studied the case of compact manifolds without boundaries (see also [1]), there exist symbols $a_{\kappa,h} \in C_0^\infty(U_\kappa \times p_\kappa^{-1}(\text{supp } \varphi))$ such that we can approximate $\chi_K \varphi(h^2 P)$ by $\sum_\kappa \text{Op}_\kappa(a_{\kappa,h})$ up to $O(h^N)$ on $L^2(M)$ for any $N \geq 0$. In particular if we obtain the following estimate:

$$(A.4) \quad \|\text{Op}_\kappa(b) \text{Op}_\kappa(a) e^{-itP} u_0\|_{L^p([0,t_0 h]; L^q(M))} \leq C \|u_0\|_{L^2(\widehat{M})}, \quad h \in (0, 1],$$

for any $a, b \in C_0^\infty(U_\kappa \times p_\kappa^{-1}(\text{supp } \varphi))$, then a same argument as that in Section 3 implies (A.2). (A.4) follows from (A.3), the TT^* -argument and the WKB parametric approximation of the propagator $e^{-itP} \text{Op}_\kappa(a)$ for $0 \leq t \leq t_0$ with sufficiently small $t_0 > 0$. The construction of the WKB parametrix is basically well known and its proof is similar as that in the case of elliptic operators on the Euclidean space. We refer to [18] for details. Dispersive estimates can be proved similarly to the case of compact manifolds without boundaries [6]. $\square$

Proof of (A.1). The Duhamel formula implies

$$\varphi(h^2 P) \chi_K e^{-itP} u_0 = e^{-itP} \varphi(h^2 P) \chi_K u_0 - i \int_0^t e^{-i(t-s)P} \varphi(h^2 P) [P, \chi_K] e^{-isP} u_0 ds.$$

By Theorem 1.1, Proposition A.1 and a same argument as that in [4, Section 5 and 6], we obtain

$$\begin{aligned} & \|\varphi(h^2 P) \chi_K e^{-itP} u_0\|_{L^p([0,1]; L^q(M))} \\ & \lesssim \|\varphi(h^2 P) \chi_K u_0\|_{L^2(M)} + h^{-1/2} \|\varphi(h^2 P) \chi_K e^{-itP} u_0\|_{L^2([0,1]; L^2(M))} \\ & \quad + h^{1/2} \|\varphi(h^2 P) [P, \chi_K] e^{-isP} u_0\|_{L^2([0,1]; L^2(M))} \\ & \lesssim \|\varphi(h^2 P) u_0\|_{L^2(M)} + h^{1/2} \|u_0\|_{L^2(M)} + \|\chi_K \varphi(h^2 P) e^{-itP} u_0\|_{L^2([0,1]; H^{1/2}(M))}, \end{aligned}$$

where $H^{1/2}(M)$ is the Sobolev space with the norm $\|(1+P)^{1/4} \cdot\|_{L^2(M)}$. In the last line we used inequalities

$$\begin{aligned} & \|[\varphi(h^2 P), \chi_K]\|_{\mathcal{L}(L^2(M))} \lesssim h, \quad \|[\varphi(h^2 P), [P, \chi_K]]\|_{\mathcal{L}(L^2(M))} \lesssim 1, \\ & \|[P, \chi_K] \tilde{\chi}_K \varphi(h^2 P)\|_{\mathcal{L}(H^{1/2}(M), H^{-1/2}(M))} \lesssim 1, \end{aligned}$$

where $\tilde{\chi}_K \in C_0^\infty(M)$ is a cut-off function so that $\tilde{\chi}_K \equiv 1$ on $\text{supp } \chi_K$. These inequalities follow from the pseudodifferential approximations of $\chi_K \varphi(h^2 P)$ and $(1 - \chi_K) \varphi(h^2 P)$ such as Lemma 2.4 (see also [1]) and the symbolic calculus.

We now use the nontrapping condition on the metric. Under the nontrapping condition, Cardoso-Vodev [7] proved that for sufficiently large $\lambda > 0$,

$$\|\chi_K(P - \lambda \pm i0)^{-1}\chi_K\|_{\mathcal{L}(L^2(M))} \lesssim \langle \lambda \rangle^{-1/2}.$$

By the abstract Kato smooth perturbation theory, this resolvent estimate implies the local smoothing effect:

$$\|\chi_K e^{-itP} u_0\|_{L^2([0,1]; H^{1/2}(M))} \lesssim \|u_0\|_{L^2(M)}.$$

Using this estimate, we obtain

$$\|\varphi(h^2 P)\chi_K e^{-itP} u_0\|_{L^p([0,1]; L^q(M))} \lesssim \|\varphi(h^2 P)u_0\|_{L^2(M)} + h^{1/2}\|u_0\|_{L^2(M)}$$

uniformly with respect to $h \in (0, 1]$. Applying the Littlewood-Paley decomposition proved by [2], we obtain (A.1) and conclude the proof. $\square$

REFERENCES

- [1] Bouclet, J.- M. Semi-classical calculus on non compact manifolds with ends and weighted L^p estimates. To appear in *Ann. Inst. Fourier*.
- [2] Bouclet, J.- M. (2010). Littlewood-Paley decompositions on manifolds with ends. *Bull. Soc. Math. France*. 138:1–37
- [3] Bouclet, J.- M. Strichartz estimates on asymptotically hyperbolic manifolds. To appear in *Analysis and PDE*.
- [4] Bouclet, J.- M., Tzvetkov, N. (2007). Strichartz estimates for long range perturbations. *Amer. J. Math.* 129:1565–1609.
- [5] Bouclet, J.- M., Tzvetkov, N. (2008). On global Strichartz estimates for non trapping metrics. *J. Funct. Analysis*. 254:1661–1682.
- [6] Burq, N., Gérard, P., Tzvetkov, N. (2004). Strichartz inequalities and the nonlinear Schrödinger equation on compact manifolds. *Amer. J. Math.* 126:569–605.
- [7] Cardoso, F., Vodev, G. (2002). Uniform estimates of the resolvent of the Laplace-Beltrami operator on infinite volume Riemannian manifolds, II. *Ann. Henri Poincaré*. 3:673–691.
- [8] Doi, S. (1996). Smoothing effects of Schrödinger evolution groups on Riemannian manifolds. *Duke Math. J.* 82:679–706.
- [9] Fujiwara, D. (1980). Remarks on convergence of the Feynman path integrals. *Duke Math. J.* 47:559–600.
- [10] Ginibre, J., Velo, G. (1985). The global Cauchy problem for the non linear Schrödinger equation. *Ann. IHP-Analyse non linéaire*. 2:309–327.
- [11] Hassell, A., Tao, T., Wunsch, J. (2006). Sharp Strichartz estimates on non-trapping asymptotically conic manifolds. *Amer. J. Math.* 128:963–1024.
- [12] Isozaki, H., Kitada, H. (1985). Modified wave operators with time independent modifiers. *J. Fac. Sci. Univ. Tokyo*. 32:77–104.
- [13] Journé, J.- L., Soffer, A., Sogge, C. D. (1991). Decay estimates for Schrödinger operators. *Comm. Pure Appl. Math.* 44:573–604.
- [14] Keel, M., Tao, T. (1998). Endpoint Strichartz Estimates. *Amer. J. Math.* 120:955–980.
- [15] Kumano-go, H. (1980). *Pseudo-Differential Operators*. MIT Press.
- [16] Marzuola, J., Metcalfe, J., Tataru, D. (2008). Strichartz estimates and local smoothing estimates for asymptotically flat Schrödinger equations. *J. Funct. Analysis*. 255:1497–1553.
- [17] Robbiano, L., Zuily, C. (2005). Strichartz estimates for Schrödinger equations with variable coefficients. *Mém. SMF. Math. Fr. (N.S.)*. 101–102.
- [18] Robert, D. (1987). *Autour de l'approximation semi-classique*. Progr. Math. 68 Birkhäuser, Basel.
- [19] Schlag, W. (2007). Dispersive estimates for Schrödinger operators: a survey. *Mathematical aspects of nonlinear dispersive equations*, *Ann. of Math. Stud. Princeton Univ. Press, Princeton, NJ*. 163:255–285
- [20] Staffilani, G., Tataru, D. (2002). Strichartz estimates for a Schrödinger operator with non smooth coefficients. *Comm. PDE*. 27:1337–1372.
- [21] Strichartz, R. (1977). Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations. *Duke Math. J.* 44:705–714.
- [22] Tataru, D. (2008). Parametrix and dispersive estimates for Schrödinger operators with variable coefficients. *Amer. J. Math.* 130:571–634.
- [23] Yajima, K. (1987). Existence of solutions for Schrödinger evolution equations. *Comm Math. Phys.* 110:415–426.
- [24] Yajima, K. (1995). The $W_{k,p}$ -continuity of wave operators for Schrödinger operators. *J. Math. Soc. Japan*. 47:551–581.

- [25] Yajima, K. (2005). Dispersive estimates for Schrödinger equations with threshold resonance and eigenvalue.
Comm. Math. Phys. 259:475–509.