TITLE:

Grazing impact on the cyanobacterium *Microcystis aeruginosa* by the heterotrophic flagellate *Collodictyon triciliatum* in an experimental pond

AUTHOR(S):

Kobayashi, Yuki; Hodoki, Yoshikuni; Ohbayashi, Kako; Okuda, Noboru; Nakano, Shin-ichi

CITATION:

Kobayashi, Yuki ...[et al]. Grazing impact on the cyanobacterium *Microcystis aeruginosa* by the heterotrophic flagellate *Collodictyon triciliatum* in an experimental pond. *Limnology* 2013, 14(1): 43-49

ISSUE DATE:

2013-01

URL:

<http://hdl.handle.net/2433/169749>

RIGHT:

The final publication is available at www.springerlink.com; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

1 **Title**

2 **Grazing impact on the cyanobacterium *Microcystis aeruginosa* by the**
3 **heterotrophic flagellate *Collodictyon triciliatum* in an experimental pond**

4 Yuki Kobayashi, Yoshikuni Hodoki, Kako Ohbayashi, Noboru Okuda, Shin-ichi
5 Nakano*

6

7 Center for Ecological Research, Kyoto University, 2-509-3 Hirano, Otsu, Shiga
8 520-2113, Japan

9

10

11 *Correspondence:

12 Center for Ecological Research, Kyoto University, 2-509-3 Hirano, Otsu, Shiga
13 520-2113, Japan

14 Tel: +81-77-549-8239,

15 Fax: +81-77-549-8201,

16 *Email: nakano@ecology.kyoto-u.ac.jp

17

18

1 Abstract

2 We estimated the grazing impact of the heterotrophic flagellate *Collodictyon*
3 *triciliatum* on the harmful, bloom-forming cyanobacterium *Microcystis aeruginosa* in
4 an experimental pond during *Microcystis* bloom from summer to winter in 2010. For
5 these experiments, we calculated the grazing rates from the *C. triciliatum* digestion rate
6 and food vacuole content. During the study period, *M. aeruginosa* exhibited one
7 blooming event with a maximum density of 1.1×10^5 cells ml^{-1} . The cell density of *C.*
8 *triciliatum* fluctuated from below the detection limit to 291 cells ml^{-1} . The number of *M.*
9 *aeruginosa* cells ingested by *C. triciliatum* food vacuoles ranged between 0.4 and 10.8
10 cells flagellate $^{-1}$, and the digestion rate of *C. triciliatum* at 25°C was 0.73% cell contents
11 min^{-1} . The grazing rate of *C. triciliatum* on *M. aeruginosa* prey was 0.2–6.9 cells
12 flagellate $^{-1} \text{ h}^{-1}$, and its grazing impact was 0.0–25.3% standing stock day^{-1} . The
13 functional response of *C. triciliatum* to *M. aeruginosa* prey followed the
14 Michaelis-Menten model of significance ($r^2 = 0.873$, $p < 0.001$) in our experimental
15 systems in which the prey concentration varied from 1.0×10^4 to 2.1×10^6 cells ml^{-1} . The
16 maximum grazing rate was 6.2 prey cells grazer $^{-1} \text{ h}^{-1}$, and the half-saturation constant
17 was 1.2×10^5 cells ml^{-1} . We present evidence that *C. triciliatum* grazing explained the
18 remarkable decrease in *M. aeruginosa* cell density in the pond. The present study is the

1 first demonstration of the high potential of protistan grazing on *M. aeruginosa* to reduce
2 cyanobacterial blooms.

3

4 Key words: *Microcystis aeruginosa*, *Collodictyon triciliatum*, Functional response,

5 Ingestion, Grazing impact

6

7

8

1 Introduction

2 The major bloom-forming cyanobacterial species *Microcystis aeruginosa* is
3 distributed ubiquitously in eutrophic lakes worldwide. It forms toxic blooms and causes
4 serious environmental problems due to deterioration of water quality, deoxygenation of
5 underlying waters and their subsequent toxicity, foul odors, and an overall decrease in
6 aesthetic value (Carmichael 1992; Park et al. 1998). Many researchers have studied the
7 environmental factors favorable to inducing *M. aeruginosa* bloom formation (Reynolds
8 and Walsby 1975; Dokulil and Teubner 2000; Latour et al. 2004; Gobler et al. 2007),
9 but only limited information have been available for *M. aeruginosa* bloom depletion.

10 A previous study demonstrated that certain flagellate species graze on *M.*
11 *aeruginosa*, including *Monas guttula* (Sugiura et al. 1992), *Poterioochromonas*
12 *malhamensis* (Zhang et al. 1996), *Poterioochromonas* sp. (Zhang et al. 2009),
13 *Collodictyon triciliatum* (Nishibe et al. 2002), *Diphyllaea rotans* (Kim et al. 2006),
14 *Ochromonas danica* (Cole and Wynne 1974), and *Ochromonas* sp. (Burkert et al. 2001;
15 Yang et al. 2008; Baek et al. 2009; Van Donk et al. 2009; Yang et al. 2009; Wilken et al.
16 2010). These flagellate species retain active growth, even while feeding on toxic *M.*
17 *aeruginosa*. Nishibe et al. (2002) determined the specific grazing rate of the large
18 flagellate *C. triciliatum* on *M. aeruginosa* using a method proposed by Dolan and Šimek

(1998), and applied the rate to estimate the *in situ* grazing pressure on a cyanobacterium population in a eutrophic pond. One would expect a high impact because of the large cell size (30–50 μm in length) of the flagellate, but it in fact had a low impact on the cyanobacterium. Beyond that study, ecophysiological reports on *C. triciliatum* remain limited.

In the present study, we determined the specific grazing rate of *C. triciliatum* on *M. aeruginosa* based on the digestion rate of *M. aeruginosa* cells in the food vacuoles of *C. triciliatum* (Dolan and Šimek 1998). In addition, we examined the functional response of *C. triciliatum* to *M. aeruginosa*. Using these results, we estimated the grazing impact by the flagellate on an *M. aeruginosa* bloom that occurred in an outdoor experimental pond.

Materials and Methods

Monitoring biological abundance

We monitored a *Microcystis* bloom and *C. triciliatum* abundance in a meso-scale experimental pond (10 m $\times$ 10 m width; maximum depth, 1.7 m; water volume, ca. 70 m^3) located at the Center for Ecological Research, Kyoto University, Japan (34° 58' 2.24''N, 135° 57' 38.93''E). We artificially induced the bloom, which is

described in detail in Hodoki et al. (2011). Water samples were collected from the pond surface with a 5L plastic bucket, and poured into three sterilized polycarbonate bottles. Samples were taken at the same time of day (10:00 to 10:30 h) about once weekly from 1 July 2010 to 28 December 2010, and the water temperature was measured simultaneously with a bar thermometer. To measure the chlorophyll *a* (chl *a*) concentration, we filtered a measured portion of each water sample through a GF/F filter to retain seston, and stored the samples in a freezer at -20°C until needed for analysis. To process the samples, we placed the filter in a glass test tube along with 10 ml N, N-dimethylformamide (DMF) to extract the chl *a*. The quantity of chl *a* was determined using a spectrofluorophotometer (RF-5300 Shimadzu; Welschmeyer 1994).

Enumeration of microorganisms

To quantify the *C. triciliatum* cells and *M. aeruginosa* cells, we fixed 500 ml water samples with acidified Lugol's solution at a final concentration of 1%, and concentrated these samples using natural sedimentation. We counted the *C. triciliatum* cells in a Fuchs-Rosenthal type haemocytometer and *M. aeruginosa* in Burkner-Turk type haemocytometer under a microscope at a magnification of ×400 at least three times. For enumeration of *M. aeruginosa* cell, concentrated samples were slightly sonicated at

1 55kw for 5 min to disperse *Microcystis* cells from colonies.

2 To quantify *M. aeruginosa* cells contained within the *C. triciliatum* food vacuoles, we

3 used samples fixed with acidified Lugol's solution and the method described above. A

4 0.05ml aliquot of the concentrated sample was mounted on a glass slide, and at least 30

5 *C. triciliatum* cells were examined under a microscope at a magnification of $\times 400$. For

6 each sample, we averaged the numbers of *M. aeruginosa* cells ingested per flagellate.

7

8 **Digestion experiment**

9 We determined the digestion rate of *C. triciliatum* feeding on the *M.*

10 *aeruginosa* population using the methods of Dolan and Šimek (1998) and Nishibe et al.

11 (2002), with minor modifications. We isolated *C. triciliatum* from the surface water of

12 the experimental pond in August 2010, and maintained the clonal cultures in CT

13 medium (Watanabe and Ichimura 1997 or MCC-NIES;

14 www.nies.go.jp/biology/mcc/home_j.htm), using *M. aeruginosa* NIES-843 as the food

15 source for *C. triciliatum*. The microorganism cultures were maintained in 300–500 ml

16 polycarbonate bottles at 25°C under a light intensity of $52 \mu\text{Em}^{-2} \text{ s}^{-1}$. We used *M.*

17 *aeruginosa* samples that were in the exponential growth phase and *C. triciliatum*

18 samples that were in the early stationary phase for the digestion and functional response

experiments, respectively. A 6 ml sample of the *C. triciliatum* culture was inoculated into a 100 ml polycarbonate bottle containing 12 ml *M. aeruginosa* culture (4.0×10^6 cell ml^{-1}). Following an incubation period of 12 h at 25°C in the dark, the flagellate culture was diluted with 6 ml of the mixture to 600 ml fresh CT medium in triplicate in 1 L Erlenmeyer flasks in the dark (25°C) to halt ingestion. We removed 50 ml subsamples from each flask at 30 min intervals for 150 min, and fixed these subsamples with acidified Lugol's solution at a final concentration 1%. The fixed samples were concentrated via natural sedimentation, and a 0.05ml aliquot of concentrated sample was mounted on a glass slide. *M. aeruginosa* cells as well as those found in the food vacuoles (ingested cells) were counted in 100 *C. triciliatum* cells under a microscope at a magnification of $\times 400$.

We performed linear regression analyses and calculated the slope of the regression of $\ln$ (% initial average number of ingested *M. aeruginosa* per flagellate) versus time to yield the digestion rate. Multiplying the slope by 100 gave a digestion rate constant (% cell content min^{-1} ; Dolan and Šimek 1998; Nishibe et al. 2002).

As per Nishibe et al. (2002), we corrected the digestion rate at 25°C for the *in situ* temperature in the experimental pond using $Q_{10} = 2.1$. The specific grazing rate of *C. triciliatum* (G_c , *M. aeruginosa* cells flagellate $^{-1} \text{ h}^{-1}$) was calculated by multiplying the

corrected digestion rate by the average number of ingested *M. aeruginosa* cells per flagellate for each sampling day, respectively. The clearance rate of *C. triciliatum* (nl flagellate⁻¹ h⁻¹) was calculated by dividing the flagellate grazing rate by the corresponding *M. aeruginosa* cell density for each sampling day. The daily grazing impact of *C. triciliatum* on the standing stock of *M. aeruginosa* (G_i , % standing stock day⁻¹) was estimated as follows:

$$G_i = 100 \times (G_c \times N_c \times 24) / N_m,$$

where N_c (cell ml⁻¹) and N_m (cells ml⁻¹) are the cell densities of *C. triciliatum* and *M. aeruginosa* for each sampling day, respectively (Dolan and Šimek 1998; Nishibe et al. 2002).

Functional response experiment

Before conducting the functional response experiment, we diluted the *C. triciliatum* culture by adding 700 ml of the mixture to 700 ml fresh CT medium, and incubated this culture in the dark for 12 h to obtain flagellates starved for food and without residual growth, and to decrease *M. aeruginosa* abundance to a negligible level (<1.0×10³ cells ml⁻¹).

We fed the *C. triciliatum* (initial condition; 2.0×10³ cells ml⁻¹) on *M.*

1 *aeruginosa* (initial condition; 1.0×10^4 to 2.1×10^6 cells ml^{-1}) as prey, and examined the
2 resultant ingestion rate using the method of Jeong et al. (2005) with slight modifications.
3 We prepared various *M. aeruginosa* cell concentrations in the culture (initial condition;
4 1.0×10^4 to 2.1×10^6 cells ml^{-1}) in triplicate 300 ml Erlenmeyer flasks, followed by
5 inoculation of *C. triciliatum* at a cell density of 2.0×10^3 cells ml^{-1} . The cultures were
6 filled with CT medium to 100 ml, and were incubated in the dark at 25°C. At 0, 15, 30
7 and 45 min of incubation, a 10 ml subsample was removed from each culture,
8 transferred to a 15 ml tube, and then fixed with acidic Lugol's solution to a final
9 concentration of 1%. We used a Burkner-Turk haemocytometer to count *M. aeruginosa*
10 cells in the sample. A 0.05ml aliquot of sample was mounted on a glass slide and count
11 *M. aeruginosa* in the food vacuoles of *C. triciliatum*. To enumerate the *M. aeruginosa*
12 as prey, we inspected at least 30 flagellate cells, and produced a linear regression line
13 representing the relationship between the number of prey cells within a *C. triciliatum*
14 cell and the incubation time. We calculated ingestion rates (prey cells flagellate $^{-1}$ h $^{-1}$)
15 according to the method of Sherr et al. (1987), and fitted all derived ingestion rates to a
16 Michaelis-Menten equation, as follows:

$$17 \quad IR = \frac{I_{\max}(x)}{K_{IR} + (x)},$$

18 where $I_{\max}$ is the maximum ingestion rate (cells flagellate $^{-1}$ h $^{-1}$), and K_{IR} is the prey

1 concentration sustaining one-half $I_{\max}$.

2

3 **Results**

4 **Monitoring *Microcystis* bloom and *Collodictyon* abundance**

5 The water temperature of the pond remained at about 30°C from the initial
6 experiment day (0) to day 74, and decreased thereafter (Fig. 1A). Two relatively high
7 chl *a* concentrations were measured on days 28 (155.3 $\mu\text{g L}^{-1}$) and 70 (138.3 $\mu\text{g L}^{-1}$; Fig.
8 1B). That on day 70 roughly coincided with the initial *M. aeruginosa* bloom period (day
9 84, 10.8×10^4 cells ml^{-1} ; Fig. 1C). The abundance of *C. triciliatum* became relatively
10 high from days 22 to 35, and from days 105 to 119, with the highest abundance (291
11 cells ml^{-1}) detected on day 28 (Fig. 1D). The number of *M. aeruginosa* cells ingested by
12 each flagellate cell was high on day 84 (10.8 cells flagellate $^{-1}$), which fluctuated during
13 the study period (Fig. 1E).

14 The digestion experiment showed that the average number of *M. aeruginosa*
15 cells ingested by each flagellate cell declined over time (Fig. 2), and we determined a
16 significant correlation between the $\ln$ (% initial average numbers of ingested *M.*
17 *aeruginosa* per flagellate) and time ($r^2 = 0.909$, $p < 0.01$, $n = 6$). The digestion rate
18 constants of *C. triciliatum* feeding on *M. aeruginosa* was 0.73% cell contents min^{-1} ,

which was close to that determined in the previous study at 25°C ($0.74 \pm 0.02\%$ cell contents min^{-1} , Nishibe et al. 2002).

The *C. triciliatum* grazing rates and clearance rates were 0.2–6.9 *M. aeruginosa* cells flagellate $^{-1} \text{h}^{-1}$ and 2.2 to 162.2 nl flagellate $^{-1} \text{h}^{-1}$, respectively (Table 1). The daily grazing impact of *C. triciliatum* on *M. aeruginosa* was 0.0–25.3% of the *M. aeruginosa* standing stock (Table 1).

Functional response

The *C. triciliatum* population demonstrated a clear functional response to prey density (Fig. 3). The half-saturation constant was 1.2×10^5 cells ml^{-1} , and the maximum ingestion rate was 6.2 cells flagellate $^{-1} \text{h}^{-1}$.

Discussion

In this study, we demonstrated the high potential of a protistan species to reduce cyanobacterial blooms by herbivorous flagellate grazing. Our observations included temperature-independent pattern changes in *C. triciliatum* abundance (Fig. 1D; see Results section), in which *C. triciliatum* concentrations increased significantly on two occasions (from days 22 to 35, and from days 105 to 119, Fig. 1D). Hence, it is

likely that growth of *C. triciliatum* in the present study was independent of temperature.

The *M. aeruginosa* population attained its maximum concentration on day 84 (Fig. 1C), when the highest number of *Microcystis* cells was ingested by each *C. triciliatum* cell (Fig. 1E). There was an abundance of *Microcystis* prey for *C. triciliatum* predation during the period days 74 to 84, suggesting a high possibility for remarkable *C. triciliatum* growth. However, we did not record any noteworthy growth rates for the flagellate during this period (Fig. 1D). One reason for this observation is the possible presence of *C. triciliatum* predators, such as large rotifers and/or carnivorous/omnivorous copepods. However, lacking any evidence of *C. triciliatum* predation, we can only present this possibility as speculation and recommend it as a topic for future study. *C. triciliatum* attained its highest concentration on study day 28 (Fig. 1D), when we also recorded the highest chl *a* concentration (Fig. 1B). Previous studies have reported that *C. triciliatum* is omnivorous (Mischke 1994; Nishibe et al. 2002). Thus, it seems likely that *C. triciliatum* could proliferate actively, with plenty of food and in the absence of predators.

The highest number of *Microcystis* cells ingested by each *C. triciliatum* was recorded on study day 84 (Fig. 1E). The abundance of *M. aeruginosa* decreased significantly between days 84 and 91 (Fig. 1C). Using the specific ingestion rate of 84

day and average *C. triciliatum* number between 84 and 91 days (Heinbokel 1978), we estimated the *M. aeruginosa* loss to grazing by *C. triciliatum* between 84 and 91 days as 1.8×10^4 cells ml^{-1} . We calculated the decrease in *Microcystis* abundance during this period as 8.6×10^4 cells ml^{-1} , indicating that 21% of *Microcystis* abundance was consumed by *C. triciliatum* grazing between 84 and 91 days. According to our microscopic observations, the dominant food item of *C. triciliatum* during this period was *M. aeruginosa*. Thus, the remarkable decrease in *Microcystis* abundance recorded from days 84 to 91 was substantially due to *C. triciliatum* grazing. Grazing rates calculated from digestion rates in the present were roughly consistent with those in our previous study (Nishibe et al. 2002; Table 1). Nishibe et al. (2002) demonstrated that the grazing impact of the flagellate may be of minor importance to the decline in *M. aeruginosa* concentrations. However, in the present study, *C. triciliatum* grazing on *Microcystis* was, if only temporarily, effective for decreasing cyanobacterial abundance. The present study is thus the first to demonstrate the high potential of protistan grazing on *M. aeruginosa* to reduce the cyanobacterial bloom.

We simply calculated the grazing impact using the data on cell densities of *C. triciliatum* and *M. aeruginosa*, and in situ grazing rate of the flagellate. However, this calculation may lead to an overestimation of grazing impact. Nishibe et al. (2002)

reported that *C. triciliatum* might have preyed on small colonies (<50 cells) of *Microcystis* as well as unicellular ones. When *Microcystis* blooms attain to the stationary phase and come to senescence, colony sizes of the cyanobacteria become smaller (Ishikawa et al. 2004, Yamamoto & Nakahara 2009). During the period between 84 days to 91 days, water temperature decreased, and chlorophyll *a* concentration and *Microcystis* cell density also did (Fig. 1A, B and C). These results indicate the cyanobacterial senescence. So, it is likely that small colonies and/or unicellular *Microcystis* were abundant during the period in the present study. Considering those, we think our estimation of *Collodictyon* grazing impact on *Microcystis* abundance is valid, though there still is the possibility of overestimation.

The maximum grazing rates and half-saturation constants recorded in the present study (Table 2) are higher than those reported by Kim et al. (2006) and Wilken et al. (2010). However, these previous studies utilized small flagellate species (*Diphyllleia*, ca. 25 μm and *Ochromonas*, ca. 10 μm), whereas the *C. triciliatum* cells used in the present study are comparatively large (30–50 μm). Because the specific ingestion rates of flagellates increase with increasing cell size (Jeong et al. 2005), the results of the present study are comparable to those in the previous studies on a sliding scale. Guo et al. (2008) reported that the flagellate genus *Poterioochromonas* has a

higher half-saturation constant on *M. aeruginosa* prey, and the maximum ingestion rate of the flagellate was lower than that of *C. triciliatum* (Table 2). The genus *Poterioochromonas* is a mixotroph, and some mixotrophic algae are mainly phototrophic with low phagotrophic ingestion rates (Guo et al. 2008). This may explain the high half-saturation constant of *Poterioochromonas*.

It is interesting to note that the grazing rate estimated from the digestion rate herein was close to that of the high food concentration measured in the functional response experiment (Table 1 and Fig. 3), although these methods are different. The possible reason for this similarity is that the *C. triciliatum*, in our digestion experiment, were fed under a food-replete condition, likely similar to the high food concentration used in our functional response experiment. The half-saturation constant of *C. triciliatum* in the present study (1.2×10^5 cell ml^{-1}) suggests that flagellate grazing on *Microcystis* is initiated in the early stages of the bloom (Nishibe et al. 2004; Imai et al. 2009).

As determined in the present study, and similar to previous reports, *C. triciliatum* cannot directly ingest a large colony of *Microcystis* (Nishibe et al. 2002). There may be more effective grazers of *M. aeruginosa* than flagellates, including some fish species (Miura 1990), cladoceran zooplankton (Hanazato and Yasuno 1984),

rotifers (Snell 1980; Fulton and Pearl 1987; Jarvis et al. 1987), and amoebae (Yamamoto and Suzuki 1984; Nishibe et al. 2004). In addition to grazing, other *Microcystis* consumers may include the algicidal effects of heterotrophic bacteria (Manage et al. 2000), and viruses (Yoshida et al. 2006). We have not yet collected data on *in situ* loss of *Microcystis* concentrations due to biological processes, and such loss may be synergistic with the aforementioned *Microcystis* eradicators. Additional studies are needed to understand *Microcystis* loss due to biological processes.

Acknowledgments

This research was supported by the Environment Research and Technology Development Fund (D-0905) of the Ministry of the Environment, Japan.

Literature cited

- Baek SH, Hong SS, Song SY, Lee HO, Nakano S, Han MS (2009) Grazing effects on toxic and non-toxic *Microcystis aeruginosa* by the mixotrophic flagellate *Ochromonas* sp. J Freshwater Ecol 24:367-373
- Burkert U, Hyenstrand P, Drakare S, Blomqvist P (2001) Effects of the mixotrophic flagellate *Ochromonas* sp. on colony formation in *Microcystis aeruginosa*.

- 1 Aquat Ecol 35:9-17
- 2 Carmichael WW (1992) Cyanobacterial secondary metabolites — the cyanotoxins. J
- 3 Appl Microbiol 72:445-459
- 4 Cole GT, Wynne MJ (1974) Endocytosis of *Microcystis aeruginosa* by *Ochromonas*
- 5 *danica*. J Phycol 10:397-410
- 6 Dokulil MT, Teubner K (2000) Cyanobacterial dominance in lakes. Hydrobiologia
- 7 438:1-12
- 8 Dolan JR, Šimek K (1998) Ingestion and digestion of an autotrophic picoplankter,
- 9 *Synechococcus*, by a heterotrophic nanoflagellate, *Bodo saltans*. Limnol
- 10 Oceanogr 43:1740-1746
- 11 Fulton RS, Pearl HW (1987) Toxic and inhibitory effects of the blue-green alga
- 12 *Microcystis aeruginosa* on herbivorous zooplankton. J Plankton Res 9:837-855
- 13 Gobler CJ, Davis TW, Coyne KJ, Boyer GL (2007) Interactive influences of nutrient
- 14 loading, zooplankton grazing, and microcystin synthetase gene expression on
- 15 cyanobacterial bloom dynamics in a eutrophic New York lake. Harmful Algae
- 16 6:119-133
- 17 Guo S, Gan N, Zheng L, Song L (2008) Effect of *Microcystis aeruginosa*
- 18 (Cyanobacteria) abundance on the growth and ingestion of *Poteroiochromonas*

- 1 sp. (Chrysophyceae). Bioinformatics and Biomedical Engineering, 2008.
- 2 ICBBE 2008. The 2nd International Conference on 16-18 May: 3639-3695
- 3 Hanazato Y, Yasuno M (1984) Growth, reproduction and assimilation of *Moina*
- 4 *macropoda* fed on *Microcystis* and/or *Chlorella*. Jpn J Ecol (Otsu) 34:195-202
- 5 Heinbokel JF (1978) Studies on the functional role of Tintinnids in the Southern
- 6 California Bight. II. Grazing rates of field populations. Mar Biol 47:191-197
- 7 Hodoki Y, Ohbayashi K, Kobayashi Y, Okuda N, Nakano S (2011) Temporal variation
- 8 in cyanobacteria species composition and photosynthetic activity in
- 9 experimentally induced blooms. J Plankton Res 33:1410-1416
- 10 Imai H, Chang KH, Kusaba M, Nakano S (2009) Temperature-dependent dominance of
- 11 *Microcystis* (Cyanophyceae) species: *M. aeruginosa* and *M. wesenbergii*. J
- 12 Plankton Res 31:171-178.
- 13 Ishikawa K, Walker RF, Tsujimura S, Nakahara H, Kumagai M (2004) Estimation of
- 14 *Microcystis* colony size in developing water blooms via image analysis. J Jpn
- 15 Soc Wat Environ 27:69-72 (in Japanese with English abstract)
- 16 Jarvis AC, Hart RC, Combrink S (1987) Zooplankton feeding on size fractionated
- 17 *Microcystis* colonies and *Chlorella* in a hypertrophic lake (Hartbeespoort Dam,
- 18 South Africa): implications to resource utilization and zooplankton succession.

- 1 J Plankton Res 9:1231-1249
- 2 Jeong HJ, Park JY, Nho JH, Park MO, Ha JH, Seong KA, Jeng C, Seong CN, Lee KY,
3 Yih WH (2005) Feeding by red-tide dinoflagellates on the cyanobacterium
4 *Synechococcus*. Aquat Microb Ecol 41:131-143
- 5 Kim BR, Nakano S, Kim BH, Han MS (2006) Grazing and growth of the heterotrophic
6 flagellate *Diphyllia rotans* on the cyanobacterium *Microcystis aeruginosa*.
7 Aquat Microb Ecol 45:163-170
- 8 Latour D, Sabido O, Salencon MJ, Giraudet H (2004) Dynamics and metabolic activity
9 of the benthic cyanobacterium *Microcystis aeruginosa* in the Grangent
10 reservoir (France). J Plankton Res 26:719-726
- 11 Manage PM, Kawabata Z, Nakano S (2000) Algicidal effect of the bacterium
12 *Alcaligenes denitrificans* on *Microcystis* spp. Aquat Microb Ecol 22:111-117
- 13 Mischke U (1994) Influence of food quality and quantity on ingestion and growth rates
14 of three omnivorous heterotrophic flagellates. Mar Microbiol Food Webs 8:
15 125-143
- 16 Miura T (1990) The effects of planktivorous fishes on the plankton community in a
17 eutrophic lake. Hydrobiologia 200/201:567-579
- 18 Nishibe Y, Kawabata Z, Nakano S (2002) Grazing on *Microcystis aeruginosa* by the

- 1 heterotrophic flagellate *Collodictyon triciliatum* in a hypertrophic pond. *Aquat*
2 *Microb Ecol* 29:173-179
- 3 Nishibe Y, Manage PM, Kawabata Z, Nakano S (2004) Trophic coupling of a testate
4 amoeba and *Microcystis* species in a hypertrophic pond. *Limnology* 5:71-76
- 5 Park HD, Iwami C, Watanabe MF, Harada K, Okino T, Hayashi H (1998) Temporal
6 variabilities of the concentration of inter- and extracellular microcystin and
7 toxic *Microcystis* species in a hypertrophic lake, Lake Suwa, Japan
8 (1991-1994). *Environ Toxicol Water Qual* 13:61-72
- 9 Reynolds CS, Walsby AE (1975) Water blooms. *Biol Rev* 50:437-481
- 10 Sherr BF, Sherr EB, Fallon RD (1987) Use of monodispersed, fluorescently labeled
11 bacteria to estimate in situ protozoan bacterivory. *Appl Environ Microbiol* 53:
12 958-965
- 13 Snell T W (1980) Blue-green algae and selection in rotifer populations. *Oecologia* 46:
14 343-346
- 15 Sugiura N, Inamori Y, Ouchiyama T, Sudo R (1992) Degradation of cyanobacteria,
16 *Microcystis* by microflagellate, *Monas guttula*. *Water Sci Technol*
17 26:2173-2176
- 18 Van Donk E, Cerbin S, Wilken S, Helmsing NR, Ptacnik R, Verschoor AM (2009) The

- 1 effect of a mixotrophic chrysophyte on toxic and colony-forming cyanobacteria.
- 2 Freshwater Biol 54:1843-1855
- 3 Watanabe MM, Ichimura T (1997) Fresh- and salt-water forms of *Spirulina platensis* in
- 4 axenic cultures. Bull. Jpn. Soc. Phycol 25 Suppl (Mem Iss Yamada) 371-377
- 5 Welschmeyer NA (1994) Fluorometric analysis of chlorophyll *a* in the presence of
- 6 chlorophyll *b* and pheopigments. Limnol Oceanogr 39:1985-1992
- 7 Wilken S, Wiezer S, Huisman J, van Donk E (2010) Microcystins do not provide
- 8 anti-herbivore defense against mixotrophic flagellates. Aquat Microb Ecol
- 9 59:207-216
- 10 Yamamoto Y, Nakahara H (2009) Seasonal variations in the morphology of
- 11 bloom-forming cyanobacteria in a eutrophic pond. Limnology 10:185-193
- 12 Yamamoto Y, Suzuki K (1984) Light and electron microscope observations and prey
- 13 specificities of an algophorous amoeba from Japanese freshwater. J Gen Appl
- 14 Microbiol 30:411-417
- 15 Yang Z, Kong F, Shi X, Zhang M, Xing P, Cao H (2008) Changes in the morphology
- 16 and polysaccharide content of *Microcystis aeruginosa* (Cyanobacteria) during
- 17 flagellate grazing. J Phycol 44:716-720
- 18 Yang Z, Kong F, Yang Z, Zhang M, Yu Y, Qian S (2009) Benefits and costs of the

- 1 grazer-induced colony formation in *Microcystis aeruginosa*. Ann Limnol-Int J
2 Lim 45:203-208
- 3 Yoshida T, Takashima Y, Tomaru Y, Shirai Y, Takao Y, Hiroishi S, Nagasaki K (2006)
4 Isolation and characterization of a cyanophage infecting the toxic
5 Cyanobacterium *Microcystis aeruginosa*. Appl Environ Microb 72:1239-1247
- 6 Zhang X, Watanabe MM, Inouye I (1996) Light and electron microscopy of grazing by
7 *Poterioochromonas malhamensis* (Chrysophyceae) on a range of
8 phytoplankton taxa. J Phycol 32:37-46
- 9 Zhang X, Hu HY, Men YJ, Yang J, Christoffersen K (2009) Feeding characteristics of a
10 golden alga (*Poterioochromonas* sp.) grazing on toxic cyanobacterium
11 *Microcystis aeruginosa*. Water Res 43:2953-2960
- 12
13
14
15

- 1 Table 1. Grazing and clearance rates of *Collodictyon triciliatum* on *Microcystis aeruginosa*, and daily grazing impact of *C. triciliatum*
- 2 on the standing stock of *M. aeruginosa* in an experimental pond. Min: a minimum value, Max: a maximum value, SD: standard
- 3 deviation.

	Range (Min - Max)	Means $\pm$ SD
Grazing rate (<i>M. aeruginosa</i> cells flagellate ⁻¹ h ⁻¹)	0.2 – 6.9	1.1 $\pm$ 1.4
Clearance rate (nl flagellate ⁻¹ h ⁻¹)	2.2 – 162.2	39.6 $\pm$ 40.0
Daily grazing impact (% standing stock day ⁻¹)	0.0 – 25.3	3.4 $\pm$ 7.1

- 4
- 5
- 6
- 7
- 8
- 9

1 Table 2. Comparison of $I_{\max}$ and K_{IR} in different predators, when fed *M. aeruginosa* or *Synechococcus*. $I_{\max}$: maximum ingestion rate, K_{IR} : the prey concentration
2 sustaining one-half $I_{\max}$.

Protist	Prey	$I_{\max}$	K_{IR}	Source
		(cells flagellate ⁻¹ h ⁻¹)	(10 ⁵ cells ml ⁻¹)	
<i>Collodictyon triciliatum</i>	<i>M. aeruginosa</i> NIES843	6.2	1.2	Present study
<i>Diphylleia rotans</i>	<i>M. aeruginosa</i> NIER -10001	0.56		Kim et al. (2006)
<i>Diphylleia rotans</i>	<i>M. aeruginosa</i> NIES -298	0.73		Kim et al. (2006)
<i>Diphylleia rotans</i>	<i>M. aeruginosa</i> NIES -101 (non-toxic strain)	0.17		Kim et al. (2006)
<i>Diphylleia rotans</i>	<i>M. viridis</i> NIES -102	0.35		Kim et al. (2006)
<i>Ochromonas</i> sp.	<i>M. aeruginosa</i> PCC7806 wild type	2.21 ± 0.25	0.13 ± 0.25	Wilken et al. (2010)
<i>Ochromonas</i> sp.	<i>M. aeruginosa</i> PCC7807 mutant	1.68 ± 0.14	0.14 ± 0.25	Wilken et al. (2010)
<i>Poterioochromonas</i> sp.	<i>M. aeruginosa</i> FACHB 469	3.3	37.6	Guo et al. (2008)

Fig 1. Changes in (A) Water temperature, (B) Chlorophyll *a* concentration, abundance of (C) *M. aeruginosa* and (D) *C. triciliatum*, and (E) average number of ingested *M. aeruginosa* in the food vacuoles of *C. triciliatum*, in an experimental pond. Error bars indicate standard deviations for each mean value in (E).

Fig 2. Relationship between *M. aeruginosa* cell number in the food vacuoles of *C. triciliatum* and time. Error bars indicate $\pm$ standard error. Error bars that are not visible are hidden behind the symbols.

Fig 3. Ingestion rate (cells flagellate⁻¹ h⁻¹) of *C. triciliatum* on *M. aeruginosa* as a function of the initial prey concentration (cell ml⁻¹). Each value was calculated from a linear regression curve of the number of prey cells in flagellate vacuoles over incubation time. The curve was fitted by a Michaelis-Menten equation (in Materials and methods) using all treatments in the experiment. Ingestion rate (*IR*, cell flagellate⁻¹ h⁻¹) = $6.185 [x/(122280 + x)]$, $r^2 = 0.873$, $p < 0.001$, where *x* is *Microcystis* abundance.

Fig. 1 Kobayashi et al.

Fig. 2 Kobayashi et al.

Fig. 3 Kobayashi et al.