

Title	Constructions of metrics and dimensions (Research trends on general and geometric topology and their applications)
Author(s)	Kato, Hisao
Citation	数理解析研究所講究録 (2011), 1728: 45-50
Issue Date	2011-02
URL	http://hdl.handle.net/2433/170531
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

距離の構成と色々な次元 Constructions of metrics and dimensions

筑波大学数理物質科学研究科 加藤久男 (Hisao Kato)
Institute of Mathematics, University of Tsukuba

The key word of this study is normal sequence of finite open covers. In general topology, the notion of normal sequence of open covers is one of the most useful tools for the study. In fact, the notion is the essence of metrizable spaces. We obtain directly the numerical properties of normal sequences of finite open covers on a given separable metric space X and we give another proof of Pontrjagin-Schnirelmann theorem. Furthermore, by use of normal sequences we can construct desired metrics d which control the values of $\log N(\epsilon, d)/|\log \epsilon|$. We investigate strong relations between topological dimension $\dim X$, metrics d and lower and upper box-counting dimensions $\underline{\dim}_B(X, d)$, $\overline{\dim}_B(X, d)$ of separable metric spaces X from a point of view of general topology. In particular, we construct *chaotic metrics* with respect to the determination of the upper and lower box-counting dimensions.

For a totally bounded metric d on X and $\epsilon > 0$, let

$$N(\epsilon, d) = \min\{|\mathcal{U}| \mid \mathcal{U} \text{ is a finite open cover of } X \text{ with } \text{mesh}_d(\mathcal{U}) \leq \epsilon\},$$

where $|A|$ denotes the cardinality of a set A . Then the lower and upper box-counting dimensions of (X, d) are given by

$$\underline{\dim}_B(X, d) = \liminf_{\epsilon \rightarrow 0} \frac{\log N(\epsilon, d)}{|\log \epsilon|}, \quad \overline{\dim}_B(X, d) = \limsup_{\epsilon \rightarrow 0} \frac{\log N(\epsilon, d)}{|\log \epsilon|}.$$

If $\underline{\dim}_B(X, d) = \overline{\dim}_B(X, d)$, we define $\dim_B(X, d) = \underline{\dim}_B(X, d)$. Suppose that \mathcal{U} is an open cover of X and $A \subset X$. Let

$$St(A, \mathcal{U}) = \bigcup \{U \in \mathcal{U} \mid U \cap A \neq \emptyset\}$$

$$\mathcal{U}^* = \{St(U, \mathcal{U}) \mid U \in \mathcal{U}\} \quad \text{and} \quad \mathcal{U}^\Delta = \{St(x, \mathcal{U}) \mid x \in X\}$$

$$\mathcal{U}^{*0} = \mathcal{U}, \quad \mathcal{U}^{\Delta 0} = \mathcal{U}$$

$$\mathcal{U}^{*p+1} = (\mathcal{U}^{*p})^* = \{St(W, \mathcal{U}^{*p}) \mid W \in \mathcal{U}^{*p}\}$$

$$\mathcal{U}^{\Delta p+1} = (\mathcal{U}^{\Delta p})^\Delta = \{St(x, \mathcal{U}^{\Delta p}) \mid x \in X\}.$$

An open cover \mathcal{V} of X is a *star-refinement* of an open cover \mathcal{U} of X if \mathcal{V}^* is a refinement of \mathcal{U} . An open cover \mathcal{V} of X is a *delta-refinement* of an open cover \mathcal{U} of X if \mathcal{V}^Δ is a refinement of \mathcal{U} . Let \mathcal{U}_i ($i = 1, 2, \dots$) be open covers of X . Then the sequence $\{\mathcal{U}_i\}_{i=1}^\infty$ is called a *normal star-sequence* if \mathcal{U}_{i+1} is a star-refinement of \mathcal{U}_i ($i = 1, 2, \dots$). Also, the sequence $\{\mathcal{U}_i\}_{i=1}^\infty$ is called a *normal delta-sequence* if \mathcal{U}_{i+1} is a delta-refinement of \mathcal{U}_i ($i = 1, 2, \dots$). The sequence $\{\mathcal{U}_i\}_{i=1}^\infty$ is called a *development* of X if $\{St(x, \mathcal{U}_i) \mid i = 1, 2, \dots\}$ is a neighborhood base for each point x of X . The following theorem is well known as Alexandroff-Urysohn metrization theorem.

Theorem 0.1. (Alexandroff-Urysohn metrization theorem) *A T_1 -space X is metrizable if and only if there exists a sequence $\{\mathcal{U}_i\}_{i=1}^{\infty}$ of open covers of X such that $\{\mathcal{U}_i\}_{i=1}^{\infty}$ is a normal sequence and a development of X .*

For any normal space X ($\neq \phi$) and natural numbers k and p , we define the following indices:

(1) The index $\star_k^p(X)$ is defined as the least natural number m such that for every open cover \mathcal{U} of X with $|\mathcal{U}| = k$, there is an open cover \mathcal{V} of X such that $|\mathcal{V}| \leq m$ and $\mathcal{V}^{\star p} \leq \mathcal{U}$.

(2) The index $\Delta_k^p(X)$ is defined as the least natural number m such that for every open cover \mathcal{U} of X with $|\mathcal{U}| = k$, there is an open cover \mathcal{V} of X such that $|\mathcal{V}| \leq m$ and $\mathcal{V}^{\Delta p} \leq \mathcal{U}$.

By C_m^k , we denote the set of all m -element subsets of the set $\{1, 2, \dots, k\}$ and by $\binom{k}{m}$

its cardinality, i.e., $\binom{k}{m} = \frac{k!}{m!(k-m)!}$.

For natural numbers k, m and p with $k \geq m$, we define the following indices;

$$\tilde{\Delta}(k; m; p) = \sum_{m \geq j_1 \geq j_2 \geq \dots \geq j_p \geq 1} \binom{k}{j_1} \binom{j_1}{j_2} \dots \binom{j_{p-1}}{j_p},$$

$$\tilde{\star}(k; m; p) = \sum_{m \geq j_1 \geq j_2 \geq \dots \geq j_p \geq 1} \binom{k}{j_1} \binom{j_1}{j_2} \dots \binom{j_{p-1}}{j_p} j_p.$$

The following result follows from Bruijning-Nagata [4], Hashimoto-Hattori [8], Bogatyi-Karpov [2] and Koto-Matsumoto [10].

Theorem 0.2. *Let X be an infinite normal space with $\dim X = m < \infty$ and let k and p be natural numbers. Then*

$$\star_k^p(X) = \begin{cases} \tilde{\star}(k; k; (1/2)(3^p - 1)) = k[(1/2)(3^p - 1) + 1]^{k-1} & (k \leq m + 1) \\ \tilde{\star}(k; m + 1; (1/2)(3^p - 1)) & (k \geq m + 1), \end{cases}$$

$$\Delta_k^p(X) = \begin{cases} \tilde{\Delta}(k; k; 2^{p-1}) = (2^{p-1} + 1)^k - (2^{p-1})^k & (k \leq m + 1) \\ \tilde{\Delta}(k; m + 1; 2^{p-1}) & (k \geq m + 1). \end{cases}$$

Lemma 0.3. (Koto-Matsumoto [10]) *Let X be an infinite separable metric space with $\dim X = m \geq 0$. Then the followings hold.*

1. *If $\{\mathcal{U}_i\}_{i=1}^{\infty}$ is a normal star-sequence of finite open covers of X and a development of X , then there is some i_0 such that*

$$|\mathcal{U}_i| \geq \tilde{\star}(m + 1; m + 1; (1/2)(3^{(i-i_0)} - 1))$$

for $i \geq i_0$. In particular, $\liminf_{i \rightarrow \infty} \frac{\log_3 |\mathcal{U}_i|}{i} \geq m$.

2. *If $\{\mathcal{U}_i\}_{i=1}^{\infty}$ is a normal delta-sequence of finite open covers of X and a development of X , then there is some i_0 such that*

$$|\mathcal{U}_i| \geq \tilde{\Delta}(m + 1; m + 1; 2^{(i-i_0-1)})$$

for $i \geq i_0$. In particular, $\liminf_{i \rightarrow \infty} \frac{\log_2 |\mathcal{U}_i|}{i} \geq m$.

Theorem 0.4. (Kato-Matsumoto [10], Kato [11]) *Let X be a nonempty separable metric space. Then*

$$\begin{aligned} \dim X &= \min\left\{\liminf_{i \rightarrow \infty} \frac{\log_3 |\mathcal{U}_i|}{i} \mid \{\mathcal{U}_i\}_{i=1}^{\infty} \text{ is a normal star-sequence of} \right. \\ &\quad \left. \text{finite open covers of } X \text{ and a development of } X\right\} \\ &= \min\left\{\liminf_{i \rightarrow \infty} \frac{\log_2 |\mathcal{U}_i|}{i} \mid \{\mathcal{U}_i\}_{i=1}^{\infty} \text{ is a normal delta-sequence of} \right. \\ &\quad \left. \text{finite open covers of } X \text{ and a development of } X\right\}. \end{aligned}$$

Moreover, there exists a normal star (resp. delta)-sequence $\{\mathcal{U}_i\}_{i=1}^{\infty}$ of finite open covers of X which is a development of X such that

$$\dim X = \lim_{i \rightarrow \infty} \frac{\log_3 |\mathcal{U}_i|}{i} \quad (\text{resp. } \dim X = \lim_{i \rightarrow \infty} \frac{\log_2 |\mathcal{U}_i|}{i}).$$

Consider the following indices:

$$\star^p(X, \mathcal{U}) = \min\{|\mathcal{V}| \mid \mathcal{V} \text{ is a finite open covering of } X \text{ such that } \mathcal{V}^{\star^p} \leq \mathcal{U}\},$$

$$\Delta^p(X, \mathcal{U}) = \min\{|\mathcal{V}| \mid \mathcal{V} \text{ is a finite open covering of } X \text{ such that } \mathcal{V}^{\Delta^p} \leq \mathcal{U}\}.$$

Theorem 0.5. (Kato-Matsumoto [10]) *Let X be a normal space. Then*

$$\dim X = \sup\left\{\limsup_{p \rightarrow \infty} \frac{\log_3 \star^p(X, \mathcal{U})}{p} \mid \mathcal{U} \text{ is a finite open covering of } X\right\}$$

and

$$\dim X = \sup\left\{\limsup_{p \rightarrow \infty} \frac{\log_2 \Delta^p(X, \mathcal{U})}{p} \mid \mathcal{U} \text{ is a finite open covering of } X\right\}.$$

The next proposition implies that for any separable metric space X there is a natural bijection from the set of all totally bounded metrics on X to the set of Alexandroff-Urysohn metrics on X induced by normal sequences of finite open covers which are developments of X , up to Lipschitz equivalence.

Proposition 0.6. (Kato-Matsumoto [10]) *Let X be a separable metric space and let ρ be a totally bounded metric on X . Then there is a normal star (resp. delta)-sequence $\{\mathcal{U}_i\}_{i=1}^{\infty}$ of finite open covers of X such that $\{\mathcal{U}_i\}_{i=1}^{\infty}$ is a development of X and ρ is Lipschitz equivalent to d , where d is the Alexandroff-Urysohn metric induced by $\{\mathcal{U}_i\}_{i=1}^{\infty}$.*

For separable metric spaces, we need the Alexandroff-Urysohn metrics induced by normal sequences of finite open covers. Define the functions $D_{\star} : X \times X \rightarrow [0, 9]$ and $D_{\Delta} : X \times X \rightarrow [0, 4]$ as follows:

(\star) Let $\{\mathcal{U}_i\}_{i=1}^{\infty}$ be a normal star-sequence of finite open covers of X and a development of X . For any pair of points x, y of X , we define the function $D_{\star} : X \times X \rightarrow [0, 9]$ by

1. $D_{\star}(x, y) = 9$ if $\{x, y\}$ is not contained in any element of \mathcal{U}_1 ,

2. $D_*(x, y) = 1/3^{(i-2)}$ if $\{x, y\}$ is contained in an element of \mathcal{U}_i and $\{x, y\}$ is not contained in any element of \mathcal{U}_j for $j > i$,
3. $D_*(x, y) = 0$ if $\{x, y\}$ is contained in an element of \mathcal{U}_i for each i .

(Δ) Let $\{\mathcal{U}_i\}_{i=1}^\infty$ be a normal delta-sequence of finite open covers of X and a development of X . For any pair of points x, y of X , we define the function $D_\Delta : X \times X \rightarrow [0, 4]$ by

1. $D_\Delta(x, y) = 4$ if $\{x, y\}$ is not contained in any element of \mathcal{U}_1 ,
2. $D_\Delta(x, y) = 1/2^{(i-2)}$ if $\{x, y\}$ is contained in an element of \mathcal{U}_i and $\{x, y\}$ is not contained in any element of \mathcal{U}_j for $j > i$,
3. $D_\Delta(x, y) = 0$ if $\{x, y\}$ is contained in an element of \mathcal{U}_i for each i .

Proposition 0.7. (see Nagami's book [14] for (2)) *Let X be a T_1 -space.*

1. *If $\{\mathcal{U}_i\}_{i=1}^\infty$ is a normal star-sequence of finite open covers of X and a development of X , then $\{\mathcal{U}_i\}_{i=1}^\infty$ induces a totally bounded metric d_* on X such that for any $x, y \in X$,*

$$d_*(x, y) \leq D_*(x, y) \leq 6d_*(x, y).$$

2. *If $\{\mathcal{U}_i\}_{i=1}^\infty$ is a normal delta-sequence of finite open covers of X and a development of X , then $\{\mathcal{U}_i\}_{i=1}^\infty$ induces a totally bounded metric d_Δ on X such that for any $x, y \in X$,*

$$d_\Delta(x, y) \leq D_\Delta(x, y) \leq 4d_\Delta(x, y).$$

By use of the above results, we obtain an another proof of the following well-known theorem.

Corollary 0.8. (Pontrjagin-Schnirelmann [19], Brouwer's theorem [3] and Kato [11]) *Let X be a separable metric space. Then*

$$\dim X = \min\{\underline{\dim}_B(X, \rho) \mid \rho \text{ is a totally bounded metric for } X\}.$$

Moreover,

$$\dim X = \min\{\overline{\dim}_B(X, \rho) \mid \rho \text{ is a totally bounded metric for } X\}.$$

Lemma 0.9. (Kato [11]) *Let X be an infinite separable metric space and let $\{\mathcal{U}_i\}_{i=1}^\infty$ be a normal star-sequence of finite open covers and a development of X such that*

$$\lim_{i \rightarrow \infty} \frac{\log_3 |\mathcal{U}_i|}{i} = \dim X.$$

Then for any α, β with $\dim X \leq \alpha \leq \beta \leq \infty$, there is a subsequence $\{\mathcal{U}_{i_j}\}_{j=1}^\infty$ of $\{\mathcal{U}_i\}_{i=1}^\infty$ such that

$$[\alpha, \beta] = \left\{ \liminf_{k \rightarrow \infty} \frac{\log_3 |\mathcal{U}_{i_{n_k}}|}{n_k} \mid \{n_k\}_{k=1}^\infty \text{ is an increasing subsequence of natural numbers} \right\}.$$

Theorem 0.10. (Kato-Matsumoto [10], Kato [11]) *Let X be an infinite separable metric space. For any $\alpha, \beta \in [\dim X, \infty]$ with $\alpha \leq \beta$, there is a totally bounded metric $d = d_{\alpha\beta}$ on X such that*

$$[\alpha, \beta] = \left\{ \liminf_{k \rightarrow \infty} \frac{\log N(\epsilon_k, d)}{|\log \epsilon_k|} \mid \{\epsilon_k\}_{k=1}^{\infty} \text{ is a decreasing sequence} \right. \\ \left. \text{of positive numbers with } \lim_{k \rightarrow \infty} \epsilon_k = 0 \right\}.$$

In particular, $\underline{\dim}_B(X, d) = \alpha$ and $\overline{\dim}_B(X, d) = \beta$.

Corollary 0.11. (Keesling [12], Kato-Matsumoto [10], Kato [11]) *Let X be a separable metric space with $\dim X \geq 1$. For any $\alpha, \beta \in [\dim X, \infty]$ with $\alpha \leq \beta$, there is a totally bounded metric $d = d_{\alpha\beta}$ on X such that*

$$[\alpha, \beta] = \left\{ \liminf_{k \rightarrow \infty} \frac{\log N(\epsilon_k, d)}{|\log \epsilon_k|} \mid \{\epsilon_k\}_{k=1}^{\infty} \text{ is a decreasing sequence} \right. \\ \left. \text{of positive numbers with } \lim_{k \rightarrow \infty} \epsilon_k = 0 \right\}.$$

In particular, $\dim_H(X, d) = \underline{\dim}_B(X, d) = \alpha$ and $\overline{\dim}_B(X, d) = \beta$, where $\dim_H(X, d)$ is the Hausdorff dimension of (X, d) .

Finally, we have the following problems.

Problem 0.12. (1) *Give another proof of the following theorem of E. Marczewski (=Szpilrajn) by use of normal sequence of open covers: For a separable metric space X , $\dim X = \min\{\dim_H(X, d) \mid d \text{ is a metric on } X\}$.*

(2) *Let X be a separable metric space with $\dim X \geq 1$. For any $\alpha, \beta, \gamma \in [\dim X, \infty]$ with $\alpha \leq \beta \leq \gamma$, does there exist a totally bounded metric d on X such that $\dim_H(X, d) = \alpha \leq \underline{\dim}_B(X, d) = \beta \leq \overline{\dim}_B(X, d) = \gamma$?*

(3) *What kinds of metrics can be embedded into Euclidean spaces, up to Lipschitz equivalence? If $\overline{\dim}_B(X, d) \leq n \in \mathbb{N}$, is it true that d can be embedded into $(2n+1)$ -dimensional Euclidean space, up to Lipschitz equivalence? Note that if $\dim X = n$, there is a metric d on X such that d can be embedded into $(2n+1)$ -dimensional Euclidean space with $\overline{\dim}_B(X, d) = n$.*

References

- [1] M. Barnsley, R. Devaney, B. Mandelbrot, H. O. Peitgen, D. Saupe and R. Voss, *The Science of Fractal Images*. Springer-Verlag, Berlin-New York, 1988.
- [2] S. A. Bogatyi and A. N. Karpov, Bruijning-Nagata and Hashimoto-Hattori Characteristics of Covering Dimension Revisited, *Mathematical Notes*, 79 (2006), 327-334.
- [3] J. Bruijning, A characterization of dimension of topological spaces by totally bounded pseudometrics, *Pacific J. Math.* 84 (1979), 283-289.

- [4] J. Bruijning and J. Nagata, A characterization of covering dimension by use of $\Delta_k(X)$, *Pacific J. Math.* 80 (1979), 1-8.
- [5] R. Engelking, Theory of dimensions finite and infinite, *Heldermann Verlag, Lemgo*, 1995.
- [6] K. Falconer, Fractal Geometry, *Mathematical Foundations and Applications. John Wiley Sons, Chichester*, 1990.
- [7] C. Fujita, H. Kato and M. Matsumoto, Fractal Metrics of Ruelle Expanding Maps and Expanding Ratios, *Topology and its Applications*, 157 (2010), 615-628.
- [8] K. Hashimoto and Y. Hattori, On Nagata's star-index $\star_k(X)$, *Topology Appl.* 122 (2002), 201-204.
- [9] W. Hurewicz and H. Wallman, Dimension Theory, *Princeton Univ. Press, Princeton, New Jersey*, 1948.
- [10] H. Kato and M. Matsumoto, Characterizations of topological dimension by use of normal sequences of finite open covers and Pontrjagin-Schniremann theorem, *J. Math. Soc. Japan*, to appear.
- [11] H. Kato, Addendum: Characterizations of topological dimension by use of normal sequences of finite open covers and Pontrjagin-Schniremann theorem, *J. Math. Soc. Japan*, to appear.
- [12] J. Keesling, Hausdorff dimension, *Topology Proc.* 11 (1986), 349-384.
- [13] R. D. Mauldin and M. Urbański, Graphic directed Markov Systems: Geometry and dynamics of limit sets, *Cambridge University Press, Cambridge* 148, 2003.
- [14] K. Nagami, Dimension Theory, *Academic Press, New York-London*, 1970.
- [15] J. Nagata, Modern Dimension Theory, *Heldermann Verlag*, 1983.
- [16] J. Nagata, Modern General Topology, *North-Holland, Amsterdam*, 1985.
- [17] J. Nagata, Open problems left in my wake of research, *Topology Appl.* 146/147 (2005), 5-13.
- [18] Y. B. Pesin, Dimension Theory in Dynamical Systems, *University of Chicago Press, Chicago* 1997.
- [19] L. Pontrjagin and L. Schnirelmann, Sur une propriété métrique de la dimension, *Ann. of Math.* (2) 33 (1932), 156-162.