

|             |  |
|-------------|--|
| Title       | On equivariant homeomorphisms of boundaries of CAT(0)groups (Research trends on general and geometric topology and their applications) |
| Author(s)   | 保坂, 哲也   |
| Citation    | 数理解析研究所講究録 (2011), 1728: 27-31   |
| Issue Date  | 2011-02  |
| URL         | <a href="http://hdl.handle.net/2433/170534">http://hdl.handle.net/2433/170534</a>  |
| Right       |  |
| Type        | Departmental Bulletin Paper  |
| Textversion | publisher  |

## On equivariant homeomorphisms of boundaries of CAT(0) groups

宇都宮大学教育学部

保坂 哲也 (Tetsuya Hosaka)

本稿では, CAT(0) 群  $G$  が幾何学的に (proper [8, p.131], ココンパクト, 等長的に) 作用する 2 つの CAT(0) 空間  $X, Y$  の境界  $\partial X, \partial Y$  が  $G$ -equivariant 同相となるための十分条件を紹介する.

CAT(0) 空間の定義と詳細は [8] [20] に見られる. CAT(0) 群とは, ある CAT(0) 空間に幾何学的に作用する群のことである.

歴史的には, まず Gromov hyperbolic group  $G$  が幾何学的に作用する測地線空間  $X, Y$  の境界  $\partial X, \partial Y$  は, quasi-isometry である自然な写像  $Gx_0 \rightarrow Gy_0$  ( $gx_0 \mapsto gy_0$ ) の連続的な拡張として境界上の  $G$ -equivariant 同相  $\partial X \rightarrow \partial Y$  を導くことがよく知られている. 実際, Gromov hyperbolic group  $G$  の境界は quasi-isometric invariant である (cf. [8], [11], [20], [21], [22]).

ここで, Gromov [22] により, 「CAT(0) 群  $G$  が幾何学的に作用する 2 つの CAT(0) 空間  $X, Y$  の境界  $\partial X, \partial Y$  は  $G$ -equivariant 同相となるか?」という質問が提起された.

まず, P. L. Bowers - K. Ruane [7] によって, ある単純な CAT(0) 群  $G$  と CAT(0) 空間  $X, Y$  の例を用いて, 自然な quasi-isometry  $Gx_0 \rightarrow Gy_0$  ( $gx_0 \mapsto gy_0$ ) の連続的な拡張として, それらの境界の連続写像  $\partial X \rightarrow \partial Y$  を導くことができない例が構成されている. (この Bowers-Ruane と類似の例は, S. Yamagata [39] によって, right-angled Coxeter 群と Davis complex を用いても構成されている.)

また, C. Croke - B. Kleiner [12] によって, ある CAT(0) 群で, それが幾何学的に作用する CAT(0) 空間の境界の位相が決定されない例が構成され, 更に J. Wilson [38] により, その Croke-Kleiner の与えた CAT(0) 群は非可算無限の種類位相の境界を持つことを示した. 最近, C. Mooney [33] によっても非可算無限の境界を持つ CAT(0) 群の例が knot 群によって構成されている.

また, 一方で M. Bestvina [5] により, 与えられた CAT(0) 群のすべての境界が shape 同値となることが考察されており, 与えられた CAT(0) 群のすべての境界が cell-like 同値となるかは Bestvina により未解決問題として提出されている. この問題の部分的な解決が [2], [34] にみられる.

CAT(0) 群  $G$  が幾何学的に作用する 2 つの CAT(0) 空間  $X, Y$  の境界  $\partial X, \partial Y$  が  $G$ -equivariant 同相となるための十分条件の紹介の前に, 上述の Bowers-Ruane の例を紹介する.

**Bowers-Ruane Example.**  $G = F_2 \times \mathbb{Z}$ ,  $X = Y = T \times \mathbb{R}$  とおく. ただし,  $F_2$  は  $\{a, b\}$  によって生成される rank 2 の free group とし,  $T$  を生成集合  $\{a, b\}$  に関する  $F_2$  の Cayley graph とする.

まず, 群  $G$  の CAT(0) 空間  $X$  への作用 “ $\cdot$ ” を以下で定義する:

$$(a, 0) \cdot (t, r) = (a \cdot t, r),$$

$$(b, 0) \cdot (t, r) = (b \cdot t, r),$$

$$(1, 1) \cdot (t, r) = (t, r + 1),$$

ここで  $(t, r) \in T \times \mathbb{R} = X$ . 群  $G$  は  $\{(a, 0), (b, 0), (1, 1)\}$  によって生成されていることに注意する.

また, 群  $G$  の CAT(0) 空間  $Y$  への作用 “ $*$ ” を以下で定義する:

$$(a, 0) * (t, r) = (a \cdot t, r),$$

$$(b, 0) * (t, r) = (b \cdot t, r + 2),$$

$$(1, 1) * (t, r) = (t, r + 1),$$

ここで  $(t, r) \in T \times \mathbb{R} = Y$ .

いま, 群  $G$  は 2 つの CAT(0) 空間  $X, Y$  に幾何学的に作用する. 一方で, quasi-isometry  $g \cdot x_0 \mapsto g * y_0$  (ここで  $x_0 = (1, 0) \in X$ ,  $y_0 = (1, 0) \in Y$ ) は  $\partial X$  から  $\partial Y$  へ連続的に写像が拡張できないことを示す.

まず,  $g_i = a^i b^i \in F_2$  について,  $\partial T$  において  $\{g_i^\infty \mid i \in \mathbb{N}\} \rightarrow a^\infty$  ( $i \rightarrow \infty$ ) が成り立つ.

また,  $X \cup \partial X$  において

$$\lim_{n \rightarrow \infty} (g_i^n, 0) \cdot x_0 = [g_i^\infty, 0]$$

$$\lim_{n \rightarrow \infty} (a^n, 0) \cdot x_0 = [a^\infty, 0]$$

が成り立つ.

一方で,  $Y \cup \partial Y$  において

$$\begin{aligned}\lim_{n \rightarrow \infty} (g_i^n, 0) * y_0 &= [g_i^\infty, \frac{\pi}{4}] \\ \lim_{n \rightarrow \infty} (a^n, 0) * y_0 &= [a^\infty, 0]\end{aligned}$$

が成り立つ.

従って, もし quasi-isometry  $\phi : G \cdot x_0 \rightarrow G * y_0$  ( $g \cdot x_0 \rightarrow g * y_0$ ) が境界上に連続的に写像  $\bar{\phi} : \partial X \rightarrow \partial Y$  に拡張したと仮定すると,

$$\begin{aligned}\bar{\phi}([g_i^\infty, 0]) &= [g_i^\infty, \frac{\pi}{4}] \\ \bar{\phi}([a^\infty, 0]) &= [a^\infty, 0]\end{aligned}$$

が成り立つはずである.

しかし, これは,

$$[g_i^\infty, 0] \rightarrow [a^\infty, 0] \quad (i \rightarrow \infty)$$

より, 点  $[a^\infty, 0]$  において写像  $\bar{\phi}$  を連続的に定義することができない ([7, p.187]).

ここで, 上の例において, 次のことに着目する.

- (a)  $d_X(a^i \cdot x_0, [x_0, g^i \cdot x_0]) = 0$  for any  $i \in \mathbb{N}$  in  $X$  and
- (b) there does *not* exist a constant  $M > 0$  such that  $d_Y(a^i * y_0, [y_0, g^i * y_0]) \leq M$  for any  $i \in \mathbb{N}$  in  $Y$ .

この考察により, 群  $G$  が 2 つの CAT(0) 空間  $X, Y$  に幾何学的に作用するときの条件 (\*) を以下で定義する (ただし  $x_0 \in X, y_0 \in Y$ ).

- (\*) ある 2 つの constant  $N > 0, M > 0$  が存在して,  $GB(x_0, N) = X,$   
 $GB(y_0, M) = Y$  が成立し, 任意の  $g, a \in G$  に対して  $[x_0, gx_0] \cap$   
 $B(ax_0, N) \neq \emptyset$  in  $X$  ならば  $[y_0, gy_0] \cap B(ay_0, M) \neq \emptyset$  in  $Y$  が成立する.

ここで, 次の定理を得た.

**Main Theorem.** 群  $G$  が 2 つの CAT(0) 空間  $X, Y$  に幾何学的に作用し,  $x_0 \in X, y_0 \in Y$  とする. このとき, 条件 (\*) が成立するならば, quasi-isometry  $\phi : Gx_0 \rightarrow Gy_0$  ( $\phi(gx_0) = gy_0$ ) は境界上の  $G$ -equivariant 同相写像  $\bar{\phi} : \partial X \rightarrow \partial Y$  に連続的に拡張する.

この定理の証明の議論および上述の条件 (\*) は, 以下の問題に応用できる可能性がある.

- (1) CAT(0) 群の境界の (equivariant) rigidity に関する問題;

- (2) Coxeter 群の境界の (equivariant) rigidity に関する問題;
- (3) Coxeter 群の Davis 複体の境界の (equivariant) rigidity に関する問題;
- (4) Coxeter 群が鏡映群として作用する CAT(0) 空間の境界の (equivariant) rigidity に関する問題;
- (5) right-angled Coxeter 群が鏡映群として作用する CAT(0) 空間の境界の (equivariant) rigidity に関する問題;
- (6) CAT(0) 群が幾何学的に作用する CAT(0) cube 複体の境界の (equivariant) rigidity に関する問題;

etc.

## REFERENCES

- [1] A. D. Alexandrov, V. N. Berestovskii and I. G. Nikolaev, *Generalized Riemannian spaces*, Russ. Math. Surveys 41 (1986), 1–54.
- [2] F. D. Ancel, C. Guilbault, and J. Wilson, *The Croke-Kleiner boundaries are cell-like equivalent*, preprint.
- [3] W. Ballmann and M. Brin, *Orbihedra of nonpositive curvature*, Inst. Hautes Études Sci. Publ. Math. 82 (1995), 169–209.
- [4] W. Ballmann, M. Gromov and V. Schroeder, *Manifolds of Nonpositive Curvature*, Progr. Math. vol. 61, Birkhäuser, Boston MA, 1985.
- [5] M. Bestvina, *Local homology properties of boundaries of groups*, Michigan Math. J. 43 (1996), 123–139.
- [6] N. Bourbaki, *Groupes et Algèbres de Lie*, Chapters IV-VI, Masson, Paris, 1981.
- [7] P. Bowers and K. Ruane, *Boundaries of nonpositively curved groups of the form  $G \times \mathbb{Z}^n$* , Glasgow Math. J. 38 (1996), 177–189.
- [8] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999.
- [9] K. S. Brown, *Buildings*, Springer-Verlag, 1980.
- [10] P. Caprace and K. Fujiwara, *Rank-one isometries of buildings and quasi-morphisms of Kac-Moody groups*, arXiv:0809.0470v3, preprint.
- [11] M. Coornaert and A. Papadopoulos, *Symbolic dynamics and hyperbolic groups*, Lecture Notes in Math 1539, Springer-Verlag, 1993.
- [12] C. B. Croke and B. Kleiner, *Spaces with nonpositive curvature and their ideal boundaries*, Topology 39 (2000), 549–556.
- [13] C. B. Croke and B. Kleiner, *The geodesic flow of a nonpositively curved graph manifold*, Geom. Funct. Anal. 12 (2002), 479–545.
- [14] M. W. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. 117 (1983), 293–324.
- [15] M. W. Davis, *Nonpositive curvature and reflection groups*, in Handbook of geometric topology (Edited by R. J. Daverman and R. B. Sher), pp. 373–422, North-Holland, Amsterdam, 2002.
- [16] A. N. Dranishnikov, *On boundaries of hyperbolic Coxeter groups*, Topology Appl. 110 (2001), 29–38.
- [17] A. N. Dranishnikov, *Boundaries of Coxeter groups and simplicial complexes with given links*, J. Pure Appl. Algebra 137 (1999), 139–151.

- [18] H. Fischer, *Boundaries of right-angled Coxeter groups with manifold nerves*, *Topology* 42 (2003), 423–446.
- [19] R. Geoghegan and P. Ontaneda, *Boundaries of cocompact proper  $CAT(0)$  spaces*, *Topology* 46 (2007), 129–137.
- [20] E. Ghys and P. de la Harpe (ed), *Sur les Groupes Hyperboliques d'après Mikhael Gromov*, *Progr. Math.* vol. 83, Birkhäuser, Boston MA, 1990.
- [21] M. Gromov, *Hyperbolic groups*, *Essays in group theory* (Edited by S. M. Gersten), pp. 75–263, M.S.R.I. Publ. 8, 1987.
- [22] M. Gromov, *Asymptotic invariants for infinite groups*, *Geometric Group Theory* (G.A. Niblo and M.A. Roller, eds.), *LMS Lecture Notes*, vol. 182, Cambridge University Press, Cambridge, 1993, pp. 1–295.
- [23] U. Hamenstädt, *Rank-one isometries of proper  $CAT(0)$ -spaces*, arXiv:0810.3794v1, preprint.
- [24] T. Hosaka, *The interior of the limit set of groups*, *Houston J. Math.* 30 (2004), 705–721.
- [25] T. Hosaka, *Reflection groups of geodesic spaces and Coxeter groups*, *Topology Appl.* 153 (2006), 1860–1866.
- [26] T. Hosaka, *On splitting theorems for  $CAT(0)$  spaces and compact geodesic spaces of non-positive curvature*, arXiv:math.GR/0405551v1, preprint.
- [27] T. Hosaka, *Parabolic subgroups of Coxeter groups acting by reflections on  $CAT(0)$  spaces*, arXiv:math/0409472v1, preprint.
- [28] T. Hosaka, *On boundaries of Coxeter groups and topological fractal structures*, arXiv:0912.0061v2, preprint.
- [29] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge University Press, 1990.
- [30] M. Mihalik and K. Ruane,  *$CAT(0)$  groups with non-locally connected boundary*, *J. London Math. Soc.* (2) 60 (1999), 757–770.
- [31] M. Mihalik, K. Ruane and S. Tschantz, *Local connectivity of right-angled Coxeter group boundaries*, *J. Group Theory* 10 (2007), 531–560.
- [32] N. Monod, *Superrigidity for irreducible lattices and geometric splitting*, *J. Amer. Math. Soc.* 19 (2006), 781–814.
- [33] C. Mooney, *Examples of non-rigid  $CAT(0)$  groups from the category of knot groups*, arXiv:0706.1581v2, preprint.
- [34] C. Mooney, *All  $CAT(0)$  boundaries of a group of the form  $H \times K$  are CE equivalent*, arXiv:0707.4316, preprint.
- [35] G. Moussong, *Hyperbolic Coxeter groups*, Ph.D. thesis, Ohio State University, 1988.
- [36] P. Papasoglu and E. L. Swenson, *Boundaries and JSJ decompositions of  $CAT(0)$ -groups*, *Geom. Funct. Anal.* 19 (2009), 558–590.
- [37] E. L. Swenson, *A cut point theorem for  $CAT(0)$  groups*, *J. Differential Geom.* 53 (1999), 327–358.
- [38] J. M. Wilson, *A  $CAT(0)$  group with uncountably many distinct boundaries*, *J. Group Theory* 8 (2005), 229–238.
- [39] S. Yamagata, *On ideal boundaries of some Coxeter groups*, *Advanced Studies Pure Math.* 55 (2009), 345–352.