

## LENS SPACE SURGERIES ALONG TWO COMPONENT LINKS AND REIDEMEISTER-TURAEV TORSION

TERUHISA KADOKAMI AND YUICHI YAMADA

### 1. INTRODUCTION

This article is a short survey of the core part of the authors' joint work "Lens space surgeries along certain 2-component links, Park's rational blow down, and Reidemeister-Turaev torsion" [KTYYY, preprint]. In [KTYYY], certain two families of 2-component links, denoted by  $A_{m,n}$  and  $B_{p,q}$  are focused, and the main result is the decision of the coefficient(s) of the knotted component yielding a lens space by Dehn surgery. The links are related to rational homology 4-ball used in J. Park's (generalized) rational blow down in 4-dimensional topology (see [3, 12]). Concrete calculus on the links  $A_{m,n}$  and  $B_{p,q}$  was important. The results made the contrast between  $A_{m,n}$  (hyperbolic) and  $B_{p,q}$  (Seifert) clear, which was one of the purpose of [KTYYY] (see also [17]).

In this article, we focus another importance, the method itself to get some necessary conditions on the lens space surgery coefficients of a given link, by using Alexander polynomial and Reidemeister torsion. Our method satisfies that a result on a link  $L$  always extends to the links whose Alexander polynomials are same with that of  $L$ .

We will compare the Reidemeister torsion of the result  $M$  of Dehn surgery along a given link and that of a lens space  $L(p, q)$  (in Example 3.4). Some necessary conditions are obtained from the value  $\tau^{\psi_d}(M)$  of the Reidemeister torsion in the  $d$ -th cyclotomic field  $\mathbb{Q}(\zeta_d)$  by  $d$ -norm, where  $d(\geq 2)$  is a divisor of  $p$ . From the sequence of the equalities on  $\tau^{\psi_d}(M)$ s in  $\mathbb{Q}(\zeta_d)$  for all divisors  $d$  of  $p$  (with a fixed combinatorial Euler structure of  $M$ ), we take an *identity on symmetric Laurent polynomials*, as a lift of the equalities. We regard the identity as an equation of the surgery coefficient for  $M$  to be a lens space.

In the next section, we start with some definitions of the Reidemeister torsion. In Section 3, we review surgery formulae. In Section 4, we will study  $d$ -norms in the  $d$ -th cyclotomic field, and show a certain uniqueness of a symmetric polynomial as a lift of the sequence of the equalities in  $\mathbb{Q}(\zeta_d)$ s. In Section 5, we will explain the method to get some necessary condition of lens space surgery coefficients of a given link. In Section 6, as a demonstration, we will apply our method to Berge's link, which is one of the most famous targets in lens space surgery ([1]).

### 2. REIDEMEISTER TORSION

For a precise definition of the Reidemeister torsion, the reader refer to V. Turaev [14, 15]. Let  $X$  be a finite CW complex and  $\pi : \tilde{X} \rightarrow X$  its maximal abelian covering. Then  $\tilde{X}$  has a CW structure induced by that of  $X$  and  $\pi$ , and the cell chain complex  $C_*$  of  $\tilde{X}$  has a

$E_L$	the complement of $L$ .
$m_i, l_i$	a meridian and a longitude of the $i$ -th component $K_i$ .
$[m_i], [l_i]$	their homology classes.
$\Delta_L(t_1, \dots, t_\mu)$	the Alexander polynomial of $L$ , where $t_i$ is represented by $[m_i]$ .
$(L; r_1, \dots, r_\mu)$	the result of Dehn surgery along $L$ , where $r_i \in \mathbb{Q} \cup \{\infty, \emptyset\}$ is the surgery coefficient of $K_i$ .
$V_i$	the solid torus attached along $K_i$ in the Dehn surgery.
$m'_i, [m'_i]$	a meridian of $V_i$ , and its homology class.
$l'_i, [l'_i]$	an oriented core curve of $V_i$ , and its homology class.

TABLE 1. Notations (for manifolds)

$\mathbb{Z}[H]$ -module structure, where  $H = H_1(X; \mathbb{Z})$  is the first homology of  $X$ . For an integral domain  $R$  and a ring homomorphism  $\psi : \mathbb{Z}[H] \rightarrow R$ , “the chain complex of  $\tilde{X}$  related with  $\psi$ ”, denoted by  $\mathbf{C}_*^\psi$ , is constructed as  $\mathbf{C}_* \otimes_{\mathbb{Z}[H]} Q(R)$ , where  $Q(R)$  is the quotient field of  $R$ . The Reidemeister torsion of  $X$  related with  $\psi$ , denoted by  $\tau^\psi(X)$ , is calculated from  $\mathbf{C}_*^\psi$ , and is an element of  $Q(R)$  determined up to multiplication of  $\pm\psi(h)$  ( $h \in H$ ). If  $R = \mathbb{Z}[H]$  and  $\psi$  is the identity map, then we denote  $\tau^\psi(X)$  by  $\tau(X)$ . We note that  $\tau^\psi(X)$  is not zero if and only if  $\mathbf{C}_*^\psi$  is acyclic.

**Notation** (for manifolds and homologies) Let  $L = K_1 \cup \dots \cup K_\mu$  be an oriented  $\mu$ -component link in  $S^3$ . We will use the notations in Table 1.

**Notation** (for algebra) For a pair of elements  $A, B$  in  $Q(R)$ , if there exists an element  $h \in H$  such that  $A = \pm\psi(h)B$ , then we denote the equality by  $A \doteq B$ . We will often take a field  $F$  and a ring homomorphism  $\psi : \mathbb{Z}[H_1(M)] \rightarrow F$ . We mainly use the  $d$ -th cyclotomic fields  $\mathbb{Q}(\zeta_d)$  as  $F$ , where  $\zeta_d$  is a primitive  $d$ -th root of unity.

### 3. SURGERY FORMULAE

Let  $E$  be a compact 3-manifold whose boundary  $\partial E$  consists of tori ( $E$  is possibly not  $E_L$  for a link  $L$ ). We study the 3-manifold  $M = E \cup V_1 \cup \dots \cup V_n$  obtained by attaching solid tori  $V_i$ s to  $E$  by attaching maps  $f_i : \partial V_i \rightarrow \partial E$  ( $\text{Im}(f_i) \cap \text{Im}(f_j) = \emptyset$  for  $i \neq j$ ). By  $l'_i$  we denote the core of  $V_i$ . We let  $\iota : E \hookrightarrow M$  denote the natural inclusion.

**Lemma 3.1.** (Surgery formula I) *If  $\psi([l'_i]) \neq 1$  for every  $i = 1, \dots, n$ , then*

$$\tau^\psi(M) \doteq \tau^{\psi'}(E) \prod_{i=1}^n (\psi([l'_i]) - 1)^{-1},$$

where  $\psi' = \psi \circ \iota_*$  ( $\iota_*$  is a ring homomorphism induced by  $\iota$ ).

For the case of the complement  $E_L$  of a  $\mu$ -component link  $L$  in  $S^3$  as in Table 1. The Reidemeister torsion is closely related with the Alexander polynomial.

**Lemma 3.2.** (Milnor [11]) *Let  $\Delta_L(t_1, \dots, t_\mu)$  be the Alexander polynomial of a  $\mu$ -component link  $L = K_1 \cup \dots \cup K_\mu$  in  $S^3$ , where a variable  $t_i$  is represented by the meridian*

of  $K_i$  ( $i = 1, \dots, \mu$ ).

$$\tau(E_L) \doteq \begin{cases} \Delta_L(t_1)(t_1 - 1)^{-1} & (\mu = 1), \\ \Delta_L(t_1, \dots, t_\mu) & (\mu \geq 2). \end{cases}$$

Next, we study the result of Dehn surgery  $M = (L; p_1/q_1, \dots, p_\mu/q_\mu)$  along  $L$ . We take integers  $r_i$  and  $s_i$  satisfying  $p_i s_i - q_i r_i = -1$ .

**Lemma 3.3.** (Surgery formula II; T. Sakai [13], V. G. Turaev [14])

- (1) In the case  $M = (K; p/q)$  ( $|p| \geq 2$ ), we have  $H = H_1(M) \cong \langle T \mid T^p = 1 \rangle \cong \mathbb{Z}/|p|\mathbb{Z}$ , where  $T$  is represented by the meridian  $[m]$ . For a divisor  $d$  ( $\geq 2$ ) of  $p$ , we define a ring homomorphism  $\psi_d : \mathbb{Z}[H] \rightarrow \mathbb{Q}(\zeta_d)$  by  $\psi_d(T) = \zeta_d$ . Then we have

$$\tau^{\psi_d}(M) \doteq \Delta_K(\zeta_d)(\zeta_d - 1)^{-1}(\zeta_d^{\bar{q}} - 1)^{-1}$$

where  $q\bar{q} \equiv 1 \pmod{p}$ .

- (2) In the case  $M = (L; p_1/q_1, \dots, p_\mu/q_\mu)$  ( $\mu \geq 2$ ). Let  $F$  be a field and  $\psi : \mathbb{Z}[H_1(M)] \rightarrow F$  a ring homomorphism. If  $\psi([m_i]^{r_i}[l_i]^{s_i}) \neq 1$  for every  $i = 1, \dots, \mu$ , then we have

$$\tau^\psi(M) \doteq \Delta_L(\psi([m_1]), \dots, \psi([m_\mu])) \prod_{i=1}^{\mu} (\psi([m_i]^{r_i}[l_i]^{s_i}) - 1)^{-1}.$$

**Example 3.4.** The lens space  $L(p, q)$  is obtained as  $-p/q$ -surgery along the unknot. By Lemma 3.3 (1), for a divisor  $d \geq 2$  of  $p$ , we have

$$\tau^{\psi_d}(L(p, q)) \doteq (\zeta_d - 1)^{-1}(\zeta_d^{\bar{q}} - 1)^{-1},$$

where  $q\bar{q} \equiv 1 \pmod{p}$ .

## 4. CYCLOTOMIC FIELD AND POLYNOMIAL

### 4.1. $d$ -norm.

About algebraic fields, the reader refer to L. C. Washington [16] for example.

For an element  $x$  in the  $d$ -th cyclotomic field  $\mathbb{Q}(\zeta_d)$ , the  $d$ -norm of  $x$  is defined as

$$N_d(x) = \prod_{\sigma \in \text{Gal}(\mathbb{Q}(\zeta_d)/\mathbb{Q})} \sigma(x),$$

where  $\text{Gal}(\mathbb{Q}(\zeta_d)/\mathbb{Q})$  is the Galois group ( $\cong (\mathbb{Z}/d\mathbb{Z})^\times$ ) related with a Galois extension  $\mathbb{Q}(\zeta_d)$  over  $\mathbb{Q}$ . The following is well-known.

**Proposition 4.1.**

- (1) If  $x \in \mathbb{Q}(\zeta_d)$ , then  $N_d(x) \in \mathbb{Q}$ . The map  $N_d : \mathbb{Q}(\zeta_d) \setminus \{0\} \rightarrow \mathbb{Q} \setminus \{0\}$  is a group homomorphism.
- (2) If  $x \in \mathbb{Z}[\zeta_d]$ , then  $N_d(x) \in \mathbb{Z}$ .

By easy calculations, we have the following.

**Lemma 4.2.**

- (1)  $N_d(\pm\zeta_d) = \begin{cases} \pm 1 & (d = 2), \\ 1 & (d \geq 3). \end{cases}$
- (2)  $N_d(1 - \zeta_d) = \begin{cases} \ell & (d \text{ is a power of a prime } \ell \geq 2), \\ 1 & (\text{otherwise}). \end{cases}$

About applications of  $d$ -norms, for example, see [5, 6, 7, 8, 9, 10].

#### 4.2. Reidemeister–Turaev torsion.

Let  $M$  be a homology lens space with  $H = H_1(M) \cong \mathbb{Z}/p\mathbb{Z}$  ( $p \geq 2$ ). Then the Reidemeister torsion  $\tau^{\psi_d}(M)$  of  $M$  related with  $\psi_d$  is determined up to multiplication of  $\pm \zeta_d^m$  ( $m \in \mathbb{Z}$ ), where  $d \geq 2$  is a divisor of  $p$  and  $\psi_d$  is the same ring homomorphism as in Lemma 3.3 (1). Once we fix a basis of a cell chain complex for the maximal abelian covering of  $M$  as a  $\mathbb{Z}[H] = \mathbb{Z}[t, t^{-1}]/(t^p - 1)$ -module, the value  $\tau^{\psi_d}(M)$  is uniquely determined as an element of  $\mathbb{Q}(\zeta_d)$  for every  $d$ . The choice of the basis up to “base change equivalence” is called a *combinatorial Euler structure* of  $M$  (cf. Turaev [15]). The Reidemeister torsion of a manifold with a fixed combinatorial Euler structure is said the *Reidemeister–Turaev torsion*.

We consider the sequence of the values  $\tau^{\psi_d}(M)$  in  $\mathbb{Q}(\zeta_d)$  of the Reidemeister–Turaev torsion for every divisor  $d \geq 2$  of  $p$ , and regard them as a value sequence  $\{\tau^{\psi_d}(M)\}_{d|p, d \geq 2}$  defined as below.

**Definition 4.3.** We define that a sequence of values  $\mathbf{x} = \{x_d\}_{d|p, d \geq 2}$  is a *value sequence* (of degree  $p$ ) if  $x_d \in \mathbb{Q}(\zeta_d)$  for every  $d$ . Two value sequences  $\mathbf{x} = \{x_d\}_{d|p, d \geq 2}$  and  $\mathbf{y} = \{y_d\}_{d|p, d \geq 2}$  are *equal* ( $\mathbf{x} = \mathbf{y}$ ) if  $x_d = y_d$  for every  $d$ . We are mainly concerned with the value sequence of type  $\mathbf{x} = \{F(\zeta_d)\}_{d|p, d \geq 2}$  for a rational function  $F(t) \in \mathbb{Q}(t)$ . In such a case, we say that  $\mathbf{x}$  is *induced by*  $F(t)$  and that  $F(t)$  is a *lift* of  $\mathbf{x}$ . A *control* of  $\mathbf{x} = \{x_d\}_{d|p, d \geq 2}$  by a trivial unit  $u = \eta t^m \in \mathbb{Z}[t, t^{-1}]/(t^p - 1)$  is defined by

$$u\mathbf{x} = \{\eta \zeta_d^m x_d\}_{d|p, d \geq 2},$$

where  $\eta = 1$  or  $-1$  (constant) and  $m \in \mathbb{Z}$ . Two value sequences  $\mathbf{x} = \{x_d\}_{d|p, d \geq 2}$  and  $\mathbf{y} = \{y_d\}_{d|p, d \geq 2}$  are *control equivalent* if there is a trivial unit  $u \in \mathbb{Z}[t, t^{-1}]/(t^p - 1)$  such that  $\mathbf{y} = u\mathbf{x}$ . A value sequence  $\mathbf{x} = \{x_d\}_{d|p, d \geq 2}$  is a *real value sequence* if  $x_d$  is a real number for every  $d$ .

**Example 4.4.** A value sequence  $\mathbf{x}$  of degree 12 is in the form  $\mathbf{x} = \{x_2, x_3, x_4, x_6, x_{12}\}$ . The following two value sequences  $\mathbf{x}, \mathbf{y}$  of degree 12 are not equal, but control equivalent for  $u = t^6$ .

$$\mathbf{x} = \{2, -1, -2, -1, 1\}, \quad \mathbf{y} = \{2, -1, 2, 1, -1\}.$$

In fact,  $\mathbf{x}$  and  $\mathbf{y}$  is induced by  $t^2 + t^{-2}$  and  $t^4 + t^{-4}$ , respectively.

Let  $M$  be a homology lens space with  $H_1(M) \cong \mathbb{Z}/p\mathbb{Z}$  ( $p \geq 2$ ). Then a sequence  $\{\tau^{\psi_d}(M)\}_{d|p, d \geq 2}$  of the Reidemeister torsions of  $M$  with a combinatorial Euler structure is a value sequence of degree  $p$ . We say the value sequence a *torsion sequence* of  $M$ .

**Lemma 4.5.**

- (1) Let  $M$  and  $M'$  be homeomorphic homology lens spaces with  $H_1(M) \cong H_1(M') \cong \mathbb{Z}/p\mathbb{Z}$  ( $p \geq 2$ ). Then torsion sequences  $\{\tau^{\psi_d}(M)\}_{d|p, d \geq 2}$  and  $\{\tau^{\psi'_d}(M')\}_{d|p, d \geq 2}$  related with the corresponding ring homomorphisms  $\psi_d$  and  $\psi'_d$  (i.e.,  $\psi_d = \psi'_d \circ h_*$ , where  $h_*$  is the induced homomorphism of the homeomorphism) are control equivalent.
- (2) Let  $M$  be a homology lens space with  $H_1(M) \cong \mathbb{Z}/p\mathbb{Z}$  ( $p \geq 2$ ). Then we can control a torsion sequence of  $M$  into a real value sequence.

*Proof.* (1) It is easy to see.

(2) Here we let  $\zeta$  denote any  $d$ -th primitive root ( $\zeta_d$ ) of unity. Since  $M$  is obtained by  $p/q$ -surgery along a knot  $K$  in a homology 3-sphere for some  $q$  (cf. [2]), and we can also

apply Lemma 2.5 (1) for the case, we have

$$\tau^{\psi_d}(M) \doteq \Delta_K(\zeta)(\zeta - 1)^{-1}(\zeta^{\bar{q}} - 1)^{-1}$$

where  $q\bar{q} \equiv 1 \pmod{p}$ . By the duality of the Alexander polynomial (cf. [11, 14, 15]), we may assume

$$\Delta_K(t) = \Delta_K(t^{-1}).$$

This is also a control of the combinatorial Euler structure of the exterior of  $K$ , which induces a control of a torsion sequence of  $M$ . We take an odd integer lift of  $\bar{q}$ . Then

$$\zeta^{\frac{1+\bar{q}}{2}} \Delta_K(\zeta)(\zeta - 1)^{-1}(\zeta^{\bar{q}} - 1)^{-1}$$

is a real number for every  $d$ . □

**Lemma 4.6.** *If two real value sequences  $\mathbf{x} = \{x_d\}_{d|p, d \geq 2}$  and  $\mathbf{y} = \{y_d\}_{d|p, d \geq 2}$  of degree  $p$  are control equivalent satisfying  $\mathbf{y} = u\mathbf{x}$  for a trivial unit  $u = \eta t^m \in \mathbb{Z}[t, t^{-1}]/(t^p - 1)$ , where  $\eta = \pm 1$  and  $m \in \mathbb{Z}$ , then the possibility of  $u$  is restricted as follows:*

- (i) *If  $p$  is odd, then  $u = 1$  or  $-1$ .*
- (ii) *If  $p$  is even, then  $u = 1, -1, t^{p/2}$  or  $-t^{p/2}$ .*

*Proof.* Since the ratio  $\zeta_p^m = \pm y_p/x_p$  is a real number, we have (i)  $m \equiv 0 \pmod{p}$  if  $p$  is odd, and (ii)  $m \equiv 0$  or  $p/2 \pmod{p}$  if  $p$  is even. □

**Definition 4.7.** (Symmetric Laurent polynomial) A Laurent polynomial  $F(t) \in \mathbb{Z}[t, t^{-1}]$  is *symmetric* if it is of the form

$$F(t) = a_0 + \sum_{i=1}^{\infty} a_i(t^i + t^{-i}),$$

where  $a_i$  is an integer for all  $i = 1, 2, \dots$  and  $a_i = 0$  for every sufficiently large  $i$ . Note that, if  $F(t)$  is a symmetric Laurent polynomial, the induced value sequence  $\{F(\zeta_d)\}_{d|p, d \geq 2}$  is a real value sequence. We are concerned with symmetric Laurent polynomials that are lifts (in  $\mathbb{Z}[t, t^{-1}]$ ) of a polynomial in the quotient ring  $\mathbb{Z}[t, t^{-1}]/(t^p - 1)$ . We say that  $F(t)$  (as above) is *reduced* if  $a_i = 0$  for all  $i > [p/2]$ . We often *reduce* the symmetric polynomials by using  $t^i + t^{-i} = t^{p+i} + t^{-(p+i)}$  modulo  $(t^p - 1)$ . We let  $\text{red}(F(t))$  denote the reduction of  $F(t)$  (i.e.,  $\text{red}(F(t))$  is reduced and  $\text{red}(F(t)) = F(t)$  in  $\mathbb{Z}[t, t^{-1}]/(t^p - 1)$ ). We will use a notation  $\langle t^i \rangle = t^i + t^{-i}$ , for short.

For a Laurent polynomial  $F(t) \in \mathbb{Z}[t, t^{-1}]$ , the *span* of  $F(t)$  is the difference of the maximal degree of  $F(t)$  and the minimal degree of  $F(t)$ , and we denote it by  $\text{span}(F(t))$ .

**Lemma 4.8.** *Let  $N \geq 2$  be an integer. Let  $F(t), G(t)$  be symmetric Laurent polynomials and  $\mathbf{x} = \{F(\zeta_d)\}_{d|N, d \geq 2}$ ,  $\mathbf{y} = \{G(\zeta_d)\}_{d|N, d \geq 2}$  the induced real value sequences, respectively. If  $\mathbf{x}$  and  $\mathbf{y}$  are control equivalent, i.e.,  $u\mathbf{x} = \mathbf{y}$  for a trivial unit  $u$  (here,  $u = 1$  or  $-1$  if  $N$  is odd,  $u = 1, -1, t^{N/2}$  or  $-t^{N/2}$  if  $N$  is even, by Lemma 4.6), and  $F(1) = G(1) = 0$ , then we have a congruence*

$$uF(t) \equiv G(t) \pmod{t^N - 1}$$

Furthermore, assuming  $\text{span}(G(t)) \leq 2[N/2]$ ,

- (i) *In the case that  $u = 1$  or  $-1$  and  $\text{span}(F(t)) \leq N - 1$ , we have an identity  $uF(t) = G(t)$  in  $\mathbb{Z}[t, t^{-1}]$ .*

(ii) Otherwise (in the case that  $N$  is even and  $u = \eta t^{N/2}$  with  $\eta = 1$  or  $-1$ ), we have  $\text{red}(t^{N/2}F(t)) = \eta G(t)$  in  $\mathbb{Z}[t, t^{-1}]$ .

*Proof.* By Chinese Remainder Theorem, we have a ring isomorphism:

$$\mathbb{Q}[t, t^{-1}]/(t^N - 1) \cong \bigoplus_{d|N, d \geq 1} \mathbb{Q}(\zeta_d),$$

where  $f(t)$  in the left-hand side maps to the value sequences  $\{f(\zeta_d)\}_{d|N, d \geq 2}$  in the right-hand side. The isomorphism implies the required congruence.  $\square$

Note that  $F(t)$  and  $t^{N/2}F(t)$  induce the control equivalent real value sequences by  $u = t^{N/2}$ , but  $\text{red}(t^{N/2}F(t)) \neq F(t)$  in general, see Example 4.4. Thus we have to care the case (ii) in the lemma. Here, we study relation between the coefficients of  $F(t)$  and those of  $\text{red}(t^{N/2}F(t))$ .

**Lemma 4.9.** *Let  $N$  be an even integer.*

$$\text{If } F(t) = a_0 + \sum_{i=1}^{N/2} a_i(t^i + t^{-i}), \text{ then } \text{red}(t^{N/2}F(t)) = b_0 + \sum_{i=1}^{N/2} b_i(t^i + t^{-i})$$

with

$$b_0 = 2a_{N/2}, \quad b_{N/2} = a_0/2 \text{ and } b_j = a_{N/2-j} \quad (j = 1, 2, \dots, N/2 - 1).$$

*Proof.* It is because

$$t^{N/2}(t^j + t^{-j}) = t^{N/2+j} + t^{N/2-j} \equiv t^{(N/2-j)} + t^{-(N/2-j)} \pmod{t^N - 1}.$$

$\square$

## 5. METHOD

Let  $L = K_1 \cup K_2 \cup \dots \cup K_\mu$  be a link. We let  $M$  simply denote the result  $(L; r_1, \dots, r_\mu)$  of the Dehn surgery. We use the notations in Table 1.

**Step 1** Study the first homologies (the generators and relations), from the exterior  $E_L$  of  $L$  (Of course,  $H_1(E_L; \mathbb{Z}) \cong \bigoplus_{i=1}^\mu \mathbb{Z}[m_i]$ ) to the result  $M$ , by attaching solid tori  $V_i$  one by one.

The first (obvious) necessary condition for the result  $M$  of Dehn surgery to be a lens space  $L(p, q)$  is

$$H_1(M; \mathbb{Z}) \cong \mathbb{Z}/p\mathbb{Z}.$$

**Step 2** Calculate the Alexander polynomial  $\Delta_L(t_1, \dots, t_\mu)$  of  $L$ . Using Lemma 3.2 and Lemma 3.3, calculate the Reidemeister torsion  $\tau^\psi(M)$  related with a ring homomorphism  $\psi: \mathbb{Z}[H_1(M)] \rightarrow \mathbb{Q}(\zeta_d)$ , where  $d (\geq 2)$  is a divisor of  $p$ .

If  $M$  is homeomorphic to a lens space  $L(p, q)$  (with undecided  $q$ ), then their Reidemeister torsions are equal to each other. By Example 3.4, there exists integers  $i, j$  coprime to  $p$  with  $0 < i, j < p$  (they are lifts of  $(\mathbb{Z}/p\mathbb{Z})^\times / \{\pm 1\}$ ) such that

$$(1) \quad \tau^\psi(M) \doteq \frac{1}{(\zeta_d^i - 1)(\zeta_d^j - 1)} \quad \text{in } \mathbb{Q}(\zeta_d),$$

for each divisor  $d (\geq 2)$  of  $p$ . We can assume  $i + j$  is even by retaking  $p - j$  instead of  $j$ .

**Step 3** Using  $d$ -norm in  $\mathbb{Q}(\zeta_d)$ , studied in Subsection 4.1, to the equality (1), we have a necessary condition on the coefficient of lens space surgery.

We fix a combinatorial Euler structure (multiple of trivial unit  $\pm\zeta_d^k$ ), deform both hand-sides of the equality (1) into real values by Lemma 4.5(2). If  $M$  is homeomorphic to  $L(p, q)$ , we have a control equivalence between the real value sequence:

$$\{\tau^\psi(M)\}_{d|p, d \geq 2} = u \{ \zeta_d^{\frac{i+j}{2}} (\zeta_d^i - 1)^{-1} (\zeta_d^j - 1)^{-1} \}_{d|p, d \geq 2},$$

where  $u$  is a trivial unit  $\pm 1$ , or  $\pm t^{p/2}$  (only in the case  $p$  is even). By Lemma 4.8, we have, via a congruence mod  $(t^p - 1)$ , an identity between symmetric Laurent polynomials. We regard the identity as an equation (on  $(i, j)$ ) of the coefficients of lens space surgery.

**Step 4** By the equation, we have a necessary condition on the coefficient(s) of lens space surgery.

## 6. DEMONSTRATION

We call the link in Figure 1 *Berge's link BL*. The compliment is a hyperbolic 3-manifold, known as Berge's manifold in [1]. The component  $K_1$  is the famous pretzel knot  $P(-2, 3, 7)$ . The link, regarded as a knot in a solid torus (the exterior of the component  $K_2$ ), admits two surgery coefficients yielding solid torus itself, and it is proved that such a hyperbolic link is unique [1]. We demonstrate our method in Section 5 to Berge's link,




FIGURE 1. Berge link  $BL$

to study lens space surgeries  $M := (BL; r, 0)$ , where  $r = \alpha/\beta$  ( $\alpha, \beta \in \mathbb{Z}, \gcd(\alpha, \beta) = 1$ ). We assume that  $\beta \geq 1$ .

(Step 1)

$$H_1(M) \cong \langle [m_1], [m_2] \mid [l_1] = [m_2]^7, [l_2] = [m_1]^7, [m_1]^\alpha [l_1]^\beta = 1, [m_1]^7 = 1 \rangle.$$

It is finite cyclic  $\mathbb{Z}/p\mathbb{Z}$  if and only if  $\gcd(\alpha, 7) = 1$ , and then we have  $p = 7^2\beta = 49\beta$ . An element  $T = [m_1]^\gamma [m_2]^{\delta'}$  with  $\alpha\delta' - 7\beta\gamma' = -1$  is a generator:  $T^{49\beta} = 1$ . We also have  $[l_1'] = [m_1]^\gamma [l_1]^\delta$  with  $\alpha\delta - \beta\gamma = -1$ , and

$$[m_1] = T^{7\beta}, [m_2] = [l_2'] = T^{-\alpha}, [l_1'] = T^7.$$

(Step 2) The Alexander polynomial of Berge's link is

$$\Delta_{BL}(t, x) \doteq 1 + t^3x + t^5x^2 + t^8x^3 + t^{11}x^4 + t^{13}x^5 + t^{16}x^6 = \sum_{i=0}^6 t^{s_i} x^i,$$

where we define a sequence  $(s_0, s_1, \dots, s_6) = (0, 3, 5, 8, 11, 13, 16)$ . This is not periodic, but we regard it as "Periodicity is broken a little". We let  $M_1 = E_{BL} \cup V_1 = (BL; \alpha/\beta, -)$ . We have, up to the ambiguity (multiplication  $\pm T^k$ ),

$$\tau(M_1) \doteq \Delta_{BL}(T^{7\beta}, T^{-\alpha})(T^7 - 1)^{-1} = \left( \sum_{i=0}^6 T^{7\beta s_i - \alpha i} \right) (T^7 - 1)^{-1}.$$

We take a divisor  $d = 7$  of  $p = 49\beta$  and let  $\zeta$  denote a primitive 7-th root of unity. We use deformations

$$T^{7\beta s_i - \alpha i} = T^{-\alpha i} (T^{7\beta s_i} - 1) + T^{-\alpha i}, \quad \frac{T^{7\beta s_i} - 1}{T^7 - 1} = 1 + T^7 + T^{14} + \dots + T^{7(\beta s_i - 1)}.$$

For a ring homomorphism  $\psi$  satisfying  $\psi(T) = \zeta$  with  $\xi = \zeta^{-\alpha}$  (then  $\xi$  is still a primitive unity, since  $\gcd(\alpha, 7) = 1$ ),

$$\begin{aligned} \tau^\psi(M) &\doteq \left\{ \beta(\zeta^{-\alpha} - 1) \left( \sum_{i=0}^6 s_i \zeta^{-\alpha i} \right) - \alpha \right\} (\zeta^{-\alpha} - 1)^{-2} \\ &= \left\{ \beta(\xi - 1) \left( \sum_{i=0}^6 s_i \xi^i \right) - \alpha \right\} (\xi - 1)^{-2}. \end{aligned}$$

In the 7-th cyclotomic field  $\mathbb{Q}(\zeta_7)$ , using the equalities  $\xi^7 = 1$  and  $1 + \xi + \xi^2 + \xi^3 + \xi^4 + \xi^5 + \xi^6 = 0$ ,

$$\begin{aligned} (\xi - 1) \sum_{i=0}^6 s_i \xi^i &= -3\xi - 2\xi^2 - 3\xi^3 - 3\xi^4 - 2\xi^5 - 3\xi^6 + 16 \\ &= -3\xi - 2\xi^2 - 3\xi^3 - 3\xi^4 - 2\xi^5 - 3\xi^6 + 16 \\ &\quad + 3(1 + \xi + \xi^2 + \xi^3 + \xi^4 + \xi^5 + \xi^6) \\ &= 19 + \xi^2 + \xi^5 \\ &= 19 + \xi^2 + \xi^{-2}. \end{aligned}$$

The Reidemeister–Turaev torsion of Dehn surgery  $M = (BL; \alpha/\beta, 0)$  is

$$(2) \quad \tau^\psi(M) \doteq \{ \beta(\xi^2 + \xi^{-2}) - (\alpha - 19\beta) \} (\xi - 1)^{-2}.$$

Now, suppose that  $M$  is a lens space  $L(p, q)$  with  $p = 49\beta$  (by Step 1) and undecided  $q$ . Then there exist integers  $i, j$  coprime to  $p$  with  $0 < i, j < p$  such that

$$(3) \quad \tau^\psi(M) \doteq (\xi^i - 1)^{-1} (\xi^j - 1)^{-1}.$$

We can assume  $i + j$  is even. We treat with  $i, j \pmod{7}$  ( $i, j \in \{1, 2, 3, 4, 5, 6\}$ ), since  $d = 7$ .

(Step 3) Using Lemma 4.2 on  $d$ -norm with  $d = 7$  on (2) and (3), we have a necessary condition for the Dehn surgery  $M = (BL; \alpha/\beta, 0)$  to be a lens space:

$$N_d(\beta(\xi^2 + \xi^{-2}) - (\alpha - 19\beta)) = 1.$$

Roughly, it means  $r = \alpha/\beta$  is near 19.



(Step 4) We set  $\alpha' = \alpha - 19\beta$ . By (2) and (3), we have

$$\xi \{ \beta(\xi^2 + \xi^{-2}) - \alpha' \} (\xi - 1)^{-2} = \pm \xi^{(i+j)/2} (\xi^i - 1)^{-1} (\xi^j - 1)^{-1}.$$

We regard it as an equality between real value sequence. Without loss of generality, we assume  $0 < i < d/2$  (i.e.,  $i = 1, 2$  or  $3$ ),  $i \leq j$ , and define  $f = (i + j)/2$ ,  $e = (j - i)/2$ . The equality lifts as an identity of symmetric Laurent polynomial

$$(4) \quad (\beta \langle t^2 \rangle - \alpha') (\langle t^f \rangle - \langle t^e \rangle) = \pm (\langle t \rangle - 2),$$

in  $\mathbb{Z}[t, t^{-1}]/(t^7 - 1)$ , where  $\langle t^i \rangle = t^i + t^{-i}$ , as in Definition 4.7. The left-hand side  $F(t)$  is expanded to

$$\beta \langle t^{f+2} \rangle + \beta \langle t^{f-2} \rangle - \alpha' \langle t^f \rangle - \beta \langle t^{e+2} \rangle - \beta \langle t^{e-2} \rangle + \alpha' \langle t^e \rangle.$$

We regard the identity (4) as an equation on  $(f, e)$ : It is a necessary condition on  $(\alpha', \beta)$  for the equation to have a solution  $(f, e)$ . Since  $f \neq e$  is obvious and  $\langle t^4 \rangle = \langle t^3 \rangle$ ,  $\langle t^5 \rangle = \langle t^2 \rangle \pmod{(t^7 - 1)}$ , we only have to consider six cases

$$(f, e) = (1, 0), (2, 0), (3, 0), (2, 1), (3, 1), (3, 2).$$

Note that  $\langle t^{-x} \rangle = \langle t^x \rangle$  and  $\langle t^0 \rangle = 2$ .

$(f, e)$	$F(t)$	$(\alpha', \beta)$
(1, 0)	$\beta \langle t^3 \rangle - 2\beta \langle t^2 \rangle - (\alpha' - \beta) \langle t^1 \rangle + 2\alpha'$	No
(2, 0)	$\beta \langle t^3 \rangle - (\alpha' + 2\beta) \langle t^2 \rangle + 2(\alpha' + \beta)$	No
(3, 0)	$-\alpha' \langle t^3 \rangle - \beta \langle t^2 \rangle + \beta \langle t^1 \rangle + 2\alpha'$	No
(2, 1)	$-\alpha' \langle t^2 \rangle + (\alpha' - \beta) \langle t^1 \rangle + 2\beta$	$(\alpha', \beta) = (0, 1)$
(3, 1)	$-(\alpha' + \beta) \langle t^3 \rangle + \beta \langle t^2 \rangle + \alpha' \langle t^1 \rangle$	No
(3, 2)	$-(\alpha' + \beta) \langle t^3 \rangle + (\alpha' + \beta) \langle t^2 \rangle + \beta \langle t^1 \rangle - 2\beta$	$(\alpha', \beta) = (-1, 1)$

Since  $\alpha' = \alpha - 19\beta$ ,  $(\alpha', \beta) = (0, 1)$  (and  $(-1, 1)$ , respectively) corresponds to  $\alpha/\beta = 19$  (and 18). We have the required conclusion (pointed out in [1]):

Berge's link  $BL$  yields a lens space as  $(BL; r, 0)$  only if  $r = 19$  or  $r = 18$ .

**Acknowledgement** The authors would like to express their sincere gratitude to the organizers of the fruitful seminar "Twisted topological invariants and topology of low-dimensional manifolds". The first author was supported by a grant (No.10801021/a010402) of NSFC. The second author was supported by KAKENHI (Grant-in-Aid for Scientific Research) No.21540072.

REFERENCES

[KTTY] T. Kadokami and Y. Yamada, *Lens space surgeries along certain 2-component links, Park's rational blow down, and Reidemeister-Turaev torsion*, preprint.

[1] J. Berge, *Some knots with surgeries yielding lens spaces*, (Unpublished manuscript, 1990).

[2] S. Boyer and D. Lines, *Surgery formulae for Casson's invariant and extensions to homology lens spaces*, *J. Reine Angew. Math.* **45** (1990), 181–220.

[3] A. Casson and J. Harer, *Some homology lens spaces which bound rational homology balls*, *Pacific J. Math.* **96** (1981), no. 1, 23–36.

[4] W. Franz, *Über die Torsion einer Überdeckung*, *J. Reine Angew. Math.* **173** (1935), 245–254.

[5] T. Kadokami, *Reidemeister torsion and lens surgeries on knots in homology 3-spheres I*, *Osaka J. Math.* **43**, no.4 (2006), 823–837.

- [6] T. Kadokami, *Reidemeister torsion of Seifert fibered homology lens spaces and Dehn surgery*, *Algebr. Geom. Topol.* **7** (2007), 1509–1529.
- [7] T. Kadokami, *Reidemeister torsion and lens surgeries on knots in homology 3-spheres II*, *Top. Appl.* **155**, no.15 (2008), 1699–1707.
- [8] T. Kadokami, N. Maruyama and M. Shimozawa, *Lens surgeries along the  $n$ -twisted Whitehead link*, preprint.
- [9] T. Kadokami and Y. Yamada, *Reidemeister torsion and lens surgeries on  $(-2, m, n)$ -pretzel knots*, *Kobe J. Math.* **23** (2006), 65–78.
- [10] T. Kadokami and Y. Yamada, *A deformation of the Alexander polynomials of knots yielding lens spaces*, *Bull. of Austral. Math. Soc.* **75** (2007), 75–89.
- [11] J. W. Milnor, *A duality theorem for Reidemeister torsion*, *Ann. of Math. (2)*, **76** (1962), 137–147.
- [12] J. Park, *Seiberg-Witten invariants of generalised rational blow-downs*, *Bull. Austral. Math. Soc.* **56** (1997) no. 3, 363–384.
- [13] T. Sakai, *Reidemeister torsion of a homology lens space*, *Kobe J. Math.* **1** (1984), 47–50.
- [14] V. G. Turaev, *Reidemeister torsion in knot theory*, *Russian Math. Surveys*, **41-1** (1986), 119–182.
- [15] V. G. Turaev, *Introduction to Combinatorial Torsions*, Birkhäuser Verlag, (2001).
- [16] L. C. Washington, *Introduction to Cyclotomic Fields*, Graduate Texts in Mathematics **83**, Springer-Verlag, (1982).
- [17] Y. Yamada, *Generalized rational blow-down, torus knots and Euclidean algorithm*, *Arxiv math.GT/0708.2316*.

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY (T. KADOKAMI)  
*E-mail address:* kadokami2007@yahoo.co.jp

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF ELECTRO-COMMUNICATIONS (Y. YAMADA)  
*E-mail address:* yyyamada@e-one.uec.ac.jp