

An asymptotic comparison of differentiable dynamics and tropical geometry

Tsuyoshi Kato

Abstract

In this paper we introduce a new comparison method to give rough asymptotic estimates of different evolutionary dynamics. It uses a kind of scale transform called tropical geometry, which connects automata with real rational dynamics. By this procedure the defining equations are transformed rather than solutions themselves. Real rational dynamics is regarded as an approximation of evolutionary dynamics given by partial differential equations. Two different evolutionary dynamics can be considered when their defining equations are transformed to the same automata at infinity.

1 Introduction

1.A Asymptotic comparison between solutions to different PDEs:

Scaling limits connect several dynamics whose features are often very different mutually. One of particular properties of scaling limits is that in many cases such associations are not injective. When two dynamical systems correspond to the same one by such scaling limits, then one might say that these two dynamics behave by the same way at infinity, and so expect that they will hold some common structural similarity.

Motivated by such aspects, in this paper we study large scale analytic properties of solutions to evolutionary differential equations by use of a particular type of scaling limit. It consists of two steps, where one is to associate discrete dynamics given by real rational functions from differential equations, and the second is automata given by $(\max, +)$ -functions from the rational

dynamics, which appears in *tropical geometry*. Combination of these two steps gives a process of association of automata from PDE. As above one of the important observations for the process is that it is not one to one, and so different differential equations can correspond to the same dynamics by automata. The situation can be interpreted that large (also very small) valued solutions to these PDEs admit mutual analytic relations in some sense, which we would expect to lead us to large scale analysis of structure for classes of differential equations.

In this paper we introduce a new method for study of solutions of non linear partial differential equations. Our main interest here is to obtain relative estimates of asymptotic growth of solutions to different PDEs with respect to higher derivatives and initial conditions.

For $T_0 \in (0, \infty]$, let $u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be a function of class $C^{\alpha+1}$. Then we introduce uniform norm of u of order $\alpha + 1$ by:

$$\|u\|_{\alpha+1} = \max_{\partial_i = \partial_x, \partial_s} \left\{ \left\| \frac{\partial^{\alpha+1} u}{\partial_1 \dots \partial_{\alpha+1}} \right\|_{C^0((0, \infty) \times [0, T_0))} \right\}.$$

Let $c = \inf_{(x,s) \in (0, \infty) \times [0, T_0)} u(x, s) \geq 0$ be the infimum of u . Suppose $u(x, s) \geq c > 0$ is positive. Then we introduce the higher derivative rates by:

$$K(u) \equiv \frac{\|u\|_{\alpha+1}}{c}$$

and call them the *derivative rates of order $\alpha + 1$* . Notice that even when c is sufficiently large, still K can be small when functions u are ‘near’ polynomial of order less than α .

Let $u, v : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be two functions of class $C^{\alpha+1}$. For small $\epsilon > 0$, we introduce the *initial rates*:

$$[u : v]_{\epsilon} \equiv \sup_{(x,s) \in (0, \infty) \times [0, \epsilon^q] \cup (0, \epsilon^p) \times [0, T_0)} \left(\frac{u(x, s)}{v(x, s)} \right)^{\pm 1}.$$

Our method provides with asymptotic relative growth very explicitly for solutions to different differential equations, with respect to their higher derivative rates and initial rates. Let us consider two differential equations of order at most α , $P(u, u_x, u_s, u_{2x}, u_{xs}, \dots) = 0$ and $Q(v, v_x, v_s, v_{2x}, v_{xs}, \dots) = 0$, and take positive solutions $u, v : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ of class $C^{\alpha+1}$. When both P and Q are ‘induced from the same automaton φ ’ which we clarify below, then we verify that there exist constants $C = C(y, r, K)$ which

depend only on the ‘rough structure’ of the differential equations P and Q , which are independent of individual solutions, so that they satisfy uniform bounds:

$$\frac{u(x, s)}{v(x, s)}, \frac{v(x, s)}{u(x, s)} \leq C(x + ks, r, K)$$

when their higher derivative rates and initial rates satisfy bounds $K(u), K(v) \leq K$ and $[u : v]_{(L+1)(2CK)^{-1}} \leq r$ respectively. Here $k, L = \max(l, d)$ and C are explicit constants which arise from scaling limits of these PDE as below.

Let us pick up the required information to determine the constants $C(y, r, K)$. Our basic process is to extract very rough framework of structures of PDE. They are given by n variable rational dynamics of the form:

$$z_{N+1}^{t+1} = f(z_{N-l_0}^{t+1}, \dots, z_N^{t+1}, z_{N-l_1}^t, \dots, z_{N+k_1}^t, \dots, z_{N+k_{d+1}}^{t-d})$$

and scaling parameters $z_N^t = \epsilon^m u(x, s)$ and $(N, t) = (\epsilon^{-p}x, \epsilon^{-q}s)$.

Once such reductions are given, then automata φ are canonically associated, and at this stage, one has chosen several numbers $L = \max(l, d)$, $k, n, D = \max(p, q)$ and C , where $l = \max(l_0, \dots, l_{d+1})$, $k = \max(k_1, \dots, k_{d+1})$, and C are the coefficients of $\alpha + 1$ derivatives in the Taylor expansions, called error constants (3.B.2.2). Relative (max, +)-functions φ are piecewise linear and they are Lipschitz. So one obtains particular two data M and c , where M is the number of the components (1.B) and c is the Lipschitz constants both for φ . In total at the level of defining equations of dynamics, induction of rational functions and scaling parameters determine the above seven data. In section 4 we see that these constants are explicitly calculated or estimated in concrete cases.

On the other hand individual solutions give the constants $[u : v]_\epsilon$ and K , Now $C(y, r, K)$ are in fact given quite explicitly as below. The above numbers are all the data which we need for the above asymptotic estimates among applicable pairs of PDEs.

As a general procedure, the rational dynamics with the scaling parameters above give pairs of partial differential equations $\mathbf{F}(\epsilon, u, u_x, \dots) = 0$ as the leading terms, and the error terms $\mathbf{F}^1(\epsilon, u, u_x, \dots) = 0$ by use of Taylor expansions (1.C).

Let us state our main theorem. The following comparison method discovers very rough structural similarity among different partial differential equations:

Theorem 1.1. *Let f and g be both relatively elementary and increasing functions of n variables, which are mutually tropically equivalent. Let \mathbf{F} and \mathbf{G} be their leading terms of order at most $\alpha \geq 0$, and take positive $C^{\alpha+1}$ solutions $u, v : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ with:*

$$\mathbf{F}(\epsilon, u, u_x, u_s, \dots, u_{\alpha x}, u_{\alpha s}) = 0, \quad \mathbf{G}(\epsilon, v, v_x, v_s, \dots, v_{\alpha x}, v_{\alpha s}) = 0.$$

Assume both u and v are ϵ_0 controlled bounded by C . Then for any $0 < \epsilon \leq \min(\frac{1}{2C}, \epsilon_0)$ and $D = \max(p, q)$, the estimates hold:

$$\left(\frac{u(x, s)}{v(x, s)}\right)^{\pm 1} \leq (2M)^{8 \frac{c^{\epsilon - D(x+ks)+1} - 1}{c-1}} ([u : v]_{(L+1)\epsilon})^{c^{\epsilon - D(x+ks)+n}}.$$

From this we will induce various estimates in concrete examples with respect to their higher derivative rates below. We notice that as a general principle, double exponential growth are optimal in our setting (remark (2) in 2.C).

Now what are the rest is to find suitable pairs of PDEs which arise from the same automata, or in other words, to find suitable rational functions which produce the desired PDEs. This is the key step for our general machinery of discretization of PDE.

Let us see explicit estimates for concrete cases. Here we treat two equations, one is quasi linear equations of order 1, and the other is diffusion equations of order 2. The proofs contain two fundamental techniques, where one is cancellation, and the other is linear deformation both for rational functions. They are obtained by combinations of results in section 4 with lemma 3.3.

Firstly let us consider the quasi linear equations, and choose the uniform norm of second order:

$$\|u\|_2 = \max\left\{\left\|\frac{\partial^2 u}{\partial x^2}\right\|_{C^0}, \left\|\frac{\partial^2 u}{\partial s^2}\right\|_{C^0}, \left\|\frac{\partial^2 u}{\partial x \partial s}\right\|_{C^0}\right\}$$

We put the second derivative rates $K(u) = \frac{\|u\|_2}{\inf_{(x,s) \in (0,\infty) \times [0,T_0)} u(x,s)}$.

Let us fix any positive constant $K_0 > 0$.

Theorem 1.2. *For any $0 < \epsilon \leq 0.1K_0^{-1}$, let $v, u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be C^2 solutions to the quasi linear equations:*

$$v_s + \epsilon v v_x - \frac{1}{2} v^2 = 0, \quad 2u_s + \epsilon u(u_s + u_x) = 0.$$

Suppose their second derivative rates are bounded by $K_0 \geq K(u), K(v)$. Then they satisfy the asymptotic estimates for all $(x, s) \in (0, \infty) \times [0, T_0)$:

$$\left(\frac{u(x, s)}{v(x, s)}\right)^{\pm 1} \leq 40^{2^{\epsilon^{-1}(x+2s)+4}} ([u : v]_{2\epsilon})^{2^{\epsilon^{-1}(x+2s)+3}}.$$

In particular when $u(x, s) \equiv R > 0$ is constant, then the estimates hold:

$$R(40)^{-2^{\epsilon^{-1}(x+2s)+4}} ([v : R]_{2\epsilon})^{-2^{\epsilon^{-1}(x+2s)+3}} \quad (1)$$

$$\leq v(x, s) \leq R(40)^{2^{\epsilon^{-1}(x+2s)+4}} ([v : R]_{2\epsilon})^{2^{\epsilon^{-1}(x+2s)+3}}. \quad (2)$$

Next we treat diffusion equations. Let F be an elementary and increasing function. Here we consider the diffusion equations of the type:

$$u_s = u_{2x} + F(u).$$

There has been various studies for such type of diffusion equations, in relation with blowing up of solutions. We point out two known results.

(1) Let $F(u) = u^l$ for $l = 1, 2, \dots$. If $l = 2$, then any positive solutions to the equation blow up at finite time. For $l \geq 4$, it has global positive solutions for small initial values ([F]). The number 3 is called the *Fujita index* (for one dimensional case).

(2) For all l , if the initial functions take sufficiently large values, then such solutions blow up at finite time ([LN]).

For this case we take the uniform norm of the third derivatives:

$$\|u\|_3 = \max\left\{\left\|\frac{\partial^3 u}{\partial x^3}\right\|_{C^0}, \left\|\frac{\partial^3 u}{\partial s^3}\right\|_{C^0}, \left\|\frac{\partial^3 u}{\partial x^2 \partial s}\right\|_{C^0}, \left\|\frac{\partial^3 u}{\partial x \partial s^2}\right\|_{C^0}\right\}.$$

Then we put the third derivative rates $K(u) \equiv \frac{\|u\|_3}{\inf_{(x,s) \in (0,\infty) \times [0,T_0)} u(x,s)}$.

Firstly let us compare linear diffusion equations with advection-diffusion equations of variable exchange. For the linear case, the corresponding Lipschitz constant is equal to one, and one obtains the exponential asymptotics:

Proposition 1.3. *Let us fix $K_0 > 0$, and choose any $0 < \epsilon \leq (200K_0)^{-1}$. Let $u, v : (0, \infty) \times [0, \infty) \rightarrow (0, \infty)$ be C^3 solutions to the linear equations:*

$$\frac{7}{5}u_s - \frac{193}{40}u_{2x} = 0, \quad \frac{15}{8}\epsilon v_s + \frac{43}{32}v_x - \frac{19}{16}\epsilon^3 v_{2s} = 0.$$

Suppose the third derivative rates satisfy the bounds $K(u), K(v) \leq K_0$. Then they satisfy the exponential asymptotic estimates for all $(x, s) \in (0, \infty) \times [0, \infty)$:

$$\left(\frac{u(x, s)}{v(x, s)}\right)^{\pm 1} \leq 10^{48(\epsilon^{-2}(x+4s)+1)} [u : v]_{5\epsilon}.$$

This is obtained by applying tropical linear deformation of rational functions. Such method is also applied for non linear case as below.

For $1 < a \in \mathbb{Q}$, let us consider the diffusion equations of the form:

$$u_s = u_{2x} + u^a.$$

Let us consider the special solution $v : [0, S_0) \rightarrow (0, \infty)$ given by:

$$v(s) = \frac{c}{(1 - c^{a-1}(a-1)s)^{(a-1)^{-1}}}$$

where $S_0 = \frac{1}{c^{a-1}(a-1)}$. Both v and its third derivative are increasing functions. Thus for any $0 < s_0 < S_0$ and $\alpha = (a-1)^{-1}$, the third derivative rate $K(s_0)$ for the restriction $v : [0, s_0] \rightarrow (0, \infty)$ is bounded by:

$$K(s_0) = \frac{c^{3\alpha-1}(\alpha+1)(\alpha+2)}{\alpha^2(1 - c^{\alpha-1}\alpha^{-1}s_0)^{\alpha+3}}$$

(Remark in 4.B.3). Conversely for any $K(0) \leq K_0 < \infty$, there are unique $s_0 < S_0$ so that the equalities $K_0 = K(s_0)$ hold.

Theorem 1.4. *Let us fix any $K_0 = K(s_0)$. For any $1 < a \in \mathbb{Q}$ and $T_0 \leq s_0$, let $u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be C^3 solutions to the diffusion equations:*

$$u_s - u_{2x} = u^a.$$

Suppose their third derivative rates are bounded by K_0 . Then for any $0 < \epsilon \leq (200K_0)^{-1}$, u satisfy the asymptotic estimates:

$$\left(\frac{u(x, s)}{v(s)}\right)^{\pm 1} \leq 10^{40 \frac{a\epsilon^{-2}(2x+4s)+1}{a-1}} ([u : v]_{5\epsilon})^{a\epsilon^{-2}(2x+4s)+4}.$$

Next we treat diffusion equations of the form:

$$u_s - u_{2x} - u^a - \delta u^b = 0, \quad (1 < a < b, 0 < \delta \ll 1)$$

where we consider the equations of the types:

$$(a, b) = (2, 3), (1 + \alpha^{-1}, 1 + 2\alpha^{-1}), (3, 5)$$

and $0.5 < \alpha < 1$ are any rational numbers. Both the right and left hand side terms touch the Fujita index ($= 3$), and the middle terms cross it. For example it contains the case $(a, b) = (2.5, 4)$.

For $0.5 \leq \alpha \leq 1$ and $c > 0$, let us put:

$$K(s_0) = \frac{c^{3\alpha^{-1}} 5^3 (\alpha + 1)(\alpha + 2)}{6^3 \alpha^2 (1 - c's_0)^{\alpha+3}} \quad (0 \leq s_0 < \frac{1}{c'}), \quad c' = \frac{5c^{\alpha^{-1}}}{6\alpha}, \quad (3)$$

$$(a, b) = (1 + \alpha^{-1}, 1 + 2\alpha^{-1}), \quad \delta = \mu\epsilon^2, \quad \mu = \frac{\alpha + 1}{9\alpha}. \quad (4)$$

For any positive rational numbers $\mu = \frac{p}{q} \in \mathbb{Q}_{>0}$, where $p, q \in \mathbb{N}$ are relatively prime, we put $c_\mu \equiv pq \in \mathbb{Z}_{>0}$.

Let us compare u with the function:

$$v(s) = \frac{c}{(1 - c's)^\alpha}.$$

Let us fix any $K_0 = K(s_0) \geq K(0)$.

Theorem 1.5. *For any $0 < T_0 \leq s_0$ and any $0 < \epsilon \leq (200K_0)^{-1}$, let $u : (0, \infty) \times [0, T_0] \rightarrow (0, \infty)$ be C^3 solutions to the the diffusion equations:*

$$u_s - u_{2x} = u^a + \delta u^b.$$

Suppose their third derivative rates are bounded by K_0 . Then u satisfy the asymptotic estimates:

$$\left(\frac{u(x, s)}{v(s)}\right)^{\pm 1} \leq (2M_\mu)^{8^{\frac{b^{\epsilon^{-2}(2x+4s)+1}-1}{b-1}}} ([u : v]_{5\epsilon})^{b^{\epsilon^{-2}(2x+4s)+4}}$$

where $M_\mu = \max(2 \times 10^3 c_\mu^2, 3 \times 10^4)$.

These results come from a general procedure of comparison method which we will describe below. Our task is to seek for discrete dynamics which induce desired PDE, but such dynamics are not unique. The analytic conditions in these results are heavily depend on choices of such discrete dynamics. Particularly of interest for us is to obtain such estimates by use of $\alpha + 1$

derivatives for larger α . If one can find more suitable discrete dynamics, then one will obtain better asymptotic estimates of solutions.

On the other hand in [KT] we have constructed some examples of pairs of PDEs whose particular solutions do not have such uniform bounds mutually, and so which do not arise from the above procedure. Thus our relations on the uniform bounds for solutions are non trivial among the set of PDEs of 2 variables.

1.A.2 Uniform bounds of higher derivative rates: Functions we consider here are assumed to satisfy uniform boundedness of higher derivatives rates. Functions ‘close’ to polynomials will be particular cases.

It follows from the next lemma that there are pairs of $C^{\alpha+1}$ functions which admit uniformly bounded derivative rates of order $\alpha + 1$, and still break these double exponential estimates on any large bounded domains:

Lemma 1.6. *Let us take any $\alpha \geq 1$ and any large $C_0 \gg 0$. Then there are constants $c_{\alpha+1}$ independent of C_0 , and pairs of $C^{\alpha+1}[0, 2)$ functions u, v whose derivative rates of order $\alpha + 1$ are bounded by $c_{\alpha+1}$, so that the estimates hold:*

$$\frac{v(s)}{u(s)} \begin{cases} = 1 & 0 \leq s \leq 1, \\ \geq C_0 & 2 - (2C_0)^{-1} \leq s < 2. \end{cases}$$

Proof: Let $u : [0, 2) \rightarrow (0, 2C_0]$ be the linear function by $u(s) = C_0(2 - s)$. $\alpha + 1$ derivatives of u vanish for $\alpha \geq 1$, and so all higher derivative rates are zero. We construct $v : [0, 2) \rightarrow [0.5C_0, 2C_0]$ which satisfies:

$$v(s) = \begin{cases} u(s) & 0 \leq s \leq 1, \\ 0.5C_0 & 1.5 \leq s \leq 2. \end{cases}$$

Let $w : [0, 2) \rightarrow [0.5, 2]$ be a smooth and non increasing function which satisfies $w(s) = 2 - s$ for $0 \leq s \leq 1$, and $\equiv 0.5$ for $1.5 \leq s \leq 2$. Then there are constants $c_{\alpha+1}$ so that its derivative rates of order $\alpha + 1$ are bounded by $c_{\alpha+1}$. Let us put $v : [0, 2) \rightarrow [0.5C_0, 2C_0]$ by $v(s) = C_0w(s)$. Then v are the desired functions, since they have the same higher derivative rates as w . This completes the proof.

When the domains for (x, s) are unbounded, what we are focusing by those double exponential estimates, is not behavior at infinity for (x, s) . In fact it follows from the assumption of uniform boundedness of higher

derivative rates that u will grow at most exponentially. So constrains of the defining equations have an effective influence on bounded regions.

1.B Real rational dynamics and tropical geometry: A relative $(\max, +)$ -function φ is a piecewise linear function of the form:

$$\varphi(\bar{x}) = \max(\alpha_1 + \bar{a}_1\bar{x}, \dots, \alpha_m + \bar{a}_m\bar{x}) - \max(\beta_1 + \bar{b}_1\bar{x}, \dots, \beta_l + \bar{b}_l\bar{x})$$

where $\bar{a}_l\bar{x} = \sum_{i=1}^n a_l^i x_i$, $\bar{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$, $\bar{a}_l = (a_l^1, \dots, a_l^n)$, $\bar{b} \in \mathbb{Z}^n$ and $\alpha_i, \beta_i \in \mathbb{R}$. We say that the multiple integer $M \equiv ml$ is the *number of the components* of φ .

Correspondingly tropical geometry associates the parametrized rational function given by (see [Mi]):

$$f_t(\bar{z}) \equiv \frac{k_t(\bar{z})}{h_t(\bar{z})} = \frac{\sum_{k=1}^m t^{\alpha_k} \bar{z}^{\bar{a}_k}}{\sum_{k=1}^l t^{\beta_k} \bar{z}^{\bar{b}_k}}$$

where $\bar{z}^{\bar{a}} = \prod_{i=1}^n z_i^{a_i}$, $\bar{z} = (z_1, \dots, z_n) \in \mathbb{R}_{>0}^n$. We say that f_t above is a *relative elementary* function. We say that both terms $h_t(\bar{z}) = \sum_{k=1}^l t^{\beta_k} \bar{z}^{\bar{b}_k}$ and $k_t(\bar{z}) = \sum_{k=1}^m t^{\alpha_k} \bar{z}^{\bar{a}_k}$ are just elementary functions.

These two functions φ and f_t admit one to one correspondence between their presentations. Moreover the defining equations are transformed by two steps, firstly taking conjugates by \log_t and secondly by letting $t \rightarrow \infty$. Notice that when all a_i and b_j are zero, then the corresponding f_t are t independent.

In some cases the same $(\max, +)$ function admits different presentations, while the corresponding rational functions are mutually different. For example for $\varphi(x) \equiv \max(x, x) = x \equiv \psi(x)$, the corresponding rational functions $f_t(z) = 2z$ and $g_t(z) = z$ are mutually different. We call such a pair of rational functions *tropically equivalent*.

Let $f_t : \mathbb{R}_{>0}^n \rightarrow (0, \infty)$ be a rational function, and consider the discrete dynamics defined by:

$$z_N = f_t(z_{N-n}, \dots, z_{N-1}), \quad N \geq n$$

with initial values $(z_0, \dots, z_{n-1}) \in \mathbb{R}_{>0}^n$. One can regard that tropically equivalent rational functions determine the same dynamics at infinity.

Let us put:

$$P_N(c) = \begin{cases} \frac{c^{N-n+1}-1}{c-1} & c > 1, \\ N - n + 1 & c = 1. \end{cases}$$

For a relative elementary function f_t , let $c_f \geq 1$ be the Lipschitz constant and M_f be the number of the components with respect to the corresponding $(\max, +)$ -function.

Our basic analysis on the orbits is given by the following (corollary 2.8):

Lemma 1.7. *Let g_t be tropically equivalent to f_t , and $\{z_N\}_N$ and $\{w_N\}_N$ be the orbits for f_t and g_t with the initial values $\bar{z}_0 = (z_0, \dots, z_{n-1})$ and $\bar{w}_0 = (w_0, \dots, w_{n-1})$ respectively. Then the estimates hold:*

$$\left(\frac{z_N}{w_N}\right)^\pm \leq M^{4P_N(c)} \left[\max_{0 \leq i \leq n-1} \left(\frac{z_i}{w_i}\right)^{\pm 1} \right]^{c^N}$$

where $c = \max(c_f, c_g)$ and $M = \max(M_f, M_g)$.

If the initial values are the same, then uniform estimates hold (prop 2.3):

$$\left(\frac{z_N}{w_N}\right)^\pm \leq M^{2P_N(c)}.$$

One particular feature is that when the Lipschitz constant is equal to 1, then the above inequalities give the exponential estimates, while for $c > 1$, they are double exponential. The former is applied for the estimates of solutions to linear PDEs.

When one considers evolutionary discrete dynamics, a parallel estimates are given. An evolutionary discrete dynamics is given by flows of the form $\{z_N^t\}_{t, N \geq 0}$, where t is time parameter. A general equation of evolutionary discrete dynamics is of the form:

$$z_{N+1}^{t+1} = f(z_{N-l_0}^{t+1}, \dots, z_N^{t+1}, z_{N-l_1}^t, \dots, z_{N+k_1}^t, \dots, z_{N+k_{d+1}}^{t-d})$$

where $l_i, k_j \geq 0$, $N \geq \max(l_0, \dots, l_{d+1})$ and $t \geq d$, with initial values:

$$\bar{z}_0^0 \equiv \{z_a^t\}_{0 \leq a \leq \max(l_0, \dots, l_{d+1}), t=0,1,\dots} \cup \{z_N^h\}_{0 \leq h \leq d, N=0,1,\dots}$$

Let us take g tropically equivalent to f , and consider the dynamics $\{w_n^t\}$ defined by g with any initial values \bar{w}_0^0 . Then we put the *initial rates* by:

$$[\bar{z}_0^0 : \bar{w}_0^0] \equiv \sup_{0 \leq a \leq \max(l_0, \dots, l_{d+1}), b=0,1,\dots, \text{ or } a=0,1,\dots, 0 \leq b \leq d} \left\{ \frac{z_a^b}{w_a^b}, \frac{w_a^b}{z_a^b} \right\}.$$

Let us put $l = \max(l_0, l_1, \dots, l_{d+1})$, $k = \max(k_1, \dots, k_{d+1})$ and

$$A(N, t) \equiv (t - d - 1)k + N - l + n - 1$$

for $N \geq l + 1$ and $t \geq d + 1$.

Proposition 1.8. (1) Let f and g be tropically equivalent. Then any orbits $\{z_N^t\}_N$ and $\{w_N^t\}_N$ for f and g with the initial values \bar{z}_0^0 and \bar{w}_0^0 respectively, satisfy the estimates:

$$\left(\frac{z_N^t}{w_N^t}\right)^\pm \leq M^{4P_{A(N,t)}(c)}[\bar{z}_0^0 : \bar{w}_0^0]^{cA(N,t)}$$

where $c = \max(c_f, c_g)$ and $M = \max(M_f, M_g)$.

(2) Let f, f', g, g' be four relative elementary functions, and assume that they are all monotone increasing and all tropically equivalent. Let $\{v_N^t\}_{N,t}$ and $\{u_N^t\}_{N,t}$ be positive sequences so that these satisfy the estimates:

$$f'(v_{N-l_0}^{t+1}, \dots, v_{N+k_{d+1}}^{t-d}) \leq v_{N+1}^{t+1} \leq f(v_{N-l_0}^{t+1}, \dots, v_{N+k_{d+1}}^{t-d}), \quad (5)$$

$$g'(u_{N-l_0}^{t+1}, \dots, u_{N+k_{d+1}}^{t-d}) \leq u_{N+1}^{t+1} \leq g(u_{N-l_0}^{t+1}, \dots, u_{N+k_{d+1}}^{t-d}) \quad (6)$$

for all N, t . Then the ratios satisfy the uniform estimates:

$$\left(\frac{v_N^t}{u_N^t}\right)^{\pm 1} \leq M^{8P_{A(N,t)}(c)}[\bar{u}_0^0 : \bar{v}_0^0]^{cA(N,t)}.$$

Here also if the Lipschitz constants c are equal to 1, then the above two estimates are at most exponential, while for the case $c > 1$, they are double exponential.

Such general form will allow us to treat wider classes of PDE. But for concrete cases, we use evolutionary discrete dynamics only of the forms:

$$z_{N+1}^{t+1} = \begin{cases} f(z_{N-1}^{t+1}, z_N^t, z_{N+2}^t) & \text{for quasi linear equations,} \\ f(z_N^t, z_{N+4}^t, z_{N-4}^{t-1}, z_{N-1}^{t-4}) & \text{for diffusion equations.} \end{cases}$$

For the former $l = 1, k = 2, d = 0$, and for the latter $l = k = d = 4$. So they are given by:

$$A(N, t) = \begin{cases} 2t + N - 1 & (N \geq 2, t \geq 1), \\ 4t + N - 21 & (N \geq 5, t \geq 5). \end{cases}$$

1.C Rough approximations by discrete dynamics: Let us describe our general procedure for approximating solutions to differential equations by discrete dynamics, and outline how to verify theorems in 1.A.

Let us consider a $C^{\alpha+1}$ function $u : (0, \infty) \rightarrow (0, \infty)$, and for $1 \leq |i| \leq n-1$, take the Taylor expansions:

$$u(x+i\epsilon) = u(x) + i\epsilon u_x + \frac{(i\epsilon)^2}{2} u_{2x} + \dots + \frac{(i\epsilon)^\alpha}{\alpha!} u_{\alpha x} + \frac{(i\epsilon)^{(\alpha+1)}}{(\alpha+1)!} u_{(\alpha+1)x}(\xi_i).$$

Then for small $\epsilon > 0$ and $N = 0, 1, 2, \dots$, let us put

$$z_N \equiv \epsilon u(\epsilon N) = \epsilon u(x), \quad (N = \frac{x}{\epsilon}).$$

Let $f = \frac{k}{h} : \mathbb{R}_{>0}^n \rightarrow (0, \infty)$ be a relative elementary function of n variables, where both h and k are elementary, and consider the discrete dynamics defined by $w_{N+1} = f(w_{N-n+1}, \dots, w_N)$ with the initial value $w_i = \epsilon u(\epsilon i) > 0$ for $0 \leq i \leq n-1$. Our basic idea is to regard that the sequence $\{w_N\}_N$ would approximate the orbit $\{z_N\}_N$.

So let us consider the difference and insert the Taylor expansions:

$$z_{N+1} - f(z_{N-n+1}, \dots, z_N) = \epsilon u(x + \epsilon) - f(\epsilon u(x - (n-1)\epsilon), \dots, \epsilon u(x)) \quad (7)$$

$$= \epsilon(u + \epsilon u_x + \frac{\epsilon^2}{2} u_{2x} + \dots) - f(\epsilon(u - (n-1)\epsilon u_x + \dots), \dots, \epsilon u) \quad (8)$$

$$= \frac{\epsilon F^1(u) + \epsilon^2 F^2(u_x) + \epsilon^3 F^3(u, u_x) + \dots + \epsilon^m F^{m'}(u, \dots, u_{(\alpha+1)x}(\xi)) + \dots}{h(\epsilon u(x - (n-1)\epsilon), \dots, \epsilon u(x))} \quad (9)$$

$$(10)$$

where F^k are monomials.

For any finite subset $A \subset \{1, 2, 3, \dots\}$, let us divide the expanded sum into two terms as:

$$= \frac{\sum_{i \in A} \epsilon^{s_i} F^{s'_i}(u, u_x, \dots, u_{\alpha x})}{h(\epsilon u(x - (n-1)\epsilon), \dots)} + \frac{\sum_{j \in A^c} \epsilon^{s_j} F^{s'_j}(u, u_x, \dots, u_{(\alpha+1)x}(\xi))}{h(\epsilon u(x - (n-1)\epsilon), \dots)} \quad (11)$$

$$\equiv \mathbf{F}(\epsilon, u, u_x, \dots, u_{\alpha x}) + \epsilon^2 \mathbf{F}^1(\epsilon, u, u_x, \dots, u_{(\alpha+1)x}(\xi_1), \dots, u_{(\alpha+1)x}(\xi_{n-1})) \quad (12)$$

We say that \mathbf{F} and \mathbf{F}^1 are the *leading* and *error* terms respectively. Once one has chosen a relative elementary function f , then the above process determines a PDE defined by \mathbf{F} , while tropical geometry gives an automaton by a $(\max, +)$ function φ . So f plays a role of a bridge to connect between PDE and automaton.

Let us define ϵ variation of \mathbf{F}^1 by:

$$\|\mathbf{F}^1(\epsilon, u, u_x, \dots, u_{\alpha x}, u_{(\alpha+1)x}(\xi_1), \dots, u_{(\alpha+1)x}(\xi_{n-1}))\|_{\epsilon}(x) \equiv \quad (13)$$

$$\sup_{\mu_i - x \in I(n, \epsilon)} |\mathbf{F}^1(\epsilon, u(x - \epsilon), \dots, u_{\alpha x}(x - \epsilon), u_{(\alpha+1)x}(\mu_1), \dots, u_{(\alpha+1)x}(\mu_{n-1}))| \quad (14)$$

where $I(n, \epsilon) = [-n\epsilon, 0]$ is the fluctuation interval.

Let us say that a $C^{\alpha+1}$ function $u : (0, \infty) \rightarrow (0, \infty)$ is ϵ controlled, if there is some constant $C > 0$ so that ϵ variation of \mathbf{F}^1 satisfy the pointwise estimates for all $x \in (0, \infty)$:

$$Cu(x) \geq \|\mathbf{F}^1(\epsilon, u, u_x, \dots, u_{\alpha x}, u_{(\alpha+1)x}(\xi_1), \dots, u_{(\alpha+1)x}(\xi_{n-1}))\|_{\epsilon}(x).$$

For two functions u, v , we put their initial rates by $[u : v]_{\epsilon} \equiv \sup_{x \in (0, \epsilon]} \left(\frac{u(x)}{v(x)}\right)^{\pm 1}$.

Now we state the first estimates. Let f and g be relatively elementary functions, and \mathbf{F} and \mathbf{G} be their corresponding leading terms. Recall that associated with f are the Lipschitz constant $c_f \geq 1$ and the number of the components M_f . Let us put $c = \max(c_f, c_g)$ and $M = \max(M_f, M_g)$.

Proposition 1.9. *Let f and g be both relatively elementary and increasing functions of n variables, which are mutually tropically equivalent. Let \mathbf{F} and \mathbf{G} be their leading terms of order at most $\alpha \geq 0$, and take positive $C^{\alpha+1}$ solutions $u, v : (0, \infty) \rightarrow (0, \infty)$ with:*

$$\mathbf{F}(\epsilon, u, u_x, \dots, u_{\alpha x}) = 0, \quad \mathbf{G}(\epsilon, v, v_x, \dots, v_{\alpha x}) = 0.$$

Assume both u and v are ϵ_0 controlled bounded by C . Then for any $0 < \epsilon \leq \min(\frac{1}{2C}, \epsilon_0)$, the estimates hold:

$$\left(\frac{u(x)}{v(x)}\right)^{\pm 1} \leq (2M)^{8\frac{\epsilon^{-1}x+1}{c-1}} ([u : v]_{n\epsilon})^{\epsilon^{-1}x+1}.$$

One can proceed parallelly for evolutionary case. Let f be a relatively elementary function, and consider the evolutionary discrete dynamics defined by the equation $z_{N+1}^{t+1} = f(z_{N-l_0}^{t+1}, \dots, z_{N+k_{d+1}}^{t-d})$.

Let us take a $C^{\alpha+1}$ function $u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$, and introduce another parameters by:

$$\epsilon^m u(x, s) = z_N^t, \quad N = \frac{x}{\epsilon^p}, \quad t = \frac{s}{\epsilon^q}$$

where $p, q \geq 1$ and $m \geq 0$ are integers. By the same way as one variable case, one takes the Taylor expansion, and take the difference:

$$z_{N+1}^{t+1} - f(z_{N-l_0}^{t+1}, \dots, z_{N+k_{d+1}}^{t-d}) \quad (15)$$

$$= \epsilon^m u(x + \epsilon^p, s + \epsilon^q) - \quad (16)$$

$$f(\epsilon^m u(x - l_0 \epsilon^p, s + \epsilon^q), \dots, \epsilon^m u(x + k_{d+1} \epsilon^p, s - d \epsilon^q)) \quad (17)$$

$$= \frac{\epsilon^m F^1(u) + \epsilon^{m+p} F^2(u_x) + \epsilon^{m+q} F^3(u_s) + \epsilon^{2m+p} F^4(u, u_x) + \dots}{h(\epsilon^m u(x - l_0 \epsilon^p, s + \epsilon^q), \dots, \epsilon u(x + k_{d+1} \epsilon^p, s - d \epsilon^q))} \quad (18)$$

$$= \mathbf{F}(\epsilon, u, u_x, u_s, u_{xs}, \dots, u_{\alpha x}, u_{\alpha s}) \quad (19)$$

$$+ \epsilon^{m+1} \mathbf{F}^1(\epsilon, u(x, s), u_x(x, s), \dots, u_{\alpha x}(x, s), \quad (20)$$

$$u_s(x, s), \dots, u_{\alpha s}(x, s), \{u_{(\alpha+1)x}(\xi_{ij}), \dots, u_{(\alpha+1)s}(\xi_{ij})\}_{i,j}). \quad (21)$$

By the same way as before one defines the ϵ variation $\|\mathbf{F}^1\|_\epsilon(x, s)$ and ϵ_0 controlledness (3.B). Combining this construction with proposition 1.9, one obtains theorem 1.1.

Our basic process goes as follows. Firstly we choose a PDE $\mathbf{F}(u, u_x, u_s, \dots) = 0$, and fix scaling parameters. Then find a relative elementary function f which induces \mathbf{F} as its leading term. Next take another relative elementary g which is tropically equivalent to f . Then by use of the same scaling parameters, it induces its leading term \mathbf{G} . Finally for two solutions u and v with $\mathbf{F}(u, u_x, u_s, \dots) = 0$ and $\mathbf{G}(v, v_x, v_s, \dots) = 0$ respectively, we seek for analytic conditions to both u and v which insure ϵ_0 controlledness. Even though choice of f and g are rather flexible, whether one could find some reasonable conditions for solutions depends on choices of these functions.

2 Discrete dynamics and tropical geometry

2.A Elementary functions: A relative $(\max, +)$ -function φ is a piecewise linear function of the form:

$$\varphi(\bar{x}) = \max(\alpha_1 + \bar{a}_1 \bar{x}, \dots, \alpha_m + \bar{a}_m \bar{x}) - \max(\beta_1 + \bar{b}_1 \bar{x}, \dots, \beta_l + \bar{b}_l \bar{x})$$

where $\bar{a}_k \bar{x} = \sum_{i=1}^n a_k^i x_i$, $\bar{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$, $\bar{a}_k = (a_k^1, \dots, a_k^n)$, $\bar{b}_k \in \mathbb{Z}^n$ and $\alpha_k, \beta_k \in \mathbb{R}$.

For each relative $(\max, +)$ function φ as above, we associate a parametrized rational function by:

$$f_t(\bar{z}) = \frac{\sum_{k=1}^m t^{\alpha_k} \bar{z}^{\bar{a}_k}}{\sum_{k=1}^l t^{\beta_k} \bar{z}^{\bar{b}_k}}$$

where $\bar{z}^{\bar{a}_k} = \prod_{i=1}^n z_i^{a_k^i}$, $\bar{z} = (z_1, \dots, z_n) \in \mathbb{R}_{>0}^n = \{(w_1, \dots, w_n) : w_i > 0\}$.

We say that f_t above is a *relative elementary function*. Notice that any relative elementary functions take positive values for $\bar{z} \in \mathbb{R}_{>0}^n$.

We say that the integer:

$$M \equiv ml$$

is the number of the components.

We say that $f_t(\bar{z}) = \sum_{k=1}^m t^{\alpha_k} \bar{z}^{\bar{j}_k}$ is an elementary rational function ([K2]). The corresponding (max, +)-function is given by $\varphi(\bar{x}) = \max(\alpha_1 + \bar{j}_1 \bar{x}, \dots, \alpha_m + \bar{j}_m \bar{x})$, and in this case m is the number of the components.

These two functions φ and f_t are connected passing through some intermediate functions φ_t ([LiM], [Mi]). Let us describe it shortly below. For $t > 1$, there is a family of semirings R_t which are all the real number \mathbb{R} as sets. The multiplications and the additions are respectively given by $x \oplus_t y = \log_t(t^x + t^y)$ and $x \otimes_t y = x + y$. As $t \rightarrow \infty$ one obtains the equality:

$$x \oplus_\infty y = \max(x, y).$$

By use of R_t as coefficients, one has *relative R_t -polynomials*:

$$\varphi_t(\bar{x}) = (\alpha_1 + \bar{a}_1 \bar{x}) \oplus_t \dots \oplus_t (\alpha_m + \bar{a}_m \bar{x}) - (\beta_1 + \bar{b}_1 \bar{x}) \oplus_t \dots \oplus_t (\beta_l + \bar{b}_l \bar{x})$$

The limit is given by the relative (max, +) function above:

$$\lim_{t \rightarrow \infty} \varphi_t(\bar{x}) = \varphi(\bar{x}).$$

Let us put $\text{Log}_t : \mathbb{R}_{>0}^n \rightarrow \mathbb{R}^n$ by $(z_1, \dots, z_n) \rightarrow (\log_t z_1, \dots, \log_t z_n)$. Then φ_t and f_t satisfy the following relation:

Proposition 2.1 (LiM,V). $f_t \equiv (\log_t)^{-1} \circ \varphi_t \circ \text{Log}_t : \mathbb{R}_{>0}^n \rightarrow (0, \infty)$ is the relative elementary function $f_t(\bar{z}) = \sum_{k=1}^m t^{\alpha_k} \bar{z}^{\bar{a}_k} / \sum_{k=1}^l t^{\beta_k} \bar{z}^{\bar{b}_k}$.

These functions φ , φ_t and f_t admit one to one correspondence between their presentations. We say that φ is the corresponding (max, +)-function to f_t . Notice that any relative (max, +) functions of the form $\varphi(\bar{x}) = \max(\bar{a}_1 \bar{x}, \dots, \bar{a}_m \bar{x}) - \max(\bar{b}_1 \bar{x}, \dots, \bar{b}_l \bar{x})$ correspond to t -independent relative elementary functions f .

2.B Discrete dynamics: Let $f_t : \mathbb{R}_{>0}^n \rightarrow (0, \infty)$ be a relative elementary function, and φ be the corresponding (max, +)-function. Let us consider the discrete dynamics defined by:

$$z_N = f_t(z_{N-n}, \dots, z_{N-1}), \quad N \geq n$$

with initial values $(z_0, \dots, z_{n-1}) \in \mathbb{R}_{>0}^n$. These orbits $\{z_N\}_N$ admit some asymptotic controll passing through tropical geometry, which we describe below. Let us compare the orbits $\{x_N\}_N$ with $\{z_N\}_N$, which are determined by:

$$x_N = \varphi(x_{N-n}, \dots, x_{N-1})$$

with the initial values $x_0 = \log_t z_0, \dots, x_{n-1} = \log_t z_{n-1}$. For this, we introduce the intermediate dynamics:

$$x'_N = \varphi_t(x'_{N-n}, \dots, x'_{N-1})$$

with the same initial data $x'_0 = \log_t z_0, \dots, x'_{n-1} = \log_t z_{n-1}$.

By proposition 2.1, two orbits $\{z_N\}_N$ and $\{x'_N\}_N$ are conjugate each other as $x'_N = \log_t z_N$ for all $N = 0, 1, \dots$. Since $\lim_{t \rightarrow \infty} \varphi_t = \varphi$ holds, one may think $\{\log_t z_N\}_N$ ‘approximate’ $\{x_N\}_N$ in some sense.

Let φ and ψ be two relative $(\max, +)$ -functions with n variables. Then ψ is *equivalent* to φ , if they are the same as maps, $\varphi(x_1, \dots, x_n) = \psi(x_1, \dots, x_n)$ for all $(x_1, \dots, x_n) \in \mathbb{R}^n$ (but possibly they can have different presentations).

Definition 2.1 (K2). *Let f_t and g_t be two relative elementary functions. g_t is tropically equivalent to f_t , if the corresponding relative $(\max, +)$ -functions φ and ψ are equivalent.*

Remarks: (1) If the pointwise estimate $\varphi \geq \varphi'$ holds, then $\psi = \max(\varphi, \varphi')$ and φ are equivalent. Let f_t and g_t be the corresponding relative elementary functions to φ and φ' respectively. Then $h_t \equiv f_t + g_t$ is tropically equivalent to f_t .

(2) For any relative elementary f_t and positive rational numbers $0 < \alpha = \frac{n}{m} \in \mathbb{Q}$, αf_t is tropically equivalent to f_t . In fact let φ correspond to f_t . Then $n f_t$ are tropically equivalent to f_t , since $n f_t$ correspond to $\max(\varphi, \dots, \varphi) = \varphi$ (n times). Similarly $\frac{1}{m} f_t$ are also tropically equivalent to f_t .

(3) For any tropically equivalent pairs of relative elementary functions f_t and g_t , the corresponding $(\max, +)$ -functions φ and ψ have the same Lipschitz constant $c > 0$, since they are the same as maps. On the other hand they may have different numbers of the components M and M' since it depends on their presentations. For example if f_t has M number of the components, then $\frac{n}{m} f_t$ has nmM number of the components.

(4) For our purposes in this paper, it is enough to treat the case that the Lipschitz constants c for φ is larger or equal to 1, and later on we will assume the bounds $c \geq 1$.

2.C Basic estimates and Lipschitz constants: Let $f_t : \mathbb{R}_{>0}^n \rightarrow (0, \infty)$ be a relative elementary function. Take initial values $(z_0, \dots, z_{n-1}) \in \mathbb{R}_{>0}^n$, and consider the orbits $\{z_N\}_{N=0}^\infty$ defined by $z_N = f_t(z_{N-n}, \dots, z_{N-1})$ for $N \geq n$. Let g_t be another relative elementary function, and consider its orbit $\{w_N\}_N$ with the same initial values $w_i = z_i$ for $0 \leq i \leq n-1$.

In order to estimate their asymptotic rates $(\frac{z_N}{w_N})^{\pm 1}$ in detail, we use the metric on \mathbb{R}^n given by:

$$d((x_0, \dots, x_{n-1}), (y_0, \dots, y_{n-1})) \equiv \max_{0 \leq i \leq n-1} \{|x_i - y_i|\}$$

(which is of course equivalent to the standard one.)

Lemma 2.2. *Let $f_t = f$ be t -independent, relative elementary and linear. Then the corresponding $(\max, +)$ -function φ has its Lipschitz constant bounded by 1.*

Proof: This follows immediately, if one checks the estimates carefully. One can express $\varphi(x_0, \dots, x_{n-1}) = \max(\alpha_1 + x_{i_1}, \dots, \alpha_{n-1} + x_{i_{n-1}}) - \max(0, \dots, 0)$. Let $\varphi(x_0, \dots, x_{n-1}) = \alpha_1 + x_{i_1} \geq \varphi(y_0, \dots, y_{n-1}) = \alpha_2 + y_{i_2}$. Then the estimates hold:

$$|\varphi(x_0, \dots, x_{n-1}) - \varphi(y_0, \dots, y_{n-1})| = \alpha_1 + x_{i_1} - (\alpha_2 + y_{i_2}) \quad (22)$$

$$\leq \alpha_1 + x_{i_1} - (\alpha_1 + y_{i_1}) = x_{i_1} - y_{i_1} \leq \max_{0 \leq i \leq n-1} \{|x_i - y_i|\}. \quad (23)$$

This completes the proof.

In general we have double exponential estimates for $(\frac{z_N}{w_N})^{\pm 1}$ as below, but in a special case that the Lipschitz constants of the corresponding $(\max, +)$ -functions are equal to 1, they can be improved to be just exponential. This happens when one considers linear PDE.

Let us put

$$P_N(c) = \begin{cases} \frac{c^{N-n+1}-1}{c-1} & c > 1, \\ (N-n+1) & c = 1. \end{cases}$$

For a relative elementary function f_t , let c_f be the Lipschitz constant and M_f be the number of the components with respect to the corresponding $(\max, +)$ -function.

Proposition 2.3. f_t and g_t are tropically equivalent, if and only if any orbits with the same initial values satisfy uniformly bounded rates:

$$\left(\frac{z_N}{w_N}\right)^{\pm 1} \equiv \frac{z_N}{w_N}, \frac{w_N}{z_N} \leq M^{2P_N(c)}, \quad (N \geq n)$$

where $c = \max(c_f, c_g)$ and $M = \max(M_f, M_g)$.

For the proof, we use the next lemma.

Let $\varphi(\bar{x}) = \max(\alpha_1 + \bar{a}_1 \bar{x}, \dots, \alpha_m + \bar{a}_m \bar{x}) - \max(\beta_1 + \bar{b}_1 \bar{x}, \dots, \beta_l + \bar{b}_l \bar{x})$ and φ_t be the corresponding functions to f_t . For the same initial values $x_0 = x'_0, \dots, x_{n-1} = x'_{n-1}$, let us denote the orbits by $\{x_N\}_N$ and $\{x'_N\}_N$ for φ and φ_t respectively.

We will improve lemma 2.2 in [K2] slightly.

Lemma 2.4. Let $c \geq 1$ and M be the Lipschitz constant and the number of the components for φ respectively. Then the estimates hold:

$$|x_N - x'_N| \leq P_N(c) \log_t M.$$

Proof: One can obtain the following estimates easily ([K2] lemma 2.1(1)):

$$|\varphi(x_0, \dots, x_{n-1}) - \varphi_t(x_0, \dots, x_{n-1})| \leq \log_t M.$$

Let us denote $\bar{x}_N = (x_N, \dots, x_{N+n-1}) \in \mathbb{R}^n$. Thus $x_{N+n} = \varphi(\bar{x}_N)$ hold for all $N \geq 0$. Similar for \bar{x}'_N .

Firstly one has the estimates $|x'_n - x_n| \leq \log_t M$ as above.

Since φ is c -Lipschitz and $\bar{x}_1 - \bar{x}'_1 = (0, \dots, 0, x_n - x'_n)$, the estimates:

$$|x_{n+1} - x'_{n+1}| = |\varphi(\bar{x}_1) - \varphi_t(\bar{x}'_1)| \tag{24}$$

$$\leq |\varphi(\bar{x}_1) - \varphi(\bar{x}'_1)| + |\varphi(\bar{x}'_1) - \varphi_t(\bar{x}'_1)| \tag{25}$$

$$\leq c|\bar{x}_1 - \bar{x}'_1| + \log_t M \leq (c+1) \log_t M \tag{26}$$

hold. Next we have estimates:

$$|\varphi(\bar{x}_2) - \varphi(\bar{x}'_2)| \leq c \max(|x_{n+1} - x'_{n+1}|, |x_n - x'_n|) \leq c(c+1) \log_t M, \tag{27}$$

$$|x_{n+2} - x'_{n+2}| = |\varphi(\bar{x}_2) - \varphi_t(\bar{x}'_2)| \tag{28}$$

$$\leq |\varphi(\bar{x}_2) - \varphi(\bar{x}'_2)| + |\varphi(\bar{x}'_2) - \varphi_t(\bar{x}'_2)| \leq [c(c+1) + 1] \log_t M \tag{29}$$

The rest is just the repetition of the same process. Now suppose $c > 1$. Then by a direct calculation, one obtains the estimates:

$$|x_N - x'_N| \leq \frac{c^{N-n+1} - 1}{c - 1} \log_t M.$$

On the other hand when $c = 1$, then $|x_N - x'_N| \leq (N - n + 1) \log_t M$ hold. This completes the proof.

Proof of proposition 2.3: The proof is almost the same as theorem 2.1 in [K2], but for convenience we will include only if part.

Let φ and ψ be the relative $(\max, +)$ -functions corresponding to f_t and g_t respectively. For the same initial values $x_i = y_i = \log_t z_i$, $0 \leq i \leq n - 1$, let us denote the corresponding orbits by $\{x_N\}_N$ and $\{y_N\}_N$. We also put $x'_N = \log_t(z_N)$ and $y'_N = \log_t(w_N)$ respectively. Thus $\{x'_N\}_N$ is the orbit for φ_t and $\{y'_N\}_N$ is for ψ_t .

By lemma 2.4, the estimates:

$$|x_N - x'_N|, |y_N - y'_N| \leq P_N(c) \log_t M$$

hold. Suppose f_t and g_t are tropically equivalent, and so φ and ψ are the same as maps. Thus $x_N = y_N$ hold, and so we have the estimates:

$$\log_t \left(\frac{z_N}{w_N} \right)^\pm \leq |\log_t(z_N) - \log_t(w_N)| = |x'_N - y'_N| \leq 2P_N(c) \log_t M.$$

Thus we have the estimates:

$$\left(\frac{z_N}{w_N} \right)^\pm \leq \max \left(\frac{z_N}{w_N}, \frac{w_N}{z_N} \right) \leq M^{2P_N(c)}.$$

This completes the proof.

Remarks: (1) In order to determine z_N for $N \geq n$, one needs to iterate $N - n + 1$ times to apply function f_t . One can say that ratios between $N - n + 1$ times iterations of f_t and g_t are at most uniformly double exponential rates.

(2) Such double exponential estimates are optimal between tropically equivalent functions. Let us consider two dynamics for $l, k \geq 1$:

$$z_N = f(z_{N-1}) = z_{N-1}^l, \quad w_N = g(w_{N-1}) = 2w_{N-1}^k.$$

If $l = k$ holds, then f and g are tropically equivalent. Let $z_0 = w_0$ be initial values. Then a direct calculation gives:

$$z_N = z_0^{l^N}, \quad w_N = 2^{\frac{k^N - 1}{k - 1}} w_0^{k^N} = 2^{\frac{k^N - 1}{k - 1}} z_0^{k^N}.$$

Thus if $l = k$, then the equality:

$$\left(\frac{w_N}{z_N}\right)^{\pm 1} = 2^{\pm \frac{l^N - 1}{l - 1}}$$

holds, which satisfies the uniformly double exponential bound.

On the other hand if $k > l$, then

$$\frac{w_N}{z_N} = 2^{\frac{k^N - 1}{k - 1}} z_0^{k^N - l^N}$$

which heavily depends on the initial values.

Lemma 2.5. *Let f_t and g_t be relative elementary and assume that both are monotone increasing. Let $\{v_N\}_N$ be a positive sequence so that the estimates:*

$$g_t(v_{N-n}, \dots, v_{N-1}) \leq v_N \leq f_t(v_{N-n}, \dots, v_{N-1}), \quad N \geq n$$

hold. Let $\{z_N\}_N$ and $\{w_N\}_N$ be two dynamics defined by $z_N = f_t(z_{N-n}, \dots, z_{N-1})$ and $w_N = g_t(w_{N-n}, \dots, w_{N-1})$ with the same initial value $z_i = w_i = v_i$ for $0 \leq i \leq n - 1$ respectively. Then the estimates hold:

$$w_N \leq v_N \leq z_N \quad (N = 0, 1, \dots)$$

Proof: We proceed by induction. For $N = n$, the estimates follows by the hypothesis. Suppose the estimates $w_N \leq v_N \leq z_N$ hold for $N \leq N_0 - 1$. Then the conclusion for N_0 follows from two estimates:

$$w_{N_0} = g_t(w_{N_0-n}, \dots, w_{N_0-1}) \leq g_t(v_{N_0-n}, \dots, v_{N_0-1}), \quad (30)$$

$$z_{N_0} = f_t(z_{N_0-n}, \dots, z_{N_0-1}) \geq f_t(v_{N_0-n}, \dots, v_{N_0-1}) \quad (31)$$

and the assumption $g_t(v_{N_0-n}, \dots, v_{N_0-1}) \leq v_{N_0} \leq f_t(v_{N_0-n}, \dots, v_{N_0-1})$.

This completes the proof.

Corollary 2.6. *Let f_t and g_t be tropically equivalent, and assume the conditions in lemma 2.4 are satisfied. Then the estimates hold:*

$$\left(\frac{z_N}{v_N}\right)^{\pm 1}, \quad \left(\frac{w_N}{v_N}\right)^{\pm 1} \leq M^{2P_N(c)}$$

where $c = \max(c_f, c_g)$ and $M = \max(M_f, M_g)$.

Proof: By lemma 2.5, the estimates $w_N \leq v_N \leq z_N$ hold for all $N = 0, 1, \dots$. On the other hand by proposition 2.3, the uniform bounds $\frac{z_N}{w_N} \leq M^{2P_N(c)}$ hold. Then the conclusions follow from the estimates $\frac{z_N}{v_N} \leq \frac{z_N}{w_N} \leq M^{2P_N(c)}$ and $\frac{v_N}{w_N} \leq \frac{z_N}{w_N} \leq M^{2P_N(c)}$. This completes the proof.

For example $g_t = \frac{1}{m}f_t$ are the cases for $m \geq 1$, when f_t is monotone increasing.

2.C.2 Dependence on initial values: Let $f_t : \mathbb{R}_{>0}^n \rightarrow (0, \infty)$ be a relative elementary function. Let us take two initial values:

$$\bar{z}_0 = (z_0, \dots, z_{n-1}), \quad \bar{w}_0 = (w_0, \dots, w_{n-1}) \in \mathbb{R}_{>0}^n$$

and consider the corresponding orbits $\{z_N\}_{N=0}^\infty$ and $\{w_N\}_{N=0}^\infty$ defined by:

$$z_N = f_t(z_{N-n}, \dots, z_{N-1}), \quad w_N = f_t(w_{N-n}, \dots, w_{N-1}), \quad (N \geq n)$$

respectively. Let φ and φ_t be the functions corresponding to f_t .

Here we have more elaborate estimates:

Proposition 2.7. *Let f_t and the orbits $\{z_N\}_{N=0}^\infty, \{w_N\}_{N=0}^\infty$ be as above with initial values \bar{z}_0 and \bar{w}_0 . Then they satisfy uniformly bounded rates:*

$$\left(\frac{z_N}{w_N}\right)^\pm \leq M^{2P_N(c)} \left[\max_{0 \leq i \leq n-1} \left(\frac{z_i}{w_i}\right)^{\pm 1} \right]^{c^N} \quad (N \geq n)$$

where c and M are the Lipschitz constant and the number of the components for φ respectively.

Proof: The idea of the proof is parallel to proposition 2.3.

Let us put $x'_N = \log_t(z_N)$ and $y'_N = \log_t(w_N)$ respectively. Thus $\{x'_N\}_N$ is the orbit for φ_t with the initial value $x'_i = \log_t z_i$ for $0 \leq i \leq n-1$, and similar for $\{y'_N\}_N$.

Let $\{x_N\}_N$ be another orbit for φ with the same initial value $x_i = \log_t z_i$ for $0 \leq i \leq n-1$, and similar for $\{y_N\}_N$.

Let $c \geq 1$ be the Lipschitz constant for φ . Let us estimate $|x_N - y_N|$ for $N \geq n$. Since $x_n = \varphi(x_0, \dots, x_{n-1})$ and $y_n = \varphi(y_0, \dots, y_{n-1})$, the estimate:

$$|x_n - y_n| = |\varphi(x_0, \dots, x_{n-1}) - \varphi(y_0, \dots, y_{n-1})| \leq c \max_{0 \leq i \leq n-1} |x_i - y_i|$$

hold. Let us iterate the same estimates:

$$|x_{n+1} - y_{n+1}| = |\varphi(x_1, \dots, x_n) - \varphi(y_1, \dots, y_n)| \quad (32)$$

$$\leq c \max_{1 \leq i \leq n} |x_i - y_i| \leq c^2 \max_{0 \leq i \leq n-1} |x_i - y_i|. \quad (33)$$

The same process gives us the estimates:

$$|x_N - y_N| \leq c^{N-n+1} \max_{0 \leq i \leq n-1} |x_i - y_i| = c^{N-n+1} \max_{0 \leq i \leq n-1} \log_t \left(\frac{z_i}{w_i} \right)^{\pm 1}.$$

On the other hand by lemma 2.4, the estimates:

$$|x_N - x'_N|, |y_N - y'_N| \leq P_N(c) \log_t M$$

hold, where M is the number of the components for φ . So combining with these estimates, we have the followings:

$$\max(\log_t \frac{z_N}{w_N}, \log_t \frac{w_N}{z_N}) = |\log_t(z_N) - \log_t(w_N)| = |x'_N - y'_N| \quad (34)$$

$$\leq |x_N - x'_N| + |y_N - y'_N| + |x_N - y_N| \quad (35)$$

$$\leq 2P_N(c) \log_t M + c^N \max_{0 \leq i \leq n-1} \log_t \left(\frac{z_i}{w_i} \right)^{\pm 1} \quad (36)$$

$$= \log_t \{ M^{2P_N(c)} [\max_{0 \leq i \leq n-1} \left(\frac{z_i}{w_i} \right)^{\pm 1}]^{c^N} \}. \quad (37)$$

Thus one obtains the estimates:

$$\left(\frac{z_N}{w_N} \right)^{\pm 1} \leq M^{2P_N(c)} \max_{0 \leq i \leq n-1} \left(\frac{z_i}{w_i} \right)^{\pm c^N}.$$

This completes the proof.

Now let g_t and f_t be two relatively elementary functions, and denote the corresponding pairs of the functions by (φ, φ_t) and (ψ, ψ_t) respectively. Let (c_f, M_f) and (c_g, M_g) be the Lipschitz constants and the numbers of the components for φ and ψ respectively.

Corollary 2.8. *Let g_t be tropically equivalent to f_t , and $\{z_N\}_N$ and $\{w_N\}_N$ be the orbits for f_t and g_t with the initial values $\bar{z}_0 = (z_0, \dots, z_{n-1})$ and $\bar{w}_0 = (w_0, \dots, w_{n-1})$ respectively. Then the estimates hold:*

$$\left(\frac{z_N}{w_N} \right)^{\pm} \leq M^{4P_N(c)} [\max_{0 \leq i \leq n-1} \left(\frac{z_i}{w_i} \right)^{\pm 1}]^{c^N}$$

where $c = \max(c_f, c_g)$ and $M = \max(M_f, M_g)$.

Proof: Let $\{z'_N\}_N$ be the orbit for f_t with the initial value $\bar{w}_0 = (w_0, \dots, w_{n-1})$. By proposition 2.7, one obtains the estimates:

$$\left(\frac{z_N}{z'_N}\right)^{\pm 1} \leq M^{2P_N(c)} \left[\max_{0 \leq i \leq n-1} \left(\frac{z_i}{w_i}\right)^{\pm 1} \right]^{c^N}.$$

On the other hand by proposition 2.3, one has another estimates:

$$\left(\frac{z'_N}{w_N}\right)^{\pm 1} \leq M^{2P_N(c)}.$$

By multiplying both sides, one obtains the desired estimates:

$$\left(\frac{z_N}{w_N}\right)^{\pm 1} = \left(\frac{z_N}{z'_N}\right)^{\pm 1} \left(\frac{z'_N}{w_N}\right)^{\pm 1} \leq M^{2P_N(c)} M^{2P_N(c)} \left[\max_{0 \leq i \leq n-1} \left(\frac{z_i}{w_i}\right)^{\pm 1} \right]^{c^N} \quad (38)$$

$$= M^{4P_N(c)} \left[\max_{0 \leq i \leq n-1} \left(\frac{z_i}{w_i}\right)^{\pm 1} \right]^{c^N}. \quad (39)$$

This completes the proof.

Now we induce the main estimates:

Theorem 2.9. *Let us take four relative elementary functions, f_t, f'_t, g_t, g'_t . Assume that they are all monotone increasing and all tropically equivalent. Let $\{v_N\}_N$ and $\{u_N\}_N$ be positive sequences which satisfy the estimates:*

$$f'_t(v_{N-n}, \dots, v_{N-1}) \leq v_N \leq f_t(v_{N-n}, \dots, v_{N-1}), \quad (40)$$

$$g'_t(u_{N-n}, \dots, u_{N-1}) \leq u_N \leq g_t(u_{N-n}, \dots, u_{N-1}). \quad (41)$$

for all $N \geq n$. Then the ratios satisfy the uniform estimates:

$$\left(\frac{v_N}{u_N}\right)^{\pm 1} \leq M^{8P_N(c)} \left[\max_{0 \leq i \leq n-1} \left(\frac{u_i}{v_i}\right)^{\pm 1} \right]^{c^N}$$

where $c = \max(c_f, c_{f'}, c_g, c_{g'})$ and $M = \max(M_f, M_{f'}, M_g, M_{g'})$.

Proof: Let us consider two orbits $\{z_N\}_N$ and $\{z'_N\}_N$ defined by $z_N = f_t(z_{N-n}, \dots, z_{N-1})$ and $z'_N = f'_t(z'_{N-n}, \dots, z'_{N-1})$ with the the same initial value $z_i = z'_i = v_i$ for $0 \leq i \leq n-1$ respectively. Similarly by use of g_t and g'_t , one has orbits for $\{w_N\}_N$ and $\{w'_N\}_N$ with the initial value $w_i = w'_i = u_i$ for $0 \leq i \leq n-1$ respectively.

Then by corollary 2.6, one has the estimates:

$$\left(\frac{z_N}{v_N}\right)^{\pm 1}, \quad \left(\frac{w_N}{u_N}\right)^{\pm 1} \leq M^{2P_N(c)}.$$

On the other hand by corollary 2.8, the estimates hold:

$$\left(\frac{z_N}{w_N}\right)^{\pm 1} \leq M^{4P_N(c)} \left[\max_{0 \leq i \leq n-1} \left(\frac{z_i}{w_i}\right)^{\pm 1} \right]^{c^N} = M^{4P_N(c)} \left[\max_{0 \leq i \leq n-1} \left(\frac{v_i}{u_i}\right)^{\pm 1} \right]^{c^N}.$$

Thus from these two, one obtains the desired uniform estimates:

$$\left(\frac{v_N}{u_N}\right)^{\pm 1} = \left(\frac{v_N}{z_N}\right)^{\pm 1} \left(\frac{z_N}{w_N}\right)^{\pm 1} \left(\frac{w_N}{u_N}\right)^{\pm 1} \quad (42)$$

$$\leq M^{2P_N(c)} M^{4P_N(c)} \left[\max_{0 \leq i \leq n-1} \left(\frac{v_i}{u_i}\right)^{\pm 1} \right]^{c^N} M^{2P_N(c)} \quad (43)$$

$$= M^{8P_N(c)} \left[\max_{0 \leq i \leq n-1} \left(\frac{v_i}{u_i}\right)^{\pm 1} \right]^{c^N}. \quad (44)$$

This completes the proof.

2.D Evolutional dynamics: For simplicity of the notation, later on we will omit to denote the parameter t for f_t and just write f for any relative elementary functions.

Let f be a relative elementary function. A general equation of evolutional discrete dynamics is of the form:

$$z_{N+1}^{t+1} = f(z_{N-l_0}^{t+1}, \dots, z_N^{t+1}, z_{N-l_1}^t, \dots, z_{N+k_1}^t, z_{N-l_2}^{t-1}, \dots, z_{N+k_2}^{t-1}, \dots, z_{N+k_{d+1}}^{t-d})$$

where $l_i, k_j \geq 0$, $N \geq \max(l_0, \dots, l_{d+1})$ and $t \geq d$, with initial values:

$$\bar{z}_0^0 \equiv \{z_a^t\}_{0 \leq a \leq \max(l_0, \dots, l_{d+1}), t=0,1,\dots} \cup \{z_N^h\}_{0 \leq h \leq d, N=0,1,\dots}$$

As before one puts the Lipschitz constant and the number of the components by c_f and M_f for the corresponding $(\max, +)$ -function to f .

Let us put $l = \max(l_0, l_1, \dots, l_{d+1})$, $k = \max(k_1, \dots, k_{d+1})$ and

$$A(N, t) \equiv (t - d - 1)k + N - l + n - 1$$

for $N \geq l + 1$ and $t \geq d + 1$.

Let us take g tropically equivalent to f , and consider the dynamics $\{w_n^t\}$ defined by g with any initial value \bar{w}_0^0 .

Now we put the *initial rates* by:

$$[\bar{z}_0^0 : \bar{w}_0^0] \equiv \sup_{0 \leq a \leq \max(l_0, \dots, l_{d+1}), b=0,1, \dots, \text{ or } a=0,1, \dots, 0 \leq b \leq d} \left\{ \frac{z_a^b}{w_a^b}, \frac{w_a^b}{z_a^b} \right\}.$$

Proposition 2.10. (1) *Let f and g be tropically equivalent. Then any orbits $\{z_N^t\}_N$ and $\{w_N^t\}_N$ for f and g with the initial values \bar{z}_0^0 and \bar{w}_0^0 respectively, satisfy the estimates:*

$$\left(\frac{z_N^t}{w_N^t} \right)^\pm \leq M^{4P_{A(N,t)}(c)} [\bar{z}_0^0 : \bar{w}_0^0]^{c^{A(N,t)}}$$

where $c = \max(c_f, c_g)$ and $M = \max(M_f, M_g)$.

(2) *Let f, f', g, g' be four relative elementary functions, and assume that they are all monotone increasing and all tropically equivalent. Let $\{v_N^t\}_{N,t}$ and $\{u_N^t\}_{N,t}$ be positive sequences so that these satisfy the estimates:*

$$f'(v_{N-l_0}^{t+1}, \dots, v_{N+k_{d+1}}^{t-d}) \leq v_{N+1}^{t+1} \leq f(v_{N-l_0}^{t+1}, \dots, v_{N+k_{d+1}}^{t-d}), \quad (45)$$

$$g'(u_{N-l_0}^{t+1}, \dots, u_{N+k_{d+1}}^{t-d}) \leq u_{N+1}^{t+1} \leq g(u_{N-l_0}^{t+1}, \dots, u_{N+k_{d+1}}^{t-d}) \quad (46)$$

for $N \geq l$ and $t \geq d$. Then the ratios satisfy the uniform estimates:

$$\left(\frac{v_N^t}{u_N^t} \right)^{\pm 1} \leq M^{8P_{A(N,t)}(c)} [\bar{u}_0^0 : \bar{v}_0^0]^{c^{A(N,t)}}$$

for $N \geq l+1$ and $t \geq d+1$.

Proof: Let us check that in order to determine z_{l+N}^{d+t} , one has to iterate at most $(t-1)k + N$ times to apply f for $N, t \geq 1$. Then the conclusions follow from corollary 2.8 and theorem 2.9 (see remark (1) below the proof of proposition 2.3).

Let us denote by $\alpha(N, t)$ the number of compositions of f in order to determine z_N^t . It is an increasing function on both variables. We show the estimates $\alpha(l+N, d+t) \leq (t-1)k + N$.

Let $\Delta_0 = \{(a, b) \in \{0, 1, \dots, k+l\} \times \{0, 1, \dots, d\} \cup \{0, \dots, l\} \times \{d+1\}\}$ be the finite set. This is a basic building block in the sense that for $N, t \geq 1$, z_{N+l}^{t+d} is determined if one knows z_{N-1+a}^{t-1+b} for $(a, b) \in \Delta_0$.

We proceed by induction on t . $\alpha(l+N, d+1) \leq N$ clearly follows.

Suppose the conclusion follows for $t \leq t_0$, and so $\alpha(N+l, d+t_0) \leq (t_0-1)k + N$ hold. Then $\alpha(l+1, d+t_0+1) = \alpha(l+k, d+t_0) + 1 \leq$

$(t_0 - 1)k + k + 1 = t_0k + 1$ hold. Next $\alpha(l + 2, d + t_0 + 1) = \max(\alpha(l + 1, d + t_0 + 1), \alpha(l + k + 1, d + t_0)) + 1 \leq t_0k + 2$. By use of the estimates $\alpha(N + l, d + t_0 + 1) \leq \max(\alpha(N - 1 + l, d + t_0 + 1), \alpha(N - 1 + l + k, d + t_0)) + 1$, one can obtain the bounds $\alpha(N + l, d + t_0 + 1) \leq t_0k + N$.

This completes the proof.

3 Asymptotic comparisons

3.A Formal Taylor expansion and ODE: Let us consider a $C^{\alpha+1}$ function $u : (0, \infty) \rightarrow (0, \infty)$. Below we proceed to approximate u very roughly by discrete dynamics defined by relative elementary functions of n variables. For $1 \leq |i| \leq n - 1$, let us take the Taylor expansions around $x \in (0, \infty)$:

$$u(x + i\epsilon) = u(x) + i\epsilon u_x + \frac{(i\epsilon)^2}{2} u_{2x} + \cdots + \frac{(i\epsilon)^\alpha}{\alpha!} u_{\alpha x} + \frac{(i\epsilon)^{(\alpha+1)}}{(\alpha + 1)!} u_{(\alpha+1)x}(\xi_i)$$

for small $|\epsilon| \ll 1$, where:

$$\begin{cases} x \leq \xi_i \leq x + i\epsilon, & i \geq 0 \\ x + i\epsilon \leq \xi_i \leq x & i < 0 \end{cases}$$

(for our applications, we will choose $\alpha \leq 2$ later).

Let $f = \frac{k}{h} : \mathbb{R}_{>0}^n \rightarrow (0, \infty)$ be a relative elementary function, where h and k are both elementary. Later on we will assume positivity:

$$h(\bar{0}) > 0.$$

Let us consider the discrete dynamics defined by $z_{N+1} = f(z_{N-n+1}, \dots, z_N)$.

We put the *fluctuation intervals* by:

$$I(n, \epsilon) = [-n\epsilon, 0].$$

For $N = 0, 1, 2, \dots$, let us put change of variables:

$$z_N \equiv \epsilon u(\epsilon N) = \epsilon u(x), \quad (N = \frac{x}{\epsilon}).$$

Let us consider the difference:

$$z_{N+1} - f(z_{N-n+1}, \dots, z_N) = \epsilon u(x + \epsilon) - f(\epsilon u(x - (n - 1)\epsilon), \dots, \epsilon u(x))$$

and insert the Taylor expansions:

$$= \epsilon(u + \epsilon u_x + \frac{\epsilon^2}{2} u_{2x} + \dots) \quad (47)$$

$$- f(\epsilon(u - (n-1)\epsilon u_x + \frac{(n-1)^2 \epsilon^2}{2} u_{2x} + \dots), \dots, \epsilon u). \quad (48)$$

By reordering the expansions with respect to the exponents of ϵ , there are rational numbers $a_0, a_1, \dots \in \mathbb{Q}$ so that the equality holds:

$$\epsilon u(x + \epsilon) - f(\epsilon u(x - (n-1)\epsilon), \dots, \epsilon u(x)) \quad (49)$$

$$= \frac{\epsilon a_0 u + \epsilon^2 a_1 u_x + \epsilon^3 a_2 u u_x + \dots + \epsilon^{\alpha+1} a_s u_{\alpha x} + \epsilon^{\alpha+2} a_{s+1} u_{(\alpha+1)x}(\xi) + \dots}{h(\epsilon u(x - (n-1)\epsilon), \dots, \epsilon u(x))} \quad (50)$$

$$\equiv \frac{\epsilon F^1(u) + \epsilon^2 F^2(u_x) + \epsilon^3 F^3(u, u_x) + \dots + \epsilon^m F^{m'}(u, \dots, u_{(\alpha+1)x}(\xi)) + \dots}{h(\epsilon u(x - (n-1)\epsilon), \dots, \epsilon u(x))} \quad (51)$$

$$(52)$$

where F^k are monomials.

Let us choose finite subsets $A \subset \{1, 2, 3, \dots\}$, and divide the expanded sum into two terms as:

$$= \frac{\sum_{i \in A} \epsilon^{s_i} F^{s'_i}(u, u_x, \dots, u_{\alpha x})}{h(\epsilon u(x - (n-1)\epsilon), \dots)} + \frac{\sum_{j \in A^c} \epsilon^{s_j} F^{s'_j}(u, u_x, \dots, u_{(\alpha+1)x}(\xi))}{h(\epsilon u(x - (n-1)\epsilon), \dots)} \quad (53)$$

$$\equiv \mathbf{F}(\epsilon, u, u_x, \dots, u_{\alpha x}) + \epsilon^2 \mathbf{F}^1(\epsilon, u, u_x, \dots, u_{(\alpha+1)x}(\xi_1), \dots, u_{(\alpha+1)x}(\xi_{n-1})). \quad (54)$$

We always choose A so that two conditions are satisfied;

(1) \mathbf{F} do not contain $u_{(l+1)x}(\xi)$, and (2) $1 \in A$, i.e. F^1 is included in \mathbf{F} .

In all the concrete cases later, we choose relative elementary functions and A so that the corresponding F^1 vanish.

Now fix $\epsilon > 0$, and suppose u obeys the equation:

$$\mathbf{F}(\epsilon, u, u_x, \dots, u_{\alpha x}) = 0.$$

Then the difference satisfies the equality:

$$\epsilon u(x + \epsilon) - f(\epsilon u(x - (n-1)\epsilon), \dots, \epsilon u(x)) = \epsilon^2 \mathbf{F}^1(\epsilon, u, u_x, \dots).$$

We say that \mathbf{F} is the *leading term*, and \mathbf{F}^1 error one for u respectively.

Remark: Conversely when one starts from ODE $\mathbf{F}(\epsilon, u, u_x, u_{\alpha x}) = 0$, there will be several choices of relative elementary functions f and A with the leading term \mathbf{F} . Various choices of f will assign different error terms \mathbf{F}^1 , which reflect estimates of solutions $\mathbf{F}(\epsilon, u, u_x, \dots, u_{\alpha x}) = 0$. So ‘better’ choice of f will give us ‘better’ estimates of large scale analysis of such solutions.

Let us define ϵ variation of \mathbf{F}^1 by

$$\begin{aligned} & \|\mathbf{F}^1(\epsilon, u, u_x, \dots, u_{\alpha x}, u_{(\alpha+1)x}(\xi_1), \dots, u_{(\alpha+1)x}(\xi_{n-1}))\|_{\epsilon}(x) \equiv & (55) \\ & \sup_{\mu_i - x \in I(n, \epsilon)} |\mathbf{F}^1(\epsilon, u(x - \epsilon), \dots, u_{\alpha x}(x - \epsilon), u_{(\alpha+1)x}(\mu_1), \dots, u_{(\alpha+1)x}(\mu_{n-1}))| & (56) \end{aligned}$$

where $I(n, \epsilon)$ is the fluctuation interval.

Let us say that a $C^{\alpha+1}$ function $u : (0, \infty) \rightarrow (0, \infty)$ is ϵ_0 controlled, if there is some constant $C > 0$ so that the ϵ_0 variation of \mathbf{F}^1 satisfy the pointwise estimates for all $x \in (0, \infty)$:

$$Cu(x) \geq \|\mathbf{F}^1(\epsilon, u, u_x, \dots, u_{\alpha x}, u_{(\alpha+1)x}(\xi_1), \dots, u_{(\alpha+1)x}(\xi_{n-1}))\|_{\epsilon_0}(x)$$

3.A.2 Comparison theorem for ODE: Let us take another relatively elementary function $g = \frac{d}{\epsilon}$ which is tropically equivalent to f . Let $v : (0, \infty) \rightarrow (0, \infty)$ be another $C^{\alpha+1}$ function. By replacing f by g and choosing another subsets $B \subset \{1, 2, 3, \dots\}$ in 3.A, one has its leading and error terms \mathbf{G} and \mathbf{G}^1 respectively. Then we have the equalities:

$$\epsilon v(x + \epsilon) - g(\epsilon v(x - (n-1)\epsilon), \dots, \epsilon v(x)) \quad (57)$$

$$= \frac{\sum_{i \in B} \epsilon^{s_i} G^{s'_i}(v, v_x, \dots, v_{\alpha x})}{e(\epsilon v(x - (n-1)\epsilon), \dots)} + \frac{\sum_{j \in B^c} \epsilon^{s_j} G^{s'_j}(v, v_x, \dots, v_{(\alpha+1)x}(\xi'_j))}{e(\epsilon v(x - (n-1)\epsilon), \dots)} \quad (58)$$

$$\equiv \mathbf{G}(\epsilon, v, \dots, v_{\alpha x}) + \epsilon^2 \mathbf{G}^1(\epsilon, v, \dots, v_{\alpha x}, v_{(\alpha+1)x}(\xi'_1), \dots, v_{(\alpha+1)x}(\xi'_{n-1})). \quad (59)$$

Let us fix a small $\epsilon > 0$, and take two positive solutions $u, v : (0, \infty) \rightarrow (0, \infty)$ to the equations:

$$\mathbf{F}(\epsilon, u, u_x, \dots) = 0, \quad \mathbf{G}(\epsilon, v, v_x, \dots) = 0.$$

Now we compare their ratios:

$$\left(\frac{u(x)}{v(x)}\right)^{\pm 1} = \left\{ \frac{u(x)}{v(x)}, \frac{v(x)}{u(x)} \right\}.$$

For this we introduce the *initial rates*:

$$[u : v]_\epsilon \equiv \sup_{x \in (0, \epsilon]} \left(\frac{u(x)}{v(x)} \right)^{\pm 1}.$$

Recall that associated with f are the Lipschitz constant $c_f \geq 1$ and the number of the components M_f . Let us put $c = \max(c_f, c_g)$ and $M = \max(M_f, M_g)$.

Theorem 3.1. *Let f and g be both relatively elementary and increasing functions of n variables, which are mutually tropically equivalent. Let \mathbf{F} and \mathbf{G} be their leading terms of order at most $\alpha \geq 0$, and take positive $C^{\alpha+1}$ solutions $u, v : (0, \infty) \rightarrow (0, \infty)$ to the equations:*

$$\mathbf{F}(\epsilon, u, u_x, \dots, u_{\alpha x}) = 0, \quad \mathbf{G}(\epsilon, v, v_x, \dots, v_{\alpha x}) = 0.$$

Assume both u and v are ϵ_0 controlled bounded by C . Then for any $0 < \epsilon \leq \min(\frac{1}{2C}, \epsilon_0)$, the estimates hold:

$$\left(\frac{u(x)}{v(x)} \right)^{\pm 1} \leq (2M)^{8 \frac{c^{\epsilon^{-1}x+1}-1}{c-1}} [u : v]_{n\epsilon}^{c^{\epsilon^{-1}x+1}}.$$

Proof: Let f and g be both n variables, and $(\mathbf{F}, \mathbf{F}^1)$ and $(\mathbf{G}, \mathbf{G}^1)$ be pairs of leading and error terms respectively.

Let us choose $0 < \epsilon \leq \min(\frac{1}{2C}, \epsilon_0)$. By the assumption, the pointwise estimates hold:

$$Cu(x + \epsilon) \geq |\mathbf{F}^1(\epsilon, u(x), u_x(x), \dots, u_{\alpha x}(x), u_{(\alpha+1)x}(\xi_1), \dots, u_{(\alpha+1)x}(\xi_{n-1}))|.$$

In particular the estimates $\epsilon^2 |\mathbf{F}^1| \leq \frac{1}{2C} \epsilon |\mathbf{F}^1| \leq \frac{1}{2} \epsilon u(x + \epsilon)$ hold.

Let us consider the equalities:

$$\epsilon u(x + \epsilon) - f(\epsilon u(x - (n-1)\epsilon), \dots, \epsilon u(x)) \tag{60}$$

$$= \mathbf{F}(\epsilon, u, \dots, u_{\alpha x}) + \epsilon^2 \mathbf{F}^1(\epsilon, u, \dots, u_{\alpha x}, u_{(\alpha+1)x}(\xi_1), \dots, u_{(\alpha+1)x}(\xi_{n-1})) \tag{61}$$

$$= \epsilon^2 \mathbf{F}^1(\epsilon, u, u_x, \dots, u_{\alpha x}, u_{(\alpha+1)x}(\xi_1), \dots, u_{(\alpha+1)x}(\xi_{n-1})) \tag{62}$$

since u obeys the equation $\mathbf{F}(\epsilon, u, u_x, \dots) = 0$.

Then combining with the above inequality, one obtains the estimates:

$$\frac{1}{2} f(\epsilon u(x - (n-1)\epsilon), \dots, \epsilon u(x)) \leq \epsilon u(x + \epsilon) \tag{63}$$

$$\leq 2f(\epsilon u(x - (n-1)\epsilon), \dots, \epsilon u(x)). \tag{64}$$

By the same way one obtains the estimates by replacing f by g :

$$\frac{1}{2}g(\epsilon v(x - (n-1)\epsilon), \dots, \epsilon v(x)) \leq \epsilon v(x + \epsilon) \quad (65)$$

$$\leq 2g(\epsilon v(x - (n-1)\epsilon), \dots, \epsilon v(x)). \quad (66)$$

f , $\frac{1}{2}f$ and $2f$ are tropically equivalent, and $\frac{1}{2}f$, $2f$, $\frac{1}{2}g$, $2g$ are all so by the assumption. Notice that the number of the components for $\frac{1}{2}f$ and $2f$ are both $2M_f$.

Thus the estimates hold by theorem 2.9:

$$\left(\frac{u(N\epsilon)}{v(N\epsilon)}\right)^{\pm 1} \leq (2M)^{8P_N(c)} \sup_{0 \leq i \leq n-1} \left(\frac{u(\epsilon i)}{v(\epsilon i)}\right)^{\pm c^N} \leq (2M)^{8P_N(c)} ([u : v]_{(n-1)\epsilon})^{c^N}.$$

For any $0 \leq \mu \leq \epsilon$, let us apply the above estimates for the translations $u(x + \mu)$ and $v(x + \mu)$. Then one obtains the estimates:

$$\left(\frac{u(N\epsilon + \mu)}{v(N\epsilon + \mu)}\right)^{\pm 1} \leq (2M)^{8P_N(c)} [u : v]_{n\epsilon}^{c^N} \quad (67)$$

$$= (2M)^{8\frac{c^{N-n+1}-1}{c-1}} [u : v]_{n\epsilon}^{c^N} \leq (2M)^{8\frac{c^{\epsilon^{-1}(N\epsilon+\mu)-n+2}-1}{c-1}} [u : v]_{n\epsilon}^{c^{\epsilon^{-1}(N\epsilon+\mu)+1}} \quad (68)$$

since $P_N(c) = \frac{c^{N-n+1}-1}{c-1}$.

Such $N\epsilon + \mu$ takes all the points $x \in (0, \infty)$, and so the estimates hold:

$$\left(\frac{u(x)}{v(x)}\right)^{\pm 1} \leq (2M)^{8\frac{c^{\epsilon^{-1}x-n+2}-1}{c-1}} [u : v]_{n\epsilon}^{c^{\epsilon^{-1}x+1}} \leq (2M)^{8\frac{c^{\epsilon^{-1}x+1}-1}{c-1}} [u : v]_{n\epsilon}^{c^{\epsilon^{-1}x+1}}.$$

This completes the proof.

Example: Let us consider a simple equation:

$$\mathbf{F}(u, u_x) = u_x + u^2 = 0.$$

It has solutions $u(x) = \frac{a}{1+ax}$ with the initial values $u(0) = a > 0$. Let us put $z_N = \epsilon u(x)$ with $x = N\epsilon$ and take the Taylor expansion $\epsilon u(x + \epsilon) = \epsilon u(x) + \epsilon^2 u_x(x) + \frac{\epsilon^3}{2} u_{2x}(\xi)$. We choose the relative elementary function $f(x) = x(1+x)^{-1}$ and calculate the difference:

$$u(x + \epsilon) - f(\epsilon u(x)) \quad (69)$$

$$= \epsilon^2 \frac{u_x + u^2}{1 + \epsilon u}(x) + \epsilon^3 \frac{\frac{1}{2} u_{2x}(\xi) + u(x)u_x(x) + \frac{\epsilon}{2} u(x)u_{2x}(\xi)}{1 + \epsilon u(x)} \quad (70)$$

Thus u is ϵ_0 controlled, since $1 + \epsilon u(x) \geq 1$, and the estimates $|u_{2x}(\xi)|, |uu_x(x)|, |u(x)u_{2x}(\xi)| \leq Cu(x)$ hold uniformly in $x \in (0, \infty)$ for some $C = C(a, \epsilon_0) \geq 0$ and $|x - \xi| \leq \epsilon$.

The corresponding $(\max, +)$ function to f is given by $V_{N+1} = V_N - \max(0, V_N)$. Notice the equality $V_N - \max(0, V_N) = V_N - \max(0, V_N, V_N)$. The tropical inverse for the latter is given by $g(y) = y(1 + 2y)^{-1}$. By choosing the same scaling parameter, one obtains the leading term $\mathbf{G}(v, v_x) = v_x + 2v^2$. It has solutions $v(x) = \frac{a'}{2a'x+1}$, and the ratio is in fact uniformly bounded:

$$\left(\frac{u(x)}{v(x)}\right)^\pm = \left(\frac{a(2a'x+1)}{a'(ax+1)}\right)^{\pm 1} \leq 2\left(\frac{a}{a'}\right)^{\pm 1} \leq 2[u : v]_\epsilon.$$

3.B Evolutional dynamics: Here we treat partial differential equations. The process of 3.B is quite parallel to 3.A by introducing time parameter.

A general equation of evolutional discrete dynamics is of the form:

$$z_{N+1}^{t+1} = f(z_{N-l_0}^{t+1}, \dots, z_N^{t+1}, z_{N-l_1}^t, \dots, z_{N+k_1}^t, z_{N-l_2}^{t-1}, \dots, z_{N+k_2}^{t-1}, \dots, z_{N+k_{d+1}}^{t-d})$$

where $l_i, k_j \geq 0$, $N \geq l \equiv \max(l_0, \dots, l_{d+1})$ and $t \geq d$, with initial values:

$$\bar{z}_0^0 \equiv \{z_a^t\}_{0 \leq a \leq l, t=0,1,\dots} \cup \{z_N^h\}_{0 \leq h \leq d, N=0,1,\dots}$$

Now let us consider a $C^{\alpha+1}$ function $u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$, and introduce another parameters by

$$N = \frac{x}{\epsilon^p}, \quad t = \frac{s}{\epsilon^q}, \quad \epsilon^m u(x, s) = z_N^t$$

where $\epsilon > 0$ is a small constant, and $p, q \geq 1$, $m \geq 0$ are integers. Then we take the Taylor expansions:

$$u(x + i\epsilon^p, s + j\epsilon^q) = u + i\epsilon^p u_x + j\epsilon^q u_s + \frac{(i\epsilon^p)^2}{2} u_{2x} + \frac{(j\epsilon^q)^2}{2} u_{2s} \quad (71)$$

$$+ j\epsilon^q i\epsilon^p u_{xs} + \dots + \frac{(i\epsilon^p)^\alpha}{\alpha!} u_{\alpha x} + \frac{(j\epsilon^q)^\alpha}{\alpha!} u_{\alpha s} \quad (72)$$

$$+ \frac{(i\epsilon^p)^{(\alpha+1)}}{(\alpha+1)!} u_{(\alpha+1)x}(\xi_{ij}) + \dots + \frac{(j\epsilon^q)^{(\alpha+1)}}{(\alpha+1)!} u_{(\alpha+1)s}(\xi_{ij}) \quad (73)$$

$$\equiv u + i\epsilon^p u_x + j\epsilon^q u_s + \frac{(i\epsilon^p)^2}{2} u_{2x} + \frac{(j\epsilon^q)^2}{2} u_{2s} + j\epsilon^q i\epsilon^p u_{xs} \quad (74)$$

$$+ \dots + \frac{(i\epsilon^p)^\alpha}{\alpha!} u_{\alpha x} + \frac{(j\epsilon^q)^\alpha}{\alpha!} u_{\alpha s} + \sum_{\bar{a}} \frac{(i\epsilon^p)^a (j\epsilon^q)^{\alpha+1-a}}{(\alpha+1)!} u_{\bar{a}}(\xi_{ij}) \quad (75)$$

where $\bar{a} = (y_{i_1}, \dots, y_{i_{\alpha+1}})$, $y_j = x$ or s , and $|(x, s) - \xi_{ij}| \leq |(i\epsilon^p, j\epsilon^q)|$.

Let $f = \frac{k}{h} : \mathbb{R}_{>0}^n \rightarrow (0, \infty)$ be a relative elementary function, and consider the difference as in 3.A:

$$z_{N+1}^{t+1} - f(z_{N-l_0}^{t+1}, \dots, z_N^{t+1}, z_{N-l_1}^t, \dots, z_{N+k_1}^t, \dots, z_{N+k_{d+1}}^{t-d}) \quad (76)$$

$$= \epsilon^m u(x + \epsilon^p, s + \epsilon^q) \quad (77)$$

$$- f(\epsilon^m u(x - l_0 \epsilon^p, s + \epsilon^q), \dots, \epsilon^m u(x + k_{d+1} \epsilon^p, s - d \epsilon^q)). \quad (78)$$

By reordering the expansions with respect to the exponents of ϵ , there are rational numbers $a_0, a_1, \dots \in \mathbb{Q}$ so that the above difference is equal to the following:

$$\epsilon^m \frac{a_0 u + \epsilon^p a_1 u_x + \epsilon^q a_2 u_s + \epsilon^{m+p} a_3 u u_x + \dots + (i\epsilon^p)^h (j\epsilon^q)^{\alpha+1-h} a_h u_{\bar{h}}(\xi_{ij}) + \dots}{h(\epsilon^m u(x - l_0 \epsilon^p, s + \epsilon^q), \dots, \epsilon^m u(x + k_{d+1} \epsilon^p, s - d \epsilon^q))} \quad (79)$$

$$\equiv \frac{\epsilon^m F^1(u) + \epsilon^{m+p} F^2(u_x) + \epsilon^{m+q} F^3(u_s) + \epsilon^{2m+p} F^4(u, u_x) + \dots}{h(\epsilon^m u(x - l_0 \epsilon^p, s + \epsilon^q), \dots, \epsilon^m u(x + k_{d+1} \epsilon^p, s - d \epsilon^q))} \quad (80)$$

where F^k are monomials.

Let us choose finite subsets $A \subset \{1, 2, 3, \dots\}$, and divide the expanded sum into two terms as:

$$\epsilon^m u(x + \epsilon^p, s + \epsilon^q) \quad (81)$$

$$- f(\epsilon^m u(x - l_0 \epsilon^p, s + \epsilon^q), \dots, \epsilon^m u(x + k_{d+1} \epsilon^p, s - d \epsilon^q)) \quad (82)$$

$$= \frac{\epsilon^m F^1(u) + \epsilon^{m+p} F^2(u_x) + \epsilon^{m+q} F^3(u_s) + \epsilon^{2m+p} F^4(u, u_x) + \dots}{h(\epsilon^m u(x - l_0 \epsilon^p, s + \epsilon^q), \dots, \epsilon^m u(x + k_{d+1} \epsilon^p, s - d \epsilon^q))} \quad (83)$$

$$= \frac{\sum_{i \in A} \epsilon^{s_i} F^{s'_i}(u, u_x, u_s, \dots, u_{\alpha s}) + \sum_{j \in A^c} \epsilon^{s_j} F^{s'_j}(u, u_x, \dots, u_{\bar{a}}(\xi_{ij}))}{h(\epsilon^m u(x - l_0 \epsilon^p, s + \epsilon^q), \dots, \epsilon^m u(x + k_{d+1} \epsilon^p, s - d \epsilon^q))} \quad (84)$$

$$\equiv \mathbf{F}(\epsilon, u, u_x, u_s, \dots, u_{\alpha s}) + \epsilon^{m+1} \mathbf{F}^1(\epsilon, u, u_x, \dots, \{u_{\bar{a}}(\xi_{ij})\}_{\bar{a}, i, j}). \quad (85)$$

As in 3.A, we always choose A so that \mathbf{F} do not contain $u_{\bar{a}}(\xi)$ and $1 \in A$. We call \mathbf{F} as the leading term and \mathbf{F}^1 the error term respectively.

3.B.2 ϵ -controlledness: Now we return to the starting point. Let f be a relative elementary function, and consider the discrete dynamics $z_{N+1}^{t+1} = f(z_{N-l_0}^{t+1}, \dots, z_N^{t+1}, z_{N-l_1}^t, \dots, z_{N+k_{d+1}}^{t-d})$. After one chooses integers p, q, m for change of variables, one determines the leading and error terms \mathbf{F} and \mathbf{F}^1 respectively.

For $(a, b) \in \mathbb{Z}^2$, let $L(a, b) = \{(ta, tb) : t \in [0, 1]\} \subset \mathbb{R}^2$ be the segment. Then for the set:

$$D \equiv \{(1, 1), (-l_0, 1), \dots, (0, 1), (-l_1, 0), \dots, (k_1, 0), \\ (-l_2, -1), \dots, (k_2, -1), \dots, (-l_{d+1}, -d), \dots, (k_{d+1}, -d)\}$$

we put the *fluctuation domain* as:

$$D(\epsilon, p, q) = \{(L(\epsilon^p a, \epsilon^q b) : (a, b) \in D\} \subset \mathbb{R}^2.$$

For example $D = \{(1, 1), (2, 0), (-1, 1)\}$ for $z_{N+1}^{t+1} = f(z_N^t, z_{N+2}^t, z_{N-1}^{t+1})$.

Let us regard \mathbf{F}^1 as a function on the variables $(x, s, \{\xi_{ij}\}_{i,j})$. Then we define its ϵ variation:

$$\|\mathbf{F}^1\|_\epsilon(x, s) \equiv \sup_{\xi_{ij} - (x - \epsilon^p, s - \epsilon^q) \in D(\epsilon, p, q)} |\mathbf{F}^1(\epsilon, u(x - \epsilon^p, s - \epsilon^q), \quad (86)$$

$$u_x(x - \epsilon^p, s - \epsilon^q), u_s(x - \epsilon^p, s - \epsilon^q), \dots, u_{\alpha s}(x - \epsilon^p, s - \epsilon^q), \{u_{\bar{a}}(\xi_{ij})\})|. \quad (87)$$

Let $u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be a $C^{\alpha+1}$ function.

Definition 3.1. u is ϵ_0 controlled bounded by C , if ϵ_0 variation of \mathbf{F}^1 satisfies the pointwise estimates:

$$Cu(x, s) \geq \|\mathbf{F}^1\|_{\epsilon_0}(x, s)$$

for all $(x, s) \in (0, \infty) \times [0, T_0)$.

3.B.2.2 Higher derivative rates: Let $u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be a $C^{\alpha+1}$ function and f be a relative elementary function. let us consider the expansions of the differences in 3.B:

$$\epsilon^m u(x + \epsilon^p, s + \epsilon^q) \quad (88)$$

$$- f(\epsilon^m u(x - l_0 \epsilon^p, s + \epsilon^q), \dots, \epsilon^m u(x + k_{d+1} \epsilon^p, s - d \epsilon^q)) \quad (89)$$

$$= \mathbf{F}(\epsilon, u, u_x, u_s, \dots, u_{\alpha s}) + \epsilon^{m+1} \mathbf{F}^1(\epsilon, u, u_x, \dots, \{u_{\bar{a}}(\xi_{ij})\}_{\bar{a}, i, j}). \quad (90)$$

\mathbf{F} has order at most α , while \mathbf{F}^1 may contain derivatives of u smaller than $\alpha + 1$ in general.

Let us say that the error term \mathbf{F}^1 is *admissible*, if it is of the form:

$$\mathbf{F}^1 = \sum_{a \in A^c} c_a \epsilon^{s_a} H_a(\epsilon^m u(x - l_0 \epsilon^p, s + \epsilon^q), \dots) u_{\bar{a}}(\xi_{ij})$$

where (1) $|\bar{a}| = \alpha + 1$ and (2) $\|H_a(x_1, x_2, \dots)\|_{C^0} \leq 1$ for any $x_1, x_2 \dots \geq 0$.

For this case we put the *error constants* by:

$$C_{\mathbf{F}^1} \equiv \sum_{a \in A^c} |c_a| \in \mathbb{Q}_{>0}.$$

The error constants are determined by the coefficients of rational functions f and of the Taylor expansions. Our applications later are all admissible cases.

Let us introduce *variation of order $\alpha + 1$* of u by:

$$\|u\|_{\alpha+1}(x, s) = \max_{\partial_i = \partial_x, \partial_s} \left\{ \sup_{\xi - (x - \epsilon^p, s - \epsilon^q) \in D(\epsilon, p, q)} \left| \frac{\partial^{\alpha+1} u}{\partial_1 \dots \partial_{\alpha+1}} \right|(\xi) \right\}.$$

Let us say that u satisfies *uniform ϵ variation*, if there is a constant C so that it satisfies the estimates:

$$Cu(x, s) \geq \|u\|_{\alpha+1}(x, s)$$

for all $(x, s) \in (0, \infty) \times [0, T_0)$.

We put the *variation constant* by:

$$V(u) \equiv \sup_{(x, s) \in (0, \infty) \times [0, T_0)} \frac{\|u\|_{\alpha+1}(x, s)}{u(x, s)}.$$

Lemma 3.2. *Suppose \mathbf{F}^1 is admissible, and u satisfies uniform ϵ variation bounded by C . Then u is ϵ controlled bounded by $CC_{\mathbf{F}^1}$.*

Proof: By admissibility, the estimates hold:

$$\begin{aligned} \|\mathbf{F}^1\|_{\epsilon}(x, s) &\leq \sum_{a \in A^c} |c_a| \epsilon^{s_a} |H_a(\epsilon^m u(x - l_0 \epsilon^p, s + \epsilon^q), \dots)| \|u\|_{\alpha+1}(x, s) \\ &\leq \sum_{a \in A^c} |c_a| \|u\|_{\alpha+1}(x, s) \leq C_{\mathbf{F}^1} \|u\|_{\alpha+1}(x, s) \leq C_{\mathbf{F}^1} Cu(x, s). \end{aligned}$$

Let $u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be a $C^{\alpha+1}$ function. Here we consider classes of functions which satisfy uniform rates between higher derivatives and lowest values.

Let us introduce the derivative constants of $\alpha + 1$, which is given by:

$$\|u\|_{\alpha+1} = \max_{\partial_i = \partial_x, \partial_s} \left\{ \left\| \frac{\partial^{\alpha+1} u}{\partial_1 \dots \partial_{\alpha+1}} \right\|_{C^0((0, \infty) \times [0, T_0))} \right\}.$$

Suppose u satisfies two conditions:

(1) $\|u\|_{\alpha+1} < \infty$ is finite and (2) $c = \inf_{(x,s) \in (0,\infty) \times [0,T_0]} u(x,s) > 0$ is positive. Then we say that the ratio:

$$K(u) \equiv \frac{\|u\|_{\alpha+1}}{c}$$

is the *derivative rates of order $\alpha + 1$* . In general the estimates hold:

$$V(u) \leq K(u).$$

Now we state the following which requires more practical conditions:

Lemma 3.3. *Suppose \mathbf{F}^1 is admissible, and the derivative rates of order $\alpha + 1$ is of finite, $K(u) < \infty$. Then for any $\epsilon > 0$, u is ϵ controlled bounded by $C_{\mathbf{F}^1} K(u)$.*

Proof: By the conditions, the error term \mathbf{F}^1 satisfies the estimates:

$$\begin{aligned} \|\mathbf{F}^1\|_{\epsilon}(x,s) &\leq C_{\mathbf{F}^1} \|u\|_{\alpha+1}(x,s) \leq C_{\mathbf{F}^1} \|u\|_{\alpha+1} \\ &= C_{\mathbf{F}^1} K(u) c \leq C_{\mathbf{F}^1} K(u) u(x,s). \end{aligned}$$

So u is ϵ -controlled bounded by $C_{\mathbf{F}^1} K(u)$ for any $\epsilon > 0$.

This completes the proof.

3.B.3 Comparison theorem: Let g be tropically equivalent to f , and choose the same scaling parameters. By replacing f by g above, one obtains another leading and error terms \mathbf{G} and \mathbf{G}^1 with the equalities:

$$\epsilon^m v(x + \epsilon^p, s + \epsilon^q) - \tag{91}$$

$$g(\epsilon^m v(x - l_0 \epsilon^p, s + \epsilon^q), \dots, \epsilon^m v(x + k_{d+1} \epsilon^p, s - d \epsilon^q)) \tag{92}$$

$$= \mathbf{G}(\epsilon, v, v_x, v_s, \dots, v_{\alpha s}) + \epsilon^{m+1} \mathbf{G}^1(\epsilon, v, v_x, v_s, \dots, v_{\alpha s}, \{v_{\bar{a}}(\eta_{ij})\}_{\bar{a},i,j}). \tag{93}$$

Let us fix a small $\epsilon > 0$, and take two positive solutions $u, v : (0, \infty) \times [0, T_0] \rightarrow (0, \infty)$ to the corresponding PDEs:

$$\mathbf{F}(\epsilon, u, u_x, u_s, \dots, u_{\alpha x}, u_{\alpha s}) = 0, \quad \mathbf{G}(\epsilon, v, v_x, v_s, \dots, v_{\alpha x}, v_{\alpha s}) = 0.$$

In order to estimate their ratios $(\frac{u(x,s)}{v(x,s)})^{\pm 1}$, we introduce the *initial rates*:

$$[u : v]_{\epsilon} \equiv \sup_{(x,s) \in (0,\infty) \times [0,\epsilon^q] \cup (0,\epsilon^p] \times [0,T_0]} \left(\frac{u(x,s)}{v(x,s)} \right)^{\pm 1}.$$

Recall the Lipschitz constant c_f and the number of the components M_f for f . Let us put $c = \max(c_f, c_g)$, $M = \max(M_f, M_g)$, $k = \max(k_1, \dots, k_{d+1})$ and $L = \max(l, d)$ for $l = \max(l_0, l_1, \dots, l_{d+1})$.

Corollary 3.4. *Let f and g be both relatively elementary and increasing functions of n variables, which are mutually tropically equivalent. Let \mathbf{F} and \mathbf{G} be their leading terms of order at most $\alpha \geq 0$, and take positive $C^{\alpha+1}$ solutions $u, v : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ to the equations:*

$$\mathbf{F}(\epsilon, u, u_x, u_s, \dots, u_{\alpha x}, u_{\alpha s}) = 0, \quad \mathbf{G}(\epsilon, v, v_x, v_s, \dots, v_{\alpha x}, v_{\alpha s}) = 0.$$

Assume both u and v are ϵ_0 controlled bounded by C . Then for any $0 < \epsilon \leq \min(\frac{1}{2C}, \epsilon_0)$ and $D = \max(p, q)$, the estimates hold:

$$\left(\frac{u(x, s)}{v(x, s)}\right)^{\pm 1} \leq (2M)^{8\frac{\epsilon^{-D}(x+ks)+1}{c-1}-1} [u : v]_{(L+1)\epsilon}^{\epsilon^{-D}(x+ks)+n}.$$

Proof: Recall $A(N, t) = (t - d - 1)k + N - l + n - 1$ for $N \geq l + 1$ and $t \geq d + 1$. Let us take $0 < \epsilon \leq \min(\frac{1}{2C}, \epsilon_0)$. Combining with proposition 2.10, the parallel argument to the proof of theorem 3.1 gives the estimates:

$$\left(\frac{u(N\epsilon^p + \mu, t\epsilon^q + \chi)}{v(N\epsilon^p + \mu, t\epsilon^q + \chi)}\right)^{\pm 1} \leq (2M)^{8P_{A(N,t)}(c)} [u : v]_{(L+1)\epsilon}^{cA(N,t)}$$

for any $0 \leq \mu \leq \epsilon^p$ and $0 \leq \chi \leq \epsilon^q$. Then we have the estimates:

$$A(N, t) = (t - d - 1)k + N - l + n - 1 \tag{94}$$

$$\leq \epsilon^{-q}k(t\epsilon^q + \chi) - dk + \epsilon^{-p}(N\epsilon^p + \mu) - l + n \tag{95}$$

$$\leq \epsilon^{-q}k(t\epsilon^q + \chi) + \epsilon^{-p}(N\epsilon^p + \mu) + n \tag{96}$$

$$\leq \epsilon^{-\alpha}[k(t\epsilon^q + \chi) + (N\epsilon^p + \mu)] + n \tag{97}$$

where $\alpha = \max(p, q)$. Then:

$$(2M)^{8P_{A(N,t)}(c)} [u : v]_{(L+1)\epsilon}^{cA(N,t)} \tag{98}$$

$$\leq (2M)^{8\frac{\epsilon^{-D}[k(t\epsilon^q + \chi) + (N\epsilon^p + \mu)] + 1}{c-1} - 1} [u : v]_{(L+1)\epsilon}^{\epsilon^{-D}[k(t\epsilon^q + \chi) + (N\epsilon^p + \mu)] + n}. \tag{99}$$

Now combing with these estimates, one obtains the desired estimates:

$$\left(\frac{u(x, s)}{v(x, s)}\right)^{\pm 1} \leq (2M)^{8\frac{\epsilon^{-D}(x+ks)+1}{c-1}-1} [u : v]_{(L+1)\epsilon}^{\epsilon^{-D}(x+ks)+n}.$$

This completes the proof.

Example: Let $b > a \geq 1$ be positive integers, and consider linear PDEs $\mathbf{F}(v_x, v_s) = av_x + bv_s = 0$. For increasing and relative elementary functions f with its leading term \mathbf{F} , let us consider the discrete dynamics:

$$z_{N+1}^{t+1} = f(z_N^t, z_{N+1}^t) = \frac{1}{b}(az_N^t + (b-a)z_{N+1}^t).$$

Let $v : (0, \infty) \times [0, \infty) \rightarrow (0, \infty)$ be C^2 functions, and take the Taylor expansions up to order 2. We choose the scaling parameters by $N = \frac{x}{\epsilon}$, $t = \frac{s}{\epsilon}$ and $z_N^t = v(x, s)$, and insert the Taylor expansions:

$$v(x + \epsilon, s + \epsilon) - f(v(x, s), v(x + \epsilon, s)) \quad (100)$$

$$= \frac{\epsilon}{b}(av_x + bv_s) + \frac{\epsilon^2}{2}((v_{2x} + v_{2s} + 2v_{xs})(\eta_1) - \frac{b-a}{b}v_{2x}(\eta_2)). \quad (101)$$

f correspond to $V_{N+1}^{t+1} = \max(V_N^t, \dots, V_N^t, V_{N+1}^t, \dots, V_{N+1}^t) - \max(0, \dots, 0)$, where their terms iterate a , $b-a$ and b times respectively. Clearly this shows that f are all tropically equivalent independently of $b > a \geq 1$.

By lemma 2.2, the Lipschitz constants $c_f = 1$ are all equal to one. For the numbers of the components, $M_f = b^2$ hold. $D = 1$, $L = 0$, $n = 2$ and $k = 1$. For any positive integers a, b, a', b' , let us take two solutions $u(x, s)$ and $v(x, s)$ satisfying the equations $au_x + bu_s = 0$ and $a'v_x + b'v_s = 0$ respectively. One may assume $b \geq b'$. Suppose both are ϵ_0 controlled bounded by C . Then by corollary 3.4, for any $0 < \epsilon \leq \min(\frac{1}{2C}, \epsilon_0)$, the exponential estimates must hold:

$$\left(\frac{u(x, s)}{v(x, s)}\right)^{\pm 1} \leq (2b^2)^{8(\epsilon^{-1}(x+s)+1)}[u : v]_\epsilon.$$

Below we apply the general procedure of the previous sections to non linear partial differential equations. We treat two PDEs, where one is the quasi linear equations of order 1, and the other is diffusion equations. Given PDE, then our procedure is to find 'good' relative elementary functions f . We have to require them to be increasing. Any elementary polynomials are increasing. One of applicable form of f is:

$$f(z_1, z_2, \dots) = \frac{z_1(\alpha + P(z_1, z_2, \dots))}{1 + z_1} + Q(z_1, \dots)$$

where both P and Q are elementary polynomials and $0 \leq \alpha \leq 1$.

One may weaken the required properties, if both the range and the domain for discrete dynamics are within the regions of monotone increasing for these functions.

4 Applications

4.A Quasi linear equations: Here we introduce a cancelation method of non linear terms and use it to compare solutions between the following equations. Let us consider the equations of the form:

$$v_s + \epsilon v v_x - \frac{1}{2}v^2 = 0, \quad 2u_s + \epsilon u(u_s + u_x) = 0$$

where $\epsilon > 0$ are small constants. These two types of the equations differ from each other, in that for the right hand side, each monomial contains differentials of u , and so in particular any constants are solutions. Notice that $v(x, s) = \frac{c}{1-0.5cs}$ are degenerate solutions on $(0, \infty) \times [0, \frac{2}{c}]$ for $c > 0$.

We choose the second variation:

$$\|u\|_2(x, s) = \sup_{\xi - (x-\epsilon, s-\epsilon) \in D(\epsilon, 1, 1)} \left\{ \left| \frac{\partial^2 u}{\partial x^2} \right|(\xi), \left| \frac{\partial^2 u}{\partial s^2} \right|(\xi), \left| \frac{\partial^2 u}{\partial x \partial s} \right|(\xi) \right\}$$

and put the variation constant:

$$V(u) \equiv \sup_{(x, s) \in (0, \infty) \times [0, T_0)} \frac{\|u\|_2(x, s)}{u(x, s)}.$$

Let us fix any positive constant $V_0 > 0$.

Theorem 4.1. *For any $0 < \epsilon \leq 0.1V_0^{-1}$, let $v, u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be C^2 solutions to the quasi linear equations:*

$$v_s + \epsilon v v_x - \frac{1}{2}v^2 = 0, \quad 2u_s + \epsilon u(u_s + u_x) = 0.$$

Suppose their variation constants $V(u), V(v)$ are bounded by V_0 . Then they satisfy the asymptotic estimates for all $(x, s) \in (0, \infty) \times [0, T_0)$:

$$\left(\frac{u(x, s)}{v(x, s)} \right)^{\pm 1} \leq 40^{2^{\epsilon^{-1}(x+2s)+4}} ([u : v]_{2\epsilon})^{2^{\epsilon^{-1}(x+2s)+3}}.$$

In particular when $u(x, s) \equiv R > 0$ is constant, then the estimates hold:

$$R(40)^{-2^{\epsilon^{-1}(x+2s)+4}} ([v : R]_{2\epsilon})^{-2^{\epsilon^{-1}(x+2s)+3}} \tag{102}$$

$$\leq v(x, s) \leq R(40)^{2^{\epsilon^{-1}(x+2s)+4}} ([v : R]_{2\epsilon})^{2^{\epsilon^{-1}(x+2s)+3}}. \tag{103}$$

4.A.2 Induction of the equations: Let us consider the dynamics:

$$z_{N+1}^{t+1} = f(z_N^t, z_{N+2}^t, z_{N-1}^{t+1}) \equiv \frac{z_{N+2}^t}{2} + \frac{z_N^t(1 + 2z_{N-1}^{t+1})}{2(1 + z_N^t)}.$$

f is an increasing function. The corresponding (max, +)-function is given by $\max(V_{N+2}^t, V_{N+2}^t + V_N^t, V_N^t, V_N^t + V_{N-1}^{t+1}, V_N^t + V_{N-1}^{t+1}) - \max(0, 0, V_N^t, V_N^t)$. The number of the components is $M = 5 \times 4 = 20$, and its Lipschitz constant is equal to 2.

We choose the scaling parameters by:

$$\epsilon v(x, s) = z_N^t, \quad N = \frac{x}{\epsilon}, \quad t = \frac{s}{\epsilon}$$

where we take a small $\epsilon > 0$ so that the estimate $\epsilon \leq 0.1V_0^{-1}$ holds.

Let $v : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be a C^2 function, and take the Taylor expansions up to order 2:

$$v(x+i\epsilon, s+j\epsilon) \tag{104}$$

$$= v + i\epsilon v_x + j\epsilon v_s + \epsilon^2 \left(\frac{i^2}{2} v_{2x} + \frac{j^2}{2} v_{2s} + ij v_{xs} \right) (\xi_{ij}) \tag{105}$$

$$\equiv v + i\epsilon v_x + j\epsilon v_s + \epsilon^2 D^2 v(\xi_{ij}). \tag{106}$$

Let us insert the formal Taylor expansions:

$$\epsilon v(x + \epsilon, s + \epsilon) - f(\epsilon v(x, s), \epsilon v(x + 2\epsilon, s), \epsilon v(x - \epsilon, s + \epsilon)) \tag{107}$$

$$= \frac{1}{2(1 + \epsilon v)} [\epsilon^2 (2v_s + 2\epsilon v v_x - v^2) - 2(\epsilon v) \epsilon^2 D^2 v(\xi_{-11})] \tag{108}$$

$$+ \epsilon^2 (D^2 v(\xi_{11}) - \frac{1}{2} D^2 v(\xi_{20})) \tag{109}$$

where the leading term is given by:

$$\mathbf{F} = \epsilon^2 \frac{2v_s + 2\epsilon v v_x - v^2}{2(1 + \epsilon v)}.$$

The error term is admissible, and let us calculate the error constant $C_{\mathbf{F}^1}$. Notice the estimates $|D^2 v(\xi_{ij})| \leq (\frac{i^2+j^2}{2} + |ij|) \|v\|_2(x, s)$, where $\|v\|_2(x, s)$ is the second variation. Then the error term satisfies the estimates:

$$\|\mathbf{F}^1\|_\epsilon(x, s) \leq \frac{2(\epsilon v) \epsilon^2 |D^2 v(\xi_{-11})|}{2(1 + \epsilon v)} + \epsilon^2 (|D^2 v(\xi_{11})| + |\frac{1}{2} D^2 v(\xi_{20})|) \leq 5\epsilon^2 \|v\|_2(x, s).$$

In particular the error constant is given by:

$$C_{\mathbf{F}^1} = 5.$$

4.A.3 Deformation and cancelation: Let us introduce a cancelation method below. Let us consider the discrete dynamics:

$$w_{N+1}^{t+1} = g(w_N^t, w_{N+2}^t, w_{N-1}^{t+1}) \equiv \frac{w_{N+2}^t}{2} + \frac{w_N^t + w_N^t w_{N-1}^{t+1}}{2(1 + w_N^t)}.$$

g is also an increasing function and is tropically equivalent to f . The number of the components is 16, and the corresponding $(\max, +)$ -function has its Lipschitz constant 2.

Let $u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be a C^2 function, and choose the same scaling parameters, $\epsilon u(x, s) = z_N^t$, $N = \frac{x}{\epsilon}$ and $t = \frac{s}{\epsilon}$. Then let us insert the Taylor expansions of u up to order 2 into the difference as before. Then the direct calculation shows that unlike to the previous case, u^2 term is eliminated, and the result is given by:

$$\epsilon u(x + \epsilon, s + \epsilon) - g(\epsilon u(x, s), \epsilon u(x + 2\epsilon, s), \epsilon u(x - \epsilon, s + \epsilon)) \quad (110)$$

$$= \frac{1}{2(1 + \epsilon u)} [\epsilon^2 (2u_s + \epsilon u u_s + \epsilon u u_x) - \epsilon u \epsilon^2 D^2 u(\eta_{-11})] \quad (111)$$

$$+ \epsilon^2 (D^2 u(\eta_{11}) - \frac{1}{2} D^2 u(\eta_{20})) \quad (112)$$

where the leading term is given by:

$$\mathbf{G} = \epsilon^2 \frac{2u_s + \epsilon u u_s + \epsilon u u_x}{2(1 + \epsilon u)}.$$

In this deformation also, the error term is admissible, and satisfies the estimates:

$$\|\mathbf{G}^1\|_\epsilon(x, s) \leq \frac{\epsilon u \epsilon^2 |D^2 u(\eta_{-11})|}{2(1 + \epsilon u)} + \epsilon^2 (|D^2 u(\eta_{11})| + |\frac{1}{2} D^2 u(\eta_{20})|) \leq 4\epsilon^2 \|u\|_2(x, s).$$

So the error constant is give by $C_{\mathbf{G}^1} = 4$.

Proof of theorem 4.1: Let $u, v : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be C^2 functions which satisfy the equations $v_s + \epsilon v v_x - \frac{1}{2} v^2 = 0$ and $2u_s + \epsilon u(u_s + u_x) = 0$.

Suppose they have bounded variation constants $V(u), V(v) \leq V_0$. Then by applying corollary 3.4 and lemma 3.2, one obtains the asymptotic estimates:

$$\left(\frac{u(x, s)}{v(x, s)}\right)^{\pm 1} \leq (2M)^{8 \frac{e^{\epsilon-D(x+ks)+1}-1}{c-1}} ([u : v]_{(L+1)\epsilon})^{c^{\epsilon-D(x+ks)+n}}$$

for any $0 < \epsilon \leq (2CV_0)^{-1}$, where in this case $D = \max(p, q) = 1$, $C = 5$, $L = 1$, $M = 20$, $c = 2$, $k = 2$ and $n = 3$. Thus for any $0 < \epsilon \leq 0.1V_0^{-1}$, the estimates:

$$\left(\frac{u(x, s)}{v(x, s)}\right)^{\pm 1} \leq (40)^{2^{\epsilon^{-1}(x+2s)+4}} ([u : v]_{2\epsilon})^{2^{\epsilon^{-1}(x+2s)+3}}$$

hold. This completes the proof.

4.B Diffusion equations: Here we introduce a linear deformation of elementary functions, and use it to compare between solutions to different diffusion equations.

Let F be a relative elementary and increasing function of one variable. Here we consider the diffusion equations of the type:

$$u_s = u_{2x} + F(u).$$

We take the third variation:

$$\|u\|_3(x, s) = \sup_{\xi - (x - \epsilon^2, s - \epsilon) \in D(\epsilon, 2, 1)} \left\{ \left| \frac{\partial^3 u}{\partial x^3} \right|(\xi), \left| \frac{\partial^3 u}{\partial s^3} \right|(\xi), \left| \frac{\partial^3 u}{\partial x^2 \partial s} \right|(\xi), \left| \frac{\partial^3 u}{\partial x \partial s^2} \right|(\xi) \right\}$$

and put the variation constant:

$$V(u) \equiv \sup_{(x, s) \in (0, \infty) \times [0, T_0]} \frac{\|u\|_3(x, s)}{u(x, s)}.$$

4.B.2 Linear deformations: Let F be relative elementary and increasing, or zero. We consider the discrete dynamics of the form:

$$z_{N+1}^{t+1} = f(z_{N-1}^{t-4}, z_{N-4}^{t-1}, z_N^t, z_{N+4}^t) = \alpha z_{N-4}^{t-1} + \beta z_{N-1}^{t-4} + \gamma z_N^t + \delta z_{N+4}^t + F(z_N^t)$$

where $\alpha, \beta, \gamma, \delta > 0$ are all positive rational numbers.

We choose the scaling parameters by

$$z_N^t = \epsilon^l u(x, s), \quad N = \frac{x}{\epsilon^m}, \quad t = \frac{s}{\epsilon^{2m}}, \quad (l \geq 0, m \geq 1).$$

For a C^3 function $u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$, let us take the Taylor expansions as before:

$$u(x + i\epsilon^m, s + j\epsilon^{2m}) = u + i\epsilon^m u_x + j\epsilon^{2m} u_s \quad (113)$$

$$+ i^2 \frac{\epsilon^{2m}}{2} u_{2x} + j^2 \frac{\epsilon^{4m}}{2} u_{2s} + ij\epsilon^{3m} u_{xs} + \epsilon^{3m} D_\epsilon^3 u(\xi_{ij}). \quad (114)$$

Firstly we consider the differences:

$$z_{N+1}^{t+1} - \left(\frac{p}{4} z_{N-4}^{t-1} + \frac{1-p}{4} z_{N-1}^{t-4} \right)$$

for $0 < p < 1$. It is immediate to see that this does not contain u_{xs} term. Let us determine $p \in \mathbb{Q}$ so that it also contains no u_{2s} term. In fact for $p = \frac{4}{5}$, the difference is:

$$\frac{3}{4} \epsilon^l u + \frac{7}{5} \epsilon^{l+2m} u_s + \frac{37}{20} \epsilon^{l+m} u_x - \frac{9}{8} \epsilon^{l+2m} u_{2x} + \epsilon^{l+3m} \text{ Higher terms}$$

where Higher terms consist of linear combinations of three derivatives.

Next we eliminate u_x term by adding δz_{N+4}^t for $\delta = \frac{37}{80}$, and then finally we eliminate u terms by adding γz_N^t for $\gamma = \frac{23}{80}$:

$$z_{N+1}^{t+1} - \left(\frac{1}{5} z_{N-4}^{t-1} + \frac{1}{20} z_{N-1}^{t-4} + \frac{37}{80} z_{N+4}^t + \frac{23}{80} z_N^t \right) \quad (115)$$

$$= \epsilon^{l+2m} \left(\frac{7}{5} u_s - \frac{193}{40} u_{2x} \right) + \epsilon^{l+3m} \text{ Higher terms} . \quad (116)$$

Next if we choose constants as below, then one induces the following:

$$w_{N+1}^{t+1} - g(w_{N-1}^{t-4}, w_{N-4}^{t-1}, w_N^t, w_{N+4}^t) \quad (117)$$

$$\equiv w_{N+1}^{t+1} - \left(\frac{1}{24} w_{N-4}^{t-1} + \frac{5}{24} w_{N-1}^{t-4} + \frac{1}{128} w_{N+4}^t + \frac{95}{128} w_N^t \right) \quad (118)$$

$$= \epsilon^{l+m} \left(\frac{15}{8} \epsilon^m v_s + \frac{43}{32} v_x - \frac{19}{16} \epsilon^{3m} v_{2s} \right) + \epsilon^{l+3m} \text{ Higher terms} . \quad (119)$$

g and f above are mutually tropically equivalent. If one exchanges the role of variables and regards x as the time parameter, then the first term of the right hand side equation gives the *advection-diffusion* equation.

Proposition 4.2. *Let us fix $V_0 > 0$ and choose any $0 < \epsilon \leq (200V_0)^{-1}$. Let $u, v : (0, \infty) \times [0, \infty) \rightarrow (0, \infty)$ be C^3 solutions to the linear equations:*

$$\frac{7}{5} u_s - \frac{193}{40} u_{2x} = 0, \quad \frac{15}{8} \epsilon v_s + \frac{43}{32} v_x - \frac{19}{16} \epsilon^3 v_{2s} = 0.$$

Suppose their variation constants satisfy the bounds $V(u), V(v) \leq V_0$. Then they satisfy the exponential asymptotic estimates for all $(x, s) \in (0, \infty) \times [0, \infty)$:

$$\left(\frac{u(x, s)}{v(x, s)}\right)^{\pm 1} \leq 10^{48(\epsilon^{-2}(x+4s)+1)} [u : v]_{5\epsilon}.$$

Proof: Let us consider two linear functions:

$$f(z_{N-1}^{t-4}, z_{N-4}^{t-1}, z_N^t, z_{N+4}^t) = \frac{1}{5}z_{N-4}^{t-1} + \frac{1}{20}z_{N-1}^{t-4} + \frac{37}{80}z_{N+4}^t + \frac{23}{80}z_N^t, \quad (120)$$

$$g(w_{N-1}^{t-4}, w_{N-4}^{t-1}, w_N^t, w_{N+4}^t) = \frac{1}{24}w_{N-4}^{t-1} + \frac{5}{24}w_{N-1}^{t-4} + \frac{1}{128}w_{N+4}^t + \frac{95}{128}w_N^t \quad (121)$$

Let us choose $m = 1$. Then the estimates in corollary 3.4 and lemma 3.2 give the following for $0 < \epsilon \leq (2CV_0)^{-1}$:

$$\left(\frac{u(x, s)}{v(x, s)}\right)^{\pm 1} \leq (2M)^{8\frac{\epsilon^{-D(x+ks)+1}-1}{c-1}} ([u : v]_{(L+1)\epsilon})^{c\epsilon^{-D(x+ks)+n}}.$$

For the corresponding $(\max, +)$ -functions, their Lipschitz constants are both $c = 1$, and the numbers of the components are bounded roughly by $M \leq \frac{10^6}{2}$. For both cases, the error terms are the Higher terms above, consisted by the linear combinations of the three derivatives. So the error constants are roughly bounded by $\frac{1}{6} \times 8 \times 4^3 \leq 10^2$. $k = 4$, $D = \max(p, q) = 2$ and $L = \max(l, d) = 4$. So in this case for any $0 < \epsilon \leq (200V_0)^{-1}$, the estimates:

$$\left(\frac{u(x, s)}{v(x, s)}\right)^{\pm 1} \leq 10^{48(\epsilon^{-2}(x+4s)+1)} [u : v]_{5\epsilon}$$

hold. This completes the proof.

4.B.3 Non deforming: Let us consider the non linear diffusion equations:

$$u_s = u_{2x} + u^a, \quad 1 < a \in \mathbb{Q}.$$

In order to estimate its asymptotics, one considers $v : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$, which is a C^3 solution to the equation $v_s = v^a$. For the initial value $c > 0$, this is easily solved as:

$$v(s) = \frac{c}{(1 - c^{a-1}(a-1)s)^{(a-1)^{-1}}}.$$

The blowing up time is $S_0 = \frac{1}{c^{a-1}(a-1)}$. Its three derivative is given by $\frac{d^3 v(s)}{ds^3} = \frac{c^{1+3\alpha-1}(\alpha+1)(\alpha+2)}{\alpha^2(1-c^{\alpha-1}\alpha^{-1}s)^{\alpha+3}}$, where $\alpha = (a-1)^{-1}$. Thus for $0 \leq s \leq s_0 < S_0$, the variation constant V is bounded by:

$$V(s_0) = \frac{c^{3\alpha-1}(\alpha+1)(\alpha+2)}{\alpha^2(1-c^{\alpha-1}\alpha^{-1}s_0)^3}.$$

Conversely for any $V(0) \leq V_0 < \infty$, there are unique $s_0 < S_0$ so that the equality $V_0 = V(s_0)$ holds.

Theorem 4.3. *Let us fix any $V_0 = V(s_0)$. For any $1 < a \in \mathbb{Q}$ and $T_0 \leq s_0$, let $u : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be C^3 solutions to the diffusion equations:*

$$u_s - u_{2x} = u^a.$$

Suppose their variation constants $V(u)$ are bounded by V_0 . Then for any $0 < \epsilon \leq (200V_0)^{-1}$, u satisfies the asymptotic estimates:

$$\left(\frac{u(x, s)}{v(s)}\right)^{\pm 1} \leq 10^{40 \frac{a\epsilon^{-2}(2x+4s)+1-1}{a-1}} ([u : v]_{5\epsilon})^{a\epsilon^{-2}(2x+4s)+4}.$$

Proof: In 4.B.2, let us choose the rescaling parameters $m = 1$ and $l \in \mathbb{Q}$ so that the equality $l + 2m = la$ holds. In order to induce the above non linear diffusion equations from discrete dynamics, we add non linear term.

Let $\tilde{u} : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be a C^3 function, and consider the discrete dynamics:

$$z_{N+1}^{t+1} - f(z_{N-1}^{t-4}, z_{N-4}^{t-1}, z_N^t, z_{N+4}^t) \quad (122)$$

$$= z_{N+1}^{t+1} - \left(\frac{1}{5}z_{N-4}^{t-1} + \frac{1}{20}z_{N-1}^{t-4} + \frac{37}{80}z_{N+4}^t + \frac{23}{80}z_N^t + \frac{7}{5}(z_N^t)^a\right) \quad (123)$$

$$= \epsilon^{la} \left(\frac{7}{5}\tilde{u}_s - \frac{193}{40}\tilde{u}_{2x} - \frac{7}{5}\tilde{u}^a\right) + \epsilon^{l+3} \text{ Higher terms.} \quad (124)$$

For the corresponding (max, +)-function to f , the Lipschitz constant is $a > 1$, and the number of the components are bounded roughly by $\frac{1}{2}10^5$. The error constant is again roughly bounded by 10^2 , $k = 4$, $D = 2$ and $L = 4$.

Suppose \tilde{u} satisfies the equation $\frac{7}{5}\tilde{u}_s - \frac{193}{40}\tilde{u}_{2x} - \frac{7}{5}\tilde{u}^a = 0$ which admits bounded variation constants $V(\tilde{u}) \leq \tilde{V}$. Then by corollary 3.4 and lemma 3.2, one finds the asymptotic estimates for $0 < \epsilon \leq (200\tilde{V})^{-1}$:

$$\left(\frac{\tilde{u}(x, s)}{v(s)}\right)^{\pm 1} \leq 10^{40 \frac{a\epsilon^{-2}(x+4s)+1-1}{a-1}} ([\tilde{u} : v]_{5\epsilon})^{a\epsilon^{-2}(x+4s)+4}.$$

Let us change the variable x as $\tilde{u}(x, s) = u(px, s)$, where $\frac{1}{2} \leq p = \sqrt{\frac{40}{193} \times \frac{7}{5}} \leq 1$, and put the variation constants of u by $V(u)$. Notice that if $u_s - u_{2x} - u^a = 0$ holds, then \tilde{u} satisfies the equation $\frac{7}{5}\tilde{u}_s - \frac{193}{40}\tilde{u}_{2x} - \frac{7}{5}\tilde{u}^a = 0$. Their variation constants satisfy the estimates $V(\tilde{u}) \leq V(u)$. Thus the asymptotic estimates hold for $0 < \epsilon \leq (200V_0)^{-1}$:

$$\left(\frac{u(x, s)}{v(s)}\right)^{\pm 1} \leq 10^{40} \frac{a^{\epsilon^{-2}(2x+4s)+1-1}}{a^{-1}} ([u : v]_{5\epsilon})^{a^{\epsilon^{-2}(2x+4s)+4}}.$$

This completes the proof.

Remark: Notice that the third derivative rates for $v : [0, s_0] \rightarrow (0, \infty)$ are given by:

$$K(s_0) = \frac{c^{3\alpha^{-1}}(\alpha + 1)(\alpha + 2)}{\alpha^2(1 - c^{\alpha^{-1}}\alpha^{-1}s_0)^{\alpha+3}}.$$

4.B.4 Inhomogeneous non linear equations: The above method does not work for diffusion equations with inhomogeneous non linear terms. In order to treat such cases, we use tropical deformations for relative elementary functions. Here we treat diffusion equations of the form:

$$u_s - u_{2x} - u^a - \delta u^b = 0, \quad (1 < a < b, 0 < \delta \ll 1).$$

Here we cover the equations of the types:

$$(a, b) = (1 + \alpha^{-1}, 1 + 2\alpha^{-1}), \quad 0.5 \leq \alpha \leq 1.$$

Let $\mu = \frac{p}{q} \in \mathbb{Q}_{>0}$ be positive rational numbers, where $p, q \in \mathbb{N}$ are relatively prime numbers. We put $c_\mu \equiv pq \in \mathbb{Z}_{>0}$ and call them as the number of the components for μ .

For $\alpha \in \mathbb{Q}$ and $c > 0$, let us put:

$$V(s_0) = \frac{c^{3\alpha^{-1}}5^3(\alpha + 1)(\alpha + 2)}{6^3\alpha^2(1 - c's_0)^3}, \quad c' = \frac{5c^{\alpha^{-1}}}{6\alpha}, \quad (125)$$

$$(a, b) = (1 + \alpha^{-1}, 1 + 2\alpha^{-1}), \quad \delta = \mu\epsilon^2, \quad \mu = \frac{\alpha + 1}{9\alpha} \quad (126)$$

for $0 \leq s_0 < (c')^{-1}$. Let us compare u with the function:

$$v(s) = \frac{c}{(1 - c's)^\alpha}.$$

Let us fix any $V_0 = V(s_0) \geq V(0)$.

Theorem 4.4. For any $0 < T_0 \leq s_0$ and any $0 < \epsilon \leq (200V_0)^{-1}$, let $u : (0, \infty) \times [0, T_0] \rightarrow (0, \infty)$ be C^3 solutions to the diffusion equations:

$$u_s - u_{2x} = u^a + \delta u^b.$$

Suppose their variation constants $V(u)$ are bounded by V_0 . Then u satisfies the asymptotic estimates:

$$\left(\frac{u(x, s)}{v(s)}\right)^{\pm 1} \leq (2M_\mu)^{8\frac{b^{\epsilon^{-2}(2x+4s)+1}-1}{b-1}} ([u : v]_{5\epsilon})^{b^{\epsilon^{-2}(2x+4s)+4}}$$

where $M_\mu = \max(2 \times 10^3 c_\mu^2, 3 \times 10^4)$.

Proof: Firstly let us consider the tropical deformation:

$$w_{N+1}^{t+1} - g(w_{N-1}^{t-4}, w_{N-4}^{t-1}, w_N^t, w_{N+4}^t) = w_{N+1}^{t+1} - \quad (127)$$

$$\left(\frac{1}{25}(w_{N-4}^{t-1} + w_{N-1}^{t-4}) + \frac{1}{25}w_{N+4}^t + \frac{22}{25}w_N^t + (w_N^t)^a + \mu(w_N^t)^b\right) \quad (128)$$

$$= \epsilon^l \left(\epsilon^{2m} \left(\frac{6}{5}v_s + \frac{4}{25}\epsilon^{2m}v_{2s}\right) + \frac{26}{25}\epsilon^m v_x + \frac{17}{25}\epsilon^{3m}v_{xs} - \frac{4}{25}\epsilon^{2m}v_{2x}\right) \quad (129)$$

$$- ((w_N^t)^a + \mu(w_N^t)^b) + \epsilon^{l+3m} \text{ Higher terms} \quad (130)$$

$$= [\epsilon^{l+2m} \left(\frac{6}{5}v_s + \frac{4}{25}\epsilon^{2m}v_{2s}\right) - \epsilon^{al}v^a - \mu\epsilon^{bl}v^b] \quad (131)$$

$$+ \epsilon^l \left(\frac{26}{25}\epsilon^m v_x + \frac{17}{25}\epsilon^{3m}v_{xs} - \frac{4}{25}\epsilon^{2m}v_{2x}\right) + \epsilon^{l+3m} \text{ Higher terms.} \quad (132)$$

Since $\mu \leq \frac{1}{3}$ hold, the number of the components for g is bounded by $75 \times 25c_\mu^2 \leq 2 \times 10^3 c_\mu^2$. The corresponding Lipschitz constant is b . $D = \max(p, q) = 2m$ and $L = 4$. The error constants are bounded by $\frac{3 \times 8}{25 \times 6} \times 4^3 \leq 11$.

Sublemma 4.5. For $(a, b) = (1 + \alpha^{-1}, 1 + 2\alpha^{-1})$, $0.5 \leq \alpha \leq 1$, one can choose $l \in \mathbb{Q}$ and $m = 1$ so that both the equalities $l + 2m = al$ and $(b - a)l = 2m$ hold.

Proof: By the condition, $m = \frac{b-a}{2}l$ must hold. By inserting into the first condition, one obtains the equality $1 + (b - a) = a$, which certainly hold for the above pairs (a, b) . This completes the proof.

If one chooses $l \in \mathbb{Q}$ and $m = 1$ as above, then the equality holds:

$$\epsilon^{l+2} \left(\frac{6}{5}v_s + \frac{4}{25}\epsilon^{2m}v_{2s}\right) - \epsilon^{al}v^a - \mu\epsilon^{bl}v^b = \epsilon^{l+2} \left(\frac{6}{5}v_s + \frac{4}{25}\epsilon^{2m}v_{2s} - v^a - \mu\epsilon^{2m}v^b\right).$$

Sublemma 4.6. *Moreover let us put $\mu = \frac{\alpha+1}{9\alpha}$. Then for any $c > 0$, $v(s) = \frac{c}{(1-c's)^\alpha}$ ($c' = \frac{5c^{\alpha-1}}{6\alpha}$) satisfy the equations:*

$$\frac{6}{5}v_s + \frac{4}{25}\epsilon^2 v_{2s} - v^a - \mu\epsilon^2 v^b = 0.$$

This can be checked by direct calculations. Notice it satisfies the equation $\epsilon^{l+2m}(\frac{6}{5}v_s + \frac{4}{25}\epsilon^{2m}v_{2s}) - \epsilon^{al}v^a - \mu\epsilon^{bl}v^b + \epsilon^l(a_1\epsilon^m v_x + a_2\epsilon^{3m}v_{xs} - a_3\epsilon^{2m}v_{2x}) = 0$.

Proof of theorem: Three derivative of v is given by $\frac{d^3 v(s)}{ds^3} = \frac{c^{1+3\alpha-1}5^3(\alpha+1)(\alpha+2)}{6^3\alpha^2(1-c's)^{\alpha+3}}$. Thus for $0 \leq s \leq s_0 < S_0 = (c')^{-1}$, the variation constants are bounded by:

$$V(s_0) = \frac{c^{3\alpha-1}5^3(\alpha+1)(\alpha+2)}{6^3\alpha^2(1-c's_0)^3}.$$

As before for any $V(0) \leq V_0 < \infty$, there are unique $s_0 < S_0$ so that the equality $V_0 = V(s_0)$ holds.

For $0 < T_0 \leq s_0$, let $\tilde{u} : (0, \infty) \times [0, T_0) \rightarrow (0, \infty)$ be C^3 functions, and consider the discrete dynamics:

$$z_{N+1}^{t+1} - f(z_{N-1}^{t-4}, z_{N-4}^{t-1}, z_N^t, z_{N+4}^t) \quad (133)$$

$$= z_{N+1}^{t+1} - \left(\frac{1}{5}z_{N-4}^{t-1} + \frac{1}{20}z_{N-1}^{t-4} + \frac{37}{80}z_{N+4}^t + \frac{23}{80}z_N^t + \frac{7}{5}(z_N^t)^a + \frac{7}{5}(z_N^t)^b\right) \quad (134)$$

$$= \epsilon^{la} \left(\frac{7}{5}\tilde{u}_s - \frac{193}{40}\tilde{u}_{2x} - \frac{7}{5}\tilde{u}^a - \frac{7}{5}\epsilon^2\tilde{u}^b\right) + \epsilon^{l+3} \text{ Higher terms.} \quad (135)$$

f and g above are mutually tropically equivalent.

For the corresponding $(\max, +)$ -function to f , the Lipschitz constant is $b > 1$, and the number of the components are bounded roughly by 3×10^4 . The error constant is roughly bounded by 10^2 . $k = 4$, $D = 2$ and $L = 4$.

Suppose \tilde{u} satisfies the equation $\frac{7}{5}\tilde{u}_s - \frac{193}{40}\tilde{u}_{2x} - \frac{7}{5}\tilde{u}^a - \frac{7}{5}\delta\tilde{u}^b = 0$, and admits bounded variation constants $V(\tilde{u}) \leq \tilde{V}$.

Now let us put $M_\mu = \max(2 \times 10^3 c_\mu^2, 3 \times 10^4)$. Then by corollary 3.4 and lemma 3.2, one finds the asymptotic estimates for any $0 < \epsilon \leq (200\tilde{V})^{-1}$:

$$\left(\frac{\tilde{u}(x, s)}{v(s)}\right)^{\pm 1} \leq (2M_\mu)^{8\frac{b\epsilon^{-2}(x+4s)+1}{b-1}-1} ([\tilde{u} : v]_{5\epsilon})^{b\epsilon^{-2}(x+4s)+4}.$$

The rest of the proof is the same as theorem 4.3 just by changing the variable x as $\tilde{u}(x, s) \equiv u(px, s)$, where $p = \sqrt{\frac{40}{193}} \times \frac{7}{5}$. Then for the variation

constants $V(u) \leq V_0$ for u and for any $0 < \epsilon \leq (200V_0)^{-1}$, the estimates hold:

$$\left(\frac{u(x, s)}{v(s)}\right)^{\pm 1} \leq (2M_\mu)^{8^{\frac{b^{\epsilon^{-2}(2x+4s)+1}-1}{b-1}}} ([u : v]_{5\epsilon})^{b^{\epsilon^{-2}(2x+4s)+4}}.$$

This completes the proof.

References

- [F] H.FUJITA, *On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$* , J. Fac. Sci. Univ. Tokyo (I) **13** pp. 109-124 (1966).
- [H] R.HIROTA, *Nonlinear partial differential equations I, III*, Journal of Phys. Soc. Japan **43** pp. 2074-2078, pp.2079-2086 (1977).
- [K1] T.KATO, *Pattern formation from projectively dynamical systems and iterations by families of maps*, To appear in the Proceedings of the 1st MSJ-SI, Probabilistic Approach to Geometry.
- [K2] T.KATO, *Deformations of real rational dynamics in tropical geometry*, Geometric and Functional Analysis **19** No 3, pp 883-901 (2009).
- [KT] T.KATO AND S.TSUJIMOTO, *A rough analytic relation on partial differential equations*, arXiv:1004.4981v1 math.MG (2010)
- [LN] T.-Y.LEE AND W.-M.NI, *Global existence, large time behaviour and life span of solutions of a semilinear parabolic Cauchy problem*, Trans. Amer. Math. Soc. **333** pp. 365-378 (1992).
- [LiM] G.LITVINOV AND V.MASLOV, *The correspondence principle for idempotent calculus and some computer applications*, Idempotency, Ed. J.Gunawardena, Cambridge Univ. Press, pp420-443 (1998).
- [Lo] J.LOGAN, *An introduction to nonlinear partial differential equations*, John Wiley and Sons Inc. Publ., (2008).
- [Mi] G.MIKHALKIN, *Amoebas and tropical geometry*, in Different faces of geometry eds, S.Donaldson, Y.Eliashberg and M.Gromov, Kluwer academic plenum publ., (2004).

- [V] O.VIRO, *Dequantization of real algebraic geometry on logarithmic paper*,
Proc. of the European Congress of Math., (2000).

E-mail: tkato@math.kyoto-u.ac.jp
Department of Mathematics, Kyoto University,
Kyoto 606-8502, Japan