TITLE:

Five amino acid residues in
cysteine-rich domain of human
T1R3 were involved in the response
for sweet-tasting protein,
thaumatin.

AUTHOR(S):

Masuda, Tetsuya; Taguchi, Wakana; Sano, Ayane;
Ohta, Keisuke; Kitabatake, Naofumi; Tani, Fumito

CITATION:

Masuda, Tetsuya ...[et al]. Five amino acid residues in cysteine-rich domain of human T1R3 were involved in the response for sweet-tasting protein, thaumatin.. *Biochimie* 2013, 95(7): 1502-1505

ISSUE DATE:

2013-07

URL:

<http://hdl.handle.net/2433/175269>

RIGHT:

© 2013 Elsevier Masson SAS.; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

Five amino acid residues in cysteine-rich domain of human T1R3 were involved in the response for sweet-tasting protein, thaumatin.

**Authors: Tetsuya Masuda ^{a,*}, Wakana Taguchi ^a, Ayane Sano ^a,
Keisuke Ohta ^a, Naofumi Kitabatake ^b, and Fumito Tani ^a**

**Affiliation: ^a Division of Food Science and Biotechnology, Graduate
School of Agriculture, Kyoto University, Uji, Kyoto
611-0011, Japan**

**^b Department of Foods and Human Nutrition, Notre Dame
Seishin University, Okayama 700-8516 Japan**

*** Corresponding author: Tetsuya Masuda,
Division of Food Science and Biotechnology,
Graduate School of Agriculture, Kyoto University,
Uji, Kyoto 611-0011, Japan
Telephone: +81-774-38-3741; Fax: +81-774-38-3740**

E-mail: t2masuda@kais.kyoto-u.ac.jp

Abstract

Thaumatococcus, a sweet-tasting plant protein, elicits a sweet taste sensation at 50 nM in humans but not rodents. Although it was shown that the cysteine-rich domain (CRD) of human T1R3 (hT1R3) is important for the response to thaumatin, the amino acid residues within CRD critical for response are still unknown. A comparison of the amino acid sequence (69 amino acid residues) of CRD between hT1R3 and mouse T1R3 (mT1R3) revealed sixteen amino acids that differ.

In the present study, we converted each of these sixteen amino acids in hT1R3 to their mouse counterpart and examined the response to thaumatin and sucralose using a cell-based assay. No significant decrease in the response to sucralose was seen among any of the sixteen mutants. However, five mutants (Q504K, A537T, R556P, S559P, and R560K) exhibited a significantly diminished response to thaumatin. The five critical residues involved in the response to thaumatin were dispersed in the CRD of hT1R3 and widely distributed when compared to brazzein.

The unique intense sweet-taste of thaumatin might be attributed to the different receptor activation mechanism compared to the small molecule sweetener sucralose.

Keywords: Thaumatococcus; Sweet-tasting protein; Sweet receptor; Cysteine-rich domain

1. Introduction

Thaumatococcus is the one of the sweetest proteins known and used as a low-calorie sugar substitute as well as for medical purposes for lifestyle-related diseases. Thaumatococcus is 100,000-fold sweeter than sucrose on a molar basis and this intense sweetness makes thaumatococcus useful for unveiling the interaction between sweeteners and sweet receptors. As sweet-tasting proteins are too large to fit the cavity of the interaction sites for small sweeteners, the activation of sweet receptors by sweet-tasting proteins seems to occur in a different manner compared to other small sweeteners [1-3]. Previous mutational studies of thaumatococcus suggested that K67 and R82 are important to the sweetness of thaumatococcus, and mutations at R82 had a more deteriorative effect on sweetness than mutations at K67 [4].

The heterodimers comprising the subunits T1R2 and T1R3, which belong to a family of class C G-protein-coupled receptors, are known to function as sweet receptors [5-8]. Each subunit of sweet receptor possesses a large N-terminal domain (NTD) and a cysteine-rich domain (CRD), and followed by a seven-helix transmembrane domain (TMD). The CRD links the NTD and TMD. Previous studies have shown that sweet-tasting proteins as well as aspartame can be perceived by humans, apes, and Old

World monkeys but not New World monkeys and rodents [9, 10]. Species differences in the response to sweeteners would provide valuable information on the molecular mechanism by which sweet receptors function as well as aid the identification of interaction sites in receptors [10-14]. Recently, we have shown that the CRD within hT1R3 is important for the response toward thaumatin [14]. However, it remains unclear whether two sweet-tasting proteins, thaumatin and brazzein, interact with the same amino acid residues in the CRD of human sweet receptors.

In the present study, to clarify the amino acid residues within the CRD of hT1R3 critical for thaumatin reception and to clarify the mechanisms by which thaumatin activates sweet receptors, we performed site-directed mutagenesis in the CRD of hT1R3 relative to the CRD of mT1R3. The findings should help clarify the activation mechanisms of proteinous sweeteners and might lead to the design of new sweeteners.

2. Materials and methods

2.1. Materials

Thaumatococcus I was purified from crude thaumatin powder as described previously [15]. Sucralose were obtained from Wako Pure Chemical Industries Ltd. (Osaka, Japan).

1

2 2.2. *Site-directed mutagenesis of the CRD of hT1R3*

3 The plasmid pcDNA3.3-hT1R3 was used as the template for
4 mutagenesis [14]. Site-directed mutagenesis was performed using a KOD
5 Plus DNA polymerase (Toyobo Co. Ltd., Osaka, Japan) with two synthetic
6 complementary to opposite strands of oligonucleotide primers containing
7 the desired mutation (Operon Biotechnologies, Tokyo, Japan, Supplement
8 Table 1). The desired mutations were confirmed by DNA sequencing.

9

10 2.3. *Functional expression of human sweet receptors*

11 Human T1R2- and T1R3- or T1R3 mutant-containing plasmids were
12 transiently transfected into HEK293 cells stably expressing the chimeric
13 G-protein, $G\alpha_{16\text{gust44}}$ as described previously [14]. After the transfection,
14 cells were seeded onto polylysine-coated 96-well culture plates (1.5×10^5
15 cells/well) (BD Biosciences, Bedford, MA) and incubated for 24 h. They
16 were then loaded with 50 μL of 3 μM fluo-8 AM (ABD Bioquest Inc.,
17 Sunnyvale, CA) in Hank's balanced salt solution (HBSS) containing 20
18 mM Hepes and 1.25 mM probenecid for 30 min at 37 °C. The cells were
19 incubated with 180 μL of 20 mM Hepes-HBSS containing 0.625 mM
20 probenecid for 10 min at 37 °C. Stimulation was performed by adding 20

1 μ L of agonist solution dissolved in 20 mM Hepes-HBSS. The response to
2 sucralose (1mM) or thaumatin (50 μ M) was detected by measuring
3 fluorescence (excitation at 495 nm and emission at 514 nm) using an
4 Infinite F200 (Tecan Group Ltd., Männedorf, Switzerland) as described
5 previously [14]. The response of hT1R2- wild-type hT1R3 (no mutation)
6 was defined as 100% and the response of each hT1R3 mutant was
7 compared to that of hT1R2- wild-type hT1R3.

8 9 **3. Results and Discussion**

10
11 Since thaumatin elicits an intense sweet taste compared to other
12 artificial sweeteners, identifying the amino acid residues required for the
13 response to thaumatin and the mechanisms by which receptors are activated
14 by thaumatin would shed light on how to design new intense sweeteners.
15 Thaumatin is a basic protein (isoelectric point =12) [16], and we reported
16 previously the importance of the basicity for its sweetness [4, 17]. These
17 positively-charged residues might affect some charged residues within the
18 CRD of hT1R3. First, three mutants (Q504K, E525K, and Q531K) were
19 prepared to examine the response to thaumatin as well as sucralose. No
20 significant decrease in the response to sucralose was observed among the

three mutants, whereas the Q504K mutant showed a significantly diminished response to thaumatin (Fig. 1). Q504 is located in the N-terminal region in the CRD of hT1R3 and is an essential determinant of the specific response to thaumatin (Fig. 2). Since the N-terminal region of the CRD was found to be important for the response to thaumatin, the effect of the mutation E505D which is adjacent to Q504 was investigated. No significant decrease in the response to either sucralose or thaumatin was observed. These results suggested that only the glutamine residue at 504 is essential to the response to thaumatin. Besides the Q504K mutant, a relative decrease in the response to thaumatin was seen for the E525K mutant (Fig. 1B). These results suggest that strict amino acid positions as well as acidic and/or polar residues are important for the response to thaumatin. A relative increase in the response to thaumatin was seen for the Q531K mutant, and the N532H mutation had no effect on responses to thaumatin and sucralose (Fig. 1). The requirement of the N-terminal region of CRD in hT1R3 for the response to thaumatin may be a novel target for the activation of receptors by signal transduction.

Brazzein is a small (6.5 kDa) sweet-tasting protein [18]. Two amino acid residues, A537 and F540, located in the middle of CRD are important for the response to brazzein (Fig. 2A) [11]. Next we examined the effect of

mutating A537 and F540 on the response to thaumatin. As shown in Fig. **1B**, the A537T mutant had a significantly reduced response to thaumatin. Although a relative decrease in the response to thaumatin as well as sucralose was seen in the F540P mutant, the F540P mutant did not significantly affect the response to thaumatin or sucralose in comparison to the A537T mutant (Fig. **1**). These results suggested F540 to be involved in the response to brazzein but not thaumatin. The responses of the mutants I536F, A537T, F540P, G542N, and E545Q to brazzein and small sweeteners such as sucrose were previously examined [11]. Our results showed responses of the I536F, G542N, and E545Q to thaumatin and sucralose to be reduced in comparison with those of wild-type hT1R3 but not significantly and were similar to those of Jiang *et al* [11].

We next converted each of the remaining six amino acids of hT1R3 located in the C-terminal region of the CRD in hT1R3 to their mouse counterpart and examined the response to thaumatin and sucralose. All six mutants (R550K, R553A, F555L, R556P, S559P, and R560K) responded to sucralose, however, three mutants (R556P, S559P, and R560K) showed diminished responses to thaumatin and were located near the transmembrane domain (Fig. **1B**, **2B**). These mutants might not be directly involved in the interaction with the large sweet protein, thaumatin, because

1 of steric hindrance between receptors and large protein-ligands.
2 Interestingly, R556 and S559 resulted in a loss of response with
3 substitution to a proline residue. Previous observations by Jiang *et al*
4 indicated that the mutation A537P resulted in unresponsive to most
5 sweeteners [11]. They suggested that a change in backbone flexibility
6 might alter the formation of the predicted β -strand and thereby alter the
7 conformation of this region in a way that makes it less able to transmit the
8 signal through the receptor. Our observation is distinct in that the loss of
9 response was only to thaumatin, not to sucralose. Taken together, the
10 substitution of proline residues in the C-terminal region of the CRD in
11 hT1R3 seems to be critical for the response to thaumatin.

12 Although the three-dimensional structure of sweet receptors has not
13 yet been determined, homology modeling using a metabotropic glutamate
14 receptor as a template has provided various models for the interaction
15 between sweet receptors and sweet-tasting proteins [1-3]. In the wedge
16 model, sweet-tasting proteins fit into a large wedge-shaped cavity and
17 activate sweet receptors by binding to an external site, thus stabilizing an
18 active form of the receptors in the absence of small sweeteners [3].
19 Recently, the CRD of hT1R3 was implicated in the response to
20 sweet-tasting proteins [8, 11, 14]. However, it was suggested that the CRD

1 plays a major role in the conformational change from the ligand-binding
2 domain to the transmembrane domain [19]. Dose response curves for the
3 five mutations to thaumatin were further investigated (Supplement Fig. 1).
4 These results showed that no significant increases of responses were
5 detected for five mutations up to 100 μ M, suggesting that five residues in
6 the CRD of hT1R3 were involved in the response to thaumatin. Recent
7 high-resolution structural analyses of thaumatin have revealed the
8 flexibility and fluctuation of the side chains of critical residues to be
9 suitable for interaction with sweet receptors [15, 20]. Sweet-tasting
10 proteins would be useful for unveiling how the ligand-binding site of
11 sweet receptors confers a broad and/or specific receptive range.

12 In conclusion, we found only five amino acid residues in the CRD of
13 hT1R3 to be involved in the response to thaumatin and the mechanisms of
14 receptor activation by thaumatin to be different from that for the small
15 molecule sweetener sucralose. Furthermore, the residues involved in the
16 response to thaumatin are dispersed in the CRD of hT1R3.

17 Insights into the molecular mechanism by which thaumatin activates
18 sweet receptors may help in understanding the signal transduction by
19 sweeteners.

20

Acknowledgements

This work was supported by Grants-in-aid for Young Scientists (B) (T.M., no.19780074) and Scientific Research (C) (T.M., no. 22580105) from The Japan Society for the Promotion of Science.

Figures and legends

Figure 1. Five amino acid residues in the CRD of hT1R3 affect the response to thaumatin.

Human T1R2- and T1R3 mutant containing plasmids were transiently transfected into $G\alpha_{16\text{gust44}}$ -expressing HEK cells and responses to sucralose (A) and thaumatin (B) were investigated by cell-based assay. The responses of 16 mutants were averaged and analyzed with a one-way ANOVA (analysis of variance). * $P < 0.05$, ** $P < 0.01$.

Figure 2. Alignment of amino acid sequences of the cysteine-rich domain (CRD) of T1R3 derived from humans and mice.

(A) Conserved residues are indicated with black letters, and non-conserved residues are in white. The residues important for the thaumatin response are indicated in red circles. Two residues, Ala537 and Phe540, previously

identified as important for the response to brazzein [11] are also shown in blue circles. (B) Schematic representation of the structure of the sweet receptor. CRD of T1R3 is shown in blue and the five amino acid residues involved in the response for thaumatin are shown in red. The figures were prepared using Pymol [21] and Modeller [22].

Supplement Figure 1. Dose-response of five mutants of CRD in hT1R3 to thaumatin.

The dose–response analysis of wild-type (black circle), Q504K (red diamond), A537T (brown square), R556P (green triangle), S559P (cyan circle), and R560K (purple triangle). The response of hT1R2- wild-type hT1R3 (no mutation) to thaumatin (50 μ M) was defined as 100%.

References

- [1] P.A. Temussi, Why are sweet proteins sweet? Interaction of brazzein, monellin and thaumatin with the T1R2–T1R3 receptor, FEBS Lett. 526 (2002) 1–3.
- [2] M. Cui, P. Jiang, E. Maillet, M. Max, R.F. Margolskee, R. Osman, The heterodimeric sweet taste receptor has multiple potential ligand

- 1 binding sites, *Curr. Pharm. Des.* 12 (2006) 4591–4600.
- 2 [3] P.A. Temussi, Determinants of sweetness in proteins: a topological
3 approach, *J. Mol. Recognit.* 24 (2011) 1033–1042.
- 4 [4] K. Ohta, T. Masuda, F. Tani, N. Kitabatake, Introduction of a negative
5 charge at Arg82 in thaumatin abolished responses to human
6 T1R2-T1R3 sweet receptors, *Biochem. Biophys. Res. Commun.* 413
7 (2011) 41–45.
- 8 [5] G. Nelson, M.A. Hoon, J. Chandrashekar, Y. Zhang, N.J.P. Ryba, C.S.
9 Zuker, Mammalian sweet taste receptors, *Cell* 106 (2001) 381–390.
- 10 [6] X. Li, L. Staszewski, H. Xu, K. Durick, M. Zoller, E. Adler, Human
11 receptors for sweet and umami taste, *Proc. Natl. Acad. Sci. USA* 99
12 (2002) 4692–4696.
- 13 [7] G.Q. Zhao, Y. Zhang, M.A. Hoon, J. Chandrashekar, I. Erlenbach, N.J.P.
14 Ryba, C.S. Zuker, The receptors for mammalian sweet and umami
15 taste, *Cell* 115 (2003) 255–266.
- 16 [8] F.M. Assadi-Porter, E.L. Maillet, J.T. Radek, J. Quijada, J.L. Markley,
17 M. Max, Key amino acid residues involved in multi-point binding
18 interactions between brazzein, a sweet protein, and the T1R2-T1R3
19 human sweet receptor, *J. Mol. Biol.* 398 (2010) 584–599.
- 20 [9] G. Hellekant, V. Danilova, Species differences toward sweeteners, *Food*

- 1 Chemistry, 56 (1996) 323–3328.
- 2 [10] B. Liu, M. Ha, X.-Y. Meng, T. Kaur, M. Khaleduzzaman, Z. Zhang, P.
- 3 Jiang, X. Li, M. Cui, Molecular mechanism of species-dependent
- 4 sweet taste toward artificial sweeteners, *J. Neurosci.* 31 (2011)
- 5 11070–11076.
- 6 [11] P. Jiang, Q. Ji, Z. Liu, L.A. Snyder, L.M.J. Benard, R.F. Margolskee,
- 7 M. Max, The cysteine-rich region of T1R3 determines responses to
- 8 intensely sweet proteins, *J. Biol. Chem.* 279 (2004) 45068–45075.
- 9 [12] H. Xu, L. Staszewski, H. Tang, E. Adler, M. Zoller, X. Li, Different
- 10 functional roles of T1R subunits in the heteromeric taste receptors,
- 11 *Proc. Natl. Acad. Sci. USA* 101 (2004) 14258–14263.
- 12 [13] A. Koizumi, K. Nakajima, T. Asakura, Y. Morita, K. Ito, A.
- 13 Shimizu-Ibuka, T. Misaka, K. Abe, Taste-modifying sweet protein,
- 14 neoculin, is received at human T1R3 amino terminal domain,
- 15 *Biochem. Biophys. Res. Commun.* 358 (2007) 585–589.
- 16 [14] K. Ohta, T. Masuda, F. Tani, N. Kitabatake, The cysteine-rich domain
- 17 of human T1R3 is necessary for the interaction between human
- 18 T1R2-T1R3 sweet receptors and a sweet-tasting protein, thaumatin,
- 19 *Biochem. Biophys. Res. Commun.* 406 (2011) 435–438.
- 20 [15] T. Masuda, K. Ohta, B. Mikami, N. Kitabatake, High-resolution

- 1 structure of the recombinant sweet-tasting protein thaumatin I, Acta
- 2 Crystallogr. Sect. F67 (2011) 652–658.
- 3 [16] H. van der Wel, K. Loeve, Isolation and characterization of thaumatin
- 4 I and II, the sweet-tasting proteins from *Thaumatococcus daniellii*
- 5 Benth, Eur. J. Biochem. 31 (1972) 221–225.
- 6 [17] K. Ohta, T. Masuda, N. Ide, N. Kitabatake, Critical molecular regions
- 7 for elicitation of the sweetness of the sweet-tasting protein, thaumatin
- 8 I, FEBS J. 275 (2008) 3644–3652.
- 9 [18] D. Ming, G. Hellelant, Brazzein, a new high-potency thermostable
- 10 sweet protein from *Pentadiplandra brazzeana* B. FEBS. Lett. 355
- 11 (1994) 106–108.
- 12 [19] T. Muto, D. Tsuchiya, K. Morikawa, H. Jingami, Structures of the
- 13 extracellular regions of the group II/III metabotropic glutamate
- 14 receptors, Proc. Natl. Acad. Sci. USA 104 (2007) 3759–3764.
- 15 [20] T. Masuda, K. Ohta, F. Tani, N. Kitabatake, Crystal structure of the
- 16 sweet-tasting protein thaumatin II at 1.27 Å, Biochem. Biophys. Res.
- 17 Commun. 410 (2011) 457–460.
- 18 [21] W.L. DeLano, The PyMOL Molecular Graphics System, DeLano
- 19 Scientific San Carlos CA, USA, 2002.
- 20 [22] Z. Yang, K. Lasker, D. Schneidman-Duhovny, B. Webb, C.C. Huang,

- 1 E.F. Pettersen, T.D. Goddard, E.C. Meng, A. Sali, T.E. Ferrin, UCSF
- 2 Chimera, MODELLER, and IMP: An integrated modeling system, J.
- 3 Struct. Biol. 179 (2012) 269-278.

Figure 1.

A **B**

Figure 2.