TITLE:

STUDIES ON MARINE GAMMARIDEAN AMPHIPODA OF THE SETO INLAND SEA. IV

AUTHOR(S):

Nagata, Kizo

CITATION:

Nagata, Kizo. STUDIES ON MARINE GAMMARIDEAN AMPHIPODA OF THE SETO INLAND SEA. IV. PUBLICATIONS OF THE SETO MARINE BIOLOGICAL LABORATORY 1966, 13(5): 327-348

ISSUE DATE:

1966-02-25

URL:

<http://hdl.handle.net/2433/175420>

RIGHT:

STUDIES ON MARINE GAMMARIDEAN AMPHIPODA OF THE SETO INLAND SEA. IV

KIZÔ NAGATA

Inland Sea Regional Fisheries Research Laboratory, Hiroshima

With 4 Text-figures

CONTENTS

PART II. Some Ecological Informations

1. General remarks on the biology of <i>Orchestia platensis japonica</i> (with Fig. 45 and Table 2).....	327
2. An observation on the nocturnal migration of the benthic gammaridean amphipods (with Fig. 46 and Table 3)	331
3. A note on the comparison of species composition between the two different areas (with Fig. 47 and Tables 4-5).....	335
4. Gammaridean amphipods as prey-animals, with special relation to the triglid fishes caught in the Seto Inland Sea (with Fig. 48 and Tables 6-8)	339
REFERENCES	343

PART II. SOME ECOLOGICAL INFORMATIONS

1. General Remarks on the Biology of *Orchestia platensis japonica*

Orchestia platensis japonica is well known as one of the "sand-hoppers" among the gammaridean group and commonly inhabits at the high-water marks on the beach; often found in a great abundance under damp seaweed or straw-mat washed ashore, sometimes living under dead leaves in the damp places far above the sea-shore. The animal is a scavenger, showing the feeding habit of a "biting" type, and often seen crowding together and biting at the shucked meat of oyster while the animals are kept in an experimental glass-vessel in the laboratory.

The material was collected once a month from March 1962 to March 1963, at the high-water mark on the beach of Ônoura, Saeki-gun, Hiroshima Pref., where the sampling spot was located just beneath the stone wall, submerged always in the high-tide of each lunar month and covered all over with

Fig. 45. Seasonal change of size-composition of *Orchestia platensis japonica*. (For female: ovigerous female; females whose ovary is well developed; females bearing the empty brood pouch).

empty shells of "Asari". Sampling in each month was made in the daytime of the new moon or the full moon when the tide was outgoing, keeping a constant distance from the stone wall. The specimens were put quickly into the collecting bag by a small elliptical hand shovel.

The length of animals is measured from head to telson along the natural curve of the dorsal line. All the individuals or a part of the total collection of each month were measured and sorted into size groups. Seasonal change of size-composition may be seen by comparing one another the results of analyses of the specimens scooped by shovel in respective months (Fig. 45). The individuals less than 4.5 mm long are defined as the sexually indifferenced youngs and treated together with the female. Adult females examined are classified into three groups according to the states, ovigerous, with fully developed ovary, or with the empty brood pouch. The number of embryos or youngs within the brood pouch is counted, when the animals are ovigerous.

The spawning appears to continue nearly throughout the year, and therefore it is difficult to define distinctively each spawning group, but it is

Table 2. Collecting data in reference to the investigation on the biology of *Orchestia platensis japonica*.

Date	Time	Age of moon in days	W.T. (°C)	Extracting rate	Material examined				
					Total	Indiffer- entiated young 4.5 mm >	Female		Male
							4.6~ 6.0 mm	6.1 mm <	
7/III	12.00	0.7	10.7	1/4	406	23	80	236	67
6/IV	11.45	1.3	12.0	1/4	212	7	11	154	40
4/V	11.00	29.3	18.0	1/4	208	20	2	119	67
4/VI	10.40	1.6	19.2	1/2	174	75	3	68	28
2/VII	10.30	0.1	21.6	1/2	157	52	45	37	23
2/VIII	11.00	1.6	26.8	1/1	212	125	64	5	18
1/IX	11.10	2.0	27.2	1/1	66	6	20	20	20
1/X	11.50	2.3	24.2	1/1	216	118	12	63	23
1/XI	13.00	3.6	19.9	1/1	181	48	30	71	32
12/XII	11.45	14.9	12.9	2/3	146	12	13	74	47
10/I	11.00	14.2	10.2	1/2	210	19	37	88	66
25/II	11.50	1.0	6.2	1/3	189	38	22	65	64
26/III	11.20	0.6	11.0	1/2	187	1	8	134	44

probable that two active spawnings are seen in the period from spring to early summer and in the autumn season from October to November. The spawning season in spring is more prominent than that in autumn, and the total number of spawns is considered to be most abundant in this season throughout the year, although a large number of the newly hatched small specimens are not shown in the figure; this is probably on account of either

the error of sampling or of the operation extracting samples from the total specimens collected. Similar seasonal change of size composition and the presence of two main spawning seasons, were also observed at another station on the same beach, at a certain damp place where the kitchen residues were piled up.

Ovigerous females are 6.0-6.5 mm in length at the minimum. For the external appearances of the adult females drawing toward the spawning, the marsupial plates begin to develop at first, in time the ovary comes to mature fully so that the purplish blue mass can be seen through the dorsal chitinous cuticle of the pereion, at this time the marsupial plates have fairly developed, but not yet armed fully with the marginal setae, and then the individuals bearing embryos within the brood pouch come to appear. The moulting will be taken just before the issue of mature eggs into the brood pouch, and after this moulting the so-called "riding position" will be seen again, followed then by a copulative posture immediately before or in the middle of the issue of mature eggs. It is pretty sure that the fertilization takes place outside the body. During the breeding period with embryos within the brood pouch, the marsupial plates develop fully and are densely armed with long marginal setae. After the release of young or embryos, the fully developed marsupial plates remain empty for a short period and then fall off in time. Such breeding processes are repeated several times by each adult female during the spawning season. Durations for respective processes (e.g. of riding position, purplish blue state of ovary, of incubation of embryos within the brood pouch, and of retaining the empty pouch), are not made clear as yet; this requires many careful observations in laboratory in future. It is, here, to be noted that the purplish blue state of ovary has never been seen from the outside in all of 434 ovigerous specimens. The mature eggs within the ovaries may all be issued into the brood pouch in a short time.

Thus, as shown in the figure, the females defined as "females whose ovary is well developed" always indicate to be in a state prior to the ovigerous stage, and "females bearing the empty brood pouch" always in a state after the release of young or embryos from the brood pouch. The latter is clearly distinguished from the former in having marsupial plates fully developed, but no sign of the purplish blue mass of ovary.

The size of mature ova within the ovary is 0.32 mm-0.43 mm in diameter. The embryonal stages within the brood pouch were provisionally divided into "eggs", "medium", and "advanced" stages, respectively 0.62 mm, 0.71 mm, and 0.79 mm in diameter on an average. In the "advanced" stage, the appendages of the young are more or less perceptible through the chorion, and the shape is almost ovoid, 0.83-1.05 mm in longer diameter. The newly hatched young is 1.5-2.0 mm long.

More than 50 percent of ovigerous females bear 5-9 embryos within the brood pouch respectively, but 22 at the maximum in a 10.7 mm long specimen. When a few number of embryos are included within the brood pouch, they are usually consisting of only youngs or embryos of the advanced stage, while when abundant, all or most of them are the eggs of the earlier stages. And it is often met with that embryos in some different stages of development are observed within the same brood pouch. These facts seem to show the possibility that the embryos may hatch or be released from the brood pouch by and by in the order of development. On the other hand, however, it is generally said in some members of both Isopods and Tanaidaceans that the number of the fertilized eggs is reduced by the expulsion of embryos before their development is completed (HOWES 1939, p. 290). Anyhow, the number of embryos within the brood pouch is very variable so far as I examined, and no significant correlation could be found between it and the season or the size of mother. If the fertilized eggs in the brood pouch could be counted exactly soon after the issue from the ovary, there might be found any relation between the egg number and some factors.

The gammaridean animal is usually said to live on for about a year. The present specimens fully grown up attain nearly to 15.0 mm long in both sexes. The spawning group seen in March 1963 is likely to belong to the group hatched in the season April to June 1962. Now, taking it into account that the size of the newly hatched specimens is 1.5-2.0 mm long, the average growth rate per a month may be roughly estimated to be about 1.0 mm, although the growth rate naturally differs according to the water temperature. Sex-ratio indicates no positive relation with the season, even though the specimens less than 6.0 mm are excluded.

2. An Observation on the Nocturnal Migration of Benthic Gammaridean Amphipods

The nocturnal migration of gammaridean amphipods inhabiting the sea floor or among seaweeds is one of the most interesting phenomena. The object of this observation was first to obtain the general outline of the upward movement at night, and second to distinguish the species migrating up to the surface waters at night from many benthic amphipods living abundantly both in number of individuals and species on the *Zostera* belt near the low-water mark.

It was planned to see successive changes in the number of individuals at different hours of observation from nightfall to dawn. The result is represented graphically in Figure 46. The "number of individuals" is represented by the number of gammarideans per horizontal haul of a plankton

Fig. 46. Diagram showing the diurnal change of number of individuals of the benthic gammarids per sample obtained by the horizontal towing of plankton net in the surface waters on *Zostera* belt near the low-water mark. The solid column indicates the proportion of *Paradexamine barnardi* to the total sample.

net (stretched with GG 40, mesh 39/inch), towed by a return sailing of a distance of about 250 meters in the surface water above the *Zostera* belt off the Ôno-Branch of our laboratory, Ônoura, Saeki-gun, Hiroshima Pref.

In the Seto Inland Sea, the tide range is fairly prominent, and the tidal flow is very fast, particularly in "Seto" (strait). This *Zostera*-belt is facing Ôno Seto, and narrowly spreads outside the inlet in the direction along the tidal flow; therefore, it is desirable to collect at the time when the tidal flowing is dead. Moreover, the *Zostera*-belt is very shallow being about 0.5 meter in depth in low water at the spring tide. Thus, the collecting was always done in the slack water just before the high water, and then it was impossible to carry out successive observations in the same day.

As seen in Table 3, the samples of both series A and C were collected after the moonset, while the series B was taken after the moonrise. No distinctive relations, however, could be seen between the migration and the moon light so far as the present collections were concerned. In general, the number of individuals rapidly increases as soon as it becomes completely dark after the nightfall, reaching to the maximum just before 22.00, then

Table 3. Data of the observations on the nocturnal migration of benthic gammarids.

Series	Date	Time	Age of moon in days	Moon-		Sun-		Depth (m)	W.T. (°C)
				rise	set	rise	set		
A	30/VIII	22.05	29.6	5.30	18.55	5.42	18.40	3.5	25.4
	6/IX	0.01	7.0	11.51	22.38	5.46	18.31	2.4	24.7
	9/IX	3.20	10.0	14.41	0.07	5.48	18.26	2.3	25.6
	10/IX-a	5.08	11.0	15.36	1.01	5.49	18.25	2.6	24.6
	11/IX	5.58	12.0	16.28	2.02	5.50	18.25	2.5	24.2
B	10/IX-b	18.35	11.0	15.36	1.01	5.49	18.25	3.0	25.4
	12/IX	19.35	13.0	17.16	3.07	5.50	18.23	3.0	25.0
	14/IX	21.15	15.0	18.42	5.28	5.52	18.20	3.7	25.0
	16/IX	22.15	17.0	20.01	7.50	5.53	18.17	3.2	25.5
	20/IX	0.30	21.0	22.55	12.19	5.56	18.11	3.0	24.8
	21/IX	1.35	22.0	23.45	13.18	5.56	18.10	2.6	24.5
	22/IX	2.45	23.0	—	14.12	5.57	18.08	2.4	25.0
	23/IX-a	4.50	24.0	0.39	15.00	5.58	18.07	2.2	25.1
	24/IX-a	5.45	25.0	1.34	15.43	5.59	18.05	2.6	25.1
C	23/IX-b	18.15	24.0	0.39	15.00	5.58	18.07	2.3	25.3
	24/IX-b	19.00	25.0	1.34	15.43	5.59	18.05	2.5	25.0
	27/IX	20.00	28.0	4.19	17.28	6.01	18.01	3.0	23.8
	1/X	21.45	2.3	7.55	19.29	6.04	17.56	3.2	23.4
	5/X	23.25	6.3	11.37	22.02	6.07	17.51	2.5	22.2
	8/X	1.00	9.3	14.17	—	6.10	17.46	2.1	22.5
	9/X	3.00	10.3	15.06	0.48	6.10	17.45	2.1	22.5
	10/X	5.15	11.3	15.51	1.54	6.10	17.44	2.6	22.8

gradually decreases towards the dawn, and animals wholly disappear while the sky is turning gray before dawn. The result of series A seems unusual.

Nevertheless, these observations made only in high water of different days are considered to accord with the day of lunar month. The results indicate, thus, the number of migrating individuals increases in the spring tide of the full moon or new moon. Similar phenomenon is also observed in the nocturnal migration of members of amphipod genus *Bathyporeia* carried out in the intertidal waters of Kames Bay, open to the Firth of Clyde, by WATKIN (1939). He says there, "when the samples are related to the day of lunar month it is shown that the numbers increase in the tidal waters in the periods immediately preceding the full moon and preceding and partly overlapping the new moon." My results show that the animals stop migrating upwards in the day time, even in the twilight. It is still questionable if the numbers always show the maximum at about 22.00 at night, although this time corresponds just to the high water of the new moon or full moon as it was so in my observations. In future, the nocturnal change of the number of migrating individuals throughout the same night will be clearly

explained by further investigations made at the location in a certain much deeper inshore area, deep enough even in the low water of neap tide and without any prominent tidal flow in neither spring nor neap tide, which will disturb the towing of plankton net.

The total number of individuals of each species obtained is given below, together with the frequency of occurrences shown by the number of samples for each species:

	(Number of individuals)	(Frequency)
<i>Paradexamine barnardi</i>	607	18 (all)
<i>Aoroides columbiae</i>	45	10
<i>Erichthonius pugnax</i>	41	11
<i>Corophium uenoi</i>	39	6
<i>Synopia ultramarina</i>	23	7
<i>Pontogeneia rostrata</i>	21	7
<i>Corophium</i> sp. (juv.)	19	8
<i>Byblis japonicus</i>	16	8
<i>Pontocrates altamarinus</i>	11	8
<i>Gitanopsis vilordes</i>	10	4
<i>Maera serratipalma</i>	8	5
<i>Grandidierella japonica</i>	6	5
<i>Stenothoe gallensis</i>	4	4
<i>Melita koreana</i>	4	4
<i>Orchomenella littoralis</i>	4	3
<i>Corophium acherusicum</i>	4	2
<i>Eriopisella sechellensis</i>	2	2
<i>Melita</i> sp. (juv.)	2	2
<i>Jassa falcata</i>	2	2
<i>Ampelisca miharaensis</i>	1	1
<i>Aoroides secunda</i>	1	1
<i>Ampithoe lacertosa</i>	1	1
<i>Microjassa cumbrensis</i>	1	1

Paradexamine barnardi occurred most abundantly nearly in all samples, showing 69.6 percent of the total individuals of all species. It is noticeable that such groups as the burrowing or tubicolous forms often appear in surface water at night. These facts seem to show that nearly all of the benthic gammarideans act the nocturnal migration. Unfortunately, the present observations were made in the season when the gammarideans were rather scarce, and so it is very desirable that further investigations will be carried out in the season when much larger numbers of individuals are available, for instance in spring to early summer.

There has been discussions by some authors on the factors causing this nocturnal migration, and the influence of the tide is regarded as an important factor in the tidal area. This may be supported also in the present observation by the increase of individual number in the high tide of the new and full moon. The migration may really have something to do with the

breeding cycle for each species. Nearly all the individuals obtained were small specimens; further detailed analyses are needed on the size and age compositions and sex-ratio.

3. A Note on the Comparison of Species Composition Between Two Different Areas

Nearly all of the species described in this paper are found in relatively shallow waters, from the *Zostera*-belt near low-water mark to about 60 meters deep; and the bottoms of the localities are in most cases composed of silt, mud, or sandy mud. Unfortunately, no detailed analyses of the nature of the bottom inhabited by animals have been made, so that the relations between the animals and the important factor, the soil grade, are almost unknown. But there is an interesting difference between the species obtained from *Zostera*-belt near low-water mark and those collected from the deeper area. The '*Zostera*-belt near low-water mark' mentioned here is the place where the tide does not completely go out even in the spring tide and about 5 meters at the deepest place in low-water. Such a *Zostera*-belt usually offers a habitat suitable for many young fishes as well as for small crustaceans important as prey-animals. Particularly gammarideans are very rich there in the number of both individuals and species, and occupies 60-90 percent of small crustaceans on the *Zostera*-belt of Mihara Bay (NAGATA 1960). The abundant occurrences of gammarideans on the *Zostera*-belt is obviously shown also by the stomach contents of fishes caught there (KITAMORI, NAGATA, and KOBAYASHI 1959).

The species of gammarideans very commonly found on the *Zostera*-belt near low water mark are listed below:

1) <i>Paradexamine barnardi</i>	10) <i>Corophium acherusicum</i>
2) <i>Pontogeneia rostrata</i>	11) <i>Byblis japonicus</i>
3) <i>Ampithoe lacertosa</i>	12) <i>Eriopisella sechellensis</i>
4) <i>Ampithoe valida</i>	13) <i>Pontocrates altamarinus</i>
5) <i>Grandidierella japonica</i>	14) <i>Orchomenella littoralis</i>
6) <i>Erichthonius pugnax</i>	15) <i>Aoroides columbiae</i>
7) <i>Pleustes panopla</i>	16) <i>Eurystheus japonicus</i>
8) <i>Cerapus tubularis</i>	17) <i>Corophium uenoi</i>
9) <i>Jassa falcata</i>	

Of the above-listed species, species 11, 12, and 13 are found abundantly and species 14-17 are found occasionally also in the area more deeper than the *Zostera*-belt near low-water mark. While the species 1-10 are mostly limited to the *Zostera*-belt. On the *Zostera*-belt of Mihara Bay (Area IV), species 1-5 were found very abundantly in net-collections, while species 6 was unexpectedly rather scarce. The list of the species living very abundantly on the inshore *Zostera*-belt is based on the materials obtained from Mihara Bay in

Fig. 47. Map showing station localities in two surveys of "Area XI". Marks ○ and × indicate respectively stations in 1959 (Area-a) and 1960 (Area-b). Sts. 16-22 in 1960 nearly coincide with Sts. 29-23, and so does St. 11 in 1959 with Sts. 19 and 26 in 1960.

the Seto Inland Sea and near Asamushi in Mutsu Bay. On the *Zostera*-belt of the latter locality, species 1, 3, 6-9, and 13-16 were very rich.

On the other hand, general aspects of their occurrences in the area deeper than the *Zostera*-belt near low-water mark are well represented in Tables 4 and 5 (see also Figs. 1 and 47). The collections in this area were made by drawing a bottom-layer net (NAGATA 1960, fig. 1) on the soft sea floor for 1.5-4 minutes at each station, but the drawing can never be done under the constant conditions throughout the stations and therefore the collections are not to be estimated quantitatively. However, several common species widely distributed in the littoral areas below the level of the low-water mark may be clearly shown in the catches of these two surveys. More than 90 percent of total individuals in each catch is occupied by the following 10 species: all the 7 species of Ampeliscidae, *Pontocrates altamarinus*, *Eriopisella sechellensis*, and *Bathymedon*

longimanus. Most of them have often been found in the stomach contents of benthos-feeding fishes, too. J. L. BARNARD started in the opening of his paper of 1954b, "Amphipoda of the family Ampeliscidae form an important component of any littoral marine soft-bottom fauna", and this appears to be the fact common to all amphipod faunas of the littoral sea floor along the Japanese coast deeper than about 10 meters. *Eurystheus utinomii*, *Anonyx nugax pacificus*, and two species of the genus *Photis* have been collected most abundantly at the three stations deeper than 10 m in Mihara Bay (NAGATA 1960, fig. 2), but not so numerous as expected in the two surveys.

As mentioned above, a prominent difference in their occurrences was noted between the species commonly found in respective areas, but other species occur comparatively less abundantly and less frequently, and thus it is unable to show definitely for each species the area of the richest occurrence.

Table 4. Species composition at each station, based on the bottom-layer net catches made in the east area of Suô Nada (Area XI-a) in the period June 12-16, 1959.

Total: 5673 specimens. Depth: 32-56 m. (see Figs. 1 and 47).

[illegible]

4. Gammaridean Amphipods as Prey-animals, with Special Relation to the Triglid Fishes Caught in the Seto Inland Sea

As previously mentioned again and again, amphipods are important prey-animals often found abundantly in the stomach contents of many benthos-feeding fishes. To make clear the trophic relations between higher consumers and their preys, our laboratory carried out the collecting of benthic fishes by the experimental two-boat trawler (30 tons & 80 H.P.), which had eight cruises over eleven "Nada"-areas around the coast of Shikoku in the period from Sept. 1957 to Sept. 1960 and obtained enormous catches of benthic fishes. And I have had an opportunity to observe the stomach contents of nearly all of the benthic fishes, but piscivorous fishes and plankton feeders.

For an example, a qualitative analysis of their feeding frequency by food-group of the fishes caught during the fourth cruise made in May-June 1958 is given in Table 6. Here, the feeding frequency is shown by the percentage of the number of fish feeding on a respective food-group to the total number of the fish examined. The total number of fishes examined here amounts to 2880 individuals, including 75 species, but in this table, only the results for 31 species are given, the species represented by less than 10 individuals being omitted. As seen on this table, main food items of those benthic fishes except for piscivores in the Seto Inland Sea are shrimps, small crustaceans, annelids, and fishes; although the proportions of these food animals are different with feeders, nearly all of fishes are known to show a fairly high percentage of shrimps in the stomach contents. Fishes or squids which all the feeders listed in this table preyed upon are small in size and not so significant quantitatively so far as concerned with those found in the Seto Inland Sea. Strictly saying, all the values in the table may be swayed to some extent by the number of feeding individuals examined, variation of their size, and also by the seasonal change of prey-animals themselves; although they may be available to get the general view of the food organisms of the fishes.

Relatively high percentages of small crustaceans are seen in flatfishes and triglid ones. Especially it is noteworthy that unexpectedly large amount of small crustaceans including gammaridean amphipods are eaten by the latter. Particularly as to *Lepidotrigla microptera*, a large number of individuals were examined so that the above-mentioned trend seems to be unsuspecting. It is supposed that triglid fishes bear some resemblance to flatfishes in their feeding habits. In feeding they usually lie prostrate on the bottom and feel the sea floor by means of their long finger-like pectoral filaments.

Of course, like other feeders, triglid fishes show some changes in the composition of food animals with growth. Table 7 shows the percentage of

Table 6. Feeding frequency of the benthic fishes caught during May-June 1958, by food-group.

Food group		Range of T.L. (cm)	Number of feeding individuals examined	Fish	Squid	Shrimp	Crab	Stomatopod	Small crustacean	Annelid	Molluscan	Echinoderm	Undet. contents
Feeding fish													
Triglidae													
Lepidotrigla abyssalis		8-15	11	—	9	73	—	—	64	18	—	—	—
L. guntheri		9-21	12	—	—	83	—	—	8	17	—	—	—
L. japonica		8-19	30	—	20	80	7	—	57	10	7	—	—
L. microptera		7-30	287	2	7	89	1	—	39	7	3	—	2
L. kishinouyei		14-24	37	—	3	100	—	—	27	—	3	—	—
Pachytrigla alata		11-22	15	7	—	94	7	—	—	—	—	—	—
Flatfishes													
Laeops lanceolata		8-13	41	—	—	—	—	—	—	100	—	—	—
Areliscus sp.		12-17	31	—	—	—	—	—	3	90	—	—	16
Poecilopsetta plinthus		6-11	22	—	—	—	—	—	73	64	—	—	9
Pleuronichthys cornutus		9-17	20	—	—	5	—	—	5	75	5	—	25
Limanda yokohamae		14-21	19	—	—	32	—	—	26	63	16	—	32
Areliscus interruptus		6-10	39	—	—	5	—	—	21	54	—	—	28
Psettina iijimai		5- 8	115	—	2	2	1	—	96	45	—	—	—
Scidorhombus pallidus		5- 6	13	—	—	23	—	—	92	8	—	—	—
Eopsetta grigorjewi		12-19	41	—	2	61	7	—	44	—	—	—	—
Arnoglossus japonicus		7-10	57	14	—	60	—	—	74	—	—	—	—
Pseudorhombus pentophthalmus		6-13	100	25	6	44	1	—	47	2	—	—	—
Sparidae													
Chrysophrys major		12-35	16	—	—	69	—	—	—	—	—	6	25
Branchiostegidae													
Branchiostegus spp.*		11-32	55	29	6	33	4	—	16	22	—	4	11
Nemipteridae													
Nemipterus spp.**		9-47	39	41	3	56	18	—	5	10	—	—	—
Sciaenidae													
Argyrosomus argentatus		9-23	274	37	—	68	—	—	12	1	—	—	7
Platycephalidae													
Kumococius detrusus		14-26	22	27	—	59	5	—	5	—	—	—	5
Suggrundus meerdervoorti		14-23	32	25	—	91	—	—	3	—	—	—	—
Cociella crocodila		20-44	18	33	6	50	28	33	—	—	6	—	—
Lamnina													
Mustelus griseus		42-85	11	27	—	55	36	18	9	—	—	—	—
M. manazo		40-77	63	16	—	81	41	11	6	—	—	—	—
Rajina***													
Holorhinus tobije		13-35	28	—	—	82	—	—	—	—	75	—	—
Raja kenoei		9-24	58	14	—	86	3	—	2	—	—	—	5
Dasyatis akajei		15-32	27	30	—	85	4	—	7	4	—	—	4

*) Comprising two species *B. argentatus* and *B. japonicus japonicus*.**) Comprising two species *N. virgatus* and *N. bathybus*.

***) Measured by 'Body Length'.

Table 7. Feeding frequency of *Lepidotrigla microptera* caught during June 1959, by size and by food-group.

T.L. (mm)	<120	<160	<200	<240	<280	<320	Total
Number of feeding individuals	1	12	103	69	35	5	225
Fish	—	—	3.8	5.7	11.4	—	5.3
Squid	—	—	5.8	2.9	2.9	—	4.0
Large shrimp	—	41.6	43.6	50.7	71.4	80.0	50.6
Small shrimp	100.0	66.6	56.3	55.0	71.4	40.0	58.6
Crab	—	—	0.9	—	—	—	0.4
Amphipod	—	75.0	43.6	26.0	22.8	20.0	36.0
Cumacean	—	—	1.9	—	—	—	0.9
Polychaete	—	16.7	4.9	2.9	5.7	—	4.9
Molluscan	—	—	3.9	2.9	5.7	—	3.6

the feeding frequency by food-group and by size in *Lepidotrigla microptera* caught during the cruise in June 1959. With the increase in size, the percentages of large shrimps and fishes grow higher, while those of small shrimps and amphipods gradually decrease. Small crustaceans were represented by only two groups of gammaridean amphipods and cumaceans.

Table 8. Feeding frequency of triglid fishes caught during Sept. 1957 to July-Aug. 1958, by season and by food-group.

	Sept. '57	Dec. '57	Jan.-Feb. '58	May-June '58	July-Aug. '58
Fish	5.5	11.0	2.9	1.8	9.1
Squid	6.3	1.3	—	7.3	2.5
Shrimp	70.8	91.9	80.9	87.9	84.4
Crab	3.3	2.3	10.3	1.5	3.6
Stomatopod	1.4	3.9	1.5	—	0.4
Small crustacean	19.6	13.3	19.1	37.1	13.8
Polychaete	1.4	6.1	—	6.6	1.8
Molluscan	0.3	6.1	—	3.0	—
Echinoderm	—	0.3	—	—	0.4
Others	—	1.0	—	0.5	—
Undet. contents	7.2	—	2.9	2.3	8.7
Numbers of feeding individuals	363	235	53	396	275

On the other hand, the feeding frequency of triglid fishes caught in the period from Sept. 1957 to July-Aug. 1958, by season and by food-group, is shown in Table 8. The feeding frequency for shrimps was highest throughout, and that of small crustaceans was always relatively higher than those of other prey-groups excepting shrimps. Figure 48 shows the seasonal change of the feeding frequency for small crustaceans given in Table 8, together with that for gammaridean amphipods eaten by various fishes living on the

Fig. 48. The seasonal change of the feeding frequency for small crustaceans (---) and for gammaridean amphipods (—). The dotted line refers to that of triglid fishes given in Table 8, while the solid line concerns that of the various fishes living on the *Zostera* belt near the low-water mark in Mihara Bay.

Zostera-belt of Mihara Bay (KITAMORI, NAGATA et KOBAYASHI 1959, fig. 8). Amphipods are the largest component of small crustaceans found in the stomach of triglid fishes treated in Table 8, and therefore it is very interesting that the two curves shown in Figure 48 indicate a similar tendency; one refers to the offshore areas, while the other concerns the inshore region near the low-water mark. Results of many surveys in Mihara Bay indicate that the seasonal change of the feeding frequency for gammaridean amphipods agrees fairly well with the seasonal variation in the quantity of the amphipods themselves in that habitat. And this means nothing but that triglid fishes

in the Seto Inland Sea feed relatively well on small crustaceans.

Now throughout the five-month data shown in Table 8, the percentage composition of the total small crustaceans found in the stomach contents of triglid fishes is given as follows:

Amphipoda	
Gammaridea	54.1
Caprellidea	5.4
Cumacea	24.4
Ostracoda	9.6
Mysidacea	5.6
Brachyura & Macrura larvae	5.6
Isopoda	0.09
Nebaliacea	0.09

More than half of small crustaceans consist of gammaridean amphipods, of which several dominant species are given below together with the percentage composition:

<i>Ampelisca brevicornis</i>	}	23.2
<i>Ampelisca cyclops</i>		
<i>Ampelisca bocki</i>		
<i>Ampelisca misakiensis</i>		
<i>Ampelisca miharaensis</i>		
<i>Ampelisca naikaiensis</i>		
<i>Byblis japonicus</i>		17.5
<i>Pontocrates altamarinus</i>		11.7

<i>Eurystheus utinomii</i>	}	37.5
<i>Eurystheus japonicus</i>		
<i>Photis longicaudata</i>	}	0.3
<i>Photis reihardi</i>		
<i>Liljeborgia japonica</i>		0.4
<i>Podocerus inconspicuus</i>		3.4
Other species		5.9

It is seen that 89.9 percent of the total are occupied by 10 species of the genera *Ampelisca*, *Byblis*, *Pontocrates*, and *Eurystheus*. The conjecture that these species important as prey-animals must be inhabiting abundantly on the offshore floor is supported actually by two benthos-collections shown in Tables 4 and 5, namely the above 10 species account for 78.6 percent of the total individuals in Area XI-a (Table 4) and for 96.2 percent in Area XI-b (Table 5).

REFERENCES

- ALDERMAN, A. L. 1936. Some new and little known amphipods of California. Univ. Calif. Publ. Zool., 41 (7): 53-74, 51 text-figs.
- BARNARD, J. L. 1952. Some Amphipoda from Central California. Wasmann Jour. Biol., 10 (1): 9-36, 9 pls.
- 1954. A new species of *Microjassa* (Amphipoda) from Los Angeles Harbor. Bull. So. Calif. Acad. Sci., 53 (3): 127-130, pls. 35-36.
- 1954a. Marine Amphipoda of Oregon. Oregon State Monog., Zool., 8: 1-103, pls. 1-33.
- 1954b. Amphipoda of the family Ampeliscidae collected in the eastern Pacific Ocean by the Velero III and Velero IV. Allan Hancock Pacific Expeds., 18 (1): 1-137, pls. 1-38.
- 1955. Gammaridean Amphipoda (Crustacea) in the collections of Bishop Museum. Bishop Mus., Bull. 215: 1-46, text-figs. 1-20.
- 1958. Index to the families, genera, and species of the gammaridean Amphipoda. Allan Hancock Found., Occ. Pap. No. 19: 1-148.
- 1959. The number of species of gammaridean Amphipoda (Crustacea). Bull. So. Calif. Acad. Sci., 58 (1), p. 16.
- 1959a. Liljeborgiid amphipods of Southern California coastal bottoms, with a revision of the family. Pacific Naturalist, 1 (4): 12-28, text-figs. 1-12, 1 tab., 2 charts.
- 1959b. Generic partition in the amphipod family Cheluridae, marine wood borers. Ibid., 1 (3): 1-12, text-figs. 1-5.
- 1959c. Ecology of Amphipoda and Polychaeta of Newport Bay, California. Allan Hancock Found. Publ., Occ. Pap. No. 21: 1-106, pls. 1-14.
- 1960. Common pleustid amphipods of Southern California, with a projected revision of the family. Pacific Naturalist, 1 (17): 37-48, 6 text-figs.
- 1960a. The amphipod family Phoxocephalidae in the eastern Pacific Ocean, with analyses of other species and notes for a revision of the family. Allan Hancock Pacific Expeds., 18 (3): 169-375, pls. 1-75, 14 tabs., 1 chart.
- 1961. Gammaridean Amphipoda from depths of 400 to 60000 meters in scientific results of the Danish Deep-Sea Expedition round the world 1950-52. Galathea Report, vol. 5: 23-128, text-figs. 1-83, 8 tabs.
- 1962. Benthic marine Amphipoda of Southern California: families Amphiloichidae, Leucothidae, Stenothoidae, Argissidae, Hyalidae. Pacific Naturalist, 3 (3): 116-163, text-figs. 1-23, 1 table.

- 1962a. Benthic marine Amphipoda of Southern California: families Tironidae to Gammaridae. *Ibid.*, 3 (2): 73-115, text-figs. 1-23, 2 tables.
- 1962b. Benthic marine Amphipoda of Southern California: families Aoridae Photidae, Ischyroceridae, Corophiidae, Podoceridae. *Ibid.*, 3 (1): 3-72, text-figs. 1-32.
- BARNARD, K. H. 1916. Contributions to the crustacean fauna of South Africa. 5.—The Amphipoda. *Ann. So. African Mus.*, 15 (3): 105-302, pls. 26-28.
- 1925. Contributions to the crustacean fauna of South Africa—No. 8. Further additions to the list of Amphipoda. *Ann. Soc. African Mus.*, 20 (5): 319-380, pl. 34.
- 1930. Amphipoda. British Antarctic ("Terra Nova") Expedition, 1910, London. *Nat. Hist. Repts., Zool.*, 8: 307-454, 63 text-figs.
- 1932. Amphipoda. *Discovery Repts.*, 5: 1-326, pl. 1, 174 text-figs.
- 1935. Report on some Amphipoda, Isopoda, and Tanaidacea in the collections of the Indian Museum. *Rec. Indian Mus.*, 37: 279-319, 21 text-figs.
- 1937. Amphipoda. John Murray Exped. 1933-34, *Sci. Repts. Brit. Mus. (Nat. Hist.)*, 4 (6): 131-201, 21 figs.
- 1940. Contributions to the crustacean fauna of South Africa, XII. Further additions to the Tanaidacea, Isopoda, and Amphipoda, together with keys for the identification of the hitherto recorded marine and fresh-water species. *Ann. So. African Mus.*, 32 (5): 381-543, 35 text-figs.
- 1955. Additions to the fauna-list of South African Crustacea and Pycnogonida. *Ann. So. African Mus.*, 43 (1): 80-99, text-figs. 38-49 (Amphipoda).
- BULYCHEVA, A. I. 1952. Novye vidy bokoplavov (Amphipoda, Gammaridea) iz Japonskogo Morja. *Akad. Nauk SSSR, Trudy Zool. Inst.*, 12: 195-250, 39 figs.
- 1955. Novye vidy bokoplavov (Amphipoda, Gammaridea) iz Japonskogo Morja. II. *Ibid.*, 21: 193-207, 6 text-figs.
- CHEVREUX, E. 1900. Amphipodes provenant des campagnes de l'*Hirondelle* (1885-1888). *Res. Camp. Sci. Albert Ier, Monaco*, 16: I-IV, 1-195, pls. 1-18.
- 1901. Amphipodes. Mission Scientifique.....Alluaud aux Iles Séchelles. *Soc. Zool. France, Mém.*, 14: 388-438, 65 text-figs.
- 1907. Amphipodes recueillis dans.....l'Océanie par M. Seurat. *Ibid.*, 20 (4): 470-527, 35 text-figs.
- 1908. Sur trois nouveaux amphipodes méditerranéens appartenant au genre *Corophium* Latreille. *Bull. Soc. Zool. France*, 33: 69-76, 6 text-figs.
- 1911. Campagnes de la *Melita*. Les amphipodes d'Algérie et de la Tunisie. *Mém. Soc. Zool. France*, 23 (3): 145-275, pls. 6-20, 17 text-figs.
- CHEVREUX, E. and L. FAGE. 1925. Amphipodes. *Faune de France*, 9: 1-448, 438 text-figs.
- CHILTON, C. 1911. The Crustacea of the Kermadec Islands. *Trans. N. Z. Inst.*, 43: 544-573, 4 text-figs.
- 1920. Some New Zealand Amphipoda: No. 1. *Ibid.*, 52: 1-8, 5 text-figs.
- 1921. Report on the Amphipoda obtained by the F.I.S. "Endeavour" in Australian Seas. *Biol. Res. "Endeavour"*, 5 (2): 33-92, 16 text-figs.
- 1921a. Fauna of the Chika Lake. Amphipoda. *Mem. Indian Mus.*, 5: 519-558, 12 text-figs.
- 1925. Zoological results of a tour in the Far East. The Amphipoda of Tale Sap. *Mem. Asiat. Soc. Bengal*, 6: 531-539, 3 text-figs.
- CRAWFORD, G. I. 1937. A review of the amphipod genus *Corophium*, with notes on the British species. *Journ. Mar. Biol. Assoc. United Kingdom*, 21 (2): 589-630, 4 text-figs.
- DAHL, E. 1944. Amphipoda of the family Ampeliscaidae from Professor Sixten Bock's Expedition to Japan, 1914. *Arkiv för Zool.*, 36A (1): 1-18, 10 text-figs.
- 1959. Amphipoda from depths exceeding 6000 meters. *Galathea Rep.*, 1: 211-241, 18 figs.

- DELLA VALLE, A. 1893. Gammarini del Golfo di Napoli. Fauna und Flora des Golfes von Neapel, 20: xi plus 948 pp., 61 pls.
- DUNBAR, M. J. 1954. The amphipod Crustacea of Ungava Bay, Canadian Eastern Arctic. Journ. Fish. Res. Board Canada, 11 (6): 709-798, 42 text-figs., 1 table.
- ENEQUIST, P. 1949. Studies on the soft-bottom amphipods of the Skagerrak. Zool. Bidrag, fran Uppsala, 28: 297-492, 67 text-figs., 6 charts.
- GURJANOVA, E. 1951. Bokoplavi morei SSSR i sopredel'nyx vod (Amphipoda-Gammaridea). Opred. po Faune SSSR, Izd. Zool. Inst. Akad. Nauk, 41: 1-1031, 705 figs.
- 1962. Bokoplavi severnoj chasti Tikhogo Okeana. Opred. po Faune SSSR, Izd. Zool. Inst. Akad. Nauk, 74: 1-441, 143 text-figs.
- HOWES, N. H. 1939. Observations on the biology and post-embryonic development of *Idotea viridis* (Slabber) (Isopoda, Valvifera) from New England Creek, South-East Essex. Journ. Mar. Biol. Assoc. United Kingdom, 23: 279-310, 14 figs., 2 tables.
- HURLEY, D. E. 1954. Studies on the New Zealand amphipodan fauna. No. 9. The families Acanthonotozomatidae, Pardaliscidae and Liljeborgiidae. Trans. Roy. Soc. N.Z., 82 (3): 763-802, 14 text-figs.
- 1954a. Studies on the New Zealand amphipodan fauna. No. 7. The family Corophiidae, including a new species of *Paracorophium*. Ibid., 82 (2): 431-460, 7 text-figs.
- 1957. Studies on the New Zealand amphipodan fauna. No. 14. The genera Hyale and Allorchestes (family Talitridae). Ibid., 84 (4): 903-933, 9 text-figs.
- IRIE, H. 1956. Tube-building amphipods occurring at the "wakame" (a species of brown algae: *Undaria pinnatifida*) grounds of Simabara, Nagasaki Prefecture. Bull. Fac. Fish., Nagasaki Univ., 4: 1-6, 6 figs.
- 1959. Studies on pelagic amphipods in the adjacent seas of Japan. Ibid., 8: 20-42, 4 figs., 11 tables.
- IWASA, M. 1934. A new amphipod (*Parhyale kurilensis*, n. sp.) from Urup. Journ. Fac. Sci. Hokkaido Imp. Univ., ser. 6, Zool., 3 (1): 1-7, pls. 1-2.
- 1939. Japanese Talitridae. Ibid., ser. 6, Zool., 6 (4): 255-296, pls. 9-22, 27 text-figs.
- KITAMORI, R., K. NAGATA, and S. KOBAYASHI. 1959. The ecological study on "Moba" (Zone of *Zostera marina* L.). (II) Seasonal changes. Bull. Naikai Reg. Fish. Res. Lab., 12: 187-199, 13 figs., 2 tables. (In Japanese).
- NAGATA, K. 1959. Notes on five species of the amphipod genus *Ampelisca* from the stomach contents of the triglid fishes. Publ. Seto Mar. Biol. Lab., 7 (2): 263-278, 11 text-figs.
- 1960. Preliminary notes on benthic gammaridean Amphipoda from the *Zostera* region of Mihara Bay, Seto Inland Sea, Japan. Ibid., 8 (1): 163-182, pls. 13-17, 2 text-figs.
- 1961. A new atyloid amphipod from Japan. Annot. Zool. Japon., 34 (4): 216-218, 2 text-figs.
- 1961a. Two new amphipods of the genus *Eurystheus* from Japan. Publ. Seto Mar. Biol. Lab., 9 (1): 31-36, 2 text-figs.
- OLIVEIRA, L. P. H. 1953. Crustacea Amphipoda do Rio de Janeiro. Mem. Inst. Oswaldo Cruz, 51: 289-376, 27 text-figs.
- PIRLOT, J. M. 1936. Les amphipodes de l'expédition du Siboga. Siboga-Exped., Mon. 33e: 237-328, text-figs. 101-146.
- 1938. Les amphipodes de l'expédition du Siboga. Siboga-Exped., Mon. 33f: 329-359, text-figs. 147-161.
- REID, D. M. 1951. Report on the Amphipoda (Gammaridea and Caprellidea) of the coast of tropical West Africa. Atlantide Rpt., Copenhagen, 2: 189-291, 58 text-figs.
- RUFFO, S. 1950. Studi sui crostacei Anfipodi. XXII—Anfipodi der Venezuela raccolti dal G. Marcuzzi. Mem. Mus. Civ. Stor. Nat. Verona, 2 (2): 49-65, 5 text-figs.
- 1959. Contributions to the knowledge of the Red Sea. No. 13. Bull. Sea Fish. Res. Stat., Haifa, 20: 1-26, 6 text-figs.

- SARS, G. O. 1895. Amphipoda. An account of the Crustacea of Norway with short descriptions and figures of all the species, 1: i-viii, 1-711, 240 pls., 8 suppl. pls.
- SCELLENBERG, A. 1926. Die Gammariden der Deutschen Südpolar-Expedition 1901-03. Deutsche Südpolar-Exped., Berlin, 18: 235-414, 68 text-figs.
- 1926a. Die Gammariden der Deutschen Tiefsee-Expedition. Wiss. Erg. D. Tielsee-Exp. "Valdivia", 23 (5): 195-243, 28 text-figs., pl. 5.
- 1927. Amphipoda des nordischen Plankton. Nordisches Plankton, 20 (6): 589-722, 104 text-figs.
- 1928. Amphipoda in Cambridge Expedition to Suez Canal. Trans. Zool. Soc. London, 22: 633-692, 12 text-figs.
- 1931. Gammariden und Caprelliden des Magellangebietes, Sidgeorigens und der Westantarktis. Further Zool. Res. Swedish Antarctic Exped. 1901-03, 2 (6): 1-290, 1 pl., 136 text-figs.
- 1936. Zwei Neue Amphipoden des Stillen Ozeans und zwei Berichtigungen. Zool. Anz., 116: 153-156, 1 text-fig.
- 1937. Schlüssel und Diagnosen der dem Süsswasser-Gammarus nahestehenden Einheiten ausschliesslich der Arten des Baikalsees und Australiens. Ibid., 117: 267-280, 4 text-figs.
- 1938. Litorale Amphipoden des tropischen Pazifiks. Kgl. Svenskapskad. Handl., (3) 16 (6): 1-105, 48 text-figs.
- 1942. Krebstiere oder Crustacea IV: Flohkrebse oder Amphipoda. Die Tierwelt Deutschlands, Jena, 40: 1-252, 204 text-figs.
- SEXTON, E. W. & D. M. REID 1951. The life-history of the multiform species *Jassa falcata* (Montagu) (Crustacea Amphipoda) with a review of the bibliography of the species. Linn. Soc. London, Jour. (Zool.), 42 (283): 29-91, pls. 4-30.
- SHEARD, K. 1938. The amphipod genera *Euonyx*, *Syndexamine* and *Paradexamine*. Rec. So. Austr. Mus., 6 (2): 169-186, 9 text-figs.
- 1939. Studies in Australian Gammaridea. (I) The genus *Ceradocus*. Ibid., 6 (3): 275-295, 8 text-figs.
- SHIINO, S. 1948. Studies on marine crustaceans III. On a new boring amphipod, *Chelura brevicauda*, sp. n. Misc. Rept. Res. Inst. Nat. Resources, 12: 25-28, 3 text-figs. (In Japanese).
- SHOEMAKER, C. R. 1930. The Amphipoda of the Cheticamp Expedition of 1917. Cont. Canad. Biol. Fish., 5 (10): 221-359, 54 text-figs.
- 1933. Two new genera and six new species of Amphipoda from Tortugas. Carnegie Inst. Wash., Pap. Tortugas Lab., 28: 245-256, 8 text-figs.
- 1942. Amphipoda crustaceans collected on the Presidential cruise of 1938. Smithsonian. Misc. Coll., 101 (11): 1-52, 17 text-figs.
- 1945. The amphipod genus *Photis* on the east coast of North America. Charleston Mus. Leaf., 22: 1-17, 5 text-figs.
- 1947. Further notes on the amphipod genus *Corophium* from the east coast of America. Journ. Wash. Acad. Sci., 37 (2): 47-63, 12 text-figs.
- 1949. The amphipod genus *Corophium* on the west coast of America. Ibid., 39 (2): 66-82, 8 text-figs.
- 1955. Amphipoda collected at the Arctic Laboratory, Office of Naval Research, Point Barrow, Alaska, by G. E. MacGinitie. Smithsonian. Misc. Colls., 128 (1): 1-78, 20 text-figs.
- 1956. Observation on the amphipod genus *Parhyale*. Proc. U.S. Nat. Mus., 106 (3372): 345-358, 4 figs.
- STEBBING, T. R. R. 1888. Amphipoda. Rept. "Challenger" Exped., Zool., vol. 29.
- 1897. Amphipoda from Copenhagen Museum and other sources. Trans. Linn. Soc. London, (2), Zool., 7: 25-45, pls. 6-14.

- 1899. Amphipoda from Copenhagen Museum and other sources. Part 2. *Ibid.*, 7: 395-432, pls. 30-35.
- 1906. Amphipoda I. Gammaridea. *Das Tierreich*, 21: 1-806, 127 text-figs.
- 1910. Crustacea. Part 5. Amphipoda. *Sci. Res. Trawling Exped. H.M.C.S. "Thetis"*. *Mem. Austr. Mus.*, 4, vol. 2, (12): 565-658, pls. 47-60.
- STEBBING, T. R. R. & D. ROBERTSON 1891. On four new British Amphipoda. *Trans. Zool. Soc. London*, 13: 31-42, pls. 5-6.
- STEPHENSEN, K. 1915. Isopoda, Tanaidacea, Cumacea, Amphipoda (excl. Hyperidea). *In Report of the Danish Oceanographical Expedition, 1908-10, to the Mediterranean and adjacent seas. vol. 2, Biology, D. 1*: 1-53, 33 text-figs.
- 1923. Crustacea Malacostraca. V (Amphipoda I). *The Danish Ingolf-Expedition*, 3 (8): 1-100, 22 text-figs., charts.
- 1925. Crustacea Malacostraca. VI (Amphipoda II). *Ibid.*, 3 (9): 101-178, text-figs. 23-53, charts.
- 1932. Some new amphipods from Japan. *Annot. Zool. Japon.*, 13 (5): 487-501, 5 text-figs.
- 1933. *Ceinina japonica* (n. gen., n. sp.), a new aberrant species of the amphipodan family Talitridae from Japan. *Trans. Sapporo Nat. Hist. Soc.*, 13 (2): 63-68.
- 1935. Indo-Pacific terrestrial Talitridae. *Bernice P. Bishop Mus., Occas. Pap.*, 10 (23): 1-20, 2 tables.
- 1938. *Gradidierella japonica* n. sp., a new amphipod with stridulating (?) organs from brackish water in Japan. *Annot. Zool. Japon.*, 17 (2): 179-184, 2 text-figs.
- 1938a. The Amphipoda of N. Norway and Spitsbergen with adjacent waters. *Tromsø Mus. Skrifter*, 3 (2): 141-278, text-figs. 20-31.
- 1940. Marine Amphipoda. *The Zoology of Iceland*, 3 (26): 1-111, 13 text-figs., 2 tables.
- 1940a. The Amphipoda of N. Norway and Spitsbergen with adjacent waters. *Tromsø Mus. Skrifter*, 3 (3): 279-362, text-figs. 32-52.
- 1942. The Amphipoda of N. Norway and Spitsbergen with adjacent waters. *Ibid.*, 3 (4): 363-526, text-figs. 53-78.
- 1944. Some Japanese amphipods. *Vidensk. Medd. fra Dansk naturh. Foren.*, 108: 25-88, 33 text-figs.
- 1944a. Crustacea Malacostraca. VIII (Amphipoda IV). *The Danish Ingolf-Expedition*, 3 (13): 3-51, 38 text-figs.
- 1944b. *The Zoology of East Greenland. Amphipoda. Meddel. om Grönland.*, 121 (14): 1-165, 18 text-figs., 3 tables.
- TATTERSALL, W. M. 1922. Zoological results of a tour in the Far East. Part 8. Amphipoda with notes on an additional species of Isopoda. *Mem. Asiatic Soc. Bengal*, 6: 435-459, pls. 18-21.
- THOMSON, G. M. 1879. New Zealand Crustacea, with descriptions of new species. *Trans. N.Z. Inst.*, 11: 230-248, pl. 10.
- 1879a. Addition to the amphipodous Crustacea of New Zealand. *Ann. Mag. Nat. Hist.*, (5) 4 (23): 329-333, pl. 16.
- 1880. New species of Crustacea from New Zealand. *Ibid.*, (5) 6: 1-6, pl. 1.
- 1881. Recent additions to and notes on New Zealand Crustacea. *Trans. N.Z. Inst.*, 13: 204-221, pls. 7-8.
- UENO, M. 1933. Freshwater Crustacea of Iturup. *Annot. Zool. Japon.*, 14: 109-113, 1 fig.
- 1935. Crustacea collected in the lakes of Southern Sakhalin. *Ibid.*, 15 (1): 88-94, 4 figs.
- UTINOMI, H. 1947. Caprellidae of Japan and adjacent waters. *Seibutu, Suppl.*, 1: 68-82, 8 text-figs. (In Japanese).

- WALKER, A. O. 1898. Crustacea collected by W. A. Herdman, F.L.S., in Puget Sound, Pacific coast of North America, Sept. 1897. Proc. Trans. Liverpool Biol. Soc., 12: 268-287, pls. 15-16.
- 1904. Report on the Amphipoda collected by Professor Herdman, at Ceylon, in 1902. Ceylon Pearl Oyster Fisheries, Suppl. Rept., 17: 229-300, 8 pls.
- 1909. Amphipoda Gammaridea from the Indian Ocean, British East Africa, and the Red Sea. Trans. Linn. Soc. London, (2), Zool., 12: 323-344, pls. 42-43.
- WATKIN, E. E. 1939. The pelagic phase in the life history of the amphipod genus *Bathyporeia*. Journ. Mar. Biol. Assoc. United Kingdom, 23: 467-481, 2 text-figs., 4 tables.