TITLE:

SCREENING OF TOXIC CORALS AND ISOLATION OF A TOXIC POLYPEPTIDE FROM GONIOPORA SPP

AUTHOR(S):

Hashimoto, Yoshiro; Ashida, Katsuro

CITATION:

Hashimoto, Yoshiro ...[et al]. SCREENING OF TOXIC CORALS AND ISOLATION OF A TOXIC POLYPEPTIDE FROM GONIOPORA SPP. PUBLICATIONS OF THE SETO MARINE BIOLOGICAL LABORATORY 1973, 20: 703-711

ISSUE DATE:

1973-12-19

URL:

<http://hdl.handle.net/2433/175749>

RIGHT:

SCREENING OF TOXIC CORALS AND ISOLATION OF A TOXIC POLYPEPTIDE FROM *GONIOPORA* SPP.¹⁾

YOSHIRO HASHIMOTO and KATSURO ASHIDA

Laboratory of Marine Biochemistry, Faculty of Agriculture
The University of Tokyo, Tokyo, Japan

With 3 Text-figures

Introduction

Many species of coral feeders are included among ciguatoxic fishes (HALSTEAD, 1967) and there has been an assumption that toxins which have been observed in these coral feeders might originate from the toxic corals on which they feed (HALSTEAD, 1967; HASHIMOTO *et al.*, 1969). In order to search for a possible origin of the toxin found in coral feeders, we screened toxic corals by extracting the specimens with hot 70% ethanol and by injecting the resulting extract into mice. We found 20 toxic specimens belonging to 8 species among 104 specimens collected at Ishigaki Island. Among them, *Goniopora* spp. were found to contain a toxic substance that induced characteristic symptoms in mice: hypersensitivity, stiffness of the entire body and marked cyanosis. Since this appeared to be a new marine toxin, we attempted its isolation and characterization.

This paper deals with the screening for toxic corals and with the isolation of *Goniopora* toxin, together with its chemical and pharmacological properties.

Materials and Methods

A total of 104 specimens were collected for screening at Ishigaki Island in May 1968. The soft parts were shaved off in the field with a knife and transported frozen by air to our laboratory, where they were kept at -20°C until used. The specimens belonging to 29 genera were identified by Dr. K. YAMAZATO, University of the Ryukyus.

Specimens of the stony corals *Goniopora* spp. were mainly collected in tide pools of coral reefs in the Ryukyu Islands. The soft parts as mentioned above were retained, corresponding in wet weight to approximately 25% of the whole mass of coral and containing approximately 30% of the calcified tissues. In order to check seasonal, geographical and individual variations in toxicity, 60 specimens were collected during

1) Studies on Marine Toxins—XXXVI.

1969 to 1971 between April and October, at Okinawa Island, Amami Islands, and Shirahama. For isolation of the toxin, a quantity of soft parts, weighing about 10 kg, was collected in July 1971 at Amami-Oshima Island.

Screening Test for Toxic Corals: A 10 g portion of each minced specimen was extracted with hot 70% ethanol; fat-soluble and water-soluble fractions were obtained by the method that we have used for ciguatoxic fishes (HASHIMOTO *et al.*, 1969). The water-soluble fraction was dissolved in 10 ml of distilled water and the fat-soluble one in 2 ml of 0.8% Tween 60 solution. Two mice, weighing 20 ± 2 g, were injected i.p. with a 1-ml portion each. The mice were observed for 3 days after injection. When one of a pair of mice died, a third mouse was injected. A specimen was regarded as toxic, if two out of three mice were killed.

Assay Method for Goniopora Toxin: Frozen coral, usually weighing 10 g, was ground in a mortar with an equal volume of aqueous 0.9% NaCl, sonicated at 20 kc per sec for 20 min, and centrifuged at 10,000 rpm for 15 min. The supernatant was filtered and used as the original test solution for the toxicity test. The amount of toxin was determined by the dose-time-to-death curve, as described by MCFARREN (1966). A 1-ml portion was injected i.p. into mice weighing 20 ± 2 g. From the dilution necessary to kill 2 mice within a range of 30 to 60 min and from the mean death time at that dilution, the amount of toxin was determined from the dose-response curve shown in Fig. 1. In the Figure, the amount of toxin is expressed as mouse units (MU); one mouse unit is defined as the minimum quantity of toxin required to kill a 20 g mouse. Toxicity of the coral specimens was indicated by MU per ml of the original test solution and toxicity of preparations by the minimum lethal dose in mg per kg of mouse or MU per ml. This assay method was also used for monitoring the elution of the toxin in column chromatography.

Assay Method for Hemolytic Activity: Hemolytic activity was determined by using a 2% rabbit blood cell suspension, as done routinely in our laboratory (HASHIMOTO and OSHIMA, 1972) and was expressed by a saponin unit (SU).

Dose-Response Curve in Mice: The dose-time-to-death curve was determined after the method of KONOSU *et al.* (1968). Two different preparations were used: one was a crude saline extract with a lethality of 300 mg per kg mouse, and the other a purified preparation having a lethality of 0.5 mg per kg mouse.

Purification of Goniopora Toxin: After preliminary tests the following procedure was established. One kilogram of raw material was ground with an equal volume of aqueous 0.9% NaCl in a mortar, sonicated at 20 kc per sec for 20 min, and centrifuged at 10,000 rpm, after the pH of the solution was adjusted to 5.5 with 0.5 N HCl. The supernatant was filtered and diluted with twice its volume of distilled water prior to chromatography on a column (6 × 50 cm) of CM-cellulose (Brown). The CM-cellulose had previously been washed thoroughly with 0.5% NaCl solution. The column was washed with 0.5% NaCl solution and the toxin was then eluted from the column with 2.0% NaCl solution. Effluents were collected in 20 ml portions with a

fraction collector. The toxic fractions were combined, concentrated below 30°C, and dialyzed against 0.9% NaCl solution in a cellophane membrane. The retentate was filtered and passed through a column (6 × 50 cm) of Sephadex G-50 (Fine). The toxin was eluted from the Sephadex column with 0.9% NaCl solution, and the entire step was repeated. The toxic fractions thus obtained were again concentrated, desalted by dialysis against distilled water and placed on a column (2.5 × 95 cm) of Sephadex G-50 (Fine). The toxin was then eluted with distilled water. The elution pattern was monitored by mouse bioassay and by absorption at 280 nm.

Thin Layer Chromatography: Among many supporting media and solvent systems tested, only a combination of Avicel SF (FCM Co.) with 1-butanol-acetic acid-H₂O (2:1:2) was found satisfactory. Spots were visualized with ninhydrin and Dragendorff reagents. In addition, the band was scraped off from the plate, extracted with 0.9% NaCl solution, and tested for toxicity.

Ultracentrifugal Analysis: Sedimentation analysis was carried out at 20°C on a Hitachi Model UT-1A ultracentrifuge equipped with a synthetic boundary cell at 60,000 rpm. The most highly purified test solution was dialyzed against 0.2 M NaCl solution prior to ultracentrifugation. The same solution was subjected to the approach-to-equilibrium sedimentation method of ARCHIBALD (SCHANCHMAN, 1959) in a cell described by YPHANTIS (1960).

Results

Screening of Toxic Corals: The water-soluble fraction was found to be toxic in 20 out of 104 specimens, but none of the fat-soluble fractions were toxic. Toxic corals and observed symptoms in mice are listed in Table 1. Both *Goniopora* sp. (No. 1) and *Palythoa tuberculosa* (Nos. 47-49) killed mice in a short time and induced characteristic symptoms. The other toxic corals killed mice very slowly without inducing any remarkable symptoms, except loss of activity. Although *Millepora* sp. (No. 30) did not kill mice and was considered nontoxic, it evoked remarkable symptoms, such as loss of activity, dyspnea, paralysis of the entire body, jumping, and rapid recovery.

Response of Mice to Goniopora Toxin: Symptoms in mice consisted of hypersensitivity, paralysis of hind limbs, diarrhea, stiffness of the entire body, dyspnea with cyanosis, and death. Contraction of capillary vessels and stiffness of the entire body before death were especially noticeable. The relationship between dose and time-to-death is shown in Fig. 1. It will be noted that identical dose-response curves were obtained for two preparations differing in activity by a factor of 600. At the minimum lethal dose, mice died within one day.

Toxicity of Goniopora Specimens: Toxicity of 60 specimens ranged from 8 to 64 MU per ml of the original test solution. Seasonal, geographical and individual variations in toxicity were insignificant.

Purification of Goniopora Toxin: In a typical run, a saline extract with a lethality

Table 1. Toxic corals from Ishigaki Island.

Species	Sample No.	Average death time (min)	Symptoms in mice
<i>Goniopora</i> sp.	1	49	Diarrhea, paralysis of hind limbs, hypersensitivity, stiffness of the entire body, dyspnea with cyanosis and death
<i>Coeloseris mayeri</i> VAUGHAN	33	780	Loss of activity, and death
<i>Clavularia</i> sp.	46	960	"
	47	65	
	48-1	3	
	" 2	110	
	" 3	105	
	" 4	9	
	" 5	22	
	" 6	3	
<i>Palythoa tuberculosa</i>	49-1	238	Loss of activity, paralysis of hind limbs, dyspnea, jumping, and death
	" 2	20	
	" 3	41	
	" 4	122	
	" 5	75	
	" 6	5	
<i>Acropora</i> sp.	66	1800	Loss of activity, and death
<i>Cyphastrea</i> sp. cf. <i>C. chalcidicum</i> (FORSKÅL)	69	1800	"
Unidentified	85	1740	"
<i>Pavona (Polyastra)</i> <i>ebtusata</i> (QUELCH)	90	480	"

of 300 mg per kg mouse was obtained. The toxic fraction from a CM-cellulose column had an estimated lethality of 180 mg per kg; the estimate is based on the observed weight of a fraction residue and the calculated amount of sodium chloride in a residue. Recovery of toxicity was 70%. The most purified toxin that was obtained by Sephadex G-50 column chromatography had a lethality of 0.5 mg per kg. Recovery of activity for the entire purification sequence was 65%. As shown in Fig. 2, the elution curves from the Sephadex G-50 column, checked by optical density at 280 nm and lethality to mice, show only single and identical peaks. The hemolytic activity was 0.11 SU in the saline extract and 0.19 SU per mg of solid in the toxic fraction from a CM-cellulose column. The hemolytic factor was separated from the toxin that is lethal to mice by Sephadex G-50 column chromatography. This factor is now under investigation.

Fig. 1. Dose-response curve for *Goniopora* toxin.

●—● Crude toxin (300 mg per kg mouse); ○—○ Purified toxin (0.5 mg per kg mouse).

Some Chemical Properties of Goniopora Toxin: When the crude saline extract was adjusted to pH 2.0, 7.0, and 12.0 with 0.5 N HCl or 0.5 N NaOH and kept at 70°C for 10 min, loss of toxicity was 100% at pH 12.0, 58% at pH 7.0, and 62% at pH 2.0, respectively. At 4°C, toxicity remained unchanged at pH 7.0 and pH 2.0 for 7 days, but was lost completely at pH 12.0. The toxin was nondialyzable against distilled water or 0.9% NaCl solution through cellophane or collodion membranes (Göttingen). From a 0.9% NaCl solution it could not be extracted with 1-butanol, ethyl acetate, or diethyl ether. It was soluble in saline and acidic aqueous solutions, barely so in distilled water and insoluble in all common organic solvents. In aqueous solution, it was precipitated quantitatively by trichloroacetic acid, ammonium sulfate, and acetone.

In thin layer chromatography, the final preparation gave a single spot positive to ninhydrin and Dragendorff reagents at R_f 0.70. It was negative to aniline-diphenylamine-phosphoric acid reagent (BAILY and BOURNE, 1960; BUCHAN and SAVAGE, 1952). Toxicity to mice was detected only in the band between R_f 0.65 and 0.75. The final preparation exhibited an absorption maximum at around 280 nm in

Fig. 2. Sephadex G-50 (fine) column chromatography of *Goniopora* toxin.
 ○—○ Toxicity to mice; ●—● Absorption at 280 nm.

water and a single peak with $s_{20,w}=1.10S$ in ultracentrifugation (Fig. 3). It was positive to Folin-Lowry (LOWRY *et al.*, 1951) and biuret reagents. The molecular weight was calculated to be 9,400.

Fig. 3. Ultracentrifugation of *Goniopora* toxin. The most highly purified test solution was dialyzed against 0.2 M NaCl solution prior to ultracentrifugation and the retentate was centrifuged at 60,000 rpm at 20°C. The picture was taken 20 min after constant velocity was reached.

Discussion

In the course of screening of 104 specimens of corals, water-soluble fractions of 20 specimens belonging to 8 species were found to be toxic, while all fat-soluble fractions were nontoxic. Specimens of both *Goniopora* sp. and *Palythoa tuberculosa* were found to kill mice in a short time and with characteristic symptoms. As far as our examination shows, none of the known fish toxins have been detected in these coral samples. It should be emphasized that the screening was carried out by using a hot 70% ethanolic extract in order to detect the toxins that are implicated in ciguatera intoxication. None of the toxic proteins, which might be distributed widely in corals, were detected. *Goniopora* toxin tolerates heating in neutral solution to some extent, which may be the reason why it was detected in the present study. In succeeding toxicity tests of *Goniopora* specimens and in purification of the toxin, hot 70% ethanol was replaced by cold saline as the extractant.

The *Goniopora* specimen from Amami-Oshima Island was tentatively identified as *G. gracilis* (BASSETT-SMITH) by Dr. H. UTINOMI, Seto Marine Biological Laboratory, Kyoto University, and those from Shirahama as *Rhodaræa tenuidens* and *G. planulata* (EHRENBERG). Our specimens were thought to include a few other species, and are therefore described in this paper as *Goniopora* spp. Since every specimen tested showed similar toxicity, *Goniopora* toxin is considered to be a common constituent in this genus. It is most likely that the toxin is contained in the nematocysts, but anatomical distribution of the toxin should await further detailed examination.

SUGIYAMA (1955) assigned to *Goniopora* sp. a Japanese name, "Dokusango", which means a toxic coral, without giving any description of toxicity. KAWAGUTI (1940) referred to stinging by *Millepora*, *Goniopora* and *Astropora* in Palau, and HALSTEAD (1965) listed *Goniopora* sp. among the stinging corals. However, we could not find any specific data on the toxicity of *Goniopora* spp. The present study has shown that "Dokusango" is an apt designation of *Goniopora* spp., but we experienced no stinging sensation when we handled *Goniopora* spp. with our bare hands.

Judging from its chemical and pharmacological properties, *Goniopora* toxin seems to be different from other toxic proteins found in coelenterates. Further purification and characterization of the toxin will be reported in the near future.

Summary

Corals collected at Ishigaki Island were examined for toxicity to mice by the method used for ciguatoxic fishes. The water-soluble fractions were found to be toxic in 20 out of 104 specimens, while none of the fat-soluble fractions were toxic. Specimens of *Goniopora* spp. and *Palythoa tuberculosa* were found to be strongly toxic.

Goniopora spp. were found to contain a new toxic substance that induces characteristic symptoms in mice and possesses hemolytic activity in rabbit blood cells.

The toxin that is lethal to mice could not be dialyzed through a cellophane membrane. It was heat-labile, but stable in neutral and acidic aqueous solutions at 4°C. It was soluble in saline and acidic solution, barely soluble in distilled water and insoluble in all common organic solvents. It was obtained chromatographically and ultracentrifugically homogeneous by CM-cellulose and Sephadex G-50 column chromatography. The purified toxin was estimated by ultracentrifugation to have a molecular weight of 9,400 and may be a polypeptide with a toxicity of 0.5 mg per kg mouse. A principle that causes hemolysis in rabbit blood cells has not yet been characterized.

Acknowledgements

The authors are grateful to Drs. H. UTINOMI, Seto Marine Biological Laboratory, Kyoto University and K. YAMAZATO, University of the Ryukyus, for their identification of the samples. Thanks are also due to Dr. P. J. SCHEUER, University of Hawaii, for his kind editorial assistance with the manuscript.

REFERENCES

- BAILEY, R. W. and E. J. BOURNE, 1960. Colour reactions given by sugars and diphenylamine-aniline spray reagents on paper chromatograms. *J. Chromatog.*, 4, 206.
- BUCHAN, J. L. and R. I. SAVAGE, 1952. Paper chromatography of some starch conversion products. *Analyst*, 77, 401.
- HALSTEAD, B. W., 1965. *In: Poisonous and Venomous Marine Animals of the World*, Vol. I. Washington, D. C.: U. S. Government Printing Office.
- HALSTEAD, B. W., 1967. *In: Poisonous and Venomous Marine Animals of the World*, Vol. II. Washington, D. C.: U. S. Government Printing Office.
- HASHIMOTO, Y., N. FUSEYANI, and S. KIMURA, 1969. Aluterin: a toxin of filefish, *Alutera scripta*, probably originating from a Zoantharian, *Palythoa tuberculosa*. *Bull. Jap. Soc. Sci. Fish.*, 35, 1086.
- HASHIMOTO, Y. and Y. OSHIMA, 1972. Separation of grammistins A, B and C from a soapfish *Pogonoperca punctata*. *Toxicon*, 10, 279.
- HASHIMOTO, Y., T. YASUMOTO, H. KAMIYA, and T. YOSHIDA, 1969. Occurrence of ciguatoxin and ciguaterin in ciguatoxic fishes in the Ryukyu and Amami Island. *Bull. Jap. Soc. Sci. Fish.*, 35, 327.
- KAWAGUTI, S., 1940. Stinging reef corals; nine note on reef corals, Note 5, *Kagaku Nanyo*.
- KONOSU, S., A. INOUE, T. NOGUCHI, and Y. HASHIMOTO, 1968. Comparison of crab toxin with saxitoxin and tetrodotoxin. *Toxicon*, 6, 113.
- LOWRY, O. H., N. J. ROSEBROUGH, A. L. FARR, and R. J. RANDALL, 1951. Protein measurement with Folin-phenol reagent. *J. Biol. Chem.*, 193, 265.
- McFARREN, E. F., 1966. Differentiation of poisons of fish, shellfish and plankton. *In: Animal Toxins*, p. 85, (RUSSELL, F. E. and P. R. SAUNDERS, Eds.). Oxford: Pergamon Press.
- SCHACHMAN, H. K., 1959. Ultracentrifugation in Biochemistry, p. 181. New York: Academic Press Inc.
- SUGIYAMA, T., 1955. *In: Illustrated Encyclopedia of the Fauna of Japan* (UCHIDA, K., Ed.). The Hokuryukan Co. Ltd., Tokyo.
- YPHANTIS, D. A., 1960. Rapid determination of molecular weights of peptides and proteins. *Ann. N. Y. Acad. Sci.*, 88, 586.

DISCUSSION

RANDALL (only a comment): Your work showing the high toxicity of gonioporiid corals sheds some light on why this species of reef corals is rejected by *Acanthaster planci*.