

## CHAPTER TWO

# Molecular Biology and Epidemiology of Dianthoviruses

**Tetsuro Okuno<sup>\*,1</sup> and Chuji Hiruki<sup>†</sup>**

<sup>\*</sup>Laboratory of Plant Pathology, Graduate School of Agriculture, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan

<sup>†</sup>Department of Agricultural, Food, and Nutritional Science, University of Alberta, Edmonton, Alberta T6G 2P5, Canada

1 Corresponding author: e-mail: okuno@kais.kyoto-u.ac.jp

### Contents

1. Introduction
2. Dianthoviruses
  - 2.1 Carnation ringspot virus
  - 2.2 Red clover necrotic mosaic virus
  - 2.3 Sweet clover necrotic mosaic virus
  - 2.4 Furcraea necrotic streak virus
  - 2.5 Other viruses suspected to be dianthoviruses
3. Economic Importance of Diseases Caused by Dianthoviruses
4. Molecular Biology of Dianthoviruses
  - 4.1 Genome organization
  - 4.2 Cap-independent translation
  - 4.3 Translation via -1 programmed frameshifting
  - 4.4 *cis*-Preferential requirement of replication protein
  - 4.5 RNA elements required for genome replication
  - 4.6 Template recognition mechanisms by replicase proteins
  - 4.7 Composition of RNA replicase complexes
  - 4.8 Assembly of RNA replicase complexes
  - 4.9 Viral RNA replication factory
  - 4.10 Subgenomic RNA synthesis
  - 4.11 A viral non-coding RNA
  - 4.12 Cell-to-cell and systemic movement
  - 4.13 Virion
  - 4.14 Suppression of RNA silencing

5. Transmission Studies and Epidemiology
  6. Effects on Environment
    - 6.1 Release of dianthoviruses from intact roots, decaying plants materials, and pollen grains
    - 6.2 Sewage
  7. Concluding Remarks and Perspectives
- Acknowledgements
- References

Key words: Plant virus, Positive-strand RNA virus, Bipartite genome, Tombusviridae, Dianthovirus, RNA replication, Translation, Soil, Transmission, Epidemiology

Advances in Virus Research, Volume 87, 2013

## Abstract

The genus *Dianthovirus* is one of eight genera in the family *Tombusviridae*. All the genera have monopartite positive-stranded RNA genomes, except the dianthoviruses which have bipartite genomes. The dianthoviruses are distributed worldwide. Although they share common structural features with the other *Tombusviridae* viruses in their virions and the terminal structure of the genomic RNAs, the bipartite nature of the dianthovirus genome offers an ideal experimental system with which to study basic issues of virology. The two genomic RNAs seem to use distinct strategies to regulate their translation, transcription, genome replication, genome packaging, and cell-to-cell movement during infection. This review summarizes the current state of our knowledge of the dianthoviruses, with its main emphasis on the molecular biology of the virus, including the viral and host factors required for its infection of host plants. The epidemiology of the virus and the possible viral impacts on agriculture and the environment are also discussed.

## 1. INTRODUCTION

Since the dianthoviruses were established as a new plant virus group by the International Committee on Taxonomy of Viruses (ICTV) in 1981 (Matthews, 1982), much information has been collected not only on their basic characteristics but also on the effects of their infection of various plant species and the behavior of the viruses in different environments. While members of this group are small in number, they cause characteristic necrosis, both local and systemic, which is often associated with severe dwarfing, a frequent cause of serious reductions in crop yields. The number of *Dianthovirus* species may increase as the molecular genetic characterization of suspected viruses advances. The viral particles are stable and abundant in reproduction, and are therefore suitable for experimental studies in many aspects.

Our knowledge of the molecular mechanisms underlying the infection processes of the dianthoviruses, has increased over the last two decades, after their infectious cDNA clones became available (Ge *et al.*, 1993; Sit *et al.*, 2001; Xiong and Lommel, 1991) since the complete nucleotide sequences of their genomes were determined (Ge *et al.*, 1992, 1993; Kendall and Lommel, 1992; Lommel *et al.*, 1988; Ryabov *et al.*, 1994; Xiong and Lommel, 1989).

This review summarizes the current state of our knowledge of the dianthoviruses,

with its main emphasis on recent advances in our understanding of the molecular mechanisms underlying the genome strategies used for translation, replication, and cell-to-cell movement during infection. The epidemiology and possible impacts of the dianthoviruses on agriculture and the environment are also discussed.

## **2. DIANTHOVIRUSES**

The group consists of four viruses: the type member is *Carnation ringspot virus* (CRSV), and the other members are *Red clover necrotic mosaic virus* (RCNMV), *Sweet clover necrotic mosaic virus* (SCNMV) (Hiruki, 1987); *Furcraea necrotic streak virus* (FNSV) is a tentative member of the genus at present (Morales *et al.*, 1992).

### **2.1. CRSV**

Symptoms such as leaf mottling and ringspotting, plant stunting and flower distortion persist after the infection of *Dianthus caryophyllus* and *D. barbatus* by CRSV. CRSV probably occurs worldwide, wherever *Dianthus* species are grown. Virions are distributed in all parts of the infected plant, including in the cell cytoplasm and nuclei. Several species are susceptible to CRSV, showing systemic or local symptoms (Tremaine and Dodds, 1985). Its transmission by nematodes *Longidorus elongatus*, *L. macrosoma*, and *Xiphinema diversicaudatum* has been reported, (Fritsche *et al.*, 1979; Trudgill *et al.*, 1983), but is still contentious (Brunt *et al.*, 1996).

### **2.2. RCNMV**

Symptoms such as necrotic lesions accompanied with mosaic, and leaf distortion persist in *Trifolium pratense*, *T. repens*, *Medicago sativa*, and *Melilotus officinalis* infected with RCNMV. The virus was first reported in the former Czechoslovakia (Musil *et al.*, 1967, 1969a), but is known to occur in the UK, Australia, Canada, New Zealand, Poland, Sweden and the USA. The virus was reported to be experimentally transmissible by the chytrid *Olpidium radicale*, but is not transmitted by seed (Gerhardson and Insunza, 1979).

### **2.3. SCNMV**

The main symptoms of SCNMV are mosaic, ringspots, and systemic veinal necrosis in the leaves of *Melilotus officinalis* and *M. alba* (Hiruki *et al.*, 1984). The most severe symptoms include stunting and distortion. SCNMV is widely spread in sweet clover growing in regions of dark gray soil and black soil in Alberta, Canada (Hiruki, 1986b;

Hiruki *et al.*, 1986, 1987). A different strain of the virus occurs in *Medicago sativa* (Pappu *et al.*, 1988; Pappu and Hiruki, 1989). SCNMV is transmitted by contact and by drainage water, but not by *Olpidium brassicae* (Hiruki, 1986b, 1994). Sixteen of 25 species in six dicotyledonous families were infected by inoculation with sap (Hiruki, 1986a, 1987).

#### **2.4. FNSV**

The major symptoms of FNSV are plant stunting and chlorotic streaks, which later coalesce to produce necrosis, drastically reducing the length and quality of the fiber of *Furcraea* (fique) spp., the most important fiber crops in Colombia. The virus is transmitted by mechanical inoculation and by grafting, but not transmitted by *Planococcus citri*, *Saissetia coffeae*, *Xiphinema*, or *Trichodorus* spp. (Morales *et al.*, 1992). The virus is consistently detected in the roots of infected fique plants (Morales *et al.*, 1992).

#### **2.5. Other Suspected Dianthoviruses**

In 1996, Editors of Viruses of Plant Viruses included the following statement (Brunt *et al.*, 1996, p.1069) that was apparently collated by Dr. M. Hollings in 1980 under Taxonomy and Relationships of Red clover necrotic mosaic dianthovirus. Virus(es) with serologically unrelated virions: 60 strains of 45 isometric viruses. Some isolates induce different symptoms, for example with those from England and Scotland are different. *Tetragonia expansa* is infected systemically only by RCNMV strain TpM-34.

An RNA with a gene organization very similar to that of dianthovirus RNA1 was reported in grassy stunt-diseased rice plants, although no virions were isolated nor any biological characters investigated (Miranda *et al.*, 2001).

### **3. ECONOMIC IMPORTANCE OF DISEASES CAUSED BY DIANTHOVIRUSES**

The economic importance of the diseases caused by the dianthoviruses varies according to the kind of crop affected. CRSV, the type virus of the dianthovirus group, is fairly common, infecting *Dianthus* spp. worldwide, spreading across the Central Asian region and India, the Eurasian region, and the North American region. It has been found in Australia, but with no evidence of field spread (Brant *et al.*, 1996).

RCNMV has been reported to occur in the former Czechoslovakia (Musil *et al.*, 1967; Musil, 1969a, 1969b), Sweden (Gerhardson and Lindsten, 1973), and the UK

(Bowen and Plumb, 1979). It has been found in Australia, Canada, Poland and New Zealand but with no evidence of field spread (Brunt *et al.*, 1996).

In Alberta, Canada, the production of forage seeds, as well as hay, is very important in supporting the \$3 billion cattle industry in the province, which is a major source of income in agricultural industry. Sweet clover is also important to beekeeping as a source of nectar and pollen, and beekeeping is an important enterprise in Alberta's agriculture diversification strategy. It contributes almost \$15 million to the agricultural economy. SCNMV infection of sweet clover not only hampers plant growth, but also significantly diminishes the production of nectar and pollen.

The genus *Furcraea* (Agavaceae) contains several species found primarily in the American tropics. In Colombia, South America, four varieties of *Furcraea* spp., collectively known as 'fique', constitute the major fiber crop with over 40,000 tons of 'cabuya' fiber produced annually by small farmers (Perez, 1974). Among the various biotic constraints that affect commercial *Furcraea* production, necrotic streak caused by FNSV is the most difficult to control because no reliable control methods are available (Molales *et al.*, 1992).

#### **4. MOLECULAR BIOLOGY OF DIANTHOVIRUS**

Dianthovirus belongs to the family *Tombusviridae*. The unifying feature of this family is that each member has a highly conserved RNA-dependent RNA polymerase (RdRP) motif. The family *Tombusviridae* is classified into the supergroup that includes the families *Flaviviridae* and *Luteoviridae* and the phage lineage (Koonin, 1991; Koonin and Dolja, 1993). The molecular mechanisms underlying the infection processes of *Dianthovirus*, including its translation, RNA replication, packaging, cell-to-cell movement, and suppression of RNA silencing, have been investigated predominantly in RCNMV (Basnayake *et al.*, 2006; Giesman-Cookmeyer and Lommel, 1993; Okuno, 2012; Mine and Okuno, 2012; Takeda *et al.*, 2005). Recent studies have increased our understanding of how the dianthoviruses replicate in host cells during successful infection. They use several unique strategies to regulate the expression of viral genes, to form viral replication factories by recruiting host proteins on cellular membranes, and to counteract host defense mechanisms. Here, we comprehensively review the molecular biology of *Dianthovirus*.

##### **4.1. Genome organization**

The genomes of the dianthoviruses consist of two positive-sense single-stranded RNAs,

RNA1 and RNA2 (Gould *et al.*, 1981; Hiruki, 1987; Okuno *et al.*, 1983). This bipartite nature of the genome is unique among the viruses of the family *Tombusviridae*, the genomes of which are usually monopartite. RNA1 (3.9 kb) encodes the RNA replicase components, an auxiliary 27-kDa protein (p27), and an N-terminally overlapping 88-kDa protein (p88) (Xiong and Lommel, 1989; Xiong *et al.*, 1993b) that contains RdRP motif (Koonin and Dolja, 1993). RNA1 also encodes a coat protein (CP), which is translated from subgenomic RNA (sgRNA) (Sit *et al.*, 1998; Tatsuta *et al.*, 2005; Zavriev *et al.*, 1996). RNA1 can replicate in a single cell in the absence of RNA2 (Osman and Buck, 1987; Paje-Manalo and Lommel, 1989; Pappu and Hiruki, 1988). RNA2 encodes a 35-kDa movement protein (MP), which is essential for the cell-to-cell and systemic movement in plants (Osman and Buck, 1987; Xiong *et al.*, 1993a). In addition to the genomic RNAs, a small noncoding RNA (0.4 kb) that consists of nearly the entire region of the 3'-untranslated region (UTR) of RNA1, accumulates in dianthovirus-infected plants and protoplasts (Iwakawa *et al.*, 2008; H. Nagano, K. Omote and T. Okuno, unpublished data). The genomic organization of RCNMV, several RNA elements of which are described in this review, is shown in Fig. 1.

#### **4.2. Cap-independent translation**

The genomic RNAs of RCNMV lack both a 5' cap structure (Mizumoto *et al.*, 2003) and a 3' poly(A) tail (Lommel *et al.*, 1988; Mizumoto *et al.*, 2002, 2003; Xiong and Lommel, 1989). Mutagenesis studies using a reporter mRNA that included the firefly luciferase (Luc) gene revealed that RNA1 contains an RNA element essential for the cap-independent translation in the 3'-UTR (Mizumoto *et al.*, 2003). The element designated "3'TE-DR1" is predicted to have five stem-loop (SL) structures (Mizumoto *et al.*, 2003; Sarawaneeyaruk *et al.*, 2009), and this overall structure has been confirmed by structure probing (Wang *et al.*, 2010; Y. Tajima and T. Okuno, unpublished data). Of these SLs, the 5'-proximal SL and two 3'-proximal SLs are required for its efficient translation (Mizumoto *et al.*, 2003; Sarawaneeyaruk *et al.*, 2009).

Cap-independent translation enhancers like 3'TE-DR1 have been identified in the 3'-UTRs of many viral RNA genomes (Kneller *et al.*, 2006; Nicholson and White, 2011). These RNA elements replace the cap-structure and poly(A) tail in facilitating the translation of viral proteins, and are termed 3' cap-independent translation elements (3'CITEs). The 3' CITEs of plant RNA viruses are classified into several structural classes (Miller *et al.*, 2007; Nicholson and White, 2011). The 5'-proximal SL of 3'TE-DR1 that is conserved in dianthoviruses, including SCNMV and CRSV, is very similar to an essential SL structure in the 3' CITE of *Barley yellow dwarf virus* (BYDV)

(Guo *et al.*, 2000; Mizumoto *et al.*, 2003; Shen and Miller., 2004).




Figure 2.1. Schematic representation of the genome structure of RCNMV with the predicted RNA structures. TABS, trans-activator-binding site; 3'TE-DR1, cap-independent translational enhancer; TA, trans-activator; YRE, Y-shaped RNA element. \*A-rich sequence (ARS): the PABP-binding site. \*\*Loop sequences required for a long-distance RNA-RNA-interaction that facilitates -1PFS. Nucleotide numbers from the 5' end are shown below the lines.

3' CITEs interact with the eukaryotic initiation factors eIF4F and eIFiso4F (Gazo *et al.*, 2004; Nicholson *et al.*, 2010; Treder *et al.*, 2008; Wang *et al.*, 2009) or the 60S ribosomal subunits (Stupina *et al.*, 2008). The 3'-UTR of RCNMV RNA1 interacted with the eIF4F/iso4F components in an RNA-aptamer (Strepto-Tag) affinity assay in a cell-free extract of evacuated BY-2 protoplasts (BYL) (Iwakawa *et al.*, 2012). BYL (Komoda *et al.*, 2004) efficiently supports the 3'TE-DR1-mediated cap-independent translation and negative-strand RNA synthesis of RCNMV (Iwakawa *et al.*, 2007). Mutations in the 5'-proximal SL of 3'TE-DR1 abolish the association of these translation initiation factors with the 3'-UTR.

Poly(A)-binding protein (PABP) also coprecipitates with the Strepto-Tag-fused 3'-UTR (Iwakawa *et al.*, 2012). The binding site of PABP resides in an adenine-rich sequence (ARS) (3518-AAACAGUAAAUAUGCAAAAAA-3538) located 60 nucleotides upstream from 3'TE-DR1 in the 3'-UTR. Interestingly, mutations in ARS that compromise PABP-binding also compromise the binding of the eIF4F/iso4F components and abolish 3'TE-DR1-mediated cap-independent translation, whereas mutations in 3'TE-DR1 that abolish translation, have no effect on PABP-binding. Therefore, PABP-binding seems to be required for the recruitment of the eIF4F/iso4F components to 3'TE-DR1. The binding of both the eIF4F/iso4F components and PABP to the 3' RNA elements is required for the efficient recruitment of the 48S and 80S ribosome complexes to the viral RNA (Iwakawa *et al.*, 2012).



Some form of communication must occur between the 5'-UTR and the 3' CITE of the viral RNAs to deliver translation factors for recruiting ribosomes to the 5' translation initiation sites. Such communications could be mediated by long-distance RNA-RNA base pairing, as reported for BYDV (Guo *et al.*, 2001), *Black currant reversion nepovirus* (Karetnikov and Lehto, 2008), *Carnation Italian ringspot tomosvirus* (Nicolson and White, 2008), *Cucumber leaf spot aureusvirus* (Xu and White, 2009), *Saguaro cactus virus* (Chattopadhyay *et al.*, 2011), and *Tomato bushy stunt virus* (TBSV) (Fabian and White, 2004 and 2006). Alternatively and additionally, mechanisms other than RNA-RNA base-pairing could mediate the 5'-3' interaction of viral RNAs. This is the case for RCNMV RNA1.

The requirement for the RNA1 5'-UTR in 3'TE-DR1-mediated cap-independent translation differs greatly between plant species (Sarawaneeyaruk *et al.*, 2009). The deletion of any one of the four SLs predicted in the 5'-UTR of RNA1 inhibited the 3'TE-DR1-mediated cap-independent activity of reporter mRNAs in BY-2 tobacco protoplasts, whereas their deletion had no effect in cowpea protoplasts. The RNA1 5'-UTR contributes to RNA stability in BY-2 protoplasts. A compensatory mutagenesis analysis used to identify possible interactions between the 5'- and 3'-UTRs of RNA1 in BY-2 protoplasts suggested that no long-distance RNA-RNA interaction plays is essential for 3'-5' communication during 3'TE-DR1-mediated cap-independent translation (Sarawaneeyaruk *et al.*, 2009; S. Sarawaneeyaruk and T. Okuno, unpublished data). Similarly, little or no involvement of RNA-RNA interactions between the 5'- and 3' UTRs has been reported in the 3' CITE-mediated cap-independent translation of *Pea enation mosaic virus* RNA 2 (Wang *et al.*, 2009), satellite *Tobacco necrosis virus* (Gazo *et al.*, 2004) or *Turnip crinkle virus* (TCV) (Qu and Morris, 2000; Stupina *et al.*, 2008). The mechanisms for delivering the translation initiation factors and translation initiation ribosome complexes to the 5'-UTR of the viral RNAs in these viruses, including RCNMV, remain to be resolved.

CP is translated from CPsgRNA, which is transcribed from RNA1 and co-terminates with RNA1. This implies that CP is translated in a 3'TE-DR1-mediated cap-independent manner. Indeed, uncapped chimeric CPsgRNA, in which the CP open reading frame (ORF) is substituted with the Luc ORF, is translated as efficiently as RNA1 (Sarawaneeyaruk *et al.*, 2009).

In contrast to RNA1 and CPsgRNA, RNA2 lacks RNA elements, such as ARS and 3'TE-DR1, that function efficiently in a reporter mRNA. Reporter Luc mRNAs with the 5'- and 3'-UTRs of RNA2 (R2-UTR-Luc) or with the Luc ORF inserted between the 5'-UTR and the MP ORF (RNA2-Luc) in RNA2 are not translated efficiently in

protoplasts in the absence of the 5'-cap structure (Mizumoto *et al.*, 2006). The cap-independent translational activities of R2-UTR-Luc and RNA2-Luc are less than 1% of 3'TE-DR1-mediated cap-independent translation. Instead, RNA2-Luc, which can replicate in the presence of RCNMV replicase proteins, functions as an efficient mRNA when replicated (Mizumoto *et al.*, 2006). This suggests a strong link between the cap-independent translation of RNA2 and RNA replication, and a difference in the translational mechanisms of RNA1 and RNA2. Host factors associated with the viral RNA replication process might facilitate the translation of RNA2. Alternatively, RNA2 may have a silencer element(s) that suppresses the translation of RNA2. The low translational activity of RNA2 may be comprehensible, if MP is only required in the late stage of replication to move the virus to neighboring cells. The difference in the translation strategy of RNA1 and RNA2 may be important for the temporal, spatial, and quantitative regulation of RCNMV gene expression during infection.

#### **4.3. Translation via –1 programmed ribosome frameshifting**

–1PRF is one of the translation strategies used by many RNA viruses to regulate viral gene expression (Brierley, 1995). p88 is translated by –1PRF from RCNMV RNA1 (Xiong *et al.*, 1993b). The *cis*-acting RNA elements required for –1PRF have been mapped to a shifty heptanucleotide sequence GGAUUUU at the slippage site (Kim and Lommel, 1994) and a highly-structured bulged SL structure predicted just downstream from the slippage site in RNA1 (Kim and Lommel, 1998). In addition to these elements, a third *cis*-acting RNA element that facilitates –1PRF was identified using an *in vitro* translation/replication system (Tajima *et al.*, 2011). This third RNA element is a small stable SL structure predicted between ARS and 3'TE-DR1 in the 3'-UTR of RNA1. The loop sequence of this element can base-pair with the bulge of the SL adjacent to the shifty site. Such long-distance base-pairing is possible in all dianthoviruses. Disruption and restoration mutagenesis analyses have demonstrated the importance of long-distance base-pairing for efficient –1PRF in RCNMV RNA1 (Tajima *et al.*, 2011). A similar requirement for long-distant RNA–RNA communication has been reported for the –1PRF of BYDV (Barry and Miller, 2002).

#### **4.4. *cis*-Preferential requirement of replicase protein**

An RNA1 mutant expressing p88 could be replicated efficiently in protoplasts when p27 was supplied in *trans*, but an RNA1 mutant expressing p27 alone could not be replicated when p88 was supplied in *trans* (Okamoto *et al.*, 2008). Thus, only RCNMV RNA1, from which p88 is translated, is an effective template, on which viral RNA

replicase can initiate RNA synthesis in the presence of p27. The *cis*-Preferential function of viral-encoded replication proteins or the coupling of translation and replication has been reported for several viruses, including *Alfalfa mosaic virus* (AMV) (Neeleman and Bol, 1999; van Rossum *et al.*, 1996), *Brome mosaic virus* (BMV) (Yi and Kao, 2008), *Clover yellow mosaic virus* (CYMV) (White *et al.*, 1992), Coronavirus (Chang *et al.*, 1994), *Cowpea mosaic virus* (Van Bokhoven *et al.*, 1993), Poliovirus (Hagino-Yamagishi and Nomoto, 1989; Johnson and Sarnow, 1991; Novak and Kirkegaard, 1994), *Turnip crinkle virus* (TCV) (White *et al.*, 1995), *Tobacco etch virus* (Mahajan *et al.*, 1996; Schaad *et al.*, 1996), *Tobacco mosaic virus* (TMV) (Lewandowski and Dawson, 2000), TBSV (Oster *et al.*, 1998), *Turnip yellow mosaic virus* (TYMV) (Weiland and Dreher, 1993), and *Rubella virus* (Liang and Gillam, 2001).

#### **4.5. RNA elements required for genome replication**

The *cis*-acting RNA elements required for viral RNA replication have been identified for many viruses with mutagenesis studies, when the RNA can be replicated by a viral replicase supplied in *trans*. However, it is difficult to apply this method to viral RNAs that encode replication proteins that are required in *cis* for RNA replication. This is the case for RCNMV RNA1.

The RNA elements required for the negative-strand synthesis of RNA1 have been determined in its 3'-UTR using capped viral RNA transcripts in BYL and BY-2 protoplasts (Iwakawa *et al.*, 2007). The use of capped viral RNA transcripts in BYL allowed the effects of introduced mutations on cap-independent translation and the negative-strand RNA synthesis of RNA1 to be distinguished. Two SL structures at the 3' end of RNA1 and the intervening sequence between the two SLs are essential for negative-strand synthesis (Iwakawa *et al.*, 2007) (see Fig. 1). These RNA elements are conserved between RNA1 and RNA2 and in SCNMV and CRSV. Interestingly, the core RNA element of 3'TE-DR1, which is essential for cap-independent translation of RNA1, is not essential for negative-strand RNA synthesis (Iwakawa *et al.*, 2007). Thus, the RNA elements in the 3' UTR of RCNMV RNA1 that are required for negative-strand RNA synthesis are separated from those required for cap-independent translation.

RCNMV-Australian (Aus) and RCNMV-Canadian (Can) strains have different temperature sensitivities during infection (Mizumoto *et al.*, 2002). RCNMV-Can does not replicate in protoplasts at temperatures higher than 22°C, indicating that replication processes are involved in the temperature-sensitive phenotype. Mutagenesis studies have shown that the temperature sensitivity of RCNMV is attributable to at least the 3'

terminal SL of RNA1, because a single-nucleotide substitution from U to C, which changes a U–G wobble pair to a stable C–G pair in the stem of the 3' terminal SL in RNA1-Can allows the RNA to replicate and to support RNA2 accumulation at nonpermissive temperatures (Mizumoto *et al.*, 2002). The lack of temperature sensitivity in RNA synthesis activity in the crude membrane-bound cellular fraction prepared from RCNMV-Can-infected plants or protoplasts suggests that the promoter elements are distorted in the initiation of minus-strand synthesis at the nonpermissive temperatures (K. Shimada and T. Okuno, unpublished results).

In contrast to RNA1, RNA2 is replicated efficiently by the RCNMV replicase proteins supplied in *trans*. The core promoter of negative-strand RNA synthesis is located in the 3' proximal region of RNA2, which is homologous to that of RNA1, except for the presence of 66 extra nucleotides upstream from the 3' terminal SL, which are not essential for replication (Iwakawa *et al.*, 2007; Turner and Buck, 1999). Three discontinuous nucleotides in the loop of the 3' terminal SL are thought to be involved in the interaction with the RCNMV RNA replicase (Weng and Xiong, 2009).

In addition to the core promoter, RNA2 has other unique replication elements. These elements include a *trans*-activator (TA) in the MP ORF (Tatsuta *et al.*, 2005) and a Y-shaped RNA element (YRE) in the 5' proximal region of the 3'-UTR (An *et al.*, 2010; Iwakawa *et al.*, 2011) (Fig. 1). Interestingly, TA is the RNA element that interacts with RNA1 and enhances the RNA-mediated transcription of CPsgRNA (Sit *et al.*, 1998) (see section J). Both TA and YRE play crucial roles in the negative-strand RNA synthesis of RNA2 (An *et al.*, 2010).

YRE consists of two small SLs with a short intervening region between them on the basal stem structure (An *et al.*, 2010). This structure is conserved among the dianthoviruses. The entire Y-shaped structure (84 nucleotides), including the two small SLs, is important for the negative-strand RNA synthesis of RNA2 (An *et al.*, 2010). YRE is the only RNA element of RCNMV that interacted with p27 supplied in *trans*, when assessed with the RNA-aptamer (Strepto Tag) affinity and immunoprecipitation assay in BYL (Iwakawa *et al.*, 2011). YRE also interacts with the 480-kDa replicase complex via p27 (Iwakawa *et al.*, 2011). The 480-kDa replicase complex contains p88 and host proteins, and is thought to be a key player in RCNMV RNA replication (Mine *et al.*, 2010b) (see section G and H). The interaction between YRE and p27 is required for the recruitment of RNA2 to the endoplasmic reticulum (ER) membrane (Hyodo *et al.*, 2011), which is the site of RCNMV RNA replication (Turner *et al.*, 2004; Hyodo *et al.*, 2013).

#### **4.6. Template recognition mechanisms of replicase proteins**

RNA1 lacks RNA elements that interact with p27 and p88 supplied in *trans*, as determined in the Strepto Tag affinity and immunoprecipitation assay in BYL (Iwakawa *et al.*, 2011). Interestingly, however, a protein-mediated coimmunoprecipitation analysis showed that both p27 and p88 can interact with their translating template RNA1, which are associated with ribosomes. The interaction between p27 or p88 and its translation templates is abolished or compromised, respectively, by puromycin treatment, which induces the dissociation of polyribosomes from mRNA (Blobel and Sabatini, 1971; Lehninger *et al.*, 1993). These results imply that p27 binds to the template RNA1 via a polyribosome-dependent and puromycin-sensitive mechanism, whereas p88 mainly binds to the template RNA in a translation-coupled, polyribosome-independent, and puromycin-tolerant manner. The interaction mechanism unique to p88 is maintained after the dissociation of the polyribosomes. p88 seems to bind specifically to the 3'-UTR of RNA1 in this translation-coupled mechanism, because SR1f, a degradation product of the 3'-UTR of RNA1 (see section K), is coimmunoprecipitated with p88 but not with p27. The translation-coupled binding of p88 may partly explain the strong *cis*-preferential requirement for p88 in the replication of RNA1 (Okamoto *et al.*, 2008).

#### **4.7. Composition of RNA replicase complexes**

A template-bound solubilized RNA polymerase has been isolated from RCNMV-infected *Nicotiana clevelandii* plants (Bates *et al.*, 1995). The polymerase becomes template-dependent after removing endogenous RNA templates with micrococcal nuclease. The RdRP contains p27 and p88 with several unknown host proteins, and produces double-stranded RNA in a template-specific manner (Bates *et al.*, 1995).

The *Agrobacterium*-mediated expression of RCNMV replicase proteins and RNAs in *N. benthamiana* leaves allowed large detergent-solubilized membrane-associated protein complexes to be isolated, with an apparent molecular mass of 480-kDa on blue-native polyacrylamide gel electrophoresis (BN-PAGE) (Mine *et al.*, 2010b). The 480-kDa integral membrane complex contains both p27 and p88, and is associated with possible host proteins. In sucrose gradient sedimentation, the 480-kDa complex cofractionates with both endogenous template-bound and exogenous template-dependent RdRP activities. The 480-kDa complex corresponds predominantly to the exogenous template-dependent RdRP activity, and specifically recognizes the 3' core promoter sequences of the RCNMV genomic RNAs to produce viral RNA fragments. In contrast, the endogenous template-bound RdRP produces genome-sized

RNAs without the addition of RNA templates (Mine *et al.*, 2010b).

Analysis of the affinity-purified solubilized membrane-bound RdRP complexes using two-dimensional BN/SDS-PAGE and mass spectrometry showed that the RdRP complexes contain many host proteins, including Hsp70, Hsp90, glyceraldehyde 3-phosphate dehydrogenase, ADP-ribosylation factor 1 (Arf1), histone deacetylase 1, ubiquitin, and several ribosomal proteins, in addition to viral replicase proteins (Hyodo *et al.*, 2013; Mine *et al.*, 2010b).

In addition to the 480-kDa complex, a 380-kDa complex is formed in BYL and RCNMV-infected plant tissues expressing p27 alone or both p27 and p88 (Mine *et al.*, 2010a and b). The 380-kDa complex could be a p27 oligomer with some host proteins.

#### **4.8. Assembly of the replicase complex**

The assembly of the viral replicase complexes of eukaryotic positive-strand RNA viruses is a regulated process: multiple viral and host proteins and template RNAs must be assembled on intracellular membranes and organized into quaternary complexes capable of synthesizing viral RNAs (Mine and Okuno, 2012; Nagy and Pogany, 2012).

p27 interacts with both p27 and p88 through direct protein-protein contacts. The C-terminal half of p27 is responsible for these interactions, whereas the nonoverlapping region unique to p88 is responsible for the p27-p88 interaction (Mine *et al.*, 2010a). Both the p27-p27 and p27-p88 interactions are required for the formation of the 480-kDa complex *in vitro* and *in planta* (Mine *et al.*, 2010a). A mutant p27 incapable of interacting with p27, but capable of interacting with p88, fails to form the 480-kDa complex. Another mutant p27 capable of interacting with p27, but incapable of interacting with p88 formed the 380 kDa complex but failed to form the 480-kDa complex. Thus, p27 oligomerization is a critical step in the formation of the 480-kDa complex. It appears that the formation of the 480-kDa complex is directed by the p27-oligomer (the 380-kDa complex), which interacts with the p88 protein(s).

The formation of the 480-kDa replicase complex is enhanced by the presence of the RCNMV genomic RNAs (Mine *et al.*, 2010b). The RNA binding domain of p27 was identified using an RNA-aptamer (Strepto-Tag)-fused YRE that binds p27 in an affinity and immunoprecipitation assay in BYL (Hyodo *et al.*, 2011). Deletion and alanine-scanning mutation analyses indicated that the main functional domains required for RNA binding differ from those required for protein binding, although they partially overlap.

There is a robust correlation between the RNA-binding activity of p27 and its RNA-recruiting activity to the ER membrane (Hyodo *et al.*, 2011). Interestingly, several

p27 mutants that retain the ability to bind to RNA2, to recruit RNA2 to the membrane, and to interact with both p27 and p88, fail to form the 480-kDa complex or to support RNA replication (Hyodo *et al.*, 2011). These p27 mutants might lack the ability to interact, directly or indirectly, with the host proteins that are required for the proper assembly of the 480-kDa replicase complex.

The roles in viral RNA replication of Hsp70 and Hsp90 detected in the affinity-purified solubilized membrane-bound RdRP complexes of RCNMV (Mine *et al.*, 2010a) were investigated (Mine *et al.*, 2012). A bimolecular fluorescence complementation (BiFC) assay using confocal microscopy showed that p27, but not p88, interacts directly with both Hsp70 and Hsp90 within the p27-induced large aggregated structures on the ER in the perinuclear region. Gene silencing and the pharmacological inhibition of Hsp70 and Hsp90 compromised RCNMV RNA replication in plant cells. The inhibition of p27–Hsp70 interaction by 2-phenylethanesulfonamide (a specific inhibitor of Hsp70) inhibited the formation of the 480-kDa complex, but instead induced the formation of nonfunctional large complexes (~1024 kDa) in BYL. Hsp70 appears to control the proper assembly of the viral replicase complexes by preventing the aggregation of p27. Alternatively, the large complexes of p27 could be intermediates in the assembly of the RCNMV replicase complex and Hsp70 might assist the assembly of these complexes into functional replicase complexes. In contrast, the inhibition of p27–Hsp90 interaction by geldanamycin (a specific inhibitor of Hsp90) inhibited the formation of the 480-kDa complex without inducing large complexes, and rendered p27 incapable of binding to a specific viral RNA element (YRE), which is a critical step in the assembly of the 480-kDa replicase complex. These findings suggest that Hsp70 and Hsp90 play essential roles in the assembly of the 480-kDa replicase complex mainly by regulating protein–protein interactions and protein–RNA interactions, respectively. Thus, Hsp70 and Hsp90 regulate different steps in the assembly of the RCNMV replicase complex.

#### **4.9. Viral RNA replication factory**

The replication of eukaryotic positive-strand RNA viruses occurs on membranes of selected subcellular organelles, such as the ER, chloroplasts, mitochondria, peroxisomes, or tonoplasts in infected cells (den Boon *et al.*, 2010; Mine and Okuno, 2012; Nagy and Pogany, 2012; Salonen *et al.*, 2005). Viral proteins play essential roles in targeting the viral replication complexes to these membranes and induce morphological changes in these membranes (Belov and van Kuppeveld, 2012; den Boon and Ahlquist, 2010; Miller and Krijnse-Locker, 2008).

Confocal microscopy using green fluorescent protein (GFP)-fused p27 and p88 showed that both proteins colocalize to the cortical and cytoplasmic ER, and that p27 causes the restructuring and proliferation of the ER membrane (Kusumanegara *et al.*, 2012; Mine *et al.*, 2012; Turner *et al.*, 2004). The domains and critical amino acids in p27 required for its association with and targeting of ER membranes were determined using a C-terminally GFP-fused p27 (p27-GFP), which supported viral RNA replication in the presence of p88. Membrane-flotation assays combined with microscopic observation revealed that the membrane association of p27 is mediated by a stretch of 20 amino acids located in its N-terminal region (amino acids 31–50) (Kusumanegara *et al.*, 2012). These 20 amino acids are predicted to form an amphipathic  $\alpha$ -helix, with one side having a cluster of hydrophilic, polar residues and the other side hydrophobic, nonpolar residues, and were sufficient to target the nonviral GFP to ER membranes. Mutations that impede the membrane association of p27 compromise the formation of the RCNMV RNA replication complexes and negative-strand RNA synthesis (Kusumanegara *et al.*, 2012).

The role in viral RNA replication of the Arf1 identified in the affinity-purified RCNMV RdRP fraction was investigated (Hyodo *et al.*, 2013). Arf1 is a ubiquitous, highly conserved, small GTPase that is implicated in the formation of the coat protein complex I (COPI) vesicles on the Golgi membranes and in the membrane transport from the Golgi to the ER (D'Souza-Schorey and Chavrier, 2006). GTP-bound Arf1 facilitates the formation of several types of vesicle coat complexes and activates lipid-modifying enzymes on the Golgi membranes (Donaldson and Jackson, 2011; Memon, 2004).

The pharmacological inhibition of the nucleotide-exchange activity of Arf1 using the inhibitor brefeldin A (BFA) and the expression of dominant-negative Arf1 mutants compromised RCNMV RNA replication in tobacco BY-2 protoplasts. *In vitro* pull-down and BiFC analyses showed that p27 interacts with Arf1 within the virus-induced large punctate structures of the ER membrane. Inhibition of Arf1 activity by BFA disrupted p27-mediated ER remodeling and the assembly of the 480-kDa viral replication complex. Thus, p27 interacts with Arf1, and recruits and relocalizes this protein to the aggregate structures on the ER membranes, where they colocalize (Hyodo *et al.*, 2013).

BFA treatment or the expression of dominant-negative mutants of Arf1 in plant cells not only inhibited the COPI pathway but also compromised COPII vesicle trafficking (Stefano *et al.*, 2006). Interestingly, the expression of a dominant-negative mutant of Sar1, a key regulator of the biogenesis of COPII vesicles at ER exit sites, also compromises RCNMV RNA replication (Hyodo *et al.*, 2013). These results suggest that



the replication of RCNMV depends on the host membrane trafficking machinery or that RCNMV rewires the cellular trafficking pathways to build a viral replication factory. A model for the formation of the viral replication factory in RCNMV-infected cells is presented (Fig. 2).

#### **4.10. Subgenomic RNA**

RCNMV uses a unique strategy to produce sgRNA. In addition to the core promoter that is predicted to exist within a stable SL (Zavriev *et al.*, 1996), the transcription of CPsgRNA from RNA1 requires RNA2 *in trans*, and is regulated by intermolecular interactions via base pairing between the eight-nucleotide sequence in the loop of TA in RNA2 and the complementary eight-nucleotide sequence in the TA binding site (TABS) immediately upstream from the CPsgRNA transcription start site in RNA1 (Sit *et al.*, 1998) (refer to Fig. 1). Biophysical analyses, using short synthetic oligonucleotides that mimic the TA of RNA2 and the TABS of RNA1, suggested the formation of a weak but stable bimolecular complex between these two mimics (Guenther *et al.*, 2004).

Several lines of evidence have suggested that CPsgRNA is generated by a premature termination mechanism (Guenther *et al.*, 2004; Sit *et al.*, 1998; Tatsuta *et al.*, 2005). The premature termination model is supported by the accumulation of negative-strand CPsgRNA in protoplasts (Iwakawa *et al.*, 2008), because the production of negative-sense sgRNA templates is a fundamental step in premature termination (Eckerle *et al.*, 2003; Lin and White, 2004; White, 2002). CPsgRNA is also unlikely to be an efficient RNA replicon, because its 3'-UTR is identical to that of RNA1 whose replication depends predominantly on replication proteins translated from its own RNA *in cis* (Iwakawa *et al.*, 2011; Okamoto *et al.*, 2008).

#### **4.11. A viral noncoding RNA**

A small noncoding RNA (0.4 kb), designated “SR1f”, which consists of nearly the entire 3'-UTR of RNA1, accumulates in RCNMV-infected plants and protoplasts, and is packaged into virions (Iwakawa *et al.*, 2008). SR1f is neither a subgenomic RNA nor a defective RNA replicon, but a stable degradation product generated by *cis*-RNA element-mediated protection against 5'→3' decay. A 58 nucleotide sequence in the 5' proximal region of the 3'-UTR of RNA1 is necessary and sufficient to protect against 5'→3' decay (Iwakawa *et al.*, 2008). SR1f efficiently suppresses both cap-independent and cap-dependent translation, both *in vitro* and *in vivo*. SR1f *trans*-inhibits the negative-strand RNA synthesis of RCNMV genomic RNAs *via* the repression of replicase protein production, but not *via* competition with the replicase proteins *in vitro*




Figure 2.2 Schematic representation of the formation of the viral replication factory in RCNMV-infected cell. The abundant auxiliary replicase protein p27 interacts with p27 itself, p88, and viral RNAs, and also interacts with host proteins, such as Hsp70, Hsp90, and Arf1. These interactions are essential for the recruitment of viral genomic RNAs, viral proteins and host factors to the ER membranes and for the reorganization of the membranes to form the viral replication factory.

(Iwakawa *et al.*, 2008). SR1f seems to play important roles in virus infection and survival, although its precise roles in RCNMV infection in nature remain to be resolved.

The stable decay products of viral noncoding RNAs generated by a mechanism similar to that for SR1f have been reported in flaviviruses (Moon *et al.*, 2012; Pijlman *et al.*, 2008) and *Beet necrotic yellow vein virus* (Peltier *et al.*, 2012). The *Flavivirus* noncoding RNA play important roles in viral pathogenicity by affecting host mRNA stability (Moon *et al.*, 2012).

#### **4.12. Cell-to-cell and systemic movement**

A 35-kDa MP encoded by RNA2 is essential for the cell-to-cell and systemic movement of RCNMV (Osman and Buck, 1987; Paje-Manalo and Lommel, 1989; Xiong *et al.*, 1993a). RCNMV MP has the ability to bind single-stranded nucleic acids cooperatively (Giesman-Cookmeyer and Lommel, 1993; Osman *et al.*, 1992, 1993), localizes in the cell wall (Osman and Buck, 1991; Tremblay *et al.*, 2005), and increases the size exclusion limit (SEL) of the plasmodesmata (PD) (Fujiwara *et al.*, 1993). RCNMV MP fused with GFP (RCNMV MP:GFP) can form a homopolymer and is targeted to the cell wall, and this targeting is required for viral cell-to-cell movement (Tremblay *et al.*, 2005).

Further analysis of the subcellular localization of RCNMV MP:GFP expressed from a recombinant virus in *N. benthamiana* epidermal cells and protoplasts showed that MP:GFP first appeared in the cell wall and was subsequently observed as punctate spots in the cortical ER (Kaido *et al.*, 2009). The ER-localization of RCNMV MP:GFP was associated with the replication of RNA1, but not with that of RNA2 or the viral replicase component proteins (p27 and p88) *per se* (Kaido *et al.*, 2009). This characteristic of MP:GFP is dependent on the 70 amino acids in the C-terminal region of MP and is required for the efficient cell-to-cell movement of the recombinant virus (Kaido *et al.*, 2011). Interestingly, the deletion of the C-terminal 70 amino acids of MP had no deleterious effects on its localization to the cell wall, or its ability to increase the PD SEL, to bind to single-stranded RNA or to interact with MP *in vivo* (Kaido *et al.*, 2011). These findings show that the recruitment of MP to the viral replication complexes is required for viral cell-to-cell movement. The mechanism that targets RCNMV MP:GFP to the ER, dependent on the replication of RNA1, which does not encode MP, may reflect a strategy to retain MP close to RNA1, thereby achieving its intracellular movement, followed by the intercellular movement of the viral genome.

Transgenic *N. benthamiana* expressing RCNMV MP supported both the cell-to-cell and systemic movement of movement-defective TMV and CMV (Giesman-cookmeyer

at al., 1995; Rao *et al.*, 1998). A chimeric TMV with the MP gene replaced with the RCNMV MP gene systemically infected *N. benthamiana* (Giesman-cookmeyer at al., 1995). Therefore, the movement proteins of tobamoviruses and dianthoviruses are functionally homologous.

In contrast, transgenic *N. benthamiana* expressing RCNMV MP supported the cell-to-cell movement but not the systemic movement, of movement-defective *Cowpea chlorotic mottle bromovirus* (Rao *et al.*, 1998). A chimeric *Barley stripe mosaic virus*, in which the triple gene block was replaced with the RCNMV MP gene, accumulated in the inoculated leaves of *N. benthamiana*, but not in the upper uninoculated leaves (Solovyev *et al.*, 1997). Immunocytochemical studies by Wang *et al.* (1998) showed that point mutations in RCNMV MP that do not affect the cell-to-cell movement of the virus prevent systemic viral movement, presumably by inhibiting RCNMV loading into the companion cell–sieve element complex. These findings suggest that the roles of RCNMV MP in the two processes are genetically distinct.

RCNMV CP is not essential for the cell-to-cell movement of the virus but virion formation is important for its systemic movement in *N. benthamiana* (Vaewhongs and Lommel, 1995; Xiong *et al.*, 1993a). The N-terminal lysine-rich motif of CP is involved in the systemic viral movement, virion accumulation, and symptomatology in *N. benthamiana* (Park *et al.*, 2012). A sobemovirus CP gene complemented the long-distance movement of a CP-null RCNMV (Callaway *et al.*, 2004).

#### **4.13. Virions**

The virions of *Dianthovirus* contain 180 copies of the 37-kDa CP subunit, forming an icosahedral particle with T = 3 symmetry and a diameter of about 36 nm (Sherman *et al.*, 2006; Hiruki, 1987). Analysis of the RCNMV virions with heat treatment and UV cross-linking suggested that RCNMV is composed of two virion populations (Basnayake *et al.*, 2006): one type contains RNA1 and RNA2, and the other type contains multiple copies (possibly four copies) of RNA2. The origin of assembly sequence (OAS) was identified in RNA2 using the TBSV-vector-based expression of RCNMV CP (Basnayake *et al.*, 2009). RNA1 is copackaged with RNA2 via an OAS–RNA1 interaction. Interestingly, the OAS is the TA that is required for the transcription of CPsgRNA from RNA1 via an RNA–RNA interaction (Sit *et al.*, 1998). The N-terminal lysine-rich motif of CP is involved in the formation of the virions in *N. benthamiana* (Park *et al.*, 2012). The heterologous combinations of RNA1 and RNA2 between RCNMV and SCNMV were infectious and produced progeny virions in their host plants (Okuno *et al.*, 1983; Chen *et al.*, 1984). A mixture of RNA1 from CRSV and

RNA2 from RCNMV was also infectious, and produced progeny virions in the host plants (Callaway *et al.*, 2001). However, unilateral compatibility of RNA1 and RNA2 was reported using the different strains of RCNMV (Rao and Hiruki, 1987).

The electrophoretic mobility of purified virions differ among the dianthoviruses, even in the different strains of the same virus (Pappu and Hiruki, 1989).

Analysis of the RCNMV virions by cryoelectron microscopy and three-dimensional image reconstruction showed that the structures of the CP subunits and the entire virion are similar to those observed in other viruses of the *Tombusviridae*, such as TBSV and TCV (Sherman *et al.*, 2006). Preliminary diffraction data are available for the X ray structure of RCNMV virion crystals at higher than 4 Å resolution (Martin *et al.*, 2010). Atomic absorption spectroscopic analysis of RCNMV virions showed that the virions contain significant amounts of Ca<sup>2+</sup> and Mg<sup>2+</sup> ions (Sherman *et al.*, 2006). Removal of these ions from the virions with a chelator, followed by exposure to ribonucleases, reduces viral infectivity, suggesting a role for these ions in virion stabilization (Sherman *et al.*, 2006).

Studies of the use of RCNMV virions as viral nanoparticles that function as drug carriers in human cells are in progress (Lockney *et al.*, 2011).

#### **4.14. Suppression of RNA silencing**

RNA silencing is an RNA-mediated plant defense mechanism against viral infection (Baulcombe, 2004). To counteract RNA silencing, viruses have developed a variety of strategies and evolved to encode a viral suppressor of RNA silencing (VSR) (Ding and Voinnet, 2007; Li and Ding, 2006; Csorba *et al.*, 2009). Many plant viruses encode at least one VSR and some viruses encode more than one. Viral proteins that function as VSRs include CPs, replicase proteins, MPs, and other nonstructural proteins, and their targets in RNA silencing are diverse (Ding and Voinnet, 2007; Li and Ding, 2006; Csorba *et al.*, 2009).

RCNMV has at least two strategies for suppressing RNA silencing. One strategy requires p27, p88, and viral RNAs (Takeda *et al.*, 2005). In a sense-transgene-mediated posttranscriptional gene silencing (S-PTGS) assay involving the *Agrobacterium*-mediated transient expression of viral components and GFP in a GFP-expressing transgenic plant, the RNA-silencing-suppression activity of RCNMV was linked to the ability of viral factors to initiate negative-strand RNA synthesis (Takeda *et al.*, 2005), and correlated with the formation of the 480-kDa replicase complex (Mine *et al.*, 2010a). A close relationship between negative-strand RNA synthesis and the suppression of RNA silencing implies a possible scenario, in which

RCNMV sequesters the host factors required for RNA silencing and reduces the antiviral silencing response. RCNMV replication also inhibits microRNA biogenesis, in which DCL1 plays an essential role, and *dcl1* mutant plants show reduced susceptibility to RCNMV infection (Takeda *et al.*, 2005). Therefore, it has been suggested that DCL1 or its homologues are recruited by the viral RNA replication complex (Takeda *et al.*, 2005). Alternatively, the reduced susceptibility of a *dcl1* mutant to RCNMV infection can also be explained by the observations that the disruption of the DCL1 function leads to the higher expression of DCL4 and DCL3 in *Arabidopsis* leaves (Qu *et al.*, 2008) and that the destabilization of the miRNA pathway, including the disruption of DCL1, leads to the increased accumulation of AGO1 and positively influences S-PTGS (Martinez de Alba *et al.*, 2011). Given that RCNMV infection inhibits miRNA biogenesis, reduced accumulations of miRNAs, including miR168 that is a regulator of AGO1 mRNA, could increase the accumulation levels of AGO1, which is the core component of the RNA-induced silencing complex (RISC).

MP has been identified as the second VSR of RCNMV in another assay system (Powers *et al.*, 2008a and b). This assay relies on TCV that contains a reporter GFP gene in place of the CP ORF (TCV-sGFP) (Powers *et al.*, 2008b). The TCV CP is a VSR (Qu *et al.*, 2003), and TCV requires CP for its cell-to-cell movement in *N. benthamiana* (Cohen *et al.*, 2000). TCV-sGFP moved from cell to cell in leaves infiltrated with *Agrobacterium* expressing RCNMV MP, but did not in leaves infiltrated with *Agrobacterium* expressing RCNMV p27, p88, and CP (Powers *et al.*, 2008a). The functional domains in RCNMV MP required for RNA silencing (Powers *et al.*, 2008a) differ from those required for the cell-to-cell movement of the virus and SEL modification (Fujiwara *et al.*, 1993; Tremblay *et al.*, 2005). Thus, RCNMV has evolved two strategies for suppressing RNA silencing and counteracting the host defenses.

## 5. TRANSMISSION STUDIES AND EPIDEMIOLOGY

Dianthoviruses multiply in infected leaves to high concentrations and are very stable *in vitro* (Hiruki, 1987). Soil transmission has been reported to be a general feature of dianthoviruses (Brunt *et al.*, 1996).

The transmission of CRSV by nematodes such as *Longidorus macrosoma* and *Xiphinema diversicaudatum* (Fritzsche and Schmelzer, 1967; Kegler *et al.*, 1977) and *L. elongates* (Fritzsche *et al.*, 1979) has been reported, but not confirmed (Brunt *et al.*, 1996).

The nematode transmission of FNSV is suspected but not demonstrated (Morales *et*

*al.*, 1992). No other vectors have been investigated in terms of the soil transmission of this virus.

The possible transmission of RCNMV by soil-inhabiting fungi was suggested as early as 1977 (Bowen and Plumb, 1979; Lange, 1977). A Swedish isolate of RCNMV was transmitted to *N. clevelandii* when the seedlings were planted into soil that had been infested by growing RCNMV-infected plants, or by adding virus suspension to the soil prior to planting. The concomitant presence of *Olpidium brassicae* only increased the rate of transmission, but not essential as an intrinsic vector for the soil transmission of the virus (Gerhaldson and Insunza, 1979). Independent tests by different groups did not demonstrate that *O. brassicae* serves as a vector of RCNMV (Leggat, 1981; Lyness *et al.*, 1981). Hiruki (1986) reported that the presence of a tobacco strain of *O. brassicae* was not required for the transmission of SCNMV in sand culture.

Compared with leguminous forage crops such as alfalfa, alsike clover, red clover, crown vetch, and white clover, sweet clover is highly susceptible to SCNMV, while the others remain practically resistant to the virus. It is very interesting to note that the most prevalent area of SCNMV occurrence in Alberta coincides with the area where beekeeping industry is most active and where sweet clover is commonly used for seed production. Hiruki *et al.* (1989) reported that the pollen from SCNMV-infected sweet clover is highly contaminated with the virus and that the virus, detectable by enzyme-linked immunosorbent assays, could be recovered by washing.

In 2007, an area of approximately a quarter section of land, consisting of 50% arable land and 50% popular forest, was cleared for residential development in a suburb of Edmonton. The topsoil, about 10 cm in depth, was completely removed and piled aside. However, due to unexpected economic downturn, the construction was halted for three years. To prevent erosion, sweet clover was seeded over the whole area (Fig. 3), offering a rare opportunity to observe the natural development of plant virus epidemic.

Regular visual inspections and samplings were conducted, and followed by infectivity tests on the indicator plants in the greenhouse. The results indicated that SCNMV infection occurred at a rate of about 6%. The source of the natural infection remains unknown. In the past, the natural infection of a solitary sweet clover plant in the forest, roadside, or riverbanks was observed frequently and shown to test positive (C. Hiruki, unpublished results). When infected sweet clover is not disturbed, abundant seeds are shed around the plant, and the young seedlings are subsequently infected by the soil transmission of SCNMV.

## 6. EFFECTS ON ENVIRONMENT

Although considerable attention has been directed toward the pollution of environmental waters with human and animal viruses, there have been only sporadic attempts to shed light on the contamination of rivers, lakes and sea with plant viruses (Tomlinson *et al.*, 1982a, 1982b, 1983; Koenig, 1985; Büttner, *et al.*, 1987).

### 6.1. Release of dianthoviruses from intact roots, decaying plant materials, and pollen grains

Generally, dianthoviruses occur at high concentrations in infected cells and are very stable *in vitro*, thus serve as potentially highly infective inoculum sources. The release of CRSV from infected roots to drainage water was reported in Germany (Kegler and Kegler, 1981). Actively growing plants of *N. clevelandii* released much larger amounts of RCNMV into drainage water than did plants from which the aerial parts had been removed or plants which were senescent or dead (Gerhardson and Insunza, 1979). CRSV was isolated from water samples collected from a canal near a sewage plant in Germany (Koenig *et al.*, 1988).

While sweet clover is a biennial crop, successive infections with SCNMV can occur from the soil around an infected plant to young germinating plants in the second year, and from the soil around the dead infected plants to young plants emerging from fallen seeds in the third year (Hiruki, unpublished data).

Another strong possibility is pollen transmission in the case of SCVMV, although its direct experimental evidence is lacking at present. However, significantly high degrees of SCNMV occurrence and release of the virus particles from pollen grains were detected serologically (Hiruki *et al.*, 1989). Moreover, in systematic field surveys over many years, high specific incidences of SCNMV coincide with the zones three (black soil) and four (gray wooded soil) in the Agriculture map of Alberta (Hiruki, 1987) where legume seed industry and bee keeping industries are concentrated (Hiruki, unpublished data). There are three main species of bees to pollinate legume crops such as honey bees (*Apis mellifera*), leafcutting bees (*Megachile rotundata*) and American bumble bees (*Bombus fervidus*). Honey bees are effective pollinators of a number of legume crops including sweet clover. However, honey bees are of little value as pollinator of alfalfa (D. T. Fairey, 2003). This fact may explain a minor incidence of SCNMV in alfalfa while frequent incidence of the same virus is observed in sweet clover in the same area (Hiruki *et al.*, 1984, Inouye and Hiruki, 1984, Hiruki, 1986). The occurrence of SCNMV in sweet clover in the forests, river banks, roadsides and



abandoned field may be explained by pollen transmission mediated mostly by honey bees and to lesser degrees by other bees.



Figure 2.3 A general view of the site where sweet clover plants were grown to prevent soil erosion. This site was used to observe the natural spread of SCNMV.

## 6.2. Sewage

CRSV was isolated from river water that was fed by the outlets of a city's sewage plant in Germany (Koenig *et al.*, 1988). The virus was identified by means of nucleic acid hybridization analysis and serology. CRSV and other viruses can also be released into streams or rivers. It is known that several plant viruses are capable of surviving a passage through the alimentary tracts of animals (Tomlinson *et al.*, 1982; Kegler *et al.*, 1987). SCNMV was not transmitted by *Sitona cylindricollus* from sweet clover to sweet clover but remained infectious in feces after a passage through the alimentary tract (Hiruki, 1986).

At any rate, these observations on the occurrence of plant viruses in streams, rivers and lakes illustrate the potential danger of using polluted water in horticultural and agricultural practices (Koenig and Lesemann, 1985; Koenig, 1986; Buttner *et al.*, 1987; Koenig *et al.*, 1988).

## 7. CONCLUDING REMARKS AND PERSPECTIVES

The dianthoviruses have moderately broad natural host ranges, restricted to dicot plant species, including legume plants (Hiruki, 1986; Descriptions lists from the VIDE Database; Virus Taxonomy, Ninth Report of the International Committee on Taxonomy of Viruses, 2012), and the viruses and many of their strains have been reported frequently worldwide except for the tropical and subtropical areas of the world. However, a natural host of FNSV, an apparent member of the dianthoviruses, is *Furcareia* spp., which are xerophytic monocots native to the tropical regions of Mexico, the Caribbean, Central America and northern South America. Dianthovirus RNA1-like RNA has also been reported in grassy stunt-diseased rice plants coinfecting with *Rice grassy stunt virus* in the Philippines (Miranda *et al.*, 2001). These reports imply that other unidentified *Dianthovirus* species or viruses related to the dianthoviruses might exist in tropical and subtropical areas. Therefore, a systematic search for dianthoviruses worldwide, including in wild plants in fields, forests, and areas of water such as river, ponds, and seas, should open the way to finding additional members of *Dianthovirus*.

Dianthoviruses are known to release large numbers of infective viral particles from infected roots into soil and drainage waters in the surrounding area (Gewrhardson and Insunza, 1979; Hiruki, 1986; Kegler and Kegler, 1981). FNSV is detected consistently in the roots of infected fiqué plants (Morales *et al.*, 1992). In nature, soil is a complex medium and acts as a heterogeneous environment. Its properties can be altered by seasonal climatic changes and agricultural practices in a particular area. This implies that any infective agents, such as viruses, introduced into the soil and drainage water may be affected by a variety of environmental factors or combination of several factors that are influenced by the interplay between the climate, vegetation and soil type. It is also expected that the application of sewage and sludge to agricultural land may alter some of the soil's physical and chemical properties that affect viral survival.

Our understanding of the molecular mechanisms underlying the infection processes of the dianthoviruses has advanced greatly in recent years, as described in this review. These advances have also given rise many interesting questions that are yet to be addressed. These questions include the differences in the gene expression strategies of RNA1 and RNA2. For example, RNA1 contains strong translation enhancer elements (3'TE-DR1 and ARS), whereas RNA2 has no such elements: instead, the translation of MP is linked to the replication of RNA2 (Mizumoto *et al.*, 2006). What do these differences in translation strategies mean? RNA1 and RNA2 might use distinct

translation factors. The recruitment of MP to viral RNAs, a process essential for viral cell-to-cell movement, also differs between RNA1 and RNA2. The replication of RNA1, but not RNA2, is associated with MP recruitment to the ER-associated viral replication complexes (Kaido et al., 2009). This strategy might assist the cell-to-cell movement of RNA1, which does not encode MP. These findings also suggest that the replication complexes formed on RNA1 might differ from those on RNA2, and may localize to different sites on the ER. Indeed, RNA2 has an RNA element (YRE) that recruits p27 and the replicase complex (An et al., 2010; Hyodo et al. 2011), whereas RNA1 has no such recruiting element. The replication of RNA1 is linked to the translation of the replicase proteins (Okamoto et al., 2008; Iwakawa et al., 2011).

The TA of RNA2 is a key regulator of translation, transcription, replication, virion formation, and cell-to-cell movement. The structural requirements of TA seem to differ depending on its role, because nucleotide substitutions in RNA2 that disrupt TA-mediated base pairing have no effect on the copackaging of RNA1 and RNA2 (Basnayake *et al.*, 2009). Further analyses of the temporal changes in the TA structure and the host and viral factors that interact with TA are necessary to clarify the complex regulatory mechanisms of the *Dianthovirus* infection processes.

The auxiliary replication protein p27 plays multiple roles in RNA replication. These roles include the recruitment of the viral RNA to the ER membranes, the assembly of the replication complexes, and the formation of the viral replication factory. p27 performs these roles by interacting with p27 itself, p88, and the viral RNA, and also by interacting with host proteins, such as Hsp70, Hsp90, and Arf1. How can p27 perform these roles properly? A model proposed by Nagy *et al.* (2012) for the *Tombusvirus* auxiliary replication protein p33 might help to answer this question. These researchers propose that “depending on the given interacting partners, the abundant p33 molecules are divided into many groups that perform different functions at different subcellular locations”. Further identification and analysis of the host proteins associated with the different functions of RCNMV p27 are required to answer this question.

Another interesting question to be addressed is the mechanism that delivers the translation initiation factors from the 3'-UTR of RCNMV RNA1 to the 5'-UTR, where the ribosome starts to scan the template (Sarawaneeyaruk *et al.*, 2009). In 3'TE-DR1-mediated cap-independent translation, PABP plays an essential role in the binding of eIF4F/eIFiso4F to 3'TE-DR1 (Iwakawa *et al.*, 2012). It is generally believed that the binding of eIF4F and eIFiso4F to viral 3' CITEs facilitates ribosome recruitment to the 5' end of the RNA *via* 5'–3' communication, which is mediated by a long-distance RNA–RNA interaction or a protein factor (Miller and White, 2006; Nicholson and

White, 2011). However, the factors other than eIF4F/eIFiso4F or PABP that are required for 3'CITE-mediated translation remains unclear. How the ribosome recognizes the uncapped 5' end, even when eIF4F/iso4F is delivered to the vicinity of the 5' end of RNA1 by 5'–3' communication is also unclear. Although we do not know how RCNMV RNA1 achieves the communication between its 5'- and 3'-ends in 3'TE-DR1-mediated cap-independent translation, our recent data suggest that the binding of eIF4F/iso4F and PABP to the 3'-UTR of RCNMV RNA1 is important for delivering cap-binding factors other than eIF4F/iso4F and PABP to its nonfunctional G-capped 5' end, and in enhancing the recruitment of the 40S ribosomal subunit to the viral mRNA (H. Iwakawa, and T. Okuno; unpublished data). Identification of the proteins associated with the 5' end of RNA1 will help to clarify these unknown mechanisms.

As described in this review, recent advances in our knowledge of the dianthoviruses, especially of their molecular biology, have relied on the development and availability of several experimental systems, such as an *in vitro* translation/replication system (BYL) (Komoda *et al.*, 2004), and the *Agrobacterium*-mediated expression of viral and host components in combination with a protein-mediated or RNA-aptamer-mediated pull-down assay and mass spectrometric analysis. Genome-wide screens of the host factors that affect viral replication have been performed using yeast as the model host. The lists of host proteins obtained from these studies in yeast (Kushner *et al.*, 2003; Panavas *et al.*, 2005; Serviene *et al.*, 2005), together with those obtained with other approaches, including those involving RCNMV, will enhance our understanding of the molecular mechanisms underlying viral infection. This knowledge will contribute to the development of antiviral measures that are more efficient, which will reduce the disease-mediated damage caused by the virus infections. However, we must consider that both the viral and host factors that affect viral infection might differ in different combinations of viruses and hosts (Li and Wong, 2007; Sarawaneeyaruk *et al.*, 2009).

## **ACKNOWLEDGMENTS**

The authors thank all their colleagues and associates who have contributed to studies of the dianthoviruses. This work was supported in part by funds from the Ministry of Education, Culture, Sports, Science, and Technology, Japan to Tetsuro Okuno, and from Natural Sciences and Engineering Research Council of Canada (Grant No. A3843, STR G1450, IC0145) and University of Alberta Distinguished University Professor Research Grant to Chuji Hiruki.

## REFERENCES

- An, M., Iwakawa, H. O., Mine, A., Kaido, M., Mise, K., and Okuno, T. (2010). A Y-shaped RNA structure in the 3' untranslated region together with the trans-activator and core promoter of Red clover necrotic mosaic virus RNA2 is required for its negative-strand RNA synthesis. *Virology* 405:100–109.
- Barry, J. K., and Miller, W. A. (2002). A -1 ribosomal frameshift element that requires base pairing across four kilobases suggests a mechanism of regulating ribosome and replicase traffic on a viral RNA. *Proc. Natl. Acad. Sci. USA* 99:11133–11138.
- Basnayake, V. R., Sit, T. L., and Lommel, S. A. (2006). The genomic RNA packaging scheme of *Red clover necrotic mosaic virus*. *Virology* 345:532–539.
- Basnayake, V. R., Sit, T. L., and Lommel, S. A. (2009). The *Red clover necrotic mosaic virus* origin of assembly is delimited to the RNA-2 trans-activator. *Virology* 384:169–178.
- Bates, H. J., Farjah, M., Osman, T. A., and Buck, K.W. (1995). Isolation and characterization of an RNA-dependent RNA polymerase from *Nicotiana clevelandii* plants infected with red clover necrotic mosaic dianthovirus. *J. Gen. Virol.* 76:1483–1491.
- Baulcombe, D. (2004). RNA silencing in plants. *Nature* 431:356–363.
- Belov, A. G., and van Kuppeveld, J. M. (2012). (+)RNA viruses rewire cellular pathways to build replication organelles. *Curr. Opin. Virol.* 2:734-741.
- Blobel, G., and Sabatini, D. (1971). Dissociation of mammalian polyribosomes into subunits by puromycin. *Proc. Natl. Acad. Sci. USA.* 68:390–394.
- Bowen, R. and Plumb, R. T. (1979). The occurrence and effects of red clover necrotic mosaic virus in red clover (*Trifolium pretense*). *Ann. Appl. Biol.* 91:227-236.
- Brierley, I. (1995). Ribosomal frameshifting viral RNAs. *J. Gen. Virol.* 76:1885–1892.
- Brunt, A. A., Crabtree, K., Dallwitz, M. J., Gibbs, A. J., and Watson, L. (1996). *Viruses of Plants*. CAB International. Wallingford, U.K.
- Buttner, C., Jacobi, V., and Koenig, R. (1987). Isolation of carnation Italian ringspot virus from a creek in a forested area Southwest of Bonn. *J. Phytopathol.* 118:131-134.
- Callaway, A. S., George, C. G., and Lommel, S. A. (2004). A Sobemovirus coat protein gene complements long-distance movement of a coat protein-null Dianthovirus. *Virology* 330:186–195.
- Callaway, A., Giesman-Cookmeyer, D., Gillock, E. T., Sit, T. L., and Lommel, S. A. (2001). The multifunctional capsid proteins of plant RNA viruses. *Ann. Rev. Phytopathol.* 39:419-460.

- Chang, R. Y., Hofmann, M. A., Sethna, P. B., and Brian, D. A., 1994. A cis-acting function for the coronavirus leader in defective interfering RNA replication. *J. Virol.* 68:8223-8231.
- Chen, M. H., Hiruki, C., and Okuno, T. (1984). Immunosorbent electron microscopy of dianthoviruses and their pseudorecombinants. *Can. J. Plant Pathol.* 6:191-195.
- Chattopadhyay, M., Shi, K., Yuan, X., and Simon, A. E. (2011). Long-distance kissing loop interactions between a 3' proximal Y-shaped structure and apical loops of 5' hairpins enhance translation of *Saguaro cactus virus*. *Virology* 417:113-125.
- Cohen, Y., Gisel, A., and Zambryski, P. C. (2000). Cell-to-cell and systemic movement of recombinant green fluorescent protein-tagged turnip crinkle viruses. *Virology* 273:258-266.
- Csorba, T., Pantaleo, V., and Burgyan, J. (2009). RNA silencing: an antiviral mechanism. *Adv. Virus Res.* 75: 35–71.
- D'Souza-Schorey, C., and Chavrier, P. (2006). ARF proteins: roles in membrane traffic and beyond. *Nat. Rev. Mol. Cell. Biol.* 7:347-358.
- den Boon, J. A., and Ahlquist, P. (2010). Organelle-Like membrane compartmentalization of positive-strand RNA virus replication factories. *Nat. Rev. Microbiol.* 64:241-256.
- Ding, S. W., and Voinnet, O. (2007). Antiviral immunity directed by small RNAs. *Cell* 130:413–426.
- Donaldson, J. G., and Jackson, C. L. (2011). ARF family G proteins and their regulators: roles in membrane transport, development and disease. *Nat. Rev. Mol. Cell Biol.* 12:362–375.
- Dreher, T. W. (1999). Functions of the 3'-untranslated region of positive strand RNA viral genomes. *Annu. Rev. Phytopathol.* 37: 151–174.
- Eckerle, L.D., Albarino, C.G., and Ball, L.A. (2003). Flock House virus subgenomic RNA3 is replicated and its replication correlates with transactivation of RNA2. *Virology* 317:95–108.
- Fabian, M. R., and White, K. A. (2004). 5'-3' RNA–RNA interaction facilitates cap- and poly(A) tail-independent translation of tomato bushy stunt virus mRNA: A potential common mechanism for tombusviridae. *J. Biol. Chem.* 279:28862–28872.
- Fabian, M. R., and White, K. A. (2006). Analysis of a 39-translation enhancer in a tombusvirus: A dynamic model for RNA-RNA interactions of mRNA termini. *RNA* 12:1304–1314.
- Fairey, D. T. (2003). Growing forage legume for seed. *Government of Alberta, Canada, Agdex* 120/15-1.

- Fritzsche, R. and Schmelzer, K. (1967). Nematode-transmissibility of carnation ringspot virus. *Naturwissenschaften* 54:498-499.
- Fritzsche, R., Kegler, H., Thiele, S., and Gruber, G. (1979). Contribution to epidemiology and transmission of carnation ringspot virus in fruit plantations. *Arch. Phytopathol. Pflanzenschutz* 15:177-180.
- Fujiwara, T., Giesman-Cookmeyer, D., Ding, B., Lommel, S. A., and Lucas, W. J. (1993). Cell-to-cell trafficking of macromolecules through plasmodesmata potentiated by the red clover necrotic mosaic virus movement protein. *Plant Cell* 5:1783–1794.
- Gazo, B.M., Murphy, P., Gatchel, J. R., and Browning, K. S. (2004). A novel interaction of cap-binding protein complexes eukaryotic initiation factor (eIF)4F and eIF(iso)4F with a region in the 3'-untranslated region of satellite tobacco necrosis virus. *J. Biol. Chem.* 279 (4):13584-13592.
- Ge, Z., Hiruki, C., and Roy, K. L. (1992). A comparative study of the RNA-2 nucleotide sequences of two sweet clover necrotic mosaic virus strains. *J. Gen. Virol.* 73: 2483–2486.
- Ge, Z., Hiruki, C., and Roy, K. L. (1993). Nucleotide sequence of sweet clover necrotic mosaic dianthovirus RNA-1. *Virus Res.* 28:113–124.
- Gerhardson, B., and Insunza, V. (1979). Soil transmission of red clover necrotic mosaic virus. *Phytopathol. Z.* 94:67-71.
- Gerhardson, B. and Lindsten, K. (1973). Red clover mottle virus and red clover necrotic mosaic virus in Sweden. *Phytopath. Z.* 76:67-79.
- Giesman-Cookmeyer, D. and Lommel, S. A. (1993). Alanine scanning mutagenesis of a plant virus movement protein identifies three functional domains. *Plant Cell* 5:973–982.
- Giesman-Cookmeyer, D., Silver, S., Vaewhong, A. A., Lommel, S. A., & Deom, C. M. (1995). Tobamovirus and dianthovirus movement proteins are functionally homologous. *Virology*, 213, 38–45.
- Gould, A. R., Francki, R. I., Hatta, T., and Hollings, M. (1981). The bipartite genome of red clover necrotic mosaic virus. *Virology* 108:499–506.
- Guenther, R. H., Sit, T. L., Gracz, H. S., Dolan, M. A., Townsend, H. L., Liu, G., Newman, W. H., Agris, P. F., and Lommel, S. A. (2004). Structural characterization of an intermolecular RNA-RNA interaction involved in the transcription regulation element of a bipartite plant virus. *Nucleic Acids Res.* 32:2819–2828.
- Guo, L., Allen, E., and Miller, W. A. (2000). Structure and function of a

- cap-independent translation element that functions in either the 3' or the 5' untranslated region. *RNA* 6:1808–1820.
- Guo, L., Allen, E., and Miller, W. A. (2001). Base-pairing between untranslated regions facilitates translation of uncapped, nonpolyadenylated viral RNA. *Mol. Cell* 7:1103–1109.
- Hagino-Yamagishi, K., and Nomoto, A., 1989. *In vitro* construction of poliovirus defective interfering particles. *J. Virol.* 63:5386-5392.
- Hiruki, C., Rao, D. V., Chen, M. H., Okuno, T., and Figueiredo, G. C. (1984). Characterization of sweet clover necrotic mosaic virus. *Phytopathology* 74:482-486.
- Hiruki, C. (1986a). Sweet clover necrotic mosaic virus. *AAB Descriptions of Plant Viruses, Assoc. Applied Biologists, U.K.* 321:1-4.
- Hiruki, C. (1986b). Incidence and geographic distribution of sweet clover necrotic mosaic virus in Alberta. *Plant Dis.* 70: 1129-1131.
- Hiruki, C. (1987). The dianthoviruses: a distinct group of isometric plant viruses with bipartite genome. *Adv. Virus Res.* 33:257–300.
- Hiruki, C., Kudo, K., and Figueiredo, G. (1989). Transmission of sweet clover necrotic mosaic virus. *Proc. Japan Acad.* 65B:234-237.
- Hiruki, C. (1994). Transmission of dianthoviruses. *Acta Hort.* 377:341-347.
- Hyodo, K., Mine, A., Iwakawa, H.O., Kaido, M., Mise, K., and Okuno, T. (2011). Identification of amino acids in auxiliary replicase protein p27 critical for its RNA-binding activity and the assembly of the replicase complex in Red clover necrotic mosaic virus. *Virology* 413:300–309.
- Hyodo, K., Mine, A., Iwakawa, H.O., Kaido, M., Mise, K., and Okuno, T. (2013). The ADP-ribosylation factor 1 plays an essential role in the replication of a plant RNA virus. *J. Virol.* 87:163-176.
- Iwakawa, H. O., Kaido, M., Mise, K., and Okuno, T. (2007). *cis*-Acting core RNA elements required for negative-strand RNA synthesis and cap-independent translation are separated in the 3'-untranslated region of *Red clover necrotic mosaic virus* RNA1. *Virology* 369:168–181.
- Iwakawa, H. O., Mine, A., Hyodo, K., An, M., Kaido, M., Mise, K., and Okuno, T. (2011). Template recognition mechanisms by replicase proteins differ between bipartite positive-strand genomic RNAs of a plant virus. *J. Virol.* 85:497–509.
- Iwakawa, H. O., Mizumoto, H., Nagano, H., Imoto, Y., Takigawa, K., Sarawaneeyaruk, S., Kaido, M., Mise, K., and Okuno, T. (2008). A viral noncoding RNA generated by *cis*-element-mediated protection against 5'→3' RNA decay represses both



- cap-independent and cap-dependent translation. *J. Virol.* 82:10162–10174.
- Iwakawa, H. O., Tajima, Y., Taniguchi, T., Kaido, M., Mise, K., Tomari, Y., Taniguchi, H., and Okuno, T. (2012). Poly(A)-binding protein facilitates translation of an uncapped/nonpolyadenylated viral RNA by binding to the 3' untranslated region. *J. Virol.* 86 (15):7836-7849.
- Johnson, K. L., and Sarnow, P., 1991. Three poliovirus 2B mutants exhibit noncomplementable defects in viral RNA amplification and display dosage-dependent dominance over wild-type poliovirus. *J. Virol.* 65:4341-4349.
- Kaido, M., Funatsu, N., Tsuno, Y., Mise, K., and Okuno, T. (2011). Viral cell-to-cell movement requires formation of cortical punctate structures containing Red clover necrotic mosaic virus movement protein. *Virology* 413:205–215.
- Kaido, M., Tsuno, Y., Mise, K., and Okuno, T. (2009). Endoplasmic reticulum targeting of the Red clover necrotic mosaic virus movement protein is associated with the replication of viral RNA1 but not that of RNA2. *Virology* 395:232–242.
- Kendall, T. L., and Lommel, S. A. (1992). Nucleotide sequence of carnation ringspot dianthovirus RNA-2. *J. Gen. Virol.* 73:2479–2482.
- Karetnikov, A., and Lehto, K. (2008). Translation mechanisms involving long-distance base pairing interactions between the 5' and 3' non-translated regions and internal ribosomal entry are conserved for both genomic RNAs of *Blackcurrant reversion nepovirus*. *Virology* 371:292–308.
- Kim, K. H., and Lommel, S. A. (1994). Identification and analysis of the site of -1 ribosomal frameshifting in red clover necrotic mosaic virus. *Virology* 200:574–582.
- Kim, K. H., and Lommel, S. A. (1998). Sequence element required for efficient -1 ribosomal frameshifting in red clover necrotic mosaic dianthovirus. *Virology* 250, 50–59.
- Kneller, E. L., Rakotondrafara, A. M., and Miller, W. A. (2006). Cap-independent translation of plant viral RNAs. *Virus. Res.* 119:63–75.
- Koenig, R., and Lesemann, D. E. (1985). Plant viruses in German rivers and lakes. I. Tombusviruses, a potyvirus and carnation mottle virus. *Phytopathol. Z.* 112:105-116.
- Koenig, R., An, D., Lesemann, D.E., and Burgermeister, W. (1988). Isolation of carnation ringspot virus from a canal near a sewage plant: cDNA hybridization analysis, serology, and cytopathology. *J. Phytopathol.* 121: 346-356.
- Komoda, K., Naito, S., and Ishikawa, M. (2004). Replication of plant RNA virus genomes in a cell-free extract of evacuated plant protoplasts. *Proc. Natl. Acad.*

- Sci. U. S. A.* 101:1863–1867.
- Koonin, E. V. (1991). The phylogeny of RNA-dependent RNA polymerases of positive-strand RNA viruses. *J. Gen. Virol.* 72:2197–2206.
- Koonin, E. V., and Dolja, V. V. (1993). Evolution and taxonomy of positive-strand RNA viruses: implications of comparative analysis of amino acid sequences. *Crit. Rev. Biochem. Mol. Biol.* 28:375–430.
- Kushner, D. B., Lindenbach, B. D., Grdzlishvili, V. Z., Noueir, A. O., Paul, S. M., and Ahlquist, P. (2003). Systematic, genome-wide identification of host genes affecting replication of a positive-strand RNA virus. *Proc. Natl. Acad. Sci. U. S. A.* 100:15764–15769.
- Kusumanegara, K., Mine, A., Hyodo, K., Kaido, M., Mise, K., and Okuno, T. (2012). Identification of domains in p27 auxiliary replicase protein essential for its association with the endoplasmic reticulum membranes in *Red clover necrotic mosaic virus*. *Virology* 433:131-141.
- Lehninger, A. L., Nelson, D. L., and Cox, M. M. (1993). *Principles of biochemistry*, p. 927–928. Worth Publishers, New York, N.Y.
- Lewandowski, D. J., and Dawson, W. O. (2000). Functions of the 126- and 183-kDa proteins of tobacco mosaic virus. *Virology* 271:90-98.
- Li, F., and Ding, S. W. (2006). Virus counterdefense: diverse strategies for evading the RNA-silencing immunity. *Annual review of microbiology* 60:503–531.
- Li, W., and Wong, S. M. (2007). Host-dependent effects of the 3' untranslated region of turnip crinkle virus RNA on accumulation in Hibiscus and Arabidopsis. *J. Gen. Virol.* 88:680-687.
- Liang, Y., and Gillam, S. (2001). Rubella virus RNA replication is *cis*-preferential and synthesis of negative- and positive-strand RNAs is regulated by the processing of nonstructural protein. *Virology* 282:307-319.
- Lin, H. X., and White, K. A. (2004). A complex network of RNA-RNA interactions controls subgenomic mRNA transcription in a tombusvirus. *EMBO J* 23, 3365–3374.
- Lockney, D. M., Guenther, R. N., Loo, L., Overton, W., Antonelli, R., Clark, J., Hu, M., Luft, C., Lommel, S. A., and Franzen, S. (2011). The Red clover necrotic mosaic virus capsid as a multifunctional cell targeting plant viral nanoparticle. *Bioconjugate Chem.* 22:67–73.
- Lommel, S. A., Weston-Fina, M., Xiong, Z., and Lomonosoff, G. P. (1988). The nucleotide sequence and gene organization of red clover necrotic mosaic virus

- RNA-2. *Nucleic Acids Res.* 16:8587–8602.
- Lynes, E. W., Teakle, D. S., and Smith, P. R. (1981). Red clover necrotic mosaic virus isolated from *Trifolium repens* and *Medicago sativa* in Victoria. *Aust. Plant Pathol.* 10:6-7.
- Mahajan, S., Dolja, V. V., and Carrington, J. C. (1996). Roles of the sequence encoding tobacco etch virus capsid protein I genome amplification: requirements for the translation process and a cis-active element. *J. Virol.* 70:4370-4379.
- Martin, S. L., Guenther, R. H., Sit, T. L., Swartz, P. D., Meilleur, F., Lommel, S. A., and Rose, R. B. (2010). Crystallization and preliminary X-ray diffraction analysis of red clover necrotic mosaic virus. *Acta Cryst.* F66:1458-1462.
- Martinez de Alba, A. E., Jauvion, V., Mallory, A. C., Bouteiller, N., and Vaucheret, H. (2011). The miRNA pathway limits AGO1 availability during siRNA-mediated PTGS defense against exogenous RNA. *Nucleic Acids Res.* 39: 9339–9344.
- Matthews, R. E. F. (1982). “Classification and Nomenclature of Viruses,” 4<sup>th</sup> Report of the International Committee for Taxonomy of Viruses. Karger, Basel.
- McCormack, J. C., Yuan, X., Yingling, Y. G., Kasprzak, W., Zamora, R. E., Shapiro, B. A., and Simon, A. E. (2008). Structural domains within the 3' untranslated region of Turnip crinkle virus. *J. Virol.* 82:8706–8720.
- Memon, A. R. (2004). The role of ADP-ribosylation factor and SAR1 in vesicular trafficking in plants. *Biochi. Biophys. Acta* 1664;9 – 30.
- Meulewaeter, F., van Lipzig, R., Gulyaev, A. P., Pleij, C. W., Van Damme, D., Cornelissen, M., and van Eldik, G. (2004). Conservation of RNA structures enables TNV and BYDV 5' and 3' elements to cooperate synergistically in cap-independent translation. *Nucleic Acids Res.* 32:1721–1730.
- Miller, S., and Krijnse-Locker, J. (2008). Modification of intracellular membrane structure for virus replication. *Nat. Rev. Microbiol.* 6:363-374.
- Miller, W. A., Wang, Z., and Treder, K. (2007). The amazing diversity of cap-independent translation elements in the 3'-untranslated regions of plant viral RNAs. *Biochem. Soc. Trans* 35:1629–1633.
- Miller, W. A., and White, K. A. (2006). Long-distance RNA-RNA interactions in plant virus gene expression and replication. *Annu. Rev. Phytopathol.* 44:447–467.
- Mine, A., and Okuno, T. (2012). Composition of plant virus RNA replicase complexes. *Curr. Opin. Virol.* 2:663-669.
- Mine, A., Hyodo, K., Takeda, A., Kaido, M., Mise, K., and Okuno, T. (2010a). Interactions between p27 and p88 replicase proteins of *Red clover necrotic mosaic virus* play an essential role in viral RNA replication and suppression of RNA

- silencing via the 480-kDa viral replicase complex assembly. *Virology* 407:213–224.
- Mine A, Takeda, A., Taniguchi, T., Taniguchi, H., Kaido, M., Mise, K. and Okuno, T. (2010b). Identification and characterization of the 480-kilodalton template-specific RNA-dependent RNA polymerase complex of *Red clover necrotic mosaic virus*. *J. Virol.* 84:6070–6081.
- Mine, A., Hyodo, K., Tajima, Y., Taniguchi, T., Taniguchi, H., Kaido, M., Mise, K., and Okuno, T. (2012). Differential roles of Hsp70 and Hsp90 in the assembly of the replicase complex of a positive-strand RNA plant virus. *J. Virol.* 86:12091-12104.
- Miranda, G. J., Aliyar, R., and Shirako, Y. (2001). Nucleotide sequence of a Dianthovirus RNA1-like RNA found in grassy stunt-diseased rice plants. *Arch. Virol.* 146:225–238.
- Mizumoto, H., Hikichi, Y., and Okuno, T. (2002). The 3'-untranslated region of RNA1 as a primary determinant of temperature sensitivity of *Red clover necrotic mosaic virus* Canadian strain. *Virology* 293:320–327.
- Mizumoto, H., Iwakawa, H. O., Kaido, M., Mise, K., and Okuno, T. (2006). Cap-independent translation mechanism of *Red clover necrotic mosaic virus* RNA2 differs from that of RNA1 and is linked to RNA replication. *J. Virol.* 80:3781–3791.
- Mizumoto, H., Tatsuta, M., Kaido, M., Mise, K., and Okuno, T. (2003). Cap-independent translational enhancement by the 3' untranslated region of *Red clover necrotic mosaic virus* RNA1. *J. Virol.* 77:12113–12121.
- Moon, S. L., Anderson, J. R., Kumagai, Y., Wilusz, C. J., Akira, S., Khromykh, A. A., and Wilusz, J. A noncoding RNA produced by arthropod-borne flaviviruses inhibits the cellular exoribonuclease XRN1 and alters host mRNA stability. (2012). *RNA* 18:2029-2040.
- Morales, F., Castaño, M., Calvert, L., and Arroyave, J. A. (1992). *Furcraea necrotic streak virus*: an apparent new member of the dianthovirus group. *J. Phytopathol.* 134: 247–254.
- Nagy, P. D., Barajas, D., & Pogany, J. (2012). Host factors with regulatory roles in tombusvirus replication. *Curr. Opin. Virol.* 2, 691-698.
- Neeleman, L., and Bol, J. F. (1999). *Cis*-acting functions of alfalfa mosaic virus proteins involved in replication and encapsidation of viral RNA. *Virology* 254:324-333.

- Musil, M., and Matisová, J. (1967). Contribution to the knowledge of mosaic viruses of red clover in Slovakia. *Ochr. Rost.* 3:225-234.
- Musil, M. (1969). Red clover necrotic mosaic virus, a new virus infecting red clover (*Trifolium pretense*) in Czechoslovakia. *Biologia, Bratisl.*24:33-45.
- Nagy, P. D., and Pogany, J. (2012). The dependence of viral RNA replication on co-opted host factors. *Nat. Rev. Microbiol.* 10:137-149.
- Novak, J. E., and Kirkegaard, K., (1994). Coupling between genome translation and replication in an RNA virus. *Genes & Dev.* 8:1726-1737.
- Nicholson, B. L., Wu, B., Chevtchenko, I., and White, K. A. (2010). Tombusvirus recruitment of host translational machinery via the 3' UTR. *RNA* 16:1402–1419.
- Nicholson, B. L., and White, K. A. (2008). Context-influenced cap-independent translation of Tombusvirus mRNAs in vitro. *Virology* 380:203–212.
- Nicholson, B. L., and White, K. A. (2011). 3' Cap-independent translation enhancers of positive-strand RNA plant viruses. *Curr. Opin. Virol.* 1:373-380.
- Okamoto, K., Nagano, H., Iwakawa, H., Mizumoto, H., Takeda, A., Kaido, M., Mise, K., and Okuno, T. (2008). cis-Preferential requirement of a -1 frameshift product p88 for the replication of *Red clover necrotic mosaic virus* RNA1. *Virology* 375:205–212.
- Okuno, T., Hiruki, C., Rao, D. V., and Figueiredo, G. C. (1983). Genetic determinants distributed in two genomic RNAs of sweet clover necrotic mosaic, red clover necrotic mosaic and clover primary leaf necrosis viruses. *J. Gen. Virol.* 64:1907–1914.
- Okuno, T. (2012). Replication mechanisms of plant RNA viruses: current understanding and perspectives. *J. Gen. Plant Pathol.* 78:404–408
- Osman, T. A., and Buck, K. W. (1991). Detection of the movement protein of red clover necrotic mosaic virus in a cell wall fraction from infected *Nicotiana clevelandii* plants. *J. Gen. Virol.* 72:2853–2856.
- Osman, T. A., Hayes, R. J., and Buck, K. W. (1992). Cooperative binding of the red clover necrotic mosaic virus movement protein to single-stranded nucleic acids. *J. Gen. Virol.* 73:223–227.
- Osman, T. A., Thömmes, P., and Buck, K. W. (1993). Localization of a single-stranded RNA-binding domain in the movement protein of red clover necrotic mosaic dianthovirus. *J. Gen. Virol* 74:2453.
- Osman, T. A., and Buck, K.W. (1987). Replication of red clover necrotic mosaic virus RNA in cowpea protoplasts: RNA 1 replicates independently of RNA 2. *J. Gen.*

*Virology* 68:289–296.

- Oster, S. K., Wu, B., and White, K. A. (1998). Uncoupled expression of p33 and p92 permits amplification of Tomato bushy stunt virus RNAs. *J. Virol.* 72:5845-5851.
- Paje-Manalo, L. L., and Lommel, S. A. (1989). Independent replication of red clover necrotic mosaic virus RNA-1 in electroporated host and nonhost *Nicotiana* species protoplasts. *Phytopathology* 79:457–461.
- Panavas, T., Serviene, E., Brasher, J., and Nagy, P. D. (2005). Yeast genome-wide screen reveals dissimilar sets of host genes affecting replication of RNA viruses. *Proc. Natl. Acad. Sci. USA.* 102:7326–7331.
- Pappu, H. R., and Hiruki, C. (1988). Replication of sweet clover necrotic mosaic virus in cowpea protoplasts. *Can. J. Plant Pathol.* 10:110–115.
- Pappu, H. R., and Hiruki, C. (1989). Electrophoretic variability among dianthoviruses. *Phytopathology* 79:1253-1257.
- Pappu, H. R., Hiruki, C., and Inouye, N. (1988). A new serotype of sweet clover necrotic mosaic virus. *Phytopathology* 78:1343-1348.
- Park, S-H., Sit, T. L., Kim, K-H., and Lommel, S. A. (2012). The *Red clover necrotic mosaic virus* capsid protein N-terminal lysine-rich motif is a determinant of symptomatology and virion accumulation, *Mol. Plant Pathol.* 13:744-753.
- Peltier, C., Klein, E., Hleibieh, K., D'Alonzo, M., Hammann, P., Bouzoubaa, S., Ratti, C., and Gilmer, D. (2012). Beet necrotic yellow vein virus subgenomic RNA3 is a cleavage product leading to stable non-coding RNA required for long-distance movement. *J. Gen. Virol.* 93:1093-1102.
- Pijlman, G. P., Funk, A., Kondratieva, N., Leung, J., Torres, S., van der Aa, L., Liu, W. J., Palmenberg, A. C., Shi, P. Y., Hall, R. A., and Khromykh, A. A. (2008). A highly structured, nuclease-resistant, noncoding RNA produced by flaviviruses is required for pathogenicity. *Cell Host Microbe* 4:579–591.
- Powers, J. G., Sit, T. L., Heinsohn, C., George, C. G., Kim, K. H., and Lommel, S. A. (2008a). The *Red clover necrotic mosaic virus* RNA-2 encoded movement protein is a second suppressor of RNA silencing. *Virology* 381:277–286.
- Powers, J. G., Sit, T. L., Qu, F., Morris, T. J., Kim, K. H., and Lommel, S. A. (2008b). A versatile assay for the identification of RNA silencing suppressors based on complementation of viral movement. *Mol. Plant Microbe Interact.* 21:879–890.
- Qu, F., and Morris, T. J. (2000). Cap-independent translational enhancement of turnip crinkle virus genomic and subgenomic RNAs. *J. Virol.* 74:1085–1093.
- Qu, F., Ren, T., and Morris, T. J. (2003). The coat protein of turnip crinkle virus

- suppresses posttranscriptional gene silencing at an early initiation step. *J. Virol.* 77:511-522.
- Qu, F., Ye, X., and Morris, T. J. (2008). *Arabidopsis* DRB4, AGO1, AGO7, and RDR6 participate in a DCL4-initiated antiviral RNA silencing pathway negatively regulated by DCL1. *Proc. Natl. Acad. Sci. USA.* 105:14732–14737.
- Rao, A. L. N., Cooper, B., & Deom, C. M. (1998). Defective movement of viruses in the family *Bromoviridae* is differentially complemented in *Nicotiana benthamiana* expressing tobamovirus or dianthovirus movement proteins. *Phytopathology*, 88, 666-672.
- Rao, A. L. N., and Hiruki, C. (1987). Unilateral compatibility of genome segments from two distinct strains of red clover necrotic mosaic virus. *J. Gen. Virol.* 36:191-194.
- Ryabov, E. V., Generozov, E. V., Kendall, T. L., Lommel, S. A., and Zavriev, S. K. (1994). Nucleotide sequence of carnation ringspot dianthovirus RNA-1. *J. Gen. Virol.* 75:243–247.
- Salonen, A., Ahola, T., Kaariainen, L. (2005). Viral RNA replication in association with cellular membranes. *Curr. Top. Microbiol. Immunol.* 285:139–173.
- Sarawaneeyaruk, S., Iwakawa, H-O., Mizumoto, H., Murakami, H., Kaido, M., Mise, K., and Okuno, T. (2009). Host-dependent roles of the viral 5' untranslated region (UTR) in RNA stabilization and cap-independent translational enhancement mediated by the 3' UTR of *Red clover necrotic mosaic virus* RNA1. *Virology* 391: 107–118.
- Serviene, E., Shapka, N., Cheng, C. P., Panavas, T., Phuangrat, B., Baker, J., and Nagy, P. D. (2005). Genome-wide screen identifies host genes affecting viral RNA recombination. *Proc. Natl. Acad. Sci. U. S. A.* 102:10545–10550.
- Schaad, M. C., Haldeman-Cahill, R., Cronin, S., and Carrington, J. C. (1996). Analysis of the VPg-proteinase, NIa, encoded by tobacco etch potyvirus: effects of mutations on subcellular transport, proteolytic processing, and genome amplification. *J. Virol.* 70:7039-7048.
- Shen, R., and Miller, W. A. (2004). The 3' untranslated region of tobacco necrosis virus RNA contains a barley yellow dwarf virus-like cap-independent translation element. *J. Virol.* 78:4655–4664.
- Sherman, M. B., Guenther, R. H., Tama, F., Sit, T. L., Brooks, C. L., Mikhailov, A. M., Orlova, E. V., Baker, T. S., and Lommel, S. A. (2006). Removal of divalent cations induces structural transitions in *Red clover necrotic mosaic virus*, revealing a potential mechanism for RNA release. *J. Virol.* 80:10395–10406.

- Sit, T. L., Haikal, P. R., Callaway, A. S., and Lommel, S. A. (2001). A single amino acid mutation in the carnation ringspot virus capsid protein allows virion formation but prevents systemic infection. *J. Virol.* 75:9538–9542.
- Sit, T. L., Vaewhongs, A. A., and Lommel, S. A. (1998). RNA-mediated *trans*-activation of transcription from a viral RNA. *Science* 281:829–832.
- Solovyev, A. G., Zelenina, D. A., Savenkov, E. I., Grdzlishvili, V. Z., Morozov, SYu, Maiss, E., Casper, R., and Atabekov, J. G. Host-controlled cell-to-cell movement of a hybrid barley stripe mosaic virus expressing a dianthovirus movement protein. *Intervirology* (1997). 40:1-6.
- Stefano, G., Renn, a L., Chatre, L., Hanton, S. L., Moreau, P., Hawes, C., and Brandizzi, F. (2006). In tobacco leaf epidermal cells, the integrity of protein export from the endoplasmic reticulum and of ER export sites depends on active COPI machinery. *Plant J.* 46:95-110.
- Stupina, V. A., Meskauskas, A., McCormack, J. C ., Yingling, Y. G., Kasprzak, W. , Shapiro, B. A., Dinman, J. D., and Simon, A. E. (2008). The 3' proximal translational enhancer of Turnip crinkle virus binds to 60S ribosomal subunits. *RNA* 14:2379–2393.
- Tajima, Y., Iwakawa, H. O., Kaido, M., Mise, K., and Okuno, T. (2011). A long-distance RNA-RNA interaction plays an important role in programmed -1 ribosomal frameshifting in the translation of p88 replicase protein of Red clover necrotic mosaic virus. *Virology* 417:169-178.
- Takeda, A., Tsukuda, M., Mizumoto, H., Okamoto, K., Kaido, M., Mise, K., and Okuno, T. (2005). A plant RNA virus suppresses RNA silencing through viral RNA replication. *EMBO J.* 24:3147–3157.
- Tatsuta, M., Mizumoto, H., Kaido, M., Mise, K., and Okuno, T. (2005). The *Red clover necrotic mosaic virus* RNA2 *trans*-activator is also a cis-acting RNA2 replication element. *J. Virol.* 79:978–986.
- Tomlinson, J. A., Faithfull, E. M., and Seeley, N. D. (1982a). Plant viruses in rivers. *Rothamsted Experimental Station Report for 1981*, 86-87.
- Tomlinson, J. A., Faithfull, E., Flewett, T. H., and Beards, G. (1982b). Isolation of infective tomato bushy stunt virus after passage through the human alimentary tract. *Nature* 300:637-638.
- Tomlinson, J. A., Faithfull, S. M., and Fraser, R. S. S. (1983). Plant viruses in river water. 33<sup>rd</sup> Annual Report for 1982. *National Vegetable Research Station Wellesbourne, Warwick, K*, 81-82.
- Treder, K., Kneller, E. L., Allen, E. M., Wang, Z., Browning, K. S., and Miller, W. A.



- (2008). The 3' cap-independent translation element of *Barley yellow dwarf virus* binds eIF4F via the eIF4G subunit to initiate translation. *RNA* 14:134–147.
- Tremaine, J. H., and Dodds, J. A. (1985). Carnation ringspot virus. *AAB Descriptions of Plant Viruses. Set 19*.
- Tremblay, D., Vaewhongs, A. A., Turner, K. A., Sit, T. L., and Lommel, S. A. (2005). Cell wall localization of *Red clover necrotic mosaic virus* movement protein is required for cell-to-cell movement. *Virology* 333:10–21.
- Turner, R. L., and Buck, K. W. (1999). Mutational analysis of *cis*-acting sequences in the 3'- and 5'-untranslated regions of RNA2 of *Red clover necrotic mosaic virus*. *Virology* 253:115–124.
- Turner, K. A., Sit, T. L., Callaway, A. S., Allen, N. S., and Lommel, S. A. (2004). Red clover necrotic mosaic virus replication proteins accumulate at the endoplasmic reticulum. *Virology* 320:276–290.
- Vaewhongs, A. A., and Lommel, S. A. (1995). Virion formation is required for the long-distance movement of red clover necrotic mosaic virus in movement protein transgenic plants. *Virology* 212:607–613.
- Van Bokhoven, H., Le Gall, O., Kasteel, D., Verver, J., Wellink, J., and Van Kammen, A. B. (1993). *Cis*- and *trans*-acting elements in cowpea mosaic virus RNA replication. *Virology* 195:377-386.
- van Rossum, C. M., Garcia, M. L. & Bol, J. F., 1996. Accumulation of alfalfa mosaic virus RNAs 1 and 2 requires the encoded proteins in cis. *J. Virol.* 70:5100-5105.
- Wang, S., Browning, K. S., and Miller, W. A. (1997). A viral sequence in the 3'-untranslated region mimics a 5' cap in facilitating translation of uncapped mRNA. *EMBO J.* 16:4107–4116.
- Wang, Z., Kraft, J. J., Hui, A. Y., and Miller, W. A. (2010). Structural plasticity of Barley yellow dwarf virus-like cap-independent translation elements in four genera of plant viral RNAs. *Virology* 402:177–186.
- Wang, Z., Treder, K., and Miller, W. A. (2009). Structure of a viral cap-independent translation element that functions via high affinity binding to the eIF4E subunit of eIF4F. *J. Biol. Chem.* 284:14189–14202.
- Wang, H-L., Wang, Y. Giesman-Cookmeyer, D., Lommel, SA., and Lucas, W. A. (1998). Mutations in viral movement protein alter systemic infection and identify an intercellular barrier to entry into the phloem long-distance transport system. *Virology* 245:75-89.
- Weiland, J. J., and Dreher, T. W. (1993). *Cis*-preferential replication of the turnip

- yellow mosaic virus RNA genome. *Proc. Natl. Acad. Sci. USA*. 90:6095-6099.
- Weng, Z., and Xiong, Z. (2009). Three discontinuous loop nucleotides in the 3' terminal stem-loop are required for *Red clover necrotic mosaic virus* RNA-2 replication. *Virology* 393:346–354.
- White, K. A. (2002). The premature termination model: a possible third mechanism for subgenomic mRNA transcription in (+)-strand RNA viruses. *Virology* 304:47–154.
- White, K.A., Bancroft, J.B., and Mackie, G.A. (1992). Coding capacity determinants in vivo accumulation of a defective RNA of clover yellow mosaic virus. *J. Virol.* 66, 3069–3076.
- White, K. A., Skuzeski, J. M., Li, W., Wei, N., and Morris, T.J. (1995). Immunodetection, expression strategy and complementation of turnip crinkle virus p28 and p88 replication components. *Virology* 211:525-534.
- Xiong, Z., Kim, K. H., Giesman-Cookmeyer, D., and Lommel, S. A. (1993a). The roles of the red clover necrotic mosaic virus capsid and cell-to-cell movement proteins in systemic infection. *Virology* 192:27–32.
- Xiong, Z., Kim, K. H., Kendall, T. L., and Lommel, S. A. (1993b). Synthesis of the putative red clover necrotic mosaic virus RNA polymerase by ribosomal frameshifting *in vitro*. *Virology* 193:213–221.
- Xiong, Z., and Lommel, S. A. (1989). The complete nucleotide sequence and genome organization of red clover necrotic mosaic virus RNA-1. *Virology* 171:543–554.
- Xiong, Z. G., and Lommel, S. A. (1991). Red clover necrotic mosaic virus infectious transcripts synthesized in vitro. *Virology* 182:388–392.
- Xu, W., and White, K. A. (2009). RNA-based regulation of transcription and translation of aureusvirus subgenomic mRNA1. *J. Virol.* 83:10096–10105.
- Yi, G., and Kao, C. C. (2008). *cis*- and *trans*-Acting functions of brome mosaic virus protein 1a in genomic RNA1 replication. *J. Virol.* 82:3045-3053.
- Zavriev, S. K., Hickey, C. M., and Lommel, S. A. (1996). Mapping of the red clover necrotic mosaic virus subgenomic RNA. *Virology* 216:407–410.