RIMS Kôkyûroku Bessatsu B15 Mathematical analysis on the self-organization and self-similarity

edited by Takayoshi Ogawa and the editorial board

December, 2009 Research Institute for Mathematical Sciences Kyoto University RIMS Kôkyûroku Bessatsu B15

Mathematical analysis

on the self-organization and self-similarity

September 16~18, 2008

edited by Tatsuyuki Nakaki, Takasi Senba, Takayoshi Ogawa(chief), Mishio Kawashita and Masaki Kurokiba

December, 2009

Research Institute for Mathematical Sciences Kyoto University, Kyoto, Japan

The papers presented in this volume of RIMS Kôkyûroku Bessatsu are in final form and refereed. ©2009 by the Research Institute for Mathematical Sciences, Kyoto University. All rights reserved. Printed in Japan.

Preface

The international conference "Mathematical analysis on the self-organization and self-similarity" was held from 16th to 18th, September 2008 at Kyoto University, Kyoto, Japan as the project conference of the Research Institute of Mathematical Science (RIMS), Kyoto University.

The mathematical analysis on the model of the chemotaxis has been largely developed particularly during the last 15 years. One of the pioneer work on the singularity formation was done by Professor Toshitaka Nagai of Hiroshima University and after that there are many interesting works were continuously done around this area.

On the other hand, the mathematical analysis on the semiconductor device simulation has longer history and it is now well recognized that the both theory are closely related involving the mathematical analysis of the fluid mechanics.

The organizing committee of the conference (all the editorial board with Tatsuyuki Nakaki as the chair) considered that it is a good opportunity to organize an international conference in the occasion of Professor Nagai's sixtieth birthday and the conference was planned to make a bridge between the chemotaxis model and the semiconductor device simulations. The topics were selected within the mathematical research on the system of nonlinear parabolic equation mainly concentrated on the chemotaxis model and the semiconductor device simulations as well as their natural origin of the fluid mechanics. Ten invited speakers including overseas researchers gave splendid lectures on their latest research and many stimulated discussion around the talks were done. The conference had more than 80 participants.

This volume of the RIMS Kôkyûroku Bessatsu is devoted as the proceedings of the conference and the main contributors are the original plenary speakers and attendant researchers.

We hope that this volume is presenting the latest research that bring further development on the mathematical analysis of the non-local type nonlinear partial differential equations.

Takayoshi Ogawa (Editor in chief)

Editorial Board:

TatsuyukiNakaki(Hiroshima University, chair)TakasiSenba(Kyushu Institute of Technology)TakayoshiOgawa(Tohoku University)MishioKawashita(Hiroshima University)MasakiKurokiba(Fukuoka University)

Contents

Shadow system for adsorbate-induced phase transition model Kousuke Kuto and Tohru Tsujikawa1
Higher-order asymptotic expansions of solutions to a parabolic system of chemotaxis in \mathbb{R}^n
Tetsuya Yamada15
Nonlocal elliptic boundary value problems related to chemotactic movement of mobile species
Dirk Horstmann and Marcello Lucia
A nondegeneracy result for least energy solutions to a biharmonic problem with nearly critical exponent
Tomohiko Sato and Futoshi Takahashi73
Self-similar blow-up for a chemotaxis system in higher dimensional domains Yuki Naito and Takasi Senba
Global solvability for a chemotaxis system in \mathbb{R}^2 Toshitaka Nagai101
The strong maximum principle
Julián López-Gómez
Conservative numerical schemes for the Keller-Segel system and numerical results $\$
Norikazu Saito
Lipschitz semigroup approach to drift-diffusion systems Toshitaka Matsumoto and Naoki Tanaka
Stationary solutions of a hydrodynamic model for semiconductors Shinya Nishibata and Masahiro Suzuki
Asymptotic expansion of solution to the Nernst-Planck drift-diffusion equation Masakazu Yamamoto