

Talun-Huma, Swidden Agriculture, and Rural Economy in West Java, Indonesia

Mizuno Kosuke,* Siti Sugiah Mugniesyah,** Ageng Setiawan Herianto,*** and Tsujii Hiroshi†

Talun-huma is an intensified land use system for swidden agriculture in Indonesia. *Talun* is a productive fallow system that is meaningful from ecological, social, and economic perspectives. Although it is considered a typical practice in West Java, this study proves that the term *talun* was used in many places in Indonesia besides West Java during colonial times.

This study also shows that *talun-huma* and agroforestry practice in the surveyed village is closely linked to the socioeconomic structure of the rural society based on the data collected in 2000–1. *Huma* is more likely to be practiced by the lower strata owning little agricultural and forestry land. The economic development reflected in the growth of banana leaf production and income from the non-agricultural sector income has not excluded or diminished *talun-huma* and agroforestry practices. Sharecropping practices and agricultural labor relations among the villagers have established *huma* practice. The social forestry program implemented in this region also supports the continuity of *huma* practice. Increase in banana leaf production does not diminish the practice of *huma* and *talun* agroforestry, although banana leaf production tends to shorten the duration of the cycle. Moreover, the development of non-agricultural sectors and wet rice cultivation has had a positive impact on the existence of *huma* practices and the continuation of slash-and-burn practices.

Talun-huma and permanent forests with or without *talun* are good practices that keep the system sustainable from an economic and social point of view. Diversified farming, balanced rotation of land use, and diversity of plants, as well as the planting of leguminous land conservation trees such as *kaliandra* and *gamal*, play an important role in sustaining the system.

The introduction of banana leaf plants to the village in the 1990s has contributed significantly to the continuity of the system because the plants are a considerable source of income for villagers, both as farmers and as agricultural laborers. They can rely on the income from the hilly dry land. A problem that might, however, become serious in the near future is that banana leaf planting shortens the duration of the cycle and may render the land infertile.

Keywords: *talun-huma*, swidden agriculture, agroforestry, rural economy, Indonesia, West Java, sustainability, *talun*

* 水野広祐, Center for Southeast Asian Studies, Kyoto University
Corresponding author's email: mizuno@cseas.kyoto-u.ac.jp

** Department of Communication and Community Development Sciences, Faculty of Human Ecology, Bogor Agricultural University, Jl. Kamper, Kampus IPB Darmaga Bogor, Indonesia

*** Department of Agricultural Socio-Economics, Faculty of Agriculture, Gadjah Mada University, Jl. Flora, Bulaksumur, Yogyakarta 55281, Indonesia

† 辻井 博, Professor Emeritus, Kyoto University, 104-1 Higashianshin-cho, Okamedani, Fukakusa, Fushimi-ku, Kyoto 612-0032, Japan

Introduction

The importance of the *talun* system as an example of agroforestry was introduced to international academia by Otto Soemarwoto in the 1980s. He used the term *talun-kebun*, which he explained as a kind of shifting cultivation practiced in a man-made forest. It combined many species of perennials and annuals in multi-layered and single-layered arrangements, forming an often-dense canopy of vegetation that protects against soil erosion and leaching. Structurally, the *talun-kebun* is divided into two parts: the *talun*, or selected productive fallow “forest,” consisting of the overhead cover of essentially long-term perennials, and the *kebun*, comprising various areas of cleared ground within the *talun* planted with annual crops such as vegetables, fruits, and cereal crops, rather than dry rice, spices and so on, destined mainly for market sale. Upon harvest, the *kebun* is allowed to grow perennials and returned to the *talun* within five to eight years.

The *talun* is planted with a mixture of many species of trees but may be dominated by one species, typically bamboo, in which case it is named after this species, so *talun awi* is bamboo *talun*. The *talun* has four important functions: (i) subsistence production, (ii) commercial production, (iii) gene banking, and (iv) soil conservation and sustained productivity (Otto Soemarwoto *et al.* 1985, 49–50).

This *talun-kebun* is clearly different from conventional notions of swidden agriculture. For example, according to Sasaki, in the swidden agriculture of Southeast Asia, a plot is cultivated for less than three years, five at most, then left fallow. This agriculture can supply food for a population density of around 25–30 persons/km². The duration of the period in which the land is left fallow is 8–15 years before people slash-and-burn the plot again (Sasaki 1970, 86–122).

Often, however, shifting cultivation practices tend to shorten the fallow period, resulting in a degraded system with time or as the population density increases. People are then likely to leave the degraded land. Alternatively, land rehabilitation takes place when farmers invest in improved land management and care for the environment—provided they have reasonably secure land or tree tenure and if it is profitable compared with other investment options. Alternative land use intensification pathways that do not first involve severe land degradation do exist in the form of complex agroforestries that have been developed by indigenous communities (Sanchez *et al.* 2005, 6–7). Some such cases have been studied, including cacao planting in swidden agriculture land (Duguma *et al.* 2001) and swidden fallow agroforestry among the Bora Indians of Amazon, which is adaptable to varying environmental and economic situations (Padoch and de Jong 1987), but such studies still remain small in number.

The *talun* system illustrates a case of alternative land use intensification for shifting

cultivation. Otto Soemarwoto distinguished between two types of *talun*—permanent *talun* and *talun-kebun*. In the permanent *talun*, trees are typically densely spaced and the canopies are closed. Hence, little light penetrates the canopies and only a few shade-tolerant species, such as the taro-like *Xanthosoma*, are planted. These crop species and weeds form the undergrowth of the *talun*. In other *talun*, the trees are sparsely planted so that more light can reach the floor. In such cases, many annuals are grown (e.g., corn, cassava, and sweet potato). Many weeds are also found in the *talun*. Such *talun* are usually called *kebun campuran*, which means “mixed garden,” denoting a mixture of annuals and perennials. In the permanent *talun*, the practice of slash-and-burn is discontinued.

In the *talun-kebun*, a clearing is deliberately made in the *talun* at the beginning of the rainy season, either by clear cutting (e.g., bamboo), or by selective cutting (e.g., *Jeungjing*, *Albizia chinensis*) and heavy pruning of the remaining perennial trees. The trunks and large branches are taken out of the clearing and sold as construction materials and fuelwood, and twigs and leaves are spread out to dry in the sun, then piled up and burned. The ash is collected and mixed with cattle dung brought in from the villages.

A mixture of annual crops is grown in the clearing, which is then called *kebun*, literally “garden.” The major crops grown are beans at Ciwidey and Soreang in Southwest Bandung, and tobacco and onion at Paseh in Southeast Bandung in West Java. The planting of the different crops is not done all at once but in succession. Harvesting is also carried out over an extended period. By the time the last crops are harvested, which occurs about 18 months after the clearing, the trees would have resprouted. People then clear another area and repeat the process. Consequently, the *kebun* moves around inside the *talun* with a cycle of about six to eight years. Essentially it is a form of shifting cultivation (Otto Soemarwoto and Idjah Soemarwoto 1984, 274–276).

We understand that the land of shifting cultivation, here called *kebun*, is mainly planted with annuals after slash-and-burn, and then planted with perennials after the harvest of annuals, at which stage it is sometimes called *kebun campuran*, after which it enters the stage of *talun*, a productive fallow stage. Another land use after slash-and-burn and subsequent planting of annuals and perennials is to discontinue the slash-and-burn process and use the land as a permanent *talun*, or *kebun campuran* with no more slash-and-burn.

Talun and *talun kebun* have been studied substantially, especially from the ecological perspective, and intensively in the case of bamboo *talun* in West Java. Species diversity has been a typical characteristic. Johan Iskandar *et al.*'s study showed that 112 species were planted by the local people at a village in Soreang sub-district in Bandung district (Johan Iskandar *et al.* 1981, quoted by Herri Y. Hadikusumah 2005, 269–283).

Herri Y. Hadikusumah (2005, 256–259) demonstrated the variety of plants in the layers that form the *talun*. Linda Christanty *et al.* (1996a) focused on biomass accumulation, and Linda Christanty *et al.* (1996b) and Mailyly *et al.* (1996) discussed the biochemical role of the bamboo. Parikesit *et al.* (2004) discussed *kebon tatangkalan*, a form of permanent *talun*, as a habitat for various organisms living there, including birds and insects, and the decreasing area of such habitat because of growing demand for agricultural, industrial, and settlement lands.

The present study attempts to show a different *talun* system in detail—the *talun-huma*, which is associated with dry rice planting as annuals after slash-and-burn, rather than vegetables as the main planting, and mixed perennials rather than bamboo at the productive fallow stage. The study aims to enrich understanding of the *talun* agroforestry system as an alternative land use intensification. It will shed light on the economic contribution that is essential for the household economy and for the sustainability of the system from a quantitative perspective (such as amount of income), as well as qualitative viewpoint (such as stratification of rural society and the land tenure system). This study also intends to describe the directions taken by land use intensification, that is, with or without slash-and-burn, and with or without *talun*, and to analyze these directions. In this way, the impact of economic development on the *talun-huma* and agroforestry system will be studied from the perspectives of growth of the agricultural and the non-agricultural sectors. The *talun-huma* and agroforestry here means *talun-huma*, related practices with or without *talun (huma)*, and permanent forest with or without *talun*. These descriptions and analyses will be based on data and information collected by fieldwork at a village in Cianjur district (*Kabupaten* Cianjur), West Java. This study also examines the validity of the concept of *talun* outside West Java as a preliminary study.

Section I discusses previous studies of the *talun* system and, using a variety of sources, areas in which the system prevails. We proceed to describe the location where field study was conducted in the second section and elaborate upon the *talun-huma* and agroforestry system in the third. The direction of land use intensification, as well as the roles of the *talun-huma* and agroforestry system, especially in terms of economic contributions, is examined in Section IV, and Section V concludes this study.

The field survey was conducted over 10 years between 1998 and 2007. In total, 60 households were surveyed, using household survey questionnaires. The data used in this paper is based mainly on the study in 2000 and 2001.¹⁾

1) The study team consisted of staff from Bogor Agricultural University, Kyoto University, the University of Tokyo, and Gadjah Mada University. This paper has been presented at the 2nd Seminar on “Toward Harmonization between Development and Environmental Conservation in Biological Production” (JSPS-DGHE Core University Program in Applied Biosciences, the University of ↗

I Talun and Agroforestry Systems in Indonesia

I-1 Talun in Indonesia

According to Otto Soemarwoto (1983, 222), land use such as the home garden (*pekarangan*)—but with no houses on the land—is found in West Java and other areas with some variations. This land use is called *talun* in West Java. He also stated that the *talun-kebun* is typical of West Java, especially in the Priangan region (Otto Soemarwoto *et al.* 1985, 48). Such a view is shared by many researchers (Herri Y. Hadikusumah 2005, 269; Johan Iskandar 2009, 150), including Terra’s study of 1953.

Terra mentioned the *talun* in the context of a mixed garden study, explaining the *kebun* besides the home garden (*pekarangan*):

The *kebun* is situated near or around the villages, sometimes at some distance. The *kebun* is sometimes planted with fruit trees, or coconut palms, sugar palms, banana, tea, coffee, or bamboo only. They are distinguished from plantings, mostly in clumps, of fruits trees etc. on former *ladang* (fields in use for shifting cultivation). Instead of *kebun* (Malay), in Java the term used is often *talun* (Sunda) and outside of Java *dusun* (Ambon, Ceram), *mamar* (Timor), *porlak* (Batak), *peureuh* (Achin), *krakal* (Purworedjo, Java). (Terra 1953, 163–164)

Data on customary law during the colonial time in Indonesia show, however, that the term *talun* was not limited to West Java. Research carried out on customary law in North Sumatra (*Adatrechtbundel* VI 1911, 124) indicate that *taloen* meant a cleared plot where people grew sugar palm for sapping, sirih, and other crops in the Gajo, Alas, and Batak areas. The land was wasteland, but trails left by labor for cultivation and the like show that people’s rights had arisen. In Minahasa, *talun* means forest, and *mengatalun* means to become forest, or to revert to forest where people do not cultivate regularly (*Adatrechtbundel* IX 1914, 132). The Malay people living in Sumatra use the term *beloekar taloen* to signify young forestland where trails of former exploitation are still apparent, and *beloekar toea-taloen tabang* for old forests that used to be exploited and

↙ Tokyo, February 2003), and at the International Workshop on “Sustainable Agricultural Development in Southeast Asia” (Research Center for Regional Resources and the Indonesian Institute of Science, Jakarta, September 14–15, 2003). The seminar and workshop issued proceedings as the 2nd Seminar on “Toward Harmonization between Development and Environmental Conservation in Biological Production” (JSPS-DGHE Core University Program in Applied Biosciences, the University of Tokyo, February 2003). The paper has also been presented at the Kyoto Sustainability Institute (KSI) International Workshop on “Swidden Agriculture in Southeast Asia, Sustainability and Contribution towards Agroforestry and Forestry,” held at the Center for Southeast Asian Studies, Kyoto University on March 15, 2010. The authors express sincere thanks to these projects and to the participants of the seminar and workshop who made comments on this paper. The authors are also grateful to the paper’s referees who gave us valuable comments.

where trails of former exploitation are still more or less apparent (*Adatrechtbundel* X 1915, 221). In East Java, *talon*, *talun*, or *talunan* means abandoned garden, abandoned cultivated land, or unirrigated cultivated land in the mountainous area. These are planted with plants other than rice and are not located at the center of the village (*Adatrechtbundels* XIV 1917, 230). In West Java, *taloen* is newly cleaned *tegal* (cultivated dry field) (*Adatrechtbundel* VIII 1914, 198), while in Borneo, Tidoengsch, and Tinggalan, the term means forest or light forest (*Adatrechtbundels* XIII 1917, 55). In Central Java, cleared land where dry rice is planted because water cannot be brought in and which is abandoned after one or two plantings is called *taloen* or *pengalang-alangan*. *Pengalang-alangan* means land where *alang-alang* (*Imperata cylindrica*) is allowed to spread on purpose, with the weeds belonging to the exploiter, and *taloen* means land formerly planted with dry rice and now covered by *alang-alang* and other weeds. Anyone may remove the weeds and plant dry rice again (*Adatrechtbundels* XIV 1917, 40).²⁾

As we can see, the term *talun* is used in many areas in Indonesia such as West, Central and East Java, North Sumatra, Minahasa, Malay people's areas in Sumatra, and Borneo. In general, *talun* is non-irrigated land located in a hilly or mountainous area. It signifies forest that was once cultivated then abandoned. In many cases, it is thought of as fallowed area. In Central Java, the *taloen* belongs to the person who first cleared the land and who would cultivate the land again.

The term *talun* even became the name of villages, mountains, and sub-districts. A dictionary of geography published in 1869 mentioned the name of Taloen in six villages (*dorps*) in Central and East Java.³⁾ Taloen was also the name of a mountain (*berg*) in the residency (*residentie*) Rembang, division (*afdeeling*) Toeban (Veth 1869, 862). A directory

2) Dictionaries of local languages contain differing definitions of *talun*. In Sundanese during colonial times, *talun* meant newly developed dry land (*tegal*), and *doeloeh taloen* referred to new settlements consisting of farmland, houses, and gardens, arising from an extension of the village (Coolsma 1913, 611), while in present-day Sundanese spoken mainly in West Java, it refers to agricultural dry land with fruit trees that live for a long time (Lembaga Basa & Sastra Sunda 1975, 503). On the other hand, a Javanese dictionary written in colonial times defined *talun* as land already harvested and not made orderly yet (Jansz 1906, 1027). A Javanese dictionary written after Independence defined *talun* as land of shifting cultivation or dry rice land (Prawiroatmodjo 1957, 628). According to a colonial-era dictionary of Madurese, *talon* is a piece of land planted with something other than rice, a garden, hilly land, a fallowed garden, a fallowed agricultural field, not irrigated (Peninga and Hendriks 1936, 311). A dictionary of Sasak (the language in Lombok) mentioned *taloen* as cultivated land (Goris 1938, 296), while *taloen* appears in a dictionary of the Boesang language (spoken by a Dayak group in Borneo) as brushwood, or young forest that has undergone swidden agriculture (Barth 1910, 204).

3) These villages were located at *afdeeling* Toeban, Salatiga, and Pati, and also regency (*regentschap*) Temanggung, *residentie* Kediri, and *regentschap* Trenggalek. Besides these, the name of Taloenombo was found at *residentie* Banjoemas (Veth 1869, 862).

of local administrative bodies published in 1931 mentioned the name of Taloen in two sub-districts (*onderdistrict*) in Central and East Java, as well as 10 villages (*desa*) in Java island. Of these, five were found in East Java, two in Central Java, two in West Java, and one in the *gewest* (province) of Soerakarta (Schoel 1931, 373).⁴ A dictionary of geography published in 1917 cited 10 locations named Taloen in Java island, and 1 location in Borneo as Taloenliao (Dumont 1917, 565–566).⁵

Thus we can say that the term *talun*, *taloen*, or *talon* has been used since at least colonial times in many parts of Indonesia with relation to forest, or forest once cultivated then fallowed or abandoned. The term *taloen* was also used as the name of villages, sub-districts, or mountains in many places in Java island, especially in Central and East Java, and partly in Borneo. Why is *talun* thought to be characteristic of West Java? One possible answer is that other areas now have different terms for a similar system of agroforestry.

I-2 Agroforestry in Indonesia

Adatrechtbundels show different words that mean forest similar to fallowed land. In the Batak area in North Sumatra, *rimba oma*, *rimba aroeng*, and *rimba haoe* mean secondary forest that has been exploited then fallowed. *Rimba oma* is forest that has been fallowed for one–six years, and *rimba aroeng* for six–nine years. *Oma* is a kind of grass, *aroeng* or *tolong* is a kind of reed, and *haoe* is grown trees (*Adatrechtbundels* XXVII 1928, 176–177).

Many recent ecological studies on agroforestry similar to *talun* have been conducted. The *dusun* system in Central Maluku has been discussed by Monk *et al.* (1997, 718–737) and by Kaya *et al.* (2002, 232–233). Sardjono introduced the *lembo* system in East Kalimantan (Sardjono 1988, quoted in Herri Y. Hadikusumah 2005, 262–268) while Johan Iskandar has studied the *kaliwo* or *kalego* system in West Sumba similar to *kebun campuran*, and the *kaleka* system in Bangka Belitung, which is multi-layered forest on a former slash-and-burn location. The planting of *lada/sahang* (*Piper nigrum*) and rubber

4) Five villages at East Java were located in *residentie* Toeban, Ponorogo, Nganjoek, Bodjonegoro, and Blitar; two villages in Central Java were located at *residentie* Rembang, and Pekalongan; two villages in West Java were located at Bandoeng and Soemedang; one village at *gewest* Soerakarta was located at *residentie* Klaten. Besides these, the name of Taloenamba was found at *residentie* Bandjarnegara, Taloenblandong at *residentie* Modjokoerto, Taloenkidoel at *residentie* Djombang, Taloenkoelon at *residentie* Toeloenagoeng, Taloenombo at *residentie* Wonosobo, and Taloenredjo at *residentie* Lamongan (Schoel 1931, 373–374). This information does not exclude the possibility that there are places called *talun*, or something similar outside Java, because Schoel's dictionary only discussed names in Java island.

5) Taloenliao is in *afdeeling* Doesoelanden, *residentie* Zuid en Oost Afdeeling van Borneo; on the other hand Taloen is located in three locations in East Java, five locations in Central Java, and two locations in West Java.

(*Hevea brasiliensis*) is booming because of their good price in the markets. The *pelak* system in Kerinci, Jambi consists of the planting of annuals such as vegetables for two years after slash-and-burn, followed by the planting and harvest of dry rice. Perennials such as coffee are planted alongside the annuals, and after the harvesting of perennials, people will chose either to slash-and-burn and then to plant dry rice again, or to keep the perennials (Johan Iskandar 2009, 153–161). Rubber agroforestry called “jungle rubber” in South Sumatra was introduced by Gouyon *et al.* (1993).⁶⁾ Damar forest produced after slash-and-burn is also man-made forest in Lampung, Sumatra (Torquebiau 1984).⁷⁾

We see that in Indonesia, besides the *talun*, there are many agroforestry systems that involve slash-and-burn, followed by fallowing and the planting of perennials during the fallow. Cyclical systems that slash, cut, and burn the old perennials are found; on the other hand, continual plantings of the perennials with no more slash-and-burning are more common, and sometimes good prices support the spreading of some kinds of perennial planting. These agroforestries always retain the characteristics of diversified farming with a multi-layered canopy as a man-made forest during the productive fallow, or continual planting of perennials with no more slash-and-burn.

However, the question why the term *talun*, *taloen*, or *talon*, once used in many places both in Java and the outer Java islands as the term for man-made forest, fell out of use in the area outside of West Java, remains unanswered. Possible answers might be: i) land use of *talun* decreased or even disappeared outside West Java because of population and production growth, or was transformed into other forms such as *kebun campuran*; ii) people forgot the meaning of the term and do not use it any more;⁸⁾ iii) local variations in government policy such as the prohibition of slash-and-burn since the colonial time was what differentiated West Java from other areas; and iv) researchers have not found *talun* land use outside West Java. This paper does not address this question; nonetheless a detailed description and analysis of the sustainability of another form of *talun* is quite important for the investigation of these issues.

6) Rubber planting by smallholders in Indonesia starts with slash-and-burn (Gede Wibawa *et al.* 2005, 223). After annuals are grown for two to three years, rubber trees are planted but other perennials are left to grow. This diversified farming of rubber creates man-made forest that supports high productivity for both rubber and the household economy. Rubber would be replanted after 20–40 years of production, sometimes with an interval of slash-and-burn (Gouyon *et al.* 1993). The replanting of rubber for rejuvenating old rubber is preferred by local people to a cyclical system that slashes, cuts, and burns the old jungle rubber (Gede Wibawa *et al.* 2005, 225–231).

7) Rattan cultivation by smallholders in the man-made forest after slash-and-burn in Central Kalimantan (Godoy and Tan 1991) has boomed due to the good price for rattan.

8) Almost all current dictionaries of Indonesian—Indonesian-Indonesian and Indonesian-other languages—do not mention *talun* as a term relating to forest, or to agroforestry or land use in hilly areas.

II Kemang Village: A General Picture and Highland Farming

II-1 Kemang Village: A General Picture and Agriculture

Kemang village, the research site, belongs to Bojongpicung sub-district of the district of Cianjur. It is located in the hilly and mountainous Priangan highlands, about 7 km from the center of the sub-district, with a mountainous pass between the village and the center.

The village is surrounded by mountains that constitute a natural barrier, as such migration into the village is low. The population density of 174 persons/km² in the village (including the land controlled by the National Forestry Corporation, Perum Perhutani) or 297 persons/km² (excluding the area controlled by the National Forestry Corporation) in 2001 is relatively small compared to the average population density of 1,009 persons/km² in West Java in 2000 (BPS 2001). The population in 2001 was 4,384, with a village area of 2,518.63 ha. Of this, forests controlled by the National Forestry Corporation cover 1,040.6 ha, comprising 135 ha of protected forest (*hutan lindung*)⁹⁾ and 905 ha of production forest (*hutan produksi*).¹⁰⁾ The land controlled by the National Forestry Corporation has been designated as forest area (*kawasan hutan*) by the central government. The farmland located outside the *kawasan hutan* is privately owned by the locals, with 878.6 ha of it used as *pasir*, dry land where upland farming and perennial tree planting are combined. Rice fields are found in the low area of the village and occupy a relatively small area, accounting for just 83 ha (Desa Kemang 2001). *Pasir* land is not counted as forest by the government; however, the landscape at the stage of *talun*, full of perennial trees, is quite similar to the forest.

In this *pasir* land, dry rice, annual crops such as vegetables and tubers, and perennial trees are planted as a form of shifting cultivation, and *talun* forms a multi-layered canopy of vegetation, particularly perennial trees, as the final stage of the land use cycle. These land use and cultivation methods are described in detail in the following section.

In the area controlled by the National Forestry Corporation, social forestry programs have been implemented since the 1990s, enabling the local people to take part in the maintenance and cultivation of the area. Under the program, they first carry out slash-and-burn, then cultivate the land with dry rice and annual crops. At the same time, they also plant trees designated by the company, usually teak (*Tectona grandis*), and maintain

9) Protected forest is government-designated forest that is protected for water source conservation, prevention from floods and erosion, and maintenance of land fertility (Article 3, Act No. 5, 1966 concerning Basic Regulations on Forest).

10) Production forest is forest that is designated by the government for the needs of the people, and for the purpose of development, industry, and export (Article 3, Act No. 5, 1966 concerning Basic Regulations on Forest).

these alongside their own plantings of other perennial trees such as fruit trees like *nangka* (*Artocarpus integra*). All the harvests from the land are meant for the people, except that of the trees belonging to the company. In 1998, the National Forestry Corporation integrated the “Forest Village Society Program” (Program Masyarakat Desa Hutan, PMDH) and the “Social Forestry Program” (Perhutanan Sosial) into the “Integrated Forest Village Society Program” (Program Masyarakat Desa Hutan Terpadu, PMDHT). Kemang Village was chosen as a model village for PMDHT (Inoue *et al.* 2001, 73).

Because *talun* located on privately-owned *pasir* land is quite similar to forest in its landscape, we can find forestry both on the land controlled by the National Forestry Corporation and on the land owned by the people. However, according to the village head, the term *hutan* (forest) is only applied to the forest controlled by the National Forestry Corporation, so *hutan* is similar to *kawasan hutan*. Other forest on the land owned by the people is called *pasir*, or *talun*, and not *hutan*.

According to the villagers, the National Forestry Corporation boundary was established during the colonial era. National Forestry Corporation land was not open to ownership by individuals; but people privately owned *pasir* in non-National Forestry Corporation land. Much of National Forestry land was managed by the people as part of a social forestry program scheme. However, since leaf banana planting has prevailed, especially since around 2004, people have been planting these crops even though land is not allocated to them in the scheme of the social forestry program.

In any event, *pasir*, the privately-owned dry farmland planted with annual crops and perennial trees, the National Forestry Corporation’s land on the slopes of the mountains and hills, and the wet rice fields located in the lowland in the village are the major areas of cultivation for the people in the village surveyed. Many of the plants in the uplands and the National Forestry Corporation area are subsistence-oriented, but some plants are highly commercialized. The most important plant in the National Forestry Corporation’s area is teak; in the private upland areas, it is banana plants (*Musa* sp.), which have been increasingly popular since the second half of the 1990s, sugar palm (*Aren, Arenga pinnata*), chili plants (*Cabe, Capsicum annuum*), and dry rice (*Pare huma, Oryza sativa*). Recently the planting of *albizzia* has increased, especially since around 2006.

Among 60 respondent households, 44 households have their own rice field, whose average area is 0.14 ha. In addition, 46 households have dry land (*pasir*), measuring on average 0.91 ha. Many of the farmers who own dry land also have their own wet rice fields. Table 1 shows the number of households according to dry land and wet rice fields owned. Some households possess more than 3 ha of dry land; these households tend to also own wet rice fields. In contrast, 10 households own neither dry land nor wet rice fields.

Table 1 Number of Households by Area of Dry Land (*Pasir*) and Wet Rice Fields Owned in 2000

		No. of Households by Area of Dry Land Owned						Total
		0 ha	<0.5 ha	<1.0 ha	<2.0 ha	<3.0 ha	≥3.0 ha	
No. of households by wet rice fields owned	0 ha	10 (1)	1 (2)	4 (3)	1 (0)	0 (1)	0 (0)	16 (7)
	<0.5 ha	4 (1)	10 (6)	10 (18)	10 (12)	4 (7)	1 (1)	39 (45)
	<1.0 ha	0 (0)	0 (0)	0 (2)	1 (1)	2 (3)	1 (1)	4 (7)
	≥1.0 ha	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)	1 (1)
	Total	14 (2)	11 (8)	14 (23)	12 (13)	6 (11)	3 (3)	60 (60)

Source: Field survey conducted by authors.

Note: The number in parentheses represents the number of households by area of dry land farmed (including National Forestry Corporation land), and wet rice field farmed.

Of the households surveyed, 29 participated in the social forestry program, farming 0.25–1.25 ha of National Forestry Corporation land. The average area of the land farmed under the program was 0.43 ha. Households participating in the social forestry program are mainly those that own small areas of farmland or none at all. Of the 10 households without their own land, 9 joined the social forestry program. However, there are some households that own relatively large areas of dry land yet still join the social forestry program. The number of households in the parentheses in Table 1 is based on the area of farmed land both for wet rice fields and dry land, including social forestry program land. This table shows that only one household does not farm any land. Besides the social forestry program, there are arrangements of sharecropping, land lease, and mortgage contracts. So the farmed land average is somewhat larger than that of owned land, especially for dry land. Average dry land farmed, which includes social forestry program land, is 1.21 ha, and 0.20 ha for wet rice fields.

Apart from the cultivation of wet rice fields and dry land, there are non-agricultural activities including furniture manufacturing, rice milling, timber trading, grocery stores, and trading banana leaves. Also important is the supply of migrant workers, especially international migrants who work in Saudi Arabia.

The village consists of 22 hamlets forming 3 sub-villages (*dusun*)—*dusun* I, *dusun* II, and *dusun* III, which have 7, 5, and 10 hamlets respectively. The village lies at an altitude of 400–800 m above sea level, and the topography ranges from gently sloping to steep hilly terrain. Access to the nearest town is not easy. The road providing access to the town was built in the 1990s. Before that, people had to walk there.

II-2 Talun-Huma in *Kemang Village*

The slopes of the mountains and hills are used for upland agriculture and forestry. The land use system is quite complicated, but there are typical cases of land use.

The following is the explanation of each stage of the *talun-huma*. The first stage, *rarahan*, takes nearly three months, from July to September. The work consists of *nyacar* (tree slashing), *ngahuru* (the first burning of slashed trees), *ngaduruk* (the second burning of remains from the first burning process), *ngadampas* (clearing land from the remains of burning), *ngababantal* (making terraces), and *nyara* (collecting the remains from the *ngadampas*). In July, farmers usually start *nyacar* by cutting trees, bamboo, shrubs, and other horticultural trees selectively. They cut the very young and old *aren* trees that have already been tapped for a long time, say 20 years, bamboo, *awi ageung* (*Gigantochloa verticillata*) and *awi tali* (*Gigantochloa apus*), leguminous trees such as *kaliandra* (*Calliandra calothyrsus*), shrubs such as *sadagori* (*Sida retusa*), and old banana plants (*Musa paradisiaca*). Then they leave the slash to dry. For 0.25 ha of dry land, the *nyacar* takes 4–5 days, and the drying usually takes about 15–20 days, sometimes even one month depending on sunlight, temperature, and the kind of slash.

The next activity is *ngahuru*, where farmers gather the small dried branches, leaves, and litter in piles and burn them. This activity is usually conducted in the period from the second week until the end of August. *Ngahuru* usually takes one–two days and burning continues until the piles become ash. Then occurs *ngaduruk*, whereby farmers pile up the remaining slash that had not burned well during *ngahuru* and reburn it until the piles are reduced to ash. *Ngaduruk* is usually conducted one or two days after *ngahuru* and can last anything between two days and two weeks, as farmers have to collect the remaining slash spread over the plots.

Next comes *ngababantal*—making *babantal* (terraces) in the sloping plot. Farmers lay trunks, large branches, and bamboo as *babantal* to prevent erosion. The distance between each *babantal* is usually 3 m. *Ngababantal* usually takes more time, about 12 days. Farmers feel this is the most difficult work in the *talun-huma* as it involves making terraces. This process is called *ngais pasir*, meaning to carry the *pasir* (dry farming land) as if it were a baby wrapped in the traditional women's sarong (*sinjang pangais*)—a phrase that expresses love and care for the *pasir*.

The next activity is called *ngadampas*, in which farmers clean the remaining vegetation from the plot by chopping the roots using a short machete (*parang*). After *ngadampas*, the work of *nyara* follows. This consists of collecting the remains from the *ngadampas* and the ash, and putting them into the *babantal*, which in turn become seedbeds. The *nyara* practice is closely related to boosting soil fertility. Most of the farmers in Kemang village do not use manure to fertilize the seedbeds as not many raise goats or sheep and the plots are located far from their yards at home.

Some trees remain in the field during the *rarahan* period, usually the wood trees, bamboo, fruit trees such as mango (*Manggah*, *Mangifera indica*), *rambutan* (*Nephelium*

lappaceum), *petai* (*Peuteuy*, *Parkia speciosa*), *jengkol* (*Pithecolobium lobatum*), jackfruit (*Artocarpus heterophyllus*), *aren*, and leguminous trees such as *kaliandra*, *gamal* (*Gliricidia sepium*), and *dadap negeri* (*Erythrina* sp.) These leguminous trees have the role of conserving the land by fixing nitrogen.

During the *rarahan* stage, the farmers start planting two kinds of young banana plants: *cau buah* (fruit bananas) and *cau manggala* or *cau daun* (leaf bananas), before the first rain when dibbling for *huma* paddy starts. As the day of the first rains approach, the seeds of horticulture commodities such as pumpkin (*Waru*, *Hibiscus similis*), cucumber (*Bonteng*, *Cucumis melo*), and watermelon (*Sumangka*, *Citrulus vulgaris*) are planted. Cucumbers and watermelon are usually planted close to the area where the slashes are burnt. Except for pumpkins, two or three seeds are put in each hole, dug close to the trunk of a tree or a *tuturus* (bamboo stick for beans or other climbing plants). Pumpkins are planted at the edge of the sloping part of the plot (*sisi gawir*). Maize (*Jagung*, *Zea mays*) and *huma* paddy are planted on the same day. *Huma* rice seeds are planted in the open area after dibbling using a simple traditional tool called *aseuk*, while *cabe rawit* (small chili, *Capsicum frutescens*) seeds are usually sowed in the area close to the *babantal*.

As the *huma* rice starts growing (usually in October), the period called *huma* starts. The name of this period/stage comes from *huma*, originally meaning dry rice growing. *Pare huma* is rice harvested from dry rice growing in Sundanese. This *huma* has become the name of a land use stage for the villagers, although, in addition to dry rice, many kinds of horticultural crops and woody plants are planted during the period. The term *rarahan* originated from the activities of slash-and-burn, but it has also come to signify a stage of land use. It can be seen that despite the varied origins of the names of the stages, people utilize these terms because of the interlinked connotations.

The *huma* stage lasts for six months. The duration of the *huma* and *rarahan* stages are quite uniform. Farmers developed the *huma* stage as a strategy to meet their consumption demand, especially for food crops in the form of cereals and vegetables. Annuals of horticultural crops, particularly vegetables such as long beans (*Kacang panjang*, *Vigna sinensis*), basil (*Selasi*, *Ocimum basilicum*), eggplants (*Terong*, *Solanum* sp.), big cucumbers locally called *herbis* or *ketimun suri*, chilies (*Capsicum* sp.), and maize are harvested during the *huma* stage before the *huma* paddy is harvested in February and March. At this stage, farmers usually start to plant young woody plants such as *jeungjing* or *albizzia* (*Albizia falcataria*) and pepper (*Marica*, *Piper nigrum*), depending on the size of land. Fig. 1 shows the kinds of plants used for each stage of land use and the area planted for each plant per stage of land use (total surveyed upland used by the 60 surveyed households is 30.65 ha).

After the *huma* rice is harvested, the next stage is called *jami*, with a duration cycle




Fig. 1 Area Planted with Useful Plants by Stage of Upland Farming (2001)

Sources: Field survey conducted by authors.

Note: For our plot-based survey, respondents were asked the kinds of plants found at the plots, and the percentage of land occupied by each plant. With this information, we calculated the area planted with trees/plants. *Palawija* means secondary crops such as maize, soybean, or the like.

of about 1.5–2 years. At this stage, the horticultural crops are continuously harvested, especially duruka (*Terubuk*, *Saccharum edule*), the tubers such as cassava (*Sampeu*, *Manihot utilissima*) and sweet potatoes (*Huwi Boled*, *Ipomoea batatas*), and others such as ginger (*Jahe*, *Zingiber officinale*) and papaya (*Gedang*, *Carica papaya*). At the end of the first year, around March or April, the farmers start to harvest the fruit bananas as well as leaf bananas. In other words, the *huma* stage reflects the farmers' strategy for obtaining cereal crops and vegetable foods, while the *jami* stage reflects their strategy to satisfy vegetable and fruit consumption needs and the necessity to generate cash income, from banana fruit and leaf as well as from fruit trees, among which mangoes, *jengkol*, and *peuteuy* are noteworthy. At the *rarahan*, *huma*, and *jami* stages, the majors are annual plants, although many perennial trees are also to be found because at the *rarahan* stage, many tall trees are left untouched during slash-and-burning, and during the *huma* stage, people start to plant woody plants.

The next stage is *reuma ngora*, which lasts for one–three years. In this stage the land is usually dominated by young *albizzias* and other wood trees as well as leaf bananas and fruit trees. Annual crops are not common in this stage. The intensity of maintenance and cultivation work is lower compared with *jami*. Farmers continue to harvest the banana leaves and other fruits at this stage, and at the end of the period, the *albizzia* trees are harvested, except when farmers decide to continue to the next stage, *reuma kolot*.

Hence, this stage is used to meet the needs for *albizzia* woods, usually for repairing houses and as fuelwood, as well as for obtaining cash income. Fig. 1 shows the importance of woody trees during the *reuma ngora* and *reuma kolot* stages. Bamboos are also harvested, especially for repairing houses and making hedges for the paddy field, and so on.

The duration of the *reuma kolot* is usually three–seven years or more. The land is less cultivated than at the stage of *reuma ngora*. Wood trees are dominant in this stage. Some farmers cut *albizzia* to earn cash by selling it; others do not cut *albizzia* as they prefer to harvest teak or other wood trees. *Aren*, teakwood, mahogany (*Swietenia* sp.), bamboos, and other trees are also scattered throughout the plot. Seasonal fruits such as mango, *pisitan* (*Lansium domesticum*), *rambutan*, *jengkol*, and *peuteuy* are harvested every year. As a consequence of the growth of woody trees, the leaf bananas gradually become smaller. Farmers usually harvest *albizzia* trees at the end of this stage.

People sometimes prefer to use land as *kebun campuran* after it has been used as *jami*. With this type of land use, the multi-layer of perennial trees consists of woody trees such as *albizzia*, and bamboo. Fruits trees such as *rambutan*, *nangka*, pineapples (*Nanas*, *Ananas comosus*) are planted, and annual plants and root plants are mixed. *Aren* is an important tree in the research area; people tap the sap (*nira*) from the *aren* palm and produce palm sugar from the sap. This is a productive use of the land, and it lasts for quite a long time, sometimes 10 years or more. Banana plants are often planted for both their fruit and their leaves during this period.

Talun can be started after the *kebun campuran* period or after the *reuma kolot* period. It can be categorized into three types: i) fruit trees, *aren*, and other natural secondary vegetation, ii) fruit trees and other natural secondary vegetation, iii) *aren* and other natural secondary vegetation (Inoue *et al.* 2001, 72). *Kaliandra* is an integral tree in all three categories. People do not cultivate the land but make use of it to obtain products for both subsistence and commercial use. In any case, the intensity of usage is far lower than during the period of *kebun campuran* or *reuma kolot*.

The *talun-huma* described above is quite different from the *talun-kebun* that has been analyzed by many authors. Firstly, in the *talun-huma*, the planting of *huma* paddy is quite important, whereas in the *talun-kebun*, almost no planting of *huma* paddy is found. In many cases of *talun-kebun*, bamboo is very important for sustaining the system, whereas in the *talun-huma*, bamboo has a somewhat minor role. Mixtures of perennial trees are common. The diversity of the plants and the planting of leguminous trees such as *kaliandra* and *gamal* play an important role in sustaining the system.

Planting the *huma* paddy with dibbling and the existence of the following stages of *jami* and *reuma* are somewhat similar to the *perladangan* (shifting cultivation) system practices in the Baduy area, Banten, adjacent to West Java. There, the practice of slash-

and-burn followed by *huma* rice is quite similar to the practice found at our research site. *Kebun campuran* practice is also found at Baduy, but there is no *talun* forest. Usually the duration of *reuma* is four years, and in some cases, old secondary forest is found (Johan Iskandar 1992, 31–38, 74–111). However, there appears to be no particular term for this old secondary forest. In contrast, the *talun-huma* at our research site has a more developed system of *reuma ngora*, *reuma kolot*, and *talun* that usually takes more time during these stages.

With regard to social forestry programs, villagers report that people have been allowed to slash-and-burn since far before the introduction of the Forest Village Society Program (PMDH) at the National Forest Corporation land. The process of land use under the social forestry program is somewhat similar to the *talun-huma*, at least until the stage of *jami*, although people are required to plant teak from an early stage and maintain the land until the teak trees grow. Teak is sometimes substituted with mahogany and so on.¹¹⁾

II-3 *Variation of Land Use Stage Cycles*

The above-described sequential pattern of stages is a typical case. In reality, many variations of stage cycles are found because there are many options for the farmers.

One option is the *reuma ngora/reuma kolot* and *kebun campuran* alternative. *Talun* is a long period with low productivity, thus some people are not keen on using this cycle. *Huma* is an important period for producing dry rice; however, some people are not interested in using the land as *huma*, preferring to plant many banana plants or have more wet rice. Yet others prefer not to use the land as *reuma ngora/reuma kolot* or *kebun campuran*. They use the land as *talun* for a long time, with no intention to slash-and-burn. Some people repeat *huma* planting every three years without entering the stage of *reuma ngora*, a strategy especially common among people who have recently obtained their land, or who have just started the social forestry program.

Once farmers have decided on the *huma* stage, they should go through *rarahan*, and after the *huma*, plant secondary crops and perennials at the *jami* stage, especially during rainy season. So this sequence constitutes one set that cannot be split. *Reuma ngora* (“young” *reuma*) and *reuma kolot* (“old” *reuma*) form another set.

From these observations, and also from discussions concerning permanent forest

11) This approach to social forestry is very different from the social forestry program implemented by the National Forestry Company in Middle Java where the National Forestry Company has not allowed people to slash-and-burn in the social forest company land, according to Dr Pujo Sumedi, researcher at Gajah Mada University (interview conducted by the authors on August 17, 2011).

and cyclical use with slash-and-burn, we have determined five types of land use sequence.

The first is the rotating sequence from *rarahan/huma/jami* to the productive fallow of *reuma ngora/reuma kolot*, and finally to *talun*, or sometimes once to *kebun campuran*, and finally *talun*. In this pattern we find *huma* and *talun* in the sequence.

The second is the cyclical sequence from *rarahan/huma/jami* to productive fallow such as *reuma ngora/reuma kolot* or *kebun campuran*, but no *talun*.

The third is the cyclical sequence among *rarahan/huma/jami*, but without leading to productive fallow such as *reuma ngora/reuma kolot* or *kebun campuran*. This type of land use includes plots that were recently acquired/sharecropped/allocated under the social forestry program, and slashed-and-burned, but where it is not clear yet whether the plot will proceed to the stage of *reuma ngora*, or be slashed-and-burned again after a fallowing interval because the cultivators are not the owners of the plots and cannot/do not answer for the land use following the *jami* stage, especially in the case of sharecropping.

The fourth type is permanent forest with a *talun* stage. The fifth is permanent forest without *talun*, such as *reuma ngora/reuma kolot* and *kebun campuran*, or *kebun campuran*, or a particular forest such as *albizzia* forest. In these cases, *talun*, *reuma ngora/reuma kolot*, or *kebun campuran* are no longer productive fallow but permanent forest. These five types are shown at Table 2.

Of the 174 plots of dry land controlled by 60 households that were surveyed, the largest number of plots had adopted the second sequence—cyclical land use with productive fallow without *talun*. This was used in 87 cases, or 50 percent of all cases. The second largest number of plots used the cyclical sequence with *talun*: 38 cases, or 21.9 percent. The number of the plots including a *talun* stage is 48, or 27.6 percent of all plots.

Table 2 Plots and Their Characteristics by Cyclical Sequence (2001)

	Number of Plots	Average Distance from House (km)	Average Area of the Plot (ha)	Average Duration of Cycle (years)	Average Duration since Acquisition (years)
1. Cyclical sequence with <i>talun</i> and <i>huma</i>	38 (21.9%)	2.2	0.49	11.8	21.9
2. Cyclical sequence with <i>huma</i> and productive fallow without <i>talun</i>	87 (50.0%)	3.5	0.49	5.3	13.8
3. Cyclical sequence with <i>huma</i> without productive fallow	32 (18.4%)	4.9	0.41	2.6	5.9
4. Permanent forest with <i>talun</i>	10 (5.7%)	1.7	0.42	–	15.3
5. Permanent forest without <i>talun</i>	7 (4.0%)	0.7	0.34	–	3.7

Source: Field survey conducted by authors.

In contrast, the number of plots including a *huma* stage is 157, or 90.2 percent of all plots. Those plots go through the process of slash-and-burn. Among the 157 plots that have a *huma* stage, 125 plots or 79.6 percent of the plots have productive fallowing such as *reuma ngora/reuma kolot*, *kebun campuran*, or *talun*. Plots of permanent forest occur in 17 cases, or 9.8 percent, and among them 10 plots, or 5.7 percent of all plots had permanent forest with *talun*. *Talun-huma* refers to the first category of the Table 2 in the narrow sense, on the other hand *talun-huma* and agroforestry can cover the whole practices discussed above.

III *Talun-Huma* in the Rural Context of Social Economy

III-1 *Talun-Huma and Socioeconomic Factors*

The *talun-huma* in Kemang village displays a range of characteristics according to the cyclical sequence pattern. The characteristics are partly geographic, partly social, and partly economic as shown at Table 2.

The *huma* cultivated plots are located in remote areas. In contrast, the nearer the plot, the more *talun* is practiced. Concerning the average duration of the cycle, greater use of the *talun* system correlates to longer cycles.

Table 3 shows that the percentage of plots owned by the respondents was far higher in cyclical sequences with *talun* and *huma*, and permanent forest with *talun*. Among the 174 plots that were managed by 60 respondents households, 105 were owned by the household surveyed, and 42 were plots allocated to farmers under the social forestry scheme, 21 were sharecropped, 5 were lease-held, and 1 case was mortgaged by the

Table 3 Plot Ownership Characteristics by Cyclical Sequence (2001)

	Percentage of Respondent's Ownership (%)	Average Price of Plot (Rp. 1,000 per ha)	Average Area of Rice Field Owned by House-hold (ha)	Average Area of Dry Land Owned by House-hold (ha)
1. Cyclical sequence with <i>talun</i> and <i>huma</i>	89.4	21,860	0.20	1.03
2. Cyclical sequence with <i>huma</i> and productive fallow without <i>talun</i>	58.6	11,950	0.17	1.06
3. Cyclical sequence with <i>huma</i> without productive fallow	9.2	15,140	0.12	0.74
4. Permanent forest with <i>talun</i>	100	22,900	0.33	2.42
5. Permanent forest without <i>talun</i>	85.7	27,280	0.15	0.74

Source: Field survey conducted by authors.

respondent households. Among 21 sharecropping practices, 12 were among family members,¹²⁾ and 9 were among non-family members. So the smaller the percentage of the plots owned by the respondents, the smaller the percentage of households who use the stage of *talun* or other productive fallows, and the shorter the average duration of the sequence cycle. People who have joined the social forestry program or sharecropped the plot tend to practice more *huma* and activities relating to *huma* (*varahan*, *huma* and *jami*).

Concerning the average plot price per hectare, plots that have a *talun* stage fetch relatively higher prices; plots of permanent forest are also relatively highly valued. Households that own a larger area of *pasir* dry farmland with perennial trees planting and wet rice fields are prone to having the *talun* stage, but the less of these lands a household possesses, the less the amount of productive fallow in the sequence.

III-2 Talun-Huma and Economic Activity

III-2-1 Agricultural Sector

Economic activities and employment opportunities in the agricultural sector are closely related to dry land farming. The dry land farming stages give rise to variation in these activities.

Fig. 2 shows the yearly income earned by agriculture-related waged laborers in




Fig. 2 Agriculture-related Wage Labor by Upland Farming Stage (2001)

Source: Field survey conducted by authors.

12) Family members here mean parents and parents-in-law, and brothers or brothers-in-law of the respondents.

relation to upland farming stages. The amount of income is for whole households. At the time of the *jami* and *reuma ngora* stages, farm-waged laborers are in peak demand for both dry land farming and wet rice farming.

Income from various activities in the agricultural sector varies according to strata. Table 4 shows the yearly income composition of surveyed households according to strata. Income from wet rice farming, agricultural waged labor, palm sugar, and leaf banana production has a close relationship with the household strata. Upper strata households have a higher proportion of total household income generated by wet rice farming and leaf banana production. For lower strata households, agricultural waged labor, palm sugar production, and dry rice production generate a higher percentage of total household income.

There are agricultural waged labor and sharecropping relations between the owners of *pasir* dry farming/wet rice farming land and agricultural waged laborers/share croppers concerning palm sugar production, leaf bananas harvesting, and *huma* cultivation. Products of agricultural waged labor or sharecropping are divided between the landowner and agricultural laborers/sharecroppers. The portion of products/harvests differs according to the distance from the settlements: if the location is far, the portion for the laborers or sharecroppers is two-thirds; if the plot is not far, the portion is a half (*maro*) (Siti Sugiah Machfud Mugniesyah *et al.* 1999).

Tables 2 and 3 show that the *talun* system is practiced by the upper strata, whereas *huma* is in many cases practiced by sharecroppers, or people who have joined the social forestry program. This tendency can be partly discerned in Table 4. Dry land farming is quite important for the lower strata, strata C (79.0 percent of income derived) and D (29.2 percent of income from dry land farming, and 61.7 percent from agricultural labor), especially considering that the agricultural labor undertaken by stratum D respondents is closely related with the dry land farming of strata A and B households. Although dry rice production is not so important for the income of respondents as a whole (2 percent), it cannot be neglected for stratum D respondents, for whom dry rice contributes 11 percent to their income.

Dry land farming is important for the upper strata farmers too because they manage a wider area of dry land and derive substantial income from it (21.5 percent for stratum A, and 25.4 percent for stratum B). These incomes support the sustainability of the agroforestry system constituting of *talun-huma*, and permanent forest with or without *talun*. Palm sugar and leaf banana production contribute substantially to this income. Palm sugar production has a long history, whereas leaf banana production only started in the middle of the 1990s. The flexibility of the *talun-huma* and agroforestry in accommodating so many kinds of plants, and the fertile soil resulting from a multi-layered, often-dense canopy of vegetation that protects against soil erosion and leaching, has

Table 4 Household Yearly Income by Economic Activity According to Strata (2001) (Units: Rp. 1,000)

	Wet Rice Farming	Dry Land Farming				Agricultural Waged Labor		Non-agricultural	Others	Average
		Total	Palm Sugar	Banana	Dry Rice	Others				
A	1,757	4,378	879	2,756	51	692	23	13,772	482	20,412
N=13	8.6%	21.5%	4.3%	13.5%	0.3%	3.4%	0.1%	67.5%	2.3%	100%
B	622	3,083	1,324	1,267	187	305	321	7,000	1,124	12,150
N=19	5.1%	25.4%	10.9%	10.4%	1.6%	2.5%	2.6%	57.6%	9.3%	100%
C	117	3,618	1,685	1,070	276	587	172	574	98	4,579
N=18	2.5%	79.0%	36.8%	23.4%	6.0%	12.8%	3.7%	12.5%	2.3%	100%
D	58	1,182	1,004	-328	451	55	2,500	311	0	4,051
N=10	1.4%	29.2%	24.8%	-8.1%	11.1%	1.4%	61.7%	7.7%	0.0%	100%
Total	622	3,208	1,288	1,260	228	432	575	5,425	489	10,319
N=60	6.0%	31.1%	12.5%	12.2%	2.2%	4.2%	5.6%	52.6%	4.7%	100%

Source: Field survey conducted by authors.

Note: Strata codes are based on scoring of socioeconomic factors of surveyed households. Agricultural land was classified according to the status of ownership: owned, mortgaged, shared, or leased. Farmhouses were classified into four groups according to type of roof, walls, and floors. Occupations were classified into four groups. According to these classifications, scores were given to each status, type, and group. Strata of households were decided according to the scores given to each household. For further explanation, refer to Siti Sugiah Machfud Mugniyah and Mizuno (2001).

enabled the planting of leaf bananas on a large scale, which is quite profitable for the respondents. It is questionable whether the widespread growth of leaf banana planting will support the *talun-huma* and agroforestry because of its profitability, or on the contrary, will hurt the system because of over-planting and declining fertility (Tsuji and Ageng forthcoming). Nevertheless, it is quite apparent that the *talun-huma* and agroforestry contributes significantly to the rural economy. Dry land cultivation, which contributes 31.1 percent of the surveyed household income, is only a part of *talun-huma* and agroforestry's economic contribution. Most of the agricultural waged labor, and the banana leaf trade that are essential parts of the *talun-huma* and agroforestry, also contribute to the rural economy.

III-2-2 Talun-Huma and the Non-Agricultural Sector

The non-agricultural sector plays an important role in the economy. The total income amongst the surveyed households coming from the non-agricultural sector is 52.6 percent. Table 4 clearly shows that the higher the household strata, the larger the percentage of non-agricultural sector income relative to total household income.

Table 5 shows the composition of the yearly income derived from the non-agricultural sector by the surveyed households. Income derived from working in the civil service, *warung* (grocery) shop management, and machinery management such as chain saw rental, are clearly related to strata level, and the higher strata derive a greater percentage of income from these activities in total household income.

Table 5 Composition of Household Non-agricultural Yearly Income (2001)
(Units: Rp. 1,000)

	Non-agriculture				Remittance	Husbandry	Rentout of Land	Average
	Total	Machinery Manage- ment	Civil Servant	<i>Warung</i> Shop				
A N=13	13,772 67.5%	3,085 15.1%	5,086 24.9%	2,946 14.4%	323 1.6%	6 0%	152 0.7%	14,253 69.8%
B N=19	7,000 57.6%	1,470 12.1%	706 5.8%	2,021 16.6%	611 5.0%	27 0.2%	486 4.0%	8,124 66.8%
C N=18	574 12.5%	0 0%	0 0%	187 4.1%	54 1.1%	0 0%	43 1%	671 14.5%
D N=10	311 7.7%	0 0%	240 5.9%	0 0%	0 0%	-1 0%	0 0%	310 7.7%
Total N=60	5,425 52.6%	1,134 11.0%	1,366 13.2%	1,3 12.9%	279 2.7%	10 0.1%	200 1.9%	5,914 57.3%

Source: Field survey conducted by authors.

Note: Figures in percentage indicate the ratio to total income of individual classes shown in Table 4.

These non-agricultural sectors may contribute to sustaining the *talun-huma* and agroforestry because people derive substantial income from the non-agricultural sector, so they need not rely heavily on the dry land management sector, which results in longer terms of productive fallow. People may invest in the dry farming sector with the income derived from the non-agricultural sector. Alternatively, the non-agricultural sector may hurt the *talun-huma* and agroforestry because people do not need substantial income from the upland, and there may be too little labor to retain the cyclical sequence of land use, or maintain permanent forest in a productive way. This issue will be addressed in the following section.

IV Analysis of the Sustainability of the *Talun-Huma*

Many factors may have an influence on the sustainability of the *talun-huma* and agroforestry. From our description and analysis above, the distance of the plot from the house, duration of holding since acquisition, duration of the cycles, ownership status of the plot, area of household holding of dry land and wet rice field, and price of the plot are considered to be important for the sustainability of the *talun-huma* and agroforestry. It is debatable whether income from banana planting and non-agriculture will support or hurt the *talun-huma* and agroforestry. In order to determine whether these factors are positively related to the sustainability of the *talun-huma* and agroforestry, correlation between these factors and sustainability, and sequential cycles were examined and analyzed by some methodologies of multivariate statistics, using the area of total *pasir*/wet rice fields land owned by surveyed households, distance of the plots, income from banana production, income from non-agricultural sectors, income from wet rice farming, ownership status, and the number of household members¹³⁾ in relation to the data of each plot as variants.

Whether the *talun* stage exists or not is qualitative data. This data can be analyzed with a probit model. We have assumed the following linear equation parameters:

$$DT = F1 (WUH, DIT, DMN, NFM, IWR, IBL, ING, OWN) (4-1)$$

In order to examine the influence of these factors on whether the *talun* stage exists or not, we have assumed the following linear equation parameters:

13) The number of household members may relate to the necessity of income and availability of household labor. Price of the land per hectare was not considered because we were concerned that the price may be more influenced by the distance of the plots from the settlement.

Table 6 Estimation of *Talun* Existence Measured by Upland Farming-related Variables

Variable	Coefficient	Std. Error	z-Statistic	Prob.
C	-0.384	0.434	-0.884	0.377
Area of farming plots owned by household	0.000	0.001	0.186	0.852
Distance from house	-0.199***	0.0629	-3.167	0.002
Duration after acquisition	0.013	0.009	1.428	0.153
Number of family members	-0.111**	0.056	-1.982	0.048
Income from wet rice farming	4.02E-08	9.63E-08	0.418	0.676
Income from banana production	-7.45E-08	6.81E-08	-1.095	0.274
Income from non-agricultural sectors	-8.07E-10	1.15E-08	-0.070	0.944
Ownership	0.979***	0.331	2.952	0.003
McFadden R-squared	0.226			
Log likelihood	-78.789			
Chi-square	26.967***	(Degrees of freedom=3)		
Obs with Dep=0	124			
Obs with Dep=1	48			
Total obs	172			

Note: ***=1%, **=5%, and *=10%, significant level

DT: Dummy variable of existence of *talun* for each plot (1: *Talun* in, 0: *Talun* out)

WUH: Area of farming plots owned by one household, which includes the wet rice farming and dry land farming (in ha)

DIT: Distance from the farmhouse surveyed to the upland farm plot (in km)

DMN: Duration of land management after acquisition by respondent (in years)

NFM: Number of family members in the household surveyed

IWR: Income from wet rice cultivation by household surveyed during 2001 (in Rupiah)

IBL: Income from banana production by household surveyed during 2001 (in Rupiah)

ING: Income from non-agricultural sector activities by household surveyed during 2001 (in Rupiah)

OWN: Ownership of plots using dummy variables (1: owned by household member, 0: not owned by household member, such as sharecropped)

From this Table 6 we can estimate that the nearer to the house, the smaller the number of family members, and whether the land is owned, the greater the likelihood of *talun* being practiced. Income from non-agricultural sectors, banana production, and wet rice are not statistically proved to be related to the question of whether or not *talun* is practiced.

Now we examine the question of whether or not *huma* is practiced, or slash-and-

Table 7 Estimation of *Huma* Existence Measured by Upland Farming Related Variables

Variable	Coefficient	Std. Error	z-Statistic	Prob.
C	-0.678	0.670	-1.012	0.312
Area of farming plots owned by household	-0.006**	0.002	-2.511	0.012
Distance from house	0.766***	0.189	4.058	0.000
Duration after acquisition	0.068***	0.021	3.291	0.001
Number of family members	0.135	0.092	1.469	0.142
Income from wet rice farming	4.14E-07**	1.97E-07	2.101	0.036
Income from banana production	-3.12E-08	1.04E-07	-0.299	0.765
Income from non-agricultural sectors	3.82E-08**	1.78E-08	2.149	0.032
Ownership	-0.872*	0.479	-1.822	0.068
McFadden R-squared	0.406			
Log likelihood	-35.650			
Chi-square	41.670***	(Degrees of freedom=5)		
Obs with Dep=0	19			
Obs with Dep=1	155			
Total obs	174			

Note: ***=1%, **=5%, and *=10%, significant level

burn is practiced, using a probit model. We have assumed the following linear equation parameters:

$$DH = F2(WUH, DIT, DMN, NFM, IWR, IBL, ING, OWN) \text{ (4-2)}$$

DH: Dummy variable of existence of *huma* practice (1: *Huma* in, 0: *Huma* out)

The other variables are the same as with equation (4-1).

From Table 7, we can see that the further away the plot, and the longer the duration after the acquisition of the plot, the more *huma* or slash-and-burn are practiced. If the plot is not owned, and the smaller the area of farming plot owned by the respondents, the more *huma* is practiced. However, more income from wet rice farming and from the non-agricultural sectors does not result in a decrease in *huma* practice. In fact, according to this estimation, they somewhat support the practice of *huma*, with more practice of *huma* or slash-and-burn.

Finally we estimate the duration of the cycle to check the sustainability of *talun-huma* and agroforestry using ordinary least squares method. We have assumed the following linear equation parameters:

$$\text{Duration of Cycle} = F3(WUH, DIT, DMN, NFM, IWR, IBL, ING, OWN) \text{ (4-3)}$$

Duration of cycle: duration of a sequence of land use cycle for each plot (in months)

The other variables are the same as with equation (4-1).

Table 8 Estimation of Duration of Cycle Measured by Upland Farming Related Variables

Variable	Coefficient	Std. Error	t-Statistic	Prob.
C	5.871***	2.017	2.910	0.004
Area of farming plots owned by household	0.010	0.006	1.537	0.127
Distance from house	-0.720***	0.256	-2.813	0.006
Duration after acquisition	0.135***	0.049	2.761	0.007
Number of family members	-0.034	0.261	-0.129	0.898
Income from wet rice farming	5.75E-07	4.91E-07	1.170	0.244
Income from banana production	-8.09E-07***	2.93E-07	-2.757	0.007
Income from non-agricultural sectors	-4.69E-08	5.71E-08	-0.820	0.414
Ownership	1.144	1.385	0.826	0.410
Adjusted R-squared	0.267			
F-statistic	6.825***			
Included observations	129			

Note: ***=1%, **=5%, and *=10%, significant level

Table 8 shows that the further away the plot, and the shorter the time since the acquisition, the shorter the duration of the cycle. We observe that with relation to banana planting; the larger the banana harvest, the shorter the duration of the cycle. We can infer somewhat that the larger the area of farming plots owned by the households, the longer the cycle, but this trend is not well supported statistically. For non-agricultural sector income, we cannot find a clear relation with the duration of the cycle.

From these three estimations, we can say that distance, and ownership have been statistically proven to be closely related with the practice of *talun* and *huma*. On the other hand, non-agriculture factors have not proved to be correlated with the existence of *talun* nor with the duration of sequence cycle. Moreover, the non-agricultural sector has proven to be positively correlated with the existence of *huma* practice. These data mean that further development of non-agricultural sectors would not damage the *talun-huma* and agroforestry, at least in the near future. Leaf banana planting is not related to the existence of *talun* and *huma*; however, it has been proven to be related to the duration of the cycle. The greater the leaf banana planting, the shorter the duration of the cycle, which may in turn lead to declining fertility of the dry land.

V Conclusion

Land use intensification for swidden agriculture has taken place in Indonesia. The most vivid case study is the *talun-kebun* in West Java, especially in relation to *talun* bamboo. *Talun* is meaningful from ecological, social, and economic perspectives, and is thought

of as being a typical practice in West Java.

This study has shown that the term *talun* was used in many places such as North Sumatra, Minahasa, and Malay people's areas in Sumatra, Borneo, Central and East Java, besides West Java at least during colonial times. *Talun* even became the name of villages, sub-districts, and mountains in many places in Java island. Besides *talun*, there are many practices of land use intensification in relation to man-made forest modified from conventional slash-and-burn. These have been studied, especially from an ecological perspective.

This study has shown that *talun-huma* and agroforestry practice in the surveyed village contributed greatly to the economy of households surveyed, through dry land farming, palm sugar production, and agricultural waged labor using the data collected at the field in 2000–1. The trading of banana leaves also constitutes as a contribution of the *talun-huma* and agroforestry to rural economy. We have demonstrated that the *talun-huma* and agroforestry is closely linked to the socioeconomic structure of the rural society. *Huma* is more likely to be practiced by the lower strata who own little farming land. The economic development reflected in the growth of leaf banana production and non-agricultural sector income has not excluded or diminished *talun-huma* and agroforestry. Sharecropping practices and agricultural labor relations among the villagers established *huma* practice amongst higher strata villagers who invest more in leaf banana production. The social forestry program implemented in this region also supports the continuity of *huma* practice. Moreover, the development of non-agricultural sectors and wet rice cultivation has had a positive impact on the existence of *huma* practices and the continuation of slash-and-burn practices. The National Forestry Corporation has enabled people to slash-and-burn for a long time at least until 2001, and that within the social forestry program, meaning that the National Forestry Company to some extent adopted the *talun-huma* and agroforestry for the planting of teak or mahogany.

Talun is practiced on nearby plots with a longer duration from the time of acquisition by families with a small number of family members. The land is owned by the respondent household.

Talun-huma and permanent forest with or without *talun* are good practices that keep the system sustainable from an economic and social point of view. Diversified farming, balanced rotation of land use, and diversity of plants, as well as the planting of land conservation trees in the form of leguminous trees such as *kaliandra* and *gamal*, play an important role in sustaining the system.

Maintaining the *rarahan-huma-jami* practice, as well as the *reuma ngora* or *kebun campuran* practices, requires effort on the part of the people and a certain amount of labor. From this point of view, the introduction of leaf banana plants to the village in the

1990s has made an important contribution to the continuity of the system because the plants supply much income to villagers both as farmers and as agricultural laborers. They can then rely on the income from the hilly dry land. However, a problem that might prove to be serious in the near future is that leaf banana planting shortens the duration of the cycle and may render the land infertile.

Accepted: July 27, 2012

References

- Adatrechtbundel* [Collection of customary law]. 1910–. Het Koninklijk Instituut Voor De Taal, Land, en Volkenkunde van Nederlandsch-Indië. S-Gravenhage: Martinus Nijhoff.
- Badan Pusat Statistik (BPS). 2001. *Statistik Indonesia* [Statistical Yearbook of Indonesia]. Jakarta: BPS.
- Barth, J. P. J. 1910. *Boesangsch-Nederlandsch Woordenboek* [Boesangnese-Dutch dictionary]. Batavia: Landsdrukkerij.
- Coolsma, S. 1913. *Soendaneesch-Hollandsch Woordenboek* [Sundanese-Dutch dictionary]. Tweede Druk, Leiden: A.W. Sijthoff's Uitgevefrs-Maatschappij.
- Desa Kemang. 2001. *Profil Desa/Kelurahan* (Buku I): *Daftar Isian Data Dasar Profil Desa/Kelurahan, Desa/Kelurahan: Kemang, Kecamatan: Bojongpicung, Kabupaten: Cianjur: Tahun 2001* [Profile of village, Book I: List of content, and basic profile of Kemang village, Bojongpicung sub-district, Cianjur district, year 2001]. Desa Kemang, Kecamatan Bojongpicung, Kabupaten Cianjur, Propinsi Jawa Barat.
- Duguma, B.; Gockowski, J.; and Bakala, J. 2001. Smallholder Cacao (*Theobroma cacao* Linn.) Cultivation in Agroforestry Systems of West and Central Africa: Challenges and Opportunities. *Agroforestry Systems* 51(3): 177–188.
- Dumont, Ch. F. H. 1917. *Aardrijkskundig Woordenboek van Nederlandsch Oost-Indië* [Geographical dictionary of the Netherlands East India]. Rotterdam: Nijgh en Van Ditmar.
- Gede Wibawa; Sinung Hendratno; and van Noordwijk, Meine. 2005. Permanent Smallholder Rubber Agroforestry Systems in Sumatra, Indonesia. In *Slash-and-Burn Agriculture: The Search for Alternatives*, edited by Cheryl A. Palm, Stephen A. Vosti, Pedro A. Sanchez, and Polly J. Ericksen, pp. 222–232. New York: Columbia University Press.
- Godoy, Ricardo A.; and Tan Ching Feaw. 1991. Agricultural Diversification among Smallholder Rattan Cultivators in Central Kalimantan, Indonesia. *Agroforestry Systems* 13(1): 27–40.
- Goris, R. 1938. *Beknopt Sasaksch-Nederlandsch Woordenboek* [Concise Sasakese-Dutch dictionary]. Singaradja: Publicatie Kirtya Lieftrinck, Van der Tuuk.
- Gouyon, A.; De Foresta, H.; and Levang, P. 1993. Does “Jungle Rubber” Deserve Its Name? An Analysis of Rubber Agroforestry Systems in Southeast Sumatra. *Agroforestry Systems* 22(3): 181–206.
- Herri Y. Hadikusumah. 2005. *Silabus Mata Kuliah: Ekologi Pedesaan* [Syllabus of subject: Rural ecology]. Bandung: Jurusan Biologi, Fakultas Matematika dan Ilmu Pengetahuan Alam, Universitas Padjadjaran.
- Inoue, Makoto; Siti Sugiah Machfud Mugniesyah; and Tsurudome, Yuuki. 2001. Tropical Forestland as a Safety Net for Local People: Significance of Sloping Land in Kemang Village, West Java, Indonesia.

- In *Proceedings of the 1st Seminar "Toward Harmonization between Development and Environmental Conservation in Biological Production,"* pp. 71–78. Graduate School of Agricultural and Life Science, The University of Tokyo.
- Jansz, P. 1906. *Practisch Javaansch-Nederlandsch Woordenboek Met Latijnsche Karakters* [Practical Javanese-Dutch dictionary with Latin characters]. Semarang, Soerabaya, Bandoeng, 's-Gravenhage: N.V. Boekhandel end Drukkerij v/h G.C.T. van Dorp & Co.
- Johan Iskandar. 2009. *Ekologi Manusia dan Pembangunan Berkelanjutan* [Human ecology and sustainable development]. Bandung: Program Studi Magister Ilmu Lingkungan, Universitas Padjadjaran.
- . 1992. *Ekologi Perladangan di Indonesia, Studi Kasus dari Daerah Baduy Banten Selatan, Jawa Barat* [Ecology of shifting cultivation in Indonesia, a case of Baduy Area, South Banteng, West Java]. Jakarta: Penerbit Djambata.
- Johan Iskandar; Isnawan, H.; H. Y. Hadikusumah; and Otto Soemarwoto. 1981. Sistem Talun-Kebun: Suatu Sistem Pertanian Hutan Tradisional [*Talun-Kebun* system: A system of traditional agroforestry]. *Proceedings Seminar Agroforestry dan Pengendalian Perladangan* [Proceedings of the seminar on agroforestry and the control of shifting cultivation], November 19–21, 1981 at Jakarta. Jakarta: Balai Penelitian Hutan, Dep. Kehutanan.
- Kaya, M.; Kammesheidt, L.; and Weidelt, H.-J. 2002. The Forest Garden System of Saparua Island, Central Maluku, Indonesia, and Its Role in Maintaining Tree Species Diversity. *Agroforestry Systems* 54(3): 225–234.
- Lembaga Basa & Sastra Sunda. 1975. *Kamus Basa Sunda* [Dictionary of Sundanese]. Bandung: Penerbit Terate.
- Linda Christanty; Maily, D.; and Kimmins, J. P. 1996a. "Without Bamboo, the Land Dies": A Conceptual Model of the Biogeochemical Role of Bamboo in an Indonesian Agroforestry System. *Forest Ecology and Management* 91(1): 83–91.
- . 1996b. "Without Bamboo, the Land Dies": Biomass, Litterfall, and Soil Organic Matter Dynamics of a Javanese Bamboo *Talun-kebum* System. *Forest Ecology and Management* 87(1–3): 75–88.
- Linda Christanty; Oekan S. Abdoellah; Marten, Gerald G.; and Johan Iskandar. 1986. Traditional Agroforestry in West Java: The *Pekarangan* (Homegarden) and *Kebun-Talun* (Perennial-Annual Rotation) Cropping Systems. In *Traditional Agriculture in Southeast Asia, a Human Ecology Perspective*, edited by Gerald G. Marten, pp. 132–170. Colorado: Westview Press.
- Linda Christanty; Priyono; and Karyono. 1980. *Penelitian Pendapatan Talun di Desa Sawangan, Jawa Tengah* [Research on the income of *talun* at the Sawangan village, Central Java]. Bandung Laporan Intern Lembaga Ekologi, Universitas Padjadjaran.
- Maily, D.; L. Christanty; and Kimmins, J. P. 1996. "Without Bamboo, the Land Dies": Nutrient Cycling and Biogeochemistry of a Javanese Bamboo *Talun-kebum* System. *Forest Ecology and Management* 91(2): 155–173.
- Mizuno, Kosuke; and Siti Sugiah Machfud Mugniesyah. 2003. Economic Crisis and Social Safety Net Programs at Kemang Village in West Java. In *Sustainable Agriculture in Rural Indonesia*, edited by Yoshihiro Hayashi, Syafrida Manuwoto, and Slamet Hartono, pp. 283–293. Yogyakarta: Gadjah Mada University Press.
- Monk, Kathryn A.; de Fretes, Yance; and Reksodiharjo-Lilley, Gayatri. 1997. *The Ecology of Nusa Tenggara and Maluku*. The Ecology of Indonesia Series, Vol. 5. Singapore: Periplus Editions.
- Otto Soemarwoto. 1983. *Ekologi Lingkungan Hidup dan Pembangunan* [Ecology of environment and development]. Jakarta: Penerbit Djambatan.
- Otto Soemarwoto; and Idjah Soemarwoto. 1984. The Javanese Rural Ecosystem. In *An Introduction to Human Ecology Research on Agricultural Systems in Southeast Asia*, edited by A. Terry Rambo and

- Percy E. Sajise, pp. 254–287. Honolulu: East-West Center; Los Baños: University of the Philippines.
- Otto Soemarwoto; Linda Christanty; Henky; Herri Y. H.; Johan Iskandar; Hadyana; and Priyono. 1985. *The Tahun-Kebun: A Man-made Forest Fitted to Family Needs. Food and Nutrition Bulletin* 7(3): 48–51.
- Padoch, Christine; and de Jong, Wil. 1987. Traditional Agroforestry Practices of Native and Ribereño Farmers in the Lowland Peruvian Amazon. In *Agroforestry: Realities, Possibilities and Potentials*, edited by H.L. Gholz, pp. 179–194. Dordrecht: Martinus Nijhoff Publishers.
- Parikesit; Takeuchi, K.; Tsunekawa, A.; and O. S. Abdoellah. 2004. Kebon Tatangkalan: A Disappearing Multi-layered Agroforest in the Upper Citarum Watershed, West Java, Indonesia. *Agroforestry Systems* 63(2): 171–182.
- Penninga, P.; and Hendriks, H. 1936. *Practisch Madurees-Nederlands Woordenboek* [Practical Madurese-Dutch dictionary]. 's-Grahenhage: Van Dorp.
- Porkas Sagala. 1994. *Mengelola Lahan Kehutanan Indonesia* [Managing the land in the Indonesian forest]. Jakarta: Yayasan Obor Indonesia.
- Prawiroatmodjo. 1957. *Bausastra Djawa-Indonesia* [Dictionary of Javanese-Indonesian]. Surabaya: Penerbit Express & Marfiah.
- Sanchez, Pedro A.; Palm, Cheryl A.; Vosti, Stephen A.; Tomich, Thomas P.; and Kasyoki, Joyce. 2005. Alternatives to Slash and Burn: Challenge and Approaches of an International Consortium. In *Slash-and-Burn Agriculture, the Search for Alternatives*, edited by Cheryl A. Palm, Stephen A. Vosti, Pedro A. Sanchez, and Polly J. Ericksen, pp. 3–37. New York: Columbia University Press.
- Sardjono, M. A. 1988. Lembo: Sistem Pemberdayaan Lahan Tradisional di Kalimantan Timur [Lembo, a system of traditional land enrichment in East Kalimantan]. *Prosiding Seminar: Agroforestry untuk Pengembangan Daerah Pedesaan di Kalimantan Timur* [Proceeding of seminar: Agroforestry for the development of rural areas at East Kalimantan], September 19–21, 1988 at Fak. Kehutanan Universitas Mulawarman, Samarinda-Deutsche Gesellschaft Für Technische Zusammenarbeit (GTZ).
- Sasaki Komei 佐々木高明. 1970. *Nettai no Yakihata* 熱帯の焼畑 [Swidden cultivation in the tropics]. Tokyo: Kokin Shoin.
- Schoel, W. F. 1931. *Alphabetisch Register van de Administratieve- (Bestuurs-) en Adatrechtelijk Indeeing van Nederlandsch-Indië* Deel I: *Java en Madoera*. [Alphabetical registration of the administrative and customary classification of the Netherlands-India, Part I: Java and Madoera]. Batavia: Landsdrukkerij.
- Siti Sugiah Machfud Mugniesyah; and Mizuno, Kosuke. 2003. Gender Relations among Upland Farming Households: The Case of Kemang Village in West Java, Indonesia. In *Sustainable Agriculture in Rural Indonesia*, edited by Yoshihiro Hayashi, Syafrida Manuwoto, and Slamet Hartono, pp. 331–344. Yogyakarta: Gadjah Mada University Press.
- . 2001. Gender, Poverty and Peasant Household Survival Strategies: A Case Study in Dry Land Village in West Java. In *Proceedings of the 1st Seminar on "Toward Harmonization between Development and Environmental Conservation in Biological Production,"* pp. 63–70. Graduate School of Agricultural and Life Science, The University of Tokyo.
- Siti Sugiah Machfud Mugniesyah; Mizuno, K; and Iwamoto, N. 1999. Agricultural Sustainable Development in Rural Indonesia: A Gender and Social Change Perspective. A Case Study in Upland Area in West Java. Paper presented at the Seminar of the JSPS-DGHE Core University Program in Kyoto, February 2, 1999.
- Terra, G. J. A. 1953. The Distribution of Mixed Gardening on Java. *Landbouw* 35(1–6): 163–224.
- Torquebiau. 1984. Man-made Dipterocarp Forest in Sumatra. *Agroforestry Systems* 2: 103–127.
- Tsujii, Hiroshi; and Ageng Setiawan Herianto. Forthcoming. An Economic Analysis of Agricultural Sustainability in a Mountainous Village of West Java: Use of Multinomial Logit Model Analysis. In

Sustainability and Crisis at a Village of Agroforestry in West Java, Indonesia: Talun-Huma System and Rural Social Economy, edited by Kosuke Mizuno and Siti Sugiah Mugniyah. Yogyakarta: Gadjadara University Press.

Veth, P. T. 1869. *Aardrijkskundig en Statistisch Woordenboek van Nederlandsch-Indië*, Derde deel: R–Z. [Geographical and statistical dictionary of the Netherlands-India, Vol. 3: R–Z]. Amsterdam: Van Kampen.

Whitten, Tony; Roehayat Emon Soeriaatmadja; and Afiff, Suraya A, eds. 1996. *The Ecology of Java and Bali*. The Ecology of Indonesia Series, Vol. 2. Singapore: Periplus Editions.