TITLE:

Highly complex mitochondrial DNA genealogy in an endemic Japanese subterranean breeding brown frog *Rana tagoi* (Amphibia, Anura, Ranidae).

AUTHOR(S):

Eto, Koshiro; Matsui, Masafumi; Sugahara, Takahiro; Tanaka-Ueno, Tomoko

CITATION:

Eto, Koshiro ...[et al]. Highly complex mitochondrial DNA genealogy in an endemic Japanese subterranean breeding brown frog *Rana tagoi* (Amphibia, Anura, Ranidae).. Zoological science 2012, 29(10): 662-671

ISSUE DATE:

2012-10

URL:

<http://hdl.handle.net/2433/182953>

RIGHT:

© 2012 Zoological Society of Japan

Highly Complex Mitochondrial Dna Genealogy in an Endemic Japanese Subterranean Breeding Brown Frog *Rana Tagoi* (Amphibia, Anura, Ranidae)

Author(s): Koshiro Eto , Masafumi Matsui , Takahiro Sugahara and Tomoko Tanaka-Ueno

Source: Zoological Science, 29(10):662-671. 2012.

Published By: Zoological Society of Japan

DOI: <http://dx.doi.org/10.2108/zsj.29.662>

URL: <http://www.bioone.org/doi/full/10.2108/zsj.29.662>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

Highly Complex Mitochondrial DNA Genealogy in an Endemic Japanese Subterranean Breeding Brown Frog *Rana tagoi* (Amphibia, Anura, Ranidae)

Koshiro Eto¹, Masafumi Matsui^{1*}, Takahiro Sugahara², and Tomoko Tanaka-Ueno³

¹Graduate School of Human and Environmental Studies, Kyoto University, Kyoto 606-8501, Japan

²Hazama-cho 1740-1, Hachioji, Tokyo 193-0941, Japan

³Center for Liberal Arts, Meiji Gakuin University, Kamikurata 1518, Totsuka, Yokohama 244-8539, Japan

The endemic Japanese frog *Rana tagoi* is unique among Holarctic brown frogs in that it breeds in small subterranean streams. Using mitochondrial 16S ribosomal RNA and NADH dehydrogenase subunit 1 genes, we investigated genealogical relationships among geographic samples of this species together with its relative *R. sakuraii*, which is also a unique stream breeder. These two species together form a monophyletic group, within which both are reciprocally paraphyletic. *Rana tagoi* is divided into two major clades (Clade A and B) that are composed of 14 genetic groups. *Rana sakuraii* is included in Clade A and split into two genetic groups, one of which forms a clade (Subclade A-2) with sympatric *R. tagoi*. This species-level paraphyly appears to be caused by incomplete taxonomy, in addition to introgressive hybridization and/or incomplete lineage sorting. *Rana tagoi* strongly differs from other Japanese anurans in its geographic pattern of genetic differentiation, most probably in relation to its unique reproductive habits. Taxonomically, *R. tagoi* surely includes many cryptic species.

Key words: *Rana tagoi*, Japan, mtDNA, paraphyly, cryptic species, subterranean breeding, genetic divergence

INTRODUCTION

The genus *Rana* historically represented a very large group of frogs that occurred almost worldwide (Boulenger, 1920; Frost, 1985; Dubois, 1992), but is now restricted to smaller number of Holarctic brown frogs (Frost et al., 2006) that are generally similar in adult morphology and ecology. Most congeners breed in still (lentic) waters, such as ponds and rice paddies (e.g., *R. temporaria* Linnaeus from Europe: Nöllert and Nöllert, 1992), and only a few (e.g., *R. graeca* Boulenger from Europe and *R. sauteri* Boulenger from Taiwan) in flowing (lotic) waters of open streams (Nöllert and Nöllert, 1992; Tanaka-Ueno et al., 1998). Compared with such species, Japanese *R. tagoi* Okada (type locality: restricted by Shibata [1988] to Kamitakara-mura, currently included in Takayama-shi, Gifu Prefecture) is unique in that it breeds in small underground streams (Maeda and Matsui, 1999). This subterranean breeding habit is highly specialized and is not known in any other congeneric species.

Rana tagoi is endemic to the main (Honshu, Shikoku, and Kyushu) and some adjacent, smaller (Yakushima, Oki, and Goto) islands of Japan. Eggs laid in subterranean streams are few in number and large in size, and once hatched tadpoles can metamorphose without feeding

(Maeda and Matsui, 1999). Such traits appear to be an adaptation to this unique breeding environment. Another brown frog, *R. sakuraii* Matsui and Matsui (type locality: Okutama-machi, Nishitama-gun, Tokyo Prefecture) occurs only on Honshu Island and breeds in wider open streams in mountain regions. Other than the difference in breeding environment, this species is generally similar to *R. tagoi* in morphology and ecology, and is thought to be a close relative of *R. tagoi*, having originated from a *R. tagoi*-like subterranean breeding ancestor (Maeda and Matsui, 1999).

Steep mountains that provide many streams and rivers occupy the larger part of the main islands of Japan. Reflecting this environmental trait, there are various amphibian species that are adapted to lotic environments (e.g., *Bufo torrenticola* Matsui; *Buergeria buergeri* [Temminck and Schlegel]). Recent extensive surveys have revealed high cryptic diversity in some lotic breeding salamanders of the genera *Hynobius* Tschudi and *Onychodactylus* Tschudi (Nishikawa et al., 2007; Yoshikawa et al., 2008). A similar situation is expected in the case of lotic breeding *R. tagoi*, as the species is unique among Japanese frogs in that it contains three distinct subspecies (*R. t. tagoi* from main islands of Japan, *R. t. okiensis* Daito from Oki Islands, and *R. t. yakushimensis* Nakatani and Okada from Yakushima Island). In addition, morphological, breeding ecological (Sugahara, 1990; Sugahara and Matsui, 1992, 1993, 1994, 1995, 1996, 1997), and karyological (Ryuzaki et al., 2006) variations reported within *R. t. tagoi* suggest that it includes cryptic species. Genetically, *R. tagoi* is also diversified as

* Corresponding author. Tel. : +81-75-753-6846;
Fax : +81-75-753-6846;
E-mail: fumi@zoo.zool.kyoto-u.ac.jp
doi:10.2108/zsj.29.662

shown by the analyses of allozymes (Nishioka et al., 1987) and mitochondrial DNA (mtDNA; Tanaka et al., 1994). In contrast, variations within *R. sakuraii* have been poorly studied.

These previous studies suggest the presence of phylogenetic and/or taxonomic problems in *R. tagoi*, while such information is lacking for *R. sakuraii*. To date, few studies (e.g., Ryuzaki et al., 2006) have compared a large number of samples from the entire distributional range of the two species, leaving the comprehensive patterns of intra- or inter-specific variations unresolved. In this study, we conducted a phylogenetic analysis using two mitochondrial genes, relatively conservative 16S ribosomal RNA (16S rRNA) and rapidly evolving NADH dehydrogenase subunit 1 (ND1; Mueller, 2006), to reveal patterns of genetic differentiation and genealogical relationships in terms of mtDNA among samples of *R. tagoi* and *R. sakuraii*.

MATERIALS AND METHODS

We collected 183 specimens of *R. t. tagoi*, including the topotypic population, from 145 localities covering its entire distributional range in Honshu, Shikoku, and Kyushu. The large and small types of *R. t. tagoi* from Kinki (Sugahara, 1990) were distinguished

according to the diagnosis of Sugahara and Matsui (1994). We also collected two specimens of *R. t. yakushimensis* from Yakushima Island and three specimens of *R. t. okiensis* from the Oki islands. Furthermore, we collected 19 specimens of *R. sakuraii*, including the topotype, from 16 localities in Honshu. Detailed sampling localities are shown in Fig. 1 and Table 1.

As outgroups, we used *R. tsushimensis* from Tsushima Islands, Japan, and *Lithobates sylvaticus* from Quebec, Canada. The latter species is morphologically and ecologically similar to members of the genus *Rana*, but has been placed recently in another ranid genus, *Lithobates* (Frost et al., 2006).

Total DNA was extracted from frozen or ethanol-preserved tissues by standard phenol-chloroform extraction procedures (Hillis et al., 1996). Fragments containing the entire 16S rRNA and ND1 sequences, approximately 2.9 kb long, were amplified by polymerase chain reaction (PCR). The PCR cycle included an initial heating at 94°C for 4 min; 33 cycles of 94°C (30 s), 50°C (30 s), and 72°C (2 min 30 s); and a final extension at 72°C for 7 min. The amplified PCR products were purified by polyethylene glycol (PEG) precipitation procedures. The cycle sequence reactions were carried out with ABI PRISM Big Dye Terminator v3.1 Cycle sequencing Kit (Applied Biosystems) and sequencing was performed on an ABI 3130 automated sequencer. We used the primers listed in Table 2 to amplify and sequence the fragments, and all samples were sequenced in both directions. The obtained sequences were depos-

Fig. 1. Map of Japan showing sampling localities of *Rana t. tagoi* (circles), *R. t. yakushimensis* (double circle), *R. t. okiensis* (stars), and *R. sakuraii* (triangles). Squares indicate localities with sympatry of *R. t. tagoi* and *R. sakuraii*. For names of localities and genetic groups, see Table 1.

Table 1. Samples used for mtDNA analysis in this study with the information of voucher and collection locality. KUHE: Graduate School of Human and Environmental Studies, Kyoto University; TMP: Temporary numbered; UN: Unnumbered.

Sample no	Locality	genetic group	Voucher (KUHE)	GenBank		Sample no	Locality	genetic group	Voucher (KUHE)	GenBank	
				16S rRNA	ND1					16S rRNA	ND1
Rana tagoi tagoi											
1	Mutsu-shi, Aomori Pref.	A-1a	44827	AB639413	AB639593	49a	Sasayama-shi, Hyogo Pref.	A-1b	10285	AB639468	AB639638
2	Goshogawara-shi, Aomori Pref.	A-1a	36949	AB639413	AB639594	49b			10307	AB639469	AB639639
3	Towada-shi, Akita Pref	A-1a	13932	AB639413	AB639603	50a	Asago-shi, Hyogo Pref.	A-1b	10319	AB639470	AB639642
4	Noda-mura, Iwate Pref.	A-1a	37028	AB639413	AB639595	50b		B-2a	36586	AB639471	AB639640
5	Kamaishi-shi, Iwate Pref.	A-1a	27750	AB639411	AB639596	51	Kobe-shi, Hyogo Pref.	A-1b	22647	AB639472	AB639641
6	Oshu-shi, Iwate Pref.	A-1a	32889	AB639413	AB639597	52	Wakasa-cho, Tottori Pref.	A-1b	34743	AB639473	AB639642
7	Ichinoseki-shi, Iwate Pref.	A-1a	35268	AB639412	AB639603	53	Nihonmatsu-shi, Fukushima Pref.	A-2	36330	AB639474	AB639643
8	Fujisawa-cho, Iwate Pref.	A-1a	36699	AB639413	AB639598	54	Hirono-machi, Fukushima Pref.	A-2	44829	AB639475	AB639644
9	Senboku-shi, Akita Pref.	A-1a	27351	AB639413	AB639603	55	Kitabaraki-shi, Ibaraki Pref.	A-2	27544	AB639476	AB639645
10	Ishinomaki-shi, Miyagi Pref.	A-1a	41545	AB639414	AB639603	56	Hitachi-shi, Ibaraki Pref.	A-2	27550	AB639477	AB639646
11	Sendai-shi, Miyagi Pref.	A-1a	37121	AB639415	AB639599	57	Hitachiomiya-shi, Ibaraki Pref.	A-2	43711	AB639478	AB639647
12	Sakata-shi, Yamagata Pref.	A-1a	37544	AB639416	AB639600	58a	Tsukuba-shi, Ibaraki Pref.	A-2	42747	AB639479	AB639648
13	Yamagata-shi, Yamagata Pref.	A-1a	37543	AB639417	AB639601	58b			42751	AB639480	AB639649
14	Kaminoyama-shi, Yamagata Pref.	A-1a	29360	AB639420	AB639602	59	Mashiko-machi, Tochigi Pref.	A-2	25968	AB639481	AB639650
15	Nishikawa-machi, Yamagata Pref.	A-1a	37548	AB639418	AB639603	60a	Akiruno-shi, Tokyo Pref.	A-2	42452	AB639483	AB639651
16	Nihonmatsu-shi, Fukushima Pref.	A-1a	29595	AB639419	AB639604	61	Ichihara-shi, Chiba Pref.	A-2	28409	AB639482	AB639652
17	Shirakawa-shi, Fukushima Pref.	A-1a	21629	AB639420	AB639605	62	Otsuki-shi, Yamanashi Pref.	A-2	28064	AB639483	AB639653
18a	Daigo-machi, Ibaraki Pref.	A-1a	42344	AB639420	AB639605	63a	Izu-shi, Shizuoka Pref.	A-2	36715	AB639484	AB639654
18b		A-2	43886	AB639421	AB639646	63b			43468	AB639485	AB639655
19	Nikko-shi, Tochigi Pref.	A-1a	36719	AB639426	AB639609	64	Fuji-shi, Shizuoka Pref.	A-2	43473	AB639486	AB639656
20a	Kanuma-shi, Tochigi Pref.	A-1a	40166	AB639422	AB639609	65	Nakanojo-machi, Gunma Pref.	A-4	22930, 22936	AB639487	AB639657
21	Minakami-machi, Gunma Pref.	A-1a	27539	AB639429	AB639612	66	Nagano-shi, Nagano Pref.	A-5	18005	AB639488	AB639658
22	Nakanojo-machi, Gunma Pref.	A-1a	27930	AB639424	AB639606	67	Hokuto-shi, Yamanashi Pref.	A-5	43483	AB639489	AB639659
23	Shibukawa-shi, Gunma Pref.	A-1a	29485	AB639425	AB639607	68a	Gujo-shi, Gifu Pref.	A-5	14228	AB639490	AB639660
24	Agano-shi, Niigata Pref.	A-1a	29600	AB639426	AB639608	68b			44832	AB639491	AB639661
25	Aga-machi, Niigata Pref.	A-1a	UN	AB639426	AB639609	69	Hayakawa-cho, Yamanashi Pref.	A-6	14208	AB639492	AB639662
26	Yahiko-mura, Niigata Pref.	A-1a	27765	AB639427	AB639610	70	Fujikawaguchiko-machi, Yamanashi Pref.	A-6	43480	AB639493	AB639663
27	Kashiwazaki-shi, Niigata Pref.	A-1a	36892	AB639428	AB639611	71a	Shizuoka-shi, Shizuoka Pref.	A-6	42977	AB639494	AB639664
28	Uonuma-shi, Niigata Pref.	A-1a	36896	AB639429	AB639612	71b			24561	AB639495	AB639665
29	Otari-mura, Nagano Pref.	A-1a	43367	AB639430	AB639613	72	Shizuoka-shi, Shizuoka Pref.	A-6	29933	AB639496	AB639666
30	Ueda-shi, Nagano Pref.	A-1a	18752	AB639431	AB639614	73	Kawanehon-cho, Shizuoka Pref.	A-6	42270	AB639497	AB639667
31	Kiso-machi, Nagano Pref.	A-1a	43382	AB639432	AB639615	74	Fujieda-shi, Shizuoka Pref.	A-6	17955	AB639498	AB639668
32	Hodatsushimizu-cho, Ishikawa Pref.	A-1a	41053	AB639433	AB639616	75	Kakegawa-shi, Shizuoka Pref.	A-6	39980	AB639499	AB639669
33	Takayama-shi, Gifu Pref.	A-1a	27613, 43018	AB639434	AB639617	76	Neba-mura, Nagano Pref.	A-6	27335	AB639500	AB639670
34	Shirakawa-mura, Gifu Pref.	A-1a	26104	AB639435	AB639618	77	Shitara-cho, Aichi Pref.	A-6	27251	AB639501	AB639671
35a	Ibigawa-cho, Gifu Pref.	A-1a	27388	AB639436	AB639619	78a	Ise-shi, Mie Pref.	A-6	42829	AB639502	AB639672
36	Ikeda-cho, Fukui Pref.	A-1a	40441	AB639438	AB639624	78b			42830	AB639503	AB639672
37	Nagahama-shi, Shiga Pref.	A-1a	41470, 41471	AB639439	AB639621	79	Minamiawaji-shi, Hyogo Pref.	A-7	43885	AB639504	AB639673
38	Maibara-shi, Shiga Pref.	A-1a	37610, 37614	AB639440	AB639622	80	Manno-cho, Kagawa Pref.	A-7	TMP_T2882	AB639505	AB639674
39a	Taga-cho, Shiga Pref.	A-1a	41287	AB639440	AB639622	81	Kamiyama-cho, Tokushima Pref.	A-7	TMP_T2876	AB639506	AB639675
39b			41551	AB639441	AB639623	82	Saijo-shi, Ehime Pref.	A-7	27679	AB639507	AB639676
40	Nagahama-shi, Shiga Pref.	A-1a	40385	AB639442	AB639624	83	Imabari-shi, Ehime Pref.	A-7	27506	AB639508	AB639677
41a	Takashima-shi, Shiga Pref.	A-1a	TMP_T3395	AB639443	AB639625	84	Seiyo-shi, Ehime Pref.	A-7	TMP_T2241	AB639509	AB639678
41b			40437	AB639444	AB639625	85	Toyo-cho, Kochi Pref.	A-7	29464	AB639510	AB639679
41c			TMP_T3402	AB639445	AB639625	86	Kochi-shi, Kochi Pref.	A-7	36184	AB639511	AB639680
41d		A-1b	TMP_T3392	AB639446	AB639626	87	Kitakyushu-shi, Fukuoka Pref.	A-9a	28614	AB639512	AB639681
42a	Takashima-shi, Shiga Pref.	A-1b	25993	AB639447	AB639627	88	Koga-shi, Fukuoka Pref.	A-9a	26841	AB639513	AB639682
42b		B-2a	43609	AB639448	AB639711	89	Fukuoka-shi, Fukuoka Pref.	A-9a	26238	AB639514	AB639683
42c		B-2a	25996	AB639453	AB639711	90	Yame-shi, Fukuoka Pref.	A-9a	26643	AB639515	AB639684
43a	Otsu-shi, Shiga Pref.	A-1b	41414, 43428	AB639449	AB639628	91	Asakura-shi, Fukuoka Pref.	A-9a	27137	AB639516	AB639685
43b		B-2a	41090	AB639450	AB639629	92	Isahaya-shi, Nagasaki Pref.	A-9a	9660	AB639517	AB639686
43c			43148	AB639451	AB639713	93	Sasebo-shi, Nagasaki Pref.	A-9a	27140	AB639518	AB639687
44a	Nantan-shi, Kyoto Pref.	A-1b	41408	AB639452	AB639630	94	Beppu-shi, Oita Pref.	A-9a	43637	AB639519	AB639688
44b		B-2a	41406	AB639453	AB639711	95	Bungo-ohno-shi, Oita Pref.	A-9a	27146	AB639520	AB639694
44c			41426	AB639457	AB639713	96	Kokonoe-machi, Oita Pref.	A-9a	26148	AB639521	AB639689
45a	Kyoto-shi, Kyoto Pref.	A-1b	43324	AB639457	AB639635	97	Gokase-cho, Miyazaki Pref.	A-9a	44834	AB639522	AB639690
45b		B-2a	41730	AB639458	AB639633	98	Ebino-shi, Miyazaki Pref.	A-9a	41284	AB639523	AB639694
45c			38698	AB639459	AB639634	99	Yatsushiro-shi, Kumamoto Pref.	A-9a	27562	AB639524	AB639691
46a	Kyoto-shi, Kyoto Pref.	A-1b	42034, 44828	AB639460	AB639635	100	Amakusa-shi, Kumamoto Pref.	A-9a	30342	AB639525	AB639692
46b		B-2a	44835	AB639462	AB639711	101	Soo-shi, Kagoshima Pref.	A-9a	42191	AB639526	AB639693
46c			42396	AB639463	AB639711	102	Izumi-shi, Kagoshima Pref.	A-9a	27564	AB639527	AB639694
46d			41439	AB639461	AB639713	103	Kanoya-shi, Kagoshima Pref.	A-9a	27295, 43404	AB639530	AB639697
46e			42319	AB639464	AB639712	104	Nobeoka-shi, Miyazaki Pref.	A-9b	27121	AB639528	AB639695
47a	Toyooka-shi, Hyogo Pref.	A-1b	25664	AB639465	AB639636	105	Nishimera-son, Miyazaki Pref.	A-9b	26088	AB639529	AB639696
47b		B-2a	25662	AB639564	AB639729	106	Aya-cho, Miyazaki Pref.	A-9b	42194	AB639531	AB639698
48a	Toyooka-shi, Hyogo Pref.	A-1b	42711	AB639466	AB639637	107	Miyakonojo-shi, Miyazaki Pref.	A-9b	30907	AB639532	AB639699
48b		B-2a	42714	AB639467	AB639729	Continued					

Cryptic Diversity of Two *Rana* Species

665

Table 1. Continued

Sample no	Locality	genetic group	Voucher (KUHE)	GenBank		Sample no	Locality	genetic group	Voucher (KUHE)	GenBank	
				16S rRNA	ND1					16S rRNA	ND1
108	Kimotsuki-cho, Kagoshima Pref.	A-9b	43397	AB639533	AB639700	141	Hatsukaichi-shi, Hiroshima Pref.	B-2b	UN	AB639571	AB639736
109a	Kanoya-shi, Kagoshima Pref.	A-9b	43401	AB639534	AB639701	142	Higashihiroshima-shi, Hiroshima Pref.	B-2b	30262	AB639572	AB639737
109b			43403	AB639535	AB639702	143	Higashihiroshima-shi, Hiroshima Pref.	B-2b	30220	AB639573	AB639738
110a	Kinko-cho, Kagoshima Pref.	A-9b	27678	AB639536	AB639703	144	Hagi-shi, Yamaguchi Pref.	B-2b	42848	AB639574	AB639739
110b			41250	AB639537	AB639704	145	Shimonoseki-shi, Yamaguchi Pref.	B-2b	34516	AB639575	AB639740
111	Goto-shi, Nagasaki Pref.	A-9c	31539	AB639538	AB639705	<i>R. t. yakushimensis</i>					
112a	Taga-cho, Shiga Pref.	B-2a	43508	AB639539	AB639706	146a	Yakushima-cho, Kagoshima Pref.	A-8	10182	AB639578	AB639741
112b			43509	AB639540	AB639707	146b			43326	AB639577	AB639741
113	Konan-shi, Shiga Pref.	B-2a	18763	AB639541	AB639708	<i>R. t. okiensis</i>					
114a	Koka-shi, Shiga Pref.	B-2a	28466	AB639542	AB639709	147a	Okinoshima-cho, Shimane Pref.	B-1	10818	AB639576	AB639742
115	Kyotango-shi, Kyoto Pref.	B-2a	24566	AB639544	AB639729	147b			22341	AB639579	AB639742
116	Maizuru-shi, Kyoto Pref.	B-2a	TMP_T3345	AB639545	AB639711	148	Nishinoshima-cho, Shimane Pref.	B-1	43647	AB639580	AB639742
117a	Kyoto-shi, Kyoto Pref.	B-2a	27168	AB639546	AB639712	<i>R. sakuraii</i>					
117b			41431	AB639547	AB639714	20b	Kanuma-shi, Tochigi Pref.	A-2	43635	AB639423	AB639744
118	Kameoka-shi, Kyoto Pref.	B-2a	41553	AB639548	AB639713	35b	Ibigawa-cho, Gifu Pref.	A-3	36297	AB639437	AB639620
119	Joyo-shi, Kyoto Pref.	B-2a	41554	AB639549	AB639714	44d	Nantan-shi, Kyoto Pref.	A-3	UN	AB639454	AB639631
120	Komono-cho, Mie Pref.	B-2a	26744	AB639550	AB639715	44e			41412	AB639455	AB639632
121	Matsuzaka-shi, Mie Pref.	B-2a	41484	AB639551	AB639716	44f			41413	AB639456	AB639632
122	Owase-shi, Mie Pref.	B-2a	26990	AB639552	AB639717	60b	Akiruno-shi, Tokyo Pref.	A-2	42450	AB639583	AB639744
123a	Odai-cho, Mie Pref.	B-2a	40190	AB639553	AB639718	114b	Koka-shi, Shiga Pref.	A-3	TMP_T2666	AB639543	AB639710
124	Izumi-shi, Osaka Pref.	B-2a	TMP_T3425	AB639556	AB639721	123b	Odai-cho, Mie Pref.	A-3	27647	AB639554	AB639719
125	Soni-mura, Nara Pref.	B-2a	24435	AB639557	AB639722	123c			40309	AB639555	AB639720
126	Sakurai-shi, Nara Pref.	B-2a	18893	AB639558	AB639723	149	Naganohara-machi, Gunma Pref.	A-2	27937	AB639581	AB639744
127	Kudoyama-cho, Wakayama Pref.	B-2a	24546	AB639559	AB639724	150	Chichibu-shi, Saitama Pref.	A-2	43736	AB639582	AB639743
128	Hongu-cho, Wakayama Pref.	B-2a	26784	AB639560	AB639725	151	Okutama-machi, Tokyo Pref.	A-2	UN	AB639583	AB639744
129	Shingu-shi, Wakayama Pref.	B-2a	24540	AB639561	AB639726	152	Kiyokawa-mura, Kanagawa Pref.	A-2	14276	AB639584	AB639745
130	Gobo-shi, Wakayama Pref.	B-2a	41229	AB639562	AB639727	153	Matsumoto-shi, Nagano Pref.	A-2	22887	AB639585	AB639746
131	Kami-cho, Hyogo Pref.	B-2a	43603	AB639563	AB639728	154	Fujikawa-cho, Yamanashi Pref.	A-2	43481	AB639586	AB639747
132	Taka-cho, Hyogo Pref.	B-2a	10330	AB639564	AB639729	155	Itoigawa-shi, Niigata Pref.	A-3	31300	AB639587	AB639748
133	Sayo-cho, Hyogo Pref.	B-2a	41021	AB639565	AB639730	156	Hamamatsu-shi, Shizuoka Pref.	A-3	UN	AB639588	AB639749
134	Kamigori-cho, Hyogo Pref.	B-2a	41022	AB639566	AB639731	157	Nakatsugawa-shi, Gifu Pref.	A-3	18201	AB639589	AB639749
135	Mimasaka-shi, Okayama Pref.	B-2a	27659	AB639567	AB639732	158	Iwakuni-shi, Yamaguchi Pref.	A-3	43893	AB639590	AB639750
136	Misasa-cho, Tottori Pref.	B-2b	24574	AB639568	AB639733	<i>R. tsushimensis</i>					
137	Daisen-cho, Tottori Pref.	B-2b	36824	AB639569	AB639734		Tsushima-shi, Nagasaki Pref.		11606	AB639592	AB639752
138	Unnan-shi, Shimane Pref.	B-2b	18877	AB639570	AB639735	<i>Lithobates sylvaticus</i>					
139a	Shobara-shi, Hiroshima Pref.	B-2b	36037				Quebec, Canada		UN	AB639591	AB639751
139b			36040								
140	Shobara-shi, Hiroshima Pref.	B-2b	24553								

ited in GenBank (Table 1).

Sequences obtained were aligned using Clustal W (Thompson et al., 1994), and gaps and ambiguous areas were excluded from alignments using Gblocks 0.91b (Castresana, 2000) with default settings. We then constructed phylogenetic trees from the combined alignments using maximum parsimony (MP), maximum likelihood (ML), and Bayesian inference (BI). The MP analysis was performed using PAUP*4.0b10 (Swofford, 2002). We used a heuristic search with the tree bisection and reconnection (TBR) branch-swapping algorithm and 100 random additions replicates, and the number of saved trees was restricted to 5,000. Transitions and transversions were equally weighted. The ML and BI analyses were respectively performed using TREEFINDER ver. Oct. 2008 (Jobb, 2008) with Phylogens 1.5.2010.03.24 (Tanabe, 2008) and MrBayes 3.1.2 (Huelsenbeck and Ronquist, 2001). Different substitution models were applied for each gene partition in both of the analyses. The optimum substitution model for each gene was selected by using Kakusan4 (Tanabe, 2010), based on the Akaike information criterion (AIC). The best model was calculated for each codon position (1st, 2nd, and 3rd positions) of the ND1 genes. In the BI analysis, two independent runs of four Markov chains were conducted for 7,000,000 generations (sampling fre-

Table 2. Primers used to amplify mtDNA in this study.

Target	Name	Sequence	Reference
16S	L1507	TACACACCGCCCGTCACCTCTT	Shimada et al. (2011)
	H1923	AAGTAGCTCGCTTAGTTTCGG	Shimada et al. (2011)
	L1879	CGTACCTTTTGCATCATGGTC	Shimada et al. (2011)
	H2315	TTCTTGTTACTAGTTCTAGCAT	Shimada et al. (2011)
	L2188	AAAGTGGGCCTAAAAGCAGCCA	Matsui et al. (2006)
	Wilkinson_6	CCCTCGTGATGCCGTTGATAC	Wilkinson et al. (2002)
	16L1	CTGACCGTGCAAAGGTAGCGTAATCACT	Hedges (1994)
	16H1	CTCCGGTCTGAACTCAGATCACGTAGG	Hedges (1994)
	L3032	CGACCTCGATGTTGGATCAGG	Shimada et al. (2011)
	ND1_Htago	GRGCRATTTGGAGTTTGARGCTCA	this study
ND1	ND1_Ltago	GACCTAAACCTCAGYATYCTATTTAT	this study
	tMet_H	AGGAAGTACAAAGGGTTTGTATC	Shimada et al. (2011)

quency: one tree per 100 generations). We used TRACER v. 1.4 (Rambaut and Drummond, 2007) to determine the burn-in size and when the log likelihood of sampled trees reached stationary distribution, and the first 7,001 trees were discarded (burn-in = 700,000).

The robustness of the MP and ML trees were tested using non-parametric bootstrap analysis (Felsenstein 1985) with 1,000 replicates. We regarded tree topologies with bootstrap value (BS) 70% or greater as sufficiently supported (Huelsenbeck and Hillis, 1993). For the BI, we regarded Bayesian posterior probability (BPP) 0.95 or greater as significant support (Huelsenbeck and Ronquist, 2001;

Leaché and Reeder 2002). Uncorrected p-distances for each gene were also calculated using PAUP* ver. 4.0b10.

RESULTS

Sequences and statistics

We obtained complete 16S rRNA (1,625 bp long) and ND1 (973 bp) sequences from 207 individuals and two out-group taxa. After excluding gaps and ambiguous areas, a combined 2,521 nucleotide sites, of which 624 were variable

Table 3. Alignment statistics for total 16S rRNA and ND1. The number of base pairs (bp), variable sites (vs), number of parsimony informative sites (pi), and transition-transversion ratio (ti/tv) are given for ingroups only.

	bp	vs	pi	ti/tv
16SrRNA	1554	310	206	6.65
ND1	967	314	250	9.38
Combined	2521	624	456	8.04

Fig. 2. Bayesian tree of total 16S rRNA and ND1 mitochondrial genes for three subspecies of *R. tagoi*, *R. sakuraii*, and outgroup taxa. Nodal values indicate bootstrap supports for MP and ML, and Bayesian posterior probability (MP-BS/ML-BS/BPP). Asterisks indicate nodes with MP-BS and ML-BS = 70% and BPP = 0.95. For locality numbers, see Table 1 and Fig. 1.

and 456 parsimoniously informative (Table 3), were used for phylogenetic analysis. We detected 190 haplotypes within the ingroup, of which 168 were in *R. t. tagoi*, two in *R. t. yakushimensis*, three in *R. t. okiensis*, and 17 in *R. sakuraii*.

The MP analysis produced 5,000 equally most parsimonious trees ($L = 2007$, $CI = 0.519$, $RI = 0.901$). For the ML analysis, the best substitution model of 16S rRNA estimated by Kakusan 4 was J2 model with a Gamma (G) shape parameter. In ND1, Hasegawa-Kishino-Yano-1985 (HKY85) model + G, HKY85 + G, and J2 + G were selected for the 1st, 2nd, and 3rd codon positions, respectively. For the BI analysis, the general time reverse (GTR) model + G was selected for 16S rRNA. In ND1, HKY85 + G, HKY85 + G, and GTR + G were selected for the 1st, 2nd, and 3rd codon positions, respectively. The likelihood values ($-\ln L$) of the ML and BI trees were 14439.77 and 15102.97, respectively.

Phylogenetic relationships

The ML and BI analyses yielded essentially identical topologies. The MP tree was also similar to these, although support values tended to be lower. The BI tree is shown in Fig. 2. *Rana tagoi* and *R. sakuraii* formed a fully supported monophyletic group, but both were paraphyletic with respect to each other. The ingroup was divided into two major clades, Clade A (MP-BS = 79%, ML-BS = 83%, BPP = 0.99) and Clade B (98%, 87%, 1.00, respectively), with uncorrected p-distances of 2.1% to 3.9% in 16S rRNA and 4.9% to 8.5% in ND1 between them. Each clade contained several subclades, some of which were further divided into two or three groups. Sequence divergences as measured by the mean uncorrected p-distances among these subclades and groups are shown in Table 4.

Clade A, which contained a subset of *R. t. tagoi*, *R. t. yakushimensis*, and *R. sakuraii* samples, was divided into nine subclades (Subclade A-1 to A-9). Subclade A-1 (94%, 98%, 1.00) contained *R. t. tagoi* samples from Tohoku, northern Chubu, and northern Kinki regions. This subclade was divided into two groups, Group A-1a (97%, 99%, 1.00) and A-1b (96%, 99%, 1.00), with sequence divergences of 0.9% to 1.9% in 16S rRNA and 3.3% to 4.9% in ND1 between them.

Group A-1a contained *R. t. tagoi* from Tohoku, northern Chubu, and northeastern Kinki (localities 1 to 41), including topotypic samples (locality 33) and a part of the *R. t. tagoi* large type (Sugahara, 1990) (locality 41). Except for samples from localities 11 to 13, which were divergent from the others, genetic variation within Group A-1a was small, despite its wide range of distribution. Group A-1b consisted of all samples of the *R. t. tagoi* small type from northern Kinki (localities 41 to 52). Within this group, genetic variation among haplotypes was significant, and four divergent subgroups were recognized.

Table 4. Mean uncorrected p-distances (%) among genetic groups of three subspecies of *R. tagoi* and *R. sakuraii* for 16S rRNA (above diagonal) and ND1 (below diagonal). Darkly shaded areas indicate distances among groups with sympatric distribution and lightly shaded areas indicate distances among groups with parapatric distribution.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
1. A-1a	—	1.3	1.6	1.9	1.6	1.4	1.5	1.9	1.9	1.7	1.8	2.0	1.7	3.0	3.3	3.0
2. A-1b	4.1	—	1.8	2.1	1.7	1.6	1.6	1.9	2.1	1.6	2.0	2.3	1.8	2.8	3.2	2.9
3. A-2 (<i>R. t. tagoi</i>)	3.9	4.3	—	1.1	1.7	1.3	1.6	1.9	1.9	1.7	1.9	2.2	1.8	2.9	3.1	2.8
4. A-2 (<i>R. sakuraii</i>)	4.2	4.2	2.1	—	1.9	1.4	1.8	2.0	2.1	1.9	2.1	2.4	2.1	3.0	3.1	2.8
5. A-3	4.7	5.0	3.7	3.6	—	1.4	1.5	1.9	2.1	1.8	1.9	2.2	1.8	2.7	3.0	2.6
6. A-4	3.9	4.3	3.0	3.3	4.1	—	1.3	1.7	1.8	1.5	1.7	2.0	1.6	2.7	2.8	2.6
7. A-5	4.4	4.9	3.5	3.8	4.4	3.0	—	1.5	2.1	1.7	2.0	2.3	1.9	2.9	3.1	2.8
8. A-6	5.0	4.7	4.3	4.3	5.4	3.4	4.1	—	2.3	2.0	2.3	2.5	2.1	2.9	3.3	2.8
9. A-7	4.7	5.2	3.8	4.4	5.1	4.1	4.7	5.3	—	1.8	2.0	2.4	1.9	3.0	3.1	2.7
10. A-8	4.0	4.4	3.1	3.4	3.9	2.8	3.6	4.0	3.1	—	1.8	2.1	1.6	2.7	2.8	2.7
11. A-9a	5.2	5.3	4.2	4.4	5.4	4.4	4.8	5.4	4.4	3.4	—	1.7	1.3	3.2	3.2	2.9
12. A-9b	4.7	5.1	3.8	4.1	4.8	3.7	4.2	4.9	4.1	2.8	3.0	—	1.4	3.3	3.5	3.1
13. A-9c	5.0	5.2	4.1	3.9	5.0	4.0	4.7	5.3	4.1	3.3	3.2	2.9	—	2.9	3.1	2.8
14. B-1	6.1	6.4	5.6	6.0	6.5	5.3	5.9	6.1	5.9	5.4	6.3	6.2	6.4	—	2.0	1.8
15. B-2a	6.9	6.7	5.9	6.3	6.5	5.9	6.3	6.7	6.3	5.8	6.7	6.6	6.6	4.1	—	1.3
16. B-2b	7.0	6.9	5.7	6.1	7.0	5.9	6.5	6.6	6.3	6.0	6.4	6.4	6.4	4.4	2.9	—

Subclade A-2 (96%, 99%, 1.00) contained *R. t. tagoi* from Kanto region (localities 18 and 53 to 64) and was divided into two divergent groups. Interestingly, *R. sakuraii* from eastern Honshu (localities 20, 60, and 149 to 154), including topotypic samples (locality 151), was completely embedded in one of these groups. Within Subclade A-2, *R. sakuraii* was not much divergent from *R. t. tagoi* (0.8% to 1.3% in 16S; 1.3% to 3.0% in ND1).

Subclade A-3 (99%, 99%, 1.00) contained *R. sakuraii* from western Honshu (localities 35, 44, 114, 123, and 155 to 158), and was divided into three groups. Subclades A-2 and A-3 tended to form a clade, but their monophyly was not supported (< 50%, 66%, 0.86).

Subclade A-4 contained only one sample of *R. t. tagoi* from Nakanojo-machi (former Kuni-mura), Gunma (locality 65), while Subclade A-5 (78%, 75%, 1.00) contained divergent haplotypes of *R. t. tagoi* from central Chubu (localities 66 to 68). Subclade A-6 (all 100%, or 1.00) contained *R. t. tagoi* from southern Chubu (localities 69 to 77) and Shima Peninsula (locality 78), where variation among haplotypes was small. This subclade included *R. t. tagoi* with $2n = 28$ chromosomes (vs. $2n = 26$ chromosomes in *R. tagoi* samples from other localities so far studied) from Neba-mura, Nagano (Ryuzaki et al., 2006; locality 76). Subclades A-4 to A-6 tended to form a clade, but their monophyly was not unambiguously supported (< 50%, 68%, 1.00). Subclades A-1 to A-6 also tended to form a clade, but the MP support of this node was low (< 50%, 77%, 1.00).

Subclade A-7 (99%, 99%, 1.00) contained *R. t. tagoi* from Shikoku (localities 80 to 86) and Awaji Island (locality 79), with small genetic variations within the group. Subclade A-8 (all 100%, or 1.00) contained *R. t. yakushimensis* from Yakushima Island (locality 146), and was close to Subclade A-7, although their monophyly was not supported (< 50%, 63%, 0.54).

Subclade A-9 (90%, 99%, 1.00) contained *R. t. tagoi* from Kyushu and tended to form a clade with A-7 and A-8 but their monophyly was not supported (< 50%, 66%, 0.98). Subclade A-9 was divided into three groups, Groups A-9a (99%, 100%, 1.00), A-9b (93%, 94%, 1.00), and A-9c (only

one sample) with divergences between them being 1.3% to 1.7% in 16S, and 2.9% to 3.2% in ND1. Group A-9a contained samples from northwestern Kyushu (localities 87 to 103), and genetic variation within the group was small. Group A-9b consisted of samples from southern Pacific side of the island (localities 104 to 110) and was divided into two subgroups. Group A-9c contained one sample from Narujima Island (locality 111).

Clade B contained *R. t. okiensis* and a part of *R. t. tagoi* samples and was divided into two subclades. One of them, Subclade B-1 (all 100% or 1.00) contained *R. t. okiensis* from Oki islands (localities 147 and 148), while another, Subclade B-2 (99%, 95%, 1.00), consisted of *R. t. tagoi* from western Honshu. Two groups, with divergences of 0.8% to 1.6% in 16S rRNA and 2.1% to 4.0% in ND1, were recognized within this subclade; Group B-2a (99%, 95%, 1.00) and Group B-2b (88%, 69%, 1.00). Group B-2a contained samples from Kinki (localities 42 to 48, 50, and 112 to 135) and was divided into three subgroups. A large portion of the *R. t. tagoi* large type (Sugahara, 1990) samples (localities 42 to 48 and 50) was included in this group. Group B-2b contained samples from Chugoku (localities 136 to 145) and was divided into two subgroups.

Geographic distribution of genetic groups

Genetic groups recognized in two major clades of *R. tagoi* (sensu lato) and *R. sakuraii* (totally 15 subclades/groups) showed a complex pattern of geographic distribution, with sympatric or parapatric occurrence in some (Figs. 1, 3 and Table 4). Only *R. t. yakushimensis* (A-8), *R. t. okiensis* (B-1), *R. t. tagoi* from Awaji Island and Shikoku (A-7), and *Rana t. tagoi* from Kyushu (A-9a, b, and c) were allopatric from the other genetic groups, although A-9a and A-9b were parapatric within Kyushu.

Rana t. tagoi Group A-1a was widely distributed throughout northeastern Honshu to the northern part of central Honshu. It was transposed by *R. t. tagoi* Groups A-1b and B-2a in northeastern Kinki, the westernmost area of its distributional range. Group A-1a and A-1b were parapatric, with the exception of one sympatric site (locality 41). Group A-1b was distributed in northern part of Kinki, and was sympatric with B-2a in almost all ranges of its distribution (localities 42 to 48 and 50).

Group A-1a was transposed by *R. t. tagoi* in Subclade A-2 in northern Kanto. They were mostly parapatric, but were sympatric in one site (locality 18). *Rana t. tagoi* in Subclade A-2 was replaced by Subclade A-6 (southern Chubu) in western Kanto. Subclades A-4 and A-5 occurred in northwestern Kanto to central Chubu, between Group A-1a in the Sea of Japan side and Subclade A-6 in the Pacific side. Subclade A-4 was sympatric with A-1a, and A-5 also seemed to overlap with A-1a. Subclade A-6 widely occurred covering southern Chubu, and was replaced by Group B-2a

Fig. 3. Distributional range of each genetic group of *Rana tagoi* (solid line) and *R. sakuraii* (dotted line). For names of genetic groups, see Table 1 and Fig. 2.

in the Shima Peninsula (locality 78).

Group B-2a of *R. t. tagoi* from Kinki, which was sympatric with the *R. t. tagoi* small type (A-1b) as shown above, was transposed in the west by B-2b, which widely occurred in Chugoku, western Honshu.

Rana sakuraii was divided into two genetic groups, eastern (A-2) and western (A-3) subclades. In western Kanto, *R. sakuraii* was sympatric with *R. t. tagoi* and together formed Subclade A-2. Also, in the northern part of its distribution, *R. sakuraii* in Subclade A-2 was sympatric with *R. t. tagoi* A-1a (locality 20) and parapatric with A-4 (localities 160 and 67), and furthermore, seemed to overlap with A-5 in central Chubu. Subclade A-2 was transposed by *R. sakuraii* Subclade A-3 in the most western range of its distribution. Subclade A-3 largely overlapped with *R. t. tagoi* genetic groups in western Honshu (e.g., A-5, A-6, and B-2b), and sympatric with A-1a (locality 35), A-1b (locality 44), and B-2a (localities 44 and 114).

DISCUSSION

Phylogenetic relationships and genetic differentiation

Using allozymes and proteins, Nishioka et al. (1987) constructed a phenogram in which *R. t. yakushimensis* (A-8 in this study) was shown to be divergent from *R. t. tagoi* from western Japan. Within the latter, populations from Kinki (B-2a), Chugoku (B-2b), and Shikoku (A-7) formed one group, and those from Kyushu (A-9a) and *R. t. okiensis* (B-1) formed another. These results are completely discordant with results obtained by us or by Tanaka et al. (1996) from the mitochondrial *cyt b* gene. Our results showed common features with those reported by Tanaka et al. (1994, 1996: i.e., paraphyly of *R. tagoi*; large differentiation between large [B-2a] and small [A-1b] types of *R. t. tagoi* from Kyoto). Although there are superficial differences between Tanaka et al. (1994, 1996) and the present study, in the relationships of *R. t. tagoi*, *R. t. yakushimensis*, and *R. t. okiensis*, such discrepancies surely resulted from insufficient sampling in the Tanaka et al. (1994, 1996) study (e.g., Tanaka et al. [1996] used seven samples from five localities of *R. t. tagoi*, one sample of *R. t. yakushimensis*, three samples of *R. t. okiensis*, and six samples from three localities of *R.*

sakuraii), and results obtained from mtDNA analyses are considered essentially similar.

Discordance between trees based on nuclear (i.e., allozymes) and mitochondrial markers is generally explained by the paralogy of genes, introgressive hybridization, and incomplete lineage sorting with ancestral polymorphism (Ballard and Whitlock, 2004). However, these factors are difficult to differentiate without additional studies, in which nuclear marker analyses are made on the samples used in the present mtDNA analysis. In contrast to mitochondrial genes, allozymes are of limited value in estimating phylogenies, as historical relationships among alleles remain unclear (Avice, 2000). Thus, phylogenetic trees based on mitochondrial genes should be more valid than the allozymic ones, although the possibility of mitochondrial gene introgression, which leads to a strongly biased gene tree, is not precluded.

The geographic pattern of genetic differentiations obtained for *R. tagoi* is quite unique among Japanese anurans in that samples from western Honshu (Clade B) first diverge from the others (Clade A). In wide-ranging Japanese anurans (e.g., *Bufo japonicus*: Matsui, 1984; Igawa et al., 2006; *R. japonica*: Sumida and Ogata, 1998; *R. rugosa*: Sekiya et al., 2010; *Buergeria buergeri*: Nishizawa et al., 2011), populations from western Honshu and those from Shikoku and Kyushu tend to form a clade, unlike in *R. tagoi*, in which populations from eastern to central Honshu, Shikoku, and Kyushu form a clade (Clade A). This unique distribution suggests that geographical and environmental factors that separated Clades A and B of *R. tagoi* differ from those that affected the distribution of other Japanese anurans. Our results do not contradict Matsui and Matsui's (1990) hypothesis that the probable common ancestor of *R. tagoi* and *R. sakuraii* would have a habit of subterranean breeding, which is quite unique among Japanese anurans. The availability of subterranean environments, which was not so critical in other anurans, may have been a major factor that caused population fragmentation and subsequent genetic divergence in the ancestor of *R. tagoi* and *R. sakuraii*.

The current wider distribution of Clade A compared to Clade B indicates the Clade A ancestor was dominant throughout Honshu, including Kinki and Chugoku, in the past, whereas Clade B now predominates. Later, ancestral Clade B appears to have arisen somewhere in western Honshu and expanded its range towards east while affecting Clade A by exclusion through competition, and/or causing gene introgression to lose its original haplotypes. *Rana sakuraii* and the small type of *R. t. tagoi* are sympatric with, and specifically distinct from Clade B in Kinki and Chugoku. It is possible that these two groups have already sufficiently differentiated ecologically to avoid competition or introgressive hybridization with Clade B for coexistence in these regions.

Taxonomic relationships

Of the many genetic groups recognized, Group A-1a should be considered true *R. t. tagoi* as it included the topotypic population from Kamitakara of the current Takayama-shi (locality 33), Gifu (Okada, 1928; Shibata, 1988). The small type of *R. t. tagoi*, one of the two types of *R. t. tagoi* from

Kinki (Sugahara, 1990), represented Group A-1b and was sympatric with the large type (parts of A-1a and B-2a). The small type differs from the large type in morphological, acoustic, and breeding ecological traits (Sugahara, 1990; Sugahara and Matsui, 1992, 1993, 1994, 1995, 1996, 1997). Thus, *R. t. tagoi* small type (A-1b) should be regarded as a distinct species. In contrast, *R. t. tagoi* morphologically identified as the large type was placed in two genetic groups (A-1a and B-2a), both with samples from the regions other than Kinki, and its taxonomic status is still unclear.

Subclade A-4 from one locality in Chubu has a unique breeding ecology and morphology different from sympatric Group A-1a (Misawa, private communication; Eto et al., 2012) and would be a distinct species. *Rana t. tagoi* from Neba-mura, Nagano, in Subclade A6 could also be another distinct species as it has $2n = 28$ chromosomes in contrast to $2n = 26$ in other *R. tagoi* and *R. sakuraii* populations (Ryuzaki et al., 2006). In our resultant tree, however, samples from Neba-mura (locality 76) were very close to and formed Subclade A6 with *R. t. tagoi* from southern Chubu and Shima Peninsula. It is thus necessary to investigate the chromosome number of the other populations in A-6 to determine taxonomic status of the Neba-mura population.

Details of morphological and ecological variations among other genetic groups of *R. t. tagoi* are generally poorly studied. Most of them are generally too similar to distinguish morphologically, but there are some exceptions. For example, representatives of Group A-1a and *R. t. tagoi* in Subclade A-2, at their range of sympatry in northern Kanto, are morphologically differentiated although slightly (Eto et al., unpublished data). Thus *R. t. tagoi* seems to include more cryptic taxa than previously suggested.

Rana t. yakushimensis formed Subclade A-8 by itself, and was split from the other *R. tagoi* subspecies and *R. sakuraii*. This result suggests its specific, rather than subspecific status, although it is morphologically very similar to *R. t. tagoi* (Maeda and Matsui, 1999). Supporting this idea, Nishioka et al. (1987) reported that *R. t. yakushimensis* was slightly isolated from *R. t. tagoi* from Chugoku (B-2b) by a low degree of hybrid inviability.

Another subspecies, *R. t. okiensis* also formed a distinct subclade (B-1) and split from other genetic groups. This subspecies is morphologically distinct from the other subspecies of *R. tagoi* and *R. sakuraii* (Maeda and Matsui, 1999), and there is little doubt to treat it as a distinct taxon. Conlon et al. (2010) suggested *R. t. okiensis* and *R. t. tagoi* to be heterospecific from antimicrobial peptide structure, and Nishioka et al. (1987) and Daito et al. (1998b) reported post-mating isolation of the two subspecies. These previous studies and present result strongly suggest that *R. t. okiensis* should be treated as a species distinct from *R. t. tagoi*.

The phylogenetic relationships obtained by our group, in which *R. tagoi* and *R. sakuraii* are revealed to be paraphyletic, are in disagreement with current taxonomy. This result may be partly due to imperfect taxonomy (i.e., insufficient detection of cryptic species), in addition to the evolutionary processes as mentioned above. *Rana sakuraii* was divided into two genetic groups (Subclade A-2 and A-3). Of these, Subclade A-2 includes topotypic samples and should be regarded as true *R. sakuraii*, in spite of the possibility of past gene introgression from *R. t. tagoi* as discussed above.

Although both subclades of *R. sakuraii* are sympatric with some genetic groups of *R. t. tagoi* in Honshu (Table 4), the two species are known to be reproductively isolated by differences in the season, site, and behavior of breeding (Maeda and Matsui, 1999). Moreover, *R. sakuraii* in A-2 is completely isolated from *R. t. tagoi* from Kinki (large type from Kyoto: B-2a) and *R. t. okiensis* (B-1) by postmating isolating mechanisms (Daito et al., 1998a; Daito, 1999). Because no obvious morphological and ecological differences have been detected between the two genetic groups of *R. sakuraii*, it seems safe at present to retain it as a single species.

It is now popular to regard uncorrected p-distances in 16S rRNA of 3–5% to be thresholds between intra- and inter-specific divergence levels in anurans (Vences et al., 2005; Fouquet et al., 2007). However, Hillis and Wilcox (2005) reported interspecific sequence divergences of 16S rRNA among American ranid frogs to be 1.2–18.7% (uncorrected p-distances calculated from GenBank data). Thus, sequence divergence alone is not an absolute indicator to draw taxonomic conclusions, though it can be considered useful in detecting candidate species. As to ND1, Vredenburg et al. (2007) separated *R. sierrae* and *R. muscosa*, with 4.6% sequence divergence in ND1 and ND2, as distinct species.

In the light of these reports, divergences among genetic groups of *R. tagoi* and *R. sakuraii* (1.3–3.5% in 16S rRNA and 2.9–7.0% in ND1) are generally not very large. Of the cryptic lineages discussed above, A-1b (small type) could be regarded as heterospecific with B-2a (large type: divergences of 3.2% in 16S rRNA and 6.7% in ND1), although its divergence from true *R. t. tagoi* (A-1a) is not large enough to indicate specific separation (1.3% and 4.1%). Of other unique groups observed, Subclade A6, including a population with extra number of chromosomes, differed from the other groups by divergences of 1.5–3.3% (16S rRNA) and 3.4–6.7% (ND1). Likewise, divergences were 1.8–3.3% and 4.1–6.4% between *R. t. okiensis* and the other groups, and 1.5–2.8% and 2.8–6.9% between *R. t. yakushimensis* and the other groups. These values partly exceed proposed thresholds or reported values for specific separation (Fouquet et al., 2007; Vredenburg et al., 2007). Other combinations produced even smaller divergences (1.4% and 3.9% between Subclade A-4 and Group A-1a; 1.7% and 4.0% between Group A-1a and *R. t. tagoi* in Subclade A-2; and 1.1% and 2.1% between *R. sakuraii* and *R. t. tagoi* in A-2), in spite of their sympatric occurrences, and posed questions about the universality of threshold values in DNA barcoding.

In frogs, sister species sometimes exhibit very small sequence divergences in spite of their distinct morphological and/or ecological differences (e.g., Matsui et al., 2006), and similar situations appear to apply to unique genetic groups recognized in *R. tagoi* and *R. sakuraii*. Small sequence divergences, like morphological and ecological similarities, suggest relatively recent separation among genetic groups of these frogs.

This study provided a large amount of new information about the complex genetic diversity and consequential taxonomic problems with respect to *R. tagoi* and *R. sakuraii*. However, mtDNA alone is not a conclusive indicator of

reproductive isolation, due to its maternal mode of inheritance. Further studies, including nuclear marker analyses, are necessary to clarify reproductive isolations among genetic groups and draw definitive taxonomic conclusions.

ACKNOWLEDGMENTS

We thank G. Aoki, K. Araya, H. Fujita, M. Hayashi, A. Hamidy, T. Hikida, S. Ikeda, K. Kasugai, M. Kato, K. Kawauchi, J. B. Kim, N. Koike, Y. Kokuryo, M. Kumagai, N. Kuraishi, N. Maeda, Mm. Matsui, Mt. Matsui, T. Matsuki, T. Matsuo, Y. Misawa, K. Mitsui, Y. Miyagata, Y. Mizuno, A. Mori, S. Mori, N. Nakahama, T. Nakano, K. Nishikawa, H. Okawa, S. Okada, J. Oki, M. Sakamoto, T. Sasaoka, Y. Sasaoka, T. Seto, T. Shimada, Z. Shimizu, R. Shimoyama, T. Sota, T. Sueyoshi, K. Sugahara, T. Sugihara, H. Takeuchi, S. Tanabe, Y. Tatewaki, H. Tokunaga, N. Tokuyama, A. Tominaga, T. Utsunomiya, Y. Utsunomiya, M. Yamagami, N. Yoshikawa, and M. Yumoto for help in collecting specimens and N. Kuraishi and N. Yoshikawa for assisting with experiments. We are also grateful to H. Ota and an anonymous reviewer for valuable comments on the manuscript. This work was partially supported by grants from The Monbukagakusho through the Japanese Society for the Promotion of Sciences (JSPS: 15370038, 20405013) to MM.

REFERENCES

- Avice JC (2000) Phylogeography: The History and Formation of Species. Harvard University Press, Cambridge, Massachusetts
- Ballard JWO, Whitlock MC (2004) The incomplete natural history of mitochondria. *Mol Ecol* 13: 729–744
- Boulenger GA (1920) A monograph of the South Asian, Papuan, Melanesian, and Australian frogs of the genus *Rana*. *Rec Ind Mus* 20: 1–126
- Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 17: 540–552
- Conlon JM, Coquet L, Jouenne T, Leprince J, Vaudry H, Iwamuro S (2010) Evidence from the primary structures of dermal antimicrobial peptides that *Rana tagoi okiensis* and *Rana tagoi tagoi* (Ranidae) are not conspecific subspecies. *Toxicon* 55: 430–435
- Daito Y (1999) Development of postmating isolation in the Tago's brown frog group. III. Offspring between Oki population, Tago's brown frog of Oki island ♀ and Stream brown frog ♂. *Amph Hist* 2: 23–29 (in Japanese)
- Daito Y, Yokota T, Inoue K, Nakata T (1998a) Development of postmating isolation in the Tago's brown frogs (*Rana*) I. Offspring between *Rana sakuraii* and *Rana tagoi tagoi*, Kurama population. *Bull Fac Hum Devel Kobe Univ* 5: 507–522 (in Japanese)
- Daito Y, Hiramoto M, Yoshimoto Y (1998b) Development of postmating isolation in the Tago's brown frogs (*Rana*) II. Offspring between *Rana tagoi tagoi* and Oki population, Tago's brown frog of Oki Isl. *Bull Fac Hum Devel Kobe Univ* 5: 523–530 (in Japanese)
- Dubois A (1992) Notes sur la classification des Ranidae (Amphibiens Anoures). *Bull Mens Soc Linn Lyon* 61: 305–352
- Eto K, Matsui M, Misawa Y (2012) On two types of *Rana tagoi tagoi* from the former Kuni-mura, Gunma Prefecture. *Bull Herpetol Soc Japan* 2012: 81 (in Japanese)
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791
- Fouquet A, Gilles A, Vences M, Marty C, Blanc M, Gemmell NJ (2007) Underestimation of species richness in Neotropical frogs revealed by mtDNA analyses. *PLoS ONE* 2: e1109
- Frost DR (1985) *Amphibian Species of the World: A Taxonomic and Geographical Reference*. Allen Press, Lawrence
- Frost DR, Grant T, Faivovich JN, Bain RH, Haas A, Haddad CFB, et al. (2006) The amphibian tree of life. *Bull Amer Mus Nat Hist* 297: 1–370

- Hedges SB (1994) Molecular evidence for the origin of birds. *Proc Nat Acad Sci USA* 91: 2621–2624
- Hillis DM, Wilcox TP (2005) Phylogeny of the New World true frogs (*Rana*). *Mol Phylogenet Evol* 34: 299–314
- Hillis DM, Mable BK, Larson A, Davis SK, Zimmer EA (1996) Nucleic acids IV: sequencing and cloning. In "Molecular Systematics" Ed by DM Hillis, BK Mable, C Moritz, Sinauer Associates, Sunderland, pp 321–378
- Huelsenbeck JP, Hillis DM (1993) Success of phylogenetic methods in the four–taxon case. *Syst Biol* 42: 247–264
- Huelsenbeck JP, Ronquist FR (2001) MrBayes: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754–755
- Igawa T, Kurabayashi A, Nishioka M, Sumida M (2006) Molecular phylogenetic relationships of toads distributed in the Far East and Europe inferred from the nucleotide sequences of mitochondrial DNA genes. *Mol Phylogenet Evol* 38: 250–260
- Jobb G (2008) TREEFINDER version of October 2008. Munich, Germany. Distributed by the author at www.treefinder.de
- Leaché AD, Reeder TW (2002) Molecular systematics of the eastern fence lizard (*Sceloporus undulatus*): a comparison of parsimony, likelihood, and Bayesian approaches. *Syst Biol* 51: 44–68
- Maeda N, Matsui M (1999) Frogs and Toads of Japan, Revised Edition. Bun-ichi Sogo Shuppan, Tokyo (in Japanese)
- Matsui M (1984) Morphometric variation analyses and revision of the Japanese toads (genus *Bufo*, Bufonidae). *Contrib Biol Lab Kyoto Univ* 26: 209–428
- Matsui T, Matsui M (1990) A new brown frog (genus *Rana*) from Honshu, Japan. *Herpetologica* 46: 78–85
- Matsui M, Shimada T, Liu WZ, Maryati M, Khonsue W, Orlov N (2006) Phylogenetic relationships of Oriental torrent frogs in the genus *Amolops* and its allies (Amphibia, Anura, Ranidae). *Mol Phylogenet Evol* 38: 659–666
- Mueller RL (2006) Evolutionary rates, divergence dates, and the performance of mitochondrial genes in Bayesian phylogenetic analysis *Syst Biol* 55: 289–300
- Nishikawa K, Matsui M, Tanabe S, Sato S (2007) Morphological and allozymic variation in *Hynobius boulengeri* and *Hynobius stejnegeri* (Amphibia: Urodela: Hynobiidae). *Zool Sci* 24: 752–766
- Nishioka M, Ohta S, Sumida M (1987) Intraspecific differentiation of *Rana tagoi* elucidated by electrophoretic analysis of enzymes and blood proteins. *Sci Rep Lab Amphib Biol Hiroshima Univ* 9: 97–133
- Nishizawa T, Kurabayashi A, Kunihara T, Sano N, Fujii T, Sumida M (2011) Mitochondrial DNA diversification, molecular phylogeny, and biogeography of the primitive rhacophorid genus *Buergeria* in East Asia. *Mol Phylogenet Evol* 59: 139–147
- Nöllert A, Nöllert C (1992) Die Amphibien Europas. Franckh-Kosmos, Stuttgart
- Okada Y (1928) Notes on Japanese frogs. *Annot Zool Jap* 11: 269–277
- Rambaut A, Drummond AJ (2007) Tracer version 1.4. Available at <http://tree.bio.ed.ac.uk/software/tracer>
- Ryuzaki M, Nishioka M, Kawamura T (2006) Karyotypes of *Rana tagoi* Okada with diploid number 28 in the Chausu Mountains of the Minamishinshu district of Nagano Prefecture, Japan (Anura:Ranidae). *Cytogenet Genome Res* 114: 56–65
- Sekiya K, Ohtani H, Ogata M, Miura I (2010) Phyletic diversity in the frog *Rana rugosa* (Anura: Ranidae) with special reference to a unique morphotype found from Sado island, Japan. *Curr Herpetol* 29: 69–78
- Shibata Y (1988) The type localities of three subspecies of a Japanese brown frog, *Rana tagoi* OKADA (Amphibia: Ranidae). *Bull Osaka Mus Nat Hist* 43: 43–46 (in Japanese)
- Shimada T, Matsui M, Yambun P, Sudin A (2011) A taxonomic study of Whitehead's torrent frog, *Meristogenys whiteheadi*, with descriptions of two new species (Amphibia: Ranidae). *Zool J Linn Soc* 161: 157–183
- Sugahara T (1990) Reproductive ecology of *Rana tagoi tagoi* at Kitayama in Kyoto. *Jpn J Herpetol* 13: 145 (in Japanese)
- Sugahara T, Matsui M (1992) Larval growth and development in two types of the Japanese brown frog, *Rana tagoi tagoi*, at Kumogahata, Kyoto. *Jpn J Herpetol* 14: 209 (in Japanese)
- Sugahara T, Matsui M (1993) Morphometric comparisons in two types of *Rana t. tagoi* from Kumogahata, Kyoto. *Jpn J Herpetol* 15: 83 (in Japanese)
- Sugahara T, Matsui M (1994) An investigation of characters that discriminate two morphotypes of *Rana tagoi tagoi*. *Jpn J Herpetol* 15: 151 (in Japanese)
- Sugahara T, Matsui M (1995) Acoustic comparisons in two types of *Rana t. tagoi* from Kumogahata, Kyoto. *Jpn J Herpetol* 16: 64–65 (in Japanese)
- Sugahara T, Matsui M (1996) Morphological variation in *Rana tagoi* from regions around Kinki. *Jpn J Herpetol* 16: 151–152
- Sugahara T, Matsui M (1997) Breeding seasons and sites in two types of *Rana tagoi tagoi* from Kumogahata, Kyoto. *Jpn J Herpetol* 17: 68–69 (in Japanese)
- Sumida M, Ogata M (1998) Intraspecific differentiation of the Japanese brown frog *Rana japonica* inferred from mitochondrial DNA sequences of the cytochrome b gene. *Zool Sci* 15: 989–1000
- Swofford DL (2002) PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods), Version 4. Sinauer Associates, Sunderland
- Tanabe AS (2008) Phylogears version 1.5.2010.03.24. Software distributed by the author at <http://www.fifthdimension.jp/>
- Tanabe AS (2010) Kakusan4 version 0.1. Software distributed by the author at <http://www.fifthdimension.jp/>
- Tanaka T, Matsui M, Takenaka O (1994) Estimation of phylogenetic relationships among Japanese brown frogs from mitochondrial cytochrome b gene (Amphibia: Anura). *Zool Sci* 11: 753–757
- Tanaka T, Matsui M, Takenaka O (1996) Phylogenetic relationships of Japanese brown frogs (*Rana*: Ranidae) assessed by mitochondrial cytochrome b gene sequences. *Biochem Syst Ecol* 24: 299–307
- Tanaka-Ueno T, Matsui M, Chen S-L, Takenaka O, Ota H (1998) Phylogenetic relationships of brown frogs from Taiwan and Japan assessed by mitochondrial cytochrome b gene sequences (*Rana*: Ranidae). *Zool Sci* 15: 283–288
- Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improvement the sensitivity of progressive multiple sequence alignment through sequence weighting, position–specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22: 4673–4680
- Vences M, Thomas M, Bonett RM, Vieites DR (2005) Deciphering amphibian diversity through DNA barcoding: chances and challenges. *Phil Trans R Soc B* 360: 1859–1868
- Vredenburg VT, Bingham R, Knapp R, Morgan JAT, Moritz C, Wake D (2007) Concordant molecular and phenotypic data delineate new taxonomy and conservation priorities for the endangered mountain yellow-legged frog. *J Zool* 271: 361–374
- Wilkinson JA, Drewes RC, Tatum OL (2002) A molecular phylogenetic analysis of the family Rhacophoridae with an emphasis on the Asian and African genera. *Mol Phylogenet Evol* 24: 265–273
- Yoshikawa N, Matsui M, Nishikawa K, Kim JB, Kryukov A (2008) Phylogenetic relationships and biogeography of the Japanese clawed salamander, *Onychodactylus japonicus* (Amphibia: Caudata: Hynobiidae), and its congener inferred from the mitochondrial cytochrome b gene. *Mol Phylogenet Evol* 49: 249–259

(Received April 14, 2012 / Accepted May 12, 2012)