

## **Efficacy and safety of switching from basal insulin to sitagliptin in Japanese type 2 diabetes patients**

Shin-ichi Harashima<sup>1</sup>, Daisuke Tanaka<sup>1</sup>, Shunsuke Yamane<sup>1</sup>, Masahito Ogura<sup>1</sup>, Yoshihito Fujita<sup>1</sup>, Yuko Murata<sup>2</sup>, Mika Seike<sup>2</sup>, Tadashi Koizumi<sup>2</sup>, Mitsuru Aono<sup>2</sup>, Yu Wang<sup>1</sup>, Nobuya Inagaki<sup>1</sup>

<sup>1</sup>Department of Diabetes and Clinical Nutrition, Graduate School of Medicine, Kyoto University, Kyoto 606-8507, Japan

<sup>2</sup>Department of Internal Medicine, Takashima General Hospital, Takashima 520-1121, Japan

\*Correspondence

Shin-ichi Harashima, M.D. Ph.D.

Department of Diabetes and Clinical Nutrition, Graduate School of Medicine,  
Kyoto University, Kyoto 606-8507, Japan

Tel: +81-75-751-3560

Fax: +81-75-771-6601

Email: harasima@metab.kuhp.kyoto-u.ac.jp

Short title: Replacing basal insulin by sitagliptin

Key words: sitagliptin, basal insulin, insulin secretion capacity

## **ABSTRACT**

Basal-supported oral therapy (BOT) is often used to treat poorly controlled type 2 diabetes. However, patients sometimes experience nocturnal and early morning hypoglycemia. Thus, maintaining targeted glycemic control by BOT is limited in some patients. We assessed the efficacy and safety of replacing basal insulin by sitagliptin therapy in Japanese type 2 diabetes patients on BOT. Forty-nine subjects were sequentially recruited for the 52-week, prospective, single arm study. Patients on BOT therapy were switched from basal insulin to sitagliptin. The primary endpoint was change in HbA1c in 52-week. The secondary endpoints were dropout rate, changes in body weight, frequency of hypoglycemia, and relationship between change in HbA1c and insulin secretion capacity evaluated by glucagon loading test. The average dose of basal insulin was  $15.0 \pm 8.4$  units. Sixteen subjects (31.3%) were dropped because replacement by sitagliptin was less effective for glycemic control. In these subjects, diabetes duration was longer, FPG and HbA1c at baseline were higher, and insulin secretion capacity was lower. Change in HbA1c in 52-week was -0.40% (95%CI -0.50 to -0.32) ( $P<0.05$ ). Change in body weight was -0.71 kg (95%CI -1.42 to -0.004) ( $P<0.05$ ). Frequency of hypoglycemia was decreased from  $1.21 \pm 1.05$  to  $0.06 \pm 0.24$  times/month. HbA1c level was improved if C-peptide index (CPI) was over 1.19. In conclusion, basal insulin in BOT can be replaced by sitagliptin with a decrease in

HbA1c level and frequency of hypoglycemia in cases where insulin secretion capacity was sufficiently preserved.

## **Introduction**

Basal insulin preparation is recommended by the American Diabetes Association (ADA) / European Association for the Study of Diabetes (EASD) consensus algorithm when lifestyle interventions and oral glucose-lowering agents no longer achieve the glycemic goal of haemoglobin A1c (HbA1c) level less than 7% [1, 2]. Recently, two long-acting insulin analogues, insulin glargine and insulin detemir, are available that attain glycemic targets more effectively and safely [3, 4]. There are no significant differences reported in glycemic control and overall hypoglycemia between the two analogues [5]. The combination of basal insulin and oral hypoglycemic agents (OHAs), known as basal-supported oral therapy (BOT), is often used to treat poorly controlled type 2 diabetes [6, 7]. Better glycemic control, fewer hypoglycemic episodes, and less weight gain are obtained by BOT than by biphasic insulin [8]. In addition, BOT is relatively cost effective with the same glycemic control level as biphasic insulin regimen [9]. BOT is also helpful in Japanese type 2 diabetes patients. In the ALOHA (Add-on to Lantus® to OHA) study, in which 5,223 Japanese type 2 diabetes patients participated, mean HbA1c was reduced from  $9.05 \pm 1.15\%$  to  $7.63 \pm 1.17\%$  in 24-week [10]. Although BOT is well-tolerated and effective for glycemic control, patients sometimes experience nocturnal and early morning hypoglycemia. In the ALOHA study,

0.97% of the patients experienced frequent hypoglycemia. In the 4T-study, 1.3% of BOT-treated patients experienced hypoglycemia with loss of consciousness [8]. Another problem of BOT is that postprandial glucose is high, although morning fasting blood glucose level is within normal range. An increase in dosage of basal insulin or SUs which are most commonly administered in BOT-treated Japanese patients is not always effective, and can result in increased hypoglycemia. In Japanese interview forms, frequency of hypoglycemia induced by SUs is reported to be 1.3% to 2.8%. Thus, maintaining targeted glycemic control by BOT is limited in some patients.

Dipeptidyl peptidase-4 (DPP-4) inhibitor is a newly developed OHA that prevents degradation of the incretin hormones, glucagon-like peptide-1 and gastric inhibitory polypeptide [11]. This compound stimulates glucose-dependent insulin secretion and suppresses glucagon release, and can improve both fasting and postprandial glucose levels. Four different DPP-4 inhibitors are available in Japan: sitagliptin, vildagliptin, alogliptin, and linagliptin. Of these, sitagliptin is most widely used, partly because it was the first approved DPP-4 inhibitor and the safety and efficacy are acceptable in Japanese clinical practice. Generally, sitagliptin is more effective for glycemic control in Japanese patients compared to Caucasian patients [12, 13]. Sitagliptin is usually combined with low dosage of SUs in Japan, less than or equal to 2 mg/day of

glimepiride and 40 mg/day of gliclazide, which is enough for glycemic control when combined with sitagliptin [14]. Patients also show improved glycemic control even if insulin secretion capacity is insufficient for oral therapy [14, 15]. The main pathophysiology of Japanese type 2 diabetes is impairment of insulin secretion [16, 17]. Insulin secretion capacity in Japanese populations is almost as half as that in Caucasians [18]. Especially, decreased basal and early phase insulin secretion is more contributed to Japanese type 2 diabetes [16]. Usually insulin therapy was needed in those whose C-peptide index (CPI) was lower than 0.8 [19]. However basal insulin therapy is not always ideal in some patients because postprandial glucose is still high, while preprandial glucose is low, resulting in large fluctuation in blood glucose. On the other hand, DPP-4 inhibitor could ameliorate decreased early phase insulin secretion. This encouraged us that basal insulin can be replaced with sitagliptin in type 2 diabetes patients treated with SUs and basal insulin in some BOT cases.

We show here that sitagliptin can be switched from basal insulin in patients with C-peptide index (CPI) and/or secretory unit of islet in transplantation (SUIT) equal to or larger than 1.19 and/or 36.4, respectively, with beneficial effects on glycemic control.

## **MATERIALS AND METHODS**

## **Study Design and Participants**

This was a prospective, 52-week, single center, single arm, intervention study to evaluate the effects on glycemic control of replacement of basal insulin to sitagliptin in type 2 diabetes patients inadequately controlled with BOT. Outpatients of Takashima General Hospital were recruited consecutively for a sample size of 45 subjects. Inclusion criteria were: type 2 diabetes treated with basal insulin (insulin glargine or detemir) and SUs (glimepiride or gliclazide)  $\pm$  metformin  $\pm$  thiazolidinedione  $\pm$   $\alpha$ -glucosidase inhibitors for more than 1 year; aged  $\geq$  20 years; HbA1c level  $\geq$  6.9%; no improvement in HbA1c  $\geq$  0.5% within 3 months in BOT; and a fasting C-peptide reactin (CPR) of  $>$  0.5 ng/mL. Exclusion criteria were: type 1 diabetes; secondary diabetes; alcoholism; severe depression or severe psychological condition; malignancy; and abnormal hemoglobinemia. The study protocol was approved by the Institutional Review Board of Takashima General Hospital, and registered on University hospital Medical Information Network in Japan (UMIN000005499). Written informed consent was obtained from all subjects.

## **Procedures and Intervention**

The duration of the study was 52 weeks. Subjects were screened for eligibility and gave basic demographic information, medical history, and frequency of hypoglycemia. Within a month before changing therapy from basal insulin to sitagliptin, glucagon loading test was performed without any OHAs or basal insulin for more than 24 h to evaluate insulin secretion capacity. When basal insulin was replaced by sitagliptin, the dosage of glimepiride or gliclazide was decreased to equal to or less than 2.0 mg/day or 40 mg/day, respectively, to prevent increased hypoglycemia if the subjects had been treated with more than 2.0 mg/day glimepiride or 40 mg/day gliclazide. If the subjects had been treated with equal to or less than 2.0 mg/day of glimepiride or 40 mg/day of gliclazide, that dosage of SUs was maintained. Metformin (Met) and thiazolidinedione (TZD) were continued without any changes during the study.  $\alpha$ -glucosidase inhibitors were discontinued. The dosage of SUs was changed depending on the frequency of hypoglycemic episodes and glycemic control level. Sitagliptin was started at 50 mg/day, the usual initial dosage in Japan, which was increased to 100 mg/day if the HbA1c level did not reach 6.9%, since titration to 100 mg/day is acceptable.

## **Measurements**



The primary endpoint was change in HbA1c in 52-week. The secondary endpoints were dropout rate due to lesser efficacy of replacement by sitagliptin of basal insulin on glycemic control, change in body weight in 52-week, change in body mass index (BMI) in 52-week, change in frequency of hypoglycemia in 52-week, adverse events, and the correlation between change in HbA1c at 8-week and insulin secretion capacity or CPI or SUII at baseline. HbA1c were expressed as NGSP values. CPI was calculated by the formula:  $[100 \times \text{fasting CPR (ng/mL)}] / [18 \times \text{FPG (mM)}]$  [19]. SUII index was calculated by the formula:  $[250 \times \text{fasting CPR (nM)}] / [(\text{FPG}-3.43) \text{ (mM)}]$  [20]. Blood glucose and C-peptide level were measured before (0 min) and 6 min after intravenous administration of 1 mg glucagon.

### **Statistical analysis**

Sample size was estimated to be 34 to detect a 0.4% change in HbA1c in 52-week with a power of 95%, alpha 0.05 two-tailed, beta 0.20, standardized effect size 0.7. To take the dropout rate of 30% into account, the aim was to include 45 subjects. IBM SPSS Statistics was used for analysis. Dependent samples Student's t-test was used to compare the means of HbA1c level, insulin secretion capacity, BMI, body weight, age, and diabetes duration of the subjects between baseline and 52-week. Person's

product-moment correlation test was used to evaluate the relationship between change in HbA1c and insulin secretion capacity or CPI or SUI. To evaluate cut-off values of diabetes duration, FPG, HbA1c, 0-min CPR, 6-min CPR, delta-CPR, CPI, and SUI, receiver operating characteristics curve (ROC) analysis was used. Independent samples Student's t-test was used to compare the mean of change in HbA1c in 52-week between subjects treated with sitagliptin + glimepiride and sitagliptin + gliclazide. Dunnett analysis was used to compare change in HbA1c in 52-week among subjects treated with sitagliptin + SUs and sitagliptin + SUs + MET and sitagliptin + SUs + TZD. *P* values <0.05 were considered as statistically significant.

## **Results**

### **Participants**

Forty-nine patients were eligible and were consecutively enrolled in the study (Table 1). Average age of subjects was  $70.0 \pm 10.2$  years; ratio of male was 60.8%; duration of diabetes was  $14.3 \pm 8.2$  years; average body weight was  $62.3 \pm 10.4$  kg; average BMI was  $24.3 \pm 3.8$  kg/m<sup>2</sup>; and HbA1c was  $7.97 \pm 0.81\%$ . All subjects were treated with SUs; 17 subjects (34.7%) were treated with glimepiride (average dose  $1.67 \pm 1.47$  mg) and 32 (65.3%) were treated with gliclazide (average dose  $33.8 \pm 12.0$  mg). Average

dosage of basal insulin analogues was  $15.0 \pm 8.4$  units. Glucagon loading test showed that 0-min CPR, 6-min CPR, CPI, and SUI were  $1.65 \pm 1.02$  ng/mL,  $3.37 \pm 1.98$  ng/mL,  $1.19 \pm 0.64$ , and  $36.5 \pm 22.1$ , respectively. Sixteen subjects (32.6%) were dropped due to an increase in HbA1c in 8-week; 6 (29.4%) and 11 (34.4%) were dropped in glimepiride- and gliclazide-treated subjects, respectively (Table 2). No subjects were dropped for other reasons. Thirty-three subjects completed the study.

### **HbA1c findings and dosage of SUs and sitagliptin**

Therapy adherence was confirmed by certified diabetes educators (nurses) in the study. Adherence of BOT therapy and the switching therapy were almost 100%, and there was no difference between both therapies (Data not shown).

HbA1c level in 52-week in final subjects was significantly decreased from  $7.75 \pm 0.70\%$  to  $7.40 \pm 0.68\%$  ( $P < 0.01$ ) (Table 2). Change in HbA1c in 52-week was  $-0.40\%$  (95%CI;  $-0.50, -0.32\%$ ) ( $P < 0.05$ ). HbA1c level in 52-week in glimepiride-treated subjects ( $n=12$ ) was significantly decreased from  $7.90 \pm 0.85\%$  to  $7.16 \pm 0.77\%$  ( $P < 0.01$ ). Change in HbA1c in 52-week was  $-0.74\%$  (95%CI;  $-1.03, -0.45\%$ ) ( $P < 0.05$ ).

HbA1c level in 52-week in gliclazide-treated subjects ( $n=21$ ) was significantly decreased from  $7.66 \pm 0.60\%$  to  $7.46 \pm 0.61\%$  ( $P < 0.05$ ). Change in HbA1c in 52-week

was -0.20% (95%CI; -0.35, -0.04%) ( $P<0.05$ ). There was a significant difference in change in HbA1c in 52-week between glimepiride-treated and gliclazide-treated subjects ( $P<0.01$ ). The original dosages of glimepiride and gliclazide before the study were  $1.58 \pm 0.93$  mg/day and  $38.2 \pm 14.0$  mg/day, respectively; the initial dosages at the beginning of the study were significantly decreased to  $0.96 \pm 0.40$  mg/day and  $24.8 \pm 8.7$  mg/day, respectively ( $P<0.05$ ); and the final dosages were significantly increased to  $1.42 \pm 0.57$  mg/day and  $31.4 \pm 12.0$  mg/day, respectively, compared to the initial dosages ( $P<0.05$ ), and were almost equal to the original dosages (Table 2). Final dosage of sitagliptin was  $74.2 \pm 25.4$  mg/day in all subjects;  $70.8 \pm 25.7$  mg/day in glimepiride-treated subjects; and  $77.3 \pm 25.5$  mg/day in gliclazide-treated subjects with no significant difference between the two groups.

Of 33 subjects who completed the study, 22 subjects were treated with sitagliptin and SUs, 6 subjects were treated with sitagliptin and SUs and MET, and 3 subjects were treated with sitagliptin and SUs and TZD; changes in HbA1c in 52-week were -0.38% (95%CI; -0.58, -0.19%) ( $P<0.05$ ), -0.54% (95%CI; -0.90, -0.20%) ( $P<0.05$ ), and -0.23% (95%CI; -0.43, -0.04%) ( $P<0.05$ ), respectively (Table 2). However, there was no significant difference among the three groups.

### **Change in body weight, BMI, and frequency of hypoglycemia**

Body weight in final subjects at baseline was  $64.2 \pm 9.5$  kg, and was decreased to  $63.5 \pm 8.7$  kg at 52-week. Change in body weight in 52-week was  $-0.71$  kg (95% CI;  $-1.42, -0.004$  kg) ( $P < 0.05$ ) (Table 3). BMI at baseline was  $24.8 \pm 3.6$  kg/m<sup>2</sup>, and decreased to  $24.5 \pm 3.4$  kg/m<sup>2</sup> at 52-week. Change in BMI in 52-week was  $-0.27$  kg/m<sup>2</sup> (95% CI;  $-0.54, 0.004$  kg/m<sup>2</sup>) ( $P > 0.05$ ).

Frequency of hypoglycemia at baseline was  $1.21 \pm 1.05$  times/month, and was significantly decreased to  $0.06 \pm 0.24$  times/month at 52-week ( $P < 0.001$ ). Change in frequency in hypoglycemia in 52-week was  $-1.21$  times/months (95% CI;  $-1.5, -0.80$ ) ( $P < 0.05$ ) (Table 3). During the study, no severe hypoglycemia was noted.

During the study, no other adverse events were not observed after replacement of basal insulin with sitagliptin.

### **Differences in HbA1c findings in 8-week in the final and dropped subjects**

Sixteen subjects of 49 recruited were dropped after 8-week due to increased HbA1c level. The remaining 33 subjects completed the study. HbA1c level at baseline (0-week) in final subjects was  $7.75 \pm 0.70\%$ , and was significantly decreased to  $7.45 \pm 0.61\%$  at 8-week ( $P < 0.001$ ) (Table 4). Change in HbA1c was  $-0.32\%$  (95% CI;  $-0.45, -0.20\%$ )

( $P < 0.05$ ). On the other hand, HbA1c level at baseline (0-week) in dropped subjects was significantly higher than that in final subjects ( $P < 0.05$ ), and was significantly increased from  $8.42 \pm 0.87\%$  to  $8.86 \pm 0.99\%$  in 8-week ( $P < 0.01$ ). Change in HbA1c was  $+0.64\%$  (95% CI; 0.29, 1.00%) ( $P < 0.05$ ).

### **Differences in clinical factors in final and dropped subjects**

There were no differences in age, sex, dosage of SUs, or dosage of basal insulin in final and dropped subjects (Table 4). Body weight and BMI also were not significantly different ( $P = 0.065$  and  $P = 0.2432$ , respectively). On the other hand, diabetes duration in dropped subjects was longer than that in final subjects ( $12.1 \pm 6.6$  vs.  $18.7 \pm 9.5$  years,  $P < 0.05$ ). FPG and HbA1c also were higher in dropped subjects than in final subjects (FPG;  $7.4 \pm 1.5$  vs.  $8.9 \pm 2.9$  mM,  $P < 0.05$ ) (HbA1c;  $7.75 \pm 0.70$  vs.  $8.42 \pm 0.87\%$ ,  $P < 0.01$ ).

Insulin secretion capacity was significantly higher in final subjects than that in dropped subjects (Table 4) ( $P < 0.05$ ). In final subjects, CPR level at 0-min, 6-min, and delta CPR (6-min CPR – 0-min CPR) were  $1.95 \pm 1.25$  ng/mL,  $3.81 \pm 2.13$  ng/mL, and  $1.98 \pm 1.35$  ng/mL, respectively. In dropped subjects, CPR level at 0-min, 6-min, and delta CPR were  $1.37 \pm 0.64$  ng/mL,  $2.42 \pm 1.21$  ng/mL, and  $1.16 \pm 0.69$  ng/mL, respectively. CPI

and SUI index also were significantly higher in final subjects than those in dropped subjects. CPI at baseline in final subjects was  $1.35 \pm 0.68$ , while that in dropped subjects was  $0.92 \pm 0.51$  ( $P < 0.05$ ). SUI at baseline was  $42.7 \pm 23.0$  in final subjects, and  $23.1 \pm 10.6$  in dropped subjects ( $P < 0.01$ ). We examined cut-off values of diabetes duration, FPG, HbA1c, 0-min CPR, 6-min CPR, delta-CPR, CPI, and SUI by analyzing ROC curves; they were 16.5 years, 8.2 mM, 7.8%, 1.25 ng/ml, 2.80 ng/ml, 1.60 ng/ml, 1.34, and 37.5, respectively (Fig. 1). This indicates that with longer diabetes duration, insulin secretion capacity becomes lower and the consequent poorer glycemic control makes switching BOT-treated patients from basal insulin to sitagliptin unsafe.

### **Correlation between efficacy of sitagliptin on glycemic control and insulin secretion capacity, CPI, and SUI**

We examined whether or not insulin secretion capacity, CPI, or SUI at baseline predicted the efficacy of replacing basal insulin with sitagliptin on glycemic control (Fig. 2). There was a correlation between change in HbA1c at 8-week and 0-min CPR ( $r = -0.281$ ), 6-min CPR ( $r = -0.326$ ), and delta CPR ( $r = -0.290$ ), assessed by glucagon loading test at baseline (Fig. 2A, B, C) ( $P < 0.05$ ). In addition, CPI ( $r = -0.360$ ) or SUI ( $r = -0.306$ ) at baseline was correlated with change in HbA1c at 8-week (Fig. 2D, E)

( $P < 0.05$ ). The value of 0-min CPR, 6-min CPR, delta CPR, CPI, and SUIIT at which the HbA1c level was not increased by replacement of basal insulin by sitagliptin were calculated to be 1.64 ng/mL, 3.36 ng/mL, 1.71 ng/mL, 1.19, and 36.4, respectively, by Pearson's product-moment correlation test (Table 5). The value of 0-min CPR, 6-min CPR, delta CPR, CPI, and SUIIT at which the HbA1c level was decreased by 0.5% in 8-week were calculated to be 1.86 ng/mL, 3.83 ng/mL, 1.98 ng/mL, 1.36, and 41.3, respectively. Other clinical characteristics of the patients such as disease duration and body weight were not significantly correlated with efficacy of replacing basal insulin with sitagliptin on glycemic control (Data not shown).

## **Discussion**

We show here that basal insulin can be switched to sitagliptin with good effects in type 2 diabetes patients treated with BOT. With this treatment, the HbA1c level decreased from  $7.75 \pm 0.70\%$  to  $7.35 \pm 0.68\%$  in 52-week ( $P < 0.01$ ). The change in HbA1c in 52-week was  $-0.40\%$  (95%CI;  $-0.50, -0.32$ ) ( $P < 0.05$ ). The efficacy of switching to sitagliptin from basal insulin was correlated with insulin secretion capacity, CPI, and SUIIT, CPI being most correlated marker in the present study. The average CPI in final



subjects was  $1.35 \pm 0.68$  ng/mL, while that of dropped subjects was  $0.92 \pm 0.51$  ng/mL. Pearson's product-moment correlation test revealed that HbA1c was improved by switching from basal insulin to sitagliptin if CPI was equal to or higher than 1.19 (Fig. 2D and Table 5). Similarly, basal insulin could be switched to sitagliptin if SUIIT was equal to or larger than 36.4 (Fig. 2E and Table 5). In the dropped subjects, diabetes duration was longer, FPG and HbA1c were worse, 0-min CPR, 6-min CPR, delta-CPR, CPI, and SUIIT were lower compared to those in final subjects. (Table 4). Cut-off values of them were 16.5 years, 8.2 mM, 7.8%, 1.25 ng/ml, 2.80 ng/ml, 1.60 ng/ml, 1.34, and 37.5, respectively (Fig. 1). This suggests that the efficacy of switching from basal insulin to sitagliptin, when combined with SUs, is dependent on basal glycemic control and the insulin secretion capacity. Baseline HbA1c of dropped subjects was higher than that of the final subjects. Dosage of basal insulin was more required to reach target HbA1c level in dropped subjects compared to that in the final subjects because of lower insulin secretion capacity. Thus, if baseline HbA1c level had been reduced by increasing dosage of basal insulin, it would be difficult to replace basal insulin to sitagliptin.

Replacement of basal insulin by sitagliptin resulted in a reduction in body weight and hypoglycemia. Body weight was reduced by 0.71 kg (95%CI; -1.41 to -0.004 kg)

( $P < 0.05$ ). Frequency of hypoglycemia was decreased from  $1.21 \pm 1.05$  to  $0.06 \pm 0.24$  times/month ( $P < 0.001$ ). Since sitagliptin is known to be body weight neutral [21, 22], discontinuation of basal insulin might contribute to body weight reduction. The combination of basal insulin and SUs often induces mild hypoglycemia by which patients feel a sense of hunger and eat between-meal snacks. This sometimes induces weight gain and poor glycemic control in BOT-treated patients. On the other hand, combination therapy with sitagliptin and low dosage SUs (less than or equal to 2 mg/day glimepiride or 40 mg/day gliclazide) was body weight neutral or led a decrease in BMI [14]. In the current study, hypoglycemia seldom occurred, and BMI was significantly decreased by  $0.38 \text{ kg/m}^2$  (95%CI -0.72 to -0.04  $\text{kg/m}^2$ ) [14]. Switching from basal insulin to sitagliptin also reduced the frequency of hypoglycemia. Although energy intake was not evaluated between baseline and 52-week in the present study, patients who had previously experienced frequent hypoglycemia reported to their physicians that the number of between-meal snacks in 52-week was fewer than at baseline. Thus, excess energy intake may be reduced after switching from basal insulin to sitagliptin to account for some of the body weight reduction and improvement in HbA1c. Another reason for improvement in the HbA1c level may be the reduced

postprandial glucose level by the combination therapy with sitagliptin and SUs compared to that by BOT.

The combination therapy of glimepiride and sitagliptin was more effective for HbA1c reduction than that of gliclazide and sitagliptin. Recently, it was reported that cAMP sensor Epac2 is a direct target of several sulfonylureas [23]. Tolbutamide, glibenclimide, and glimepiride bound Epac2 and enhanced glucose-stimulated insulin secretion. However, gliclazide did not bind Epac2. Because Epac2 also mediates the potentiation of insulin secretion by cAMP increased by endogenous incretin, the combination therapy of glimepiride and sitagliptin more enhances insulin secretion through activation of Epac2. This might be a potential mechanism why the combination therapy of glimepiride and sitagliptin was more effective for glycemic control than that of gliclazide and sitagliptin.

Generally, insulin secretion capacity of Japanese is as half as that of Caucasian [16, 17, 18]. Therefore, more than 60% of Japanese type 2 diabetes patients are treated with SUs [24]. DPP-4 inhibitor now is one of the most popular OADs, and more than 2 million patients were treated with DPP-4 inhibitors in Japan. Based on pathophysiology of Japanese patients and the mechanism of incretin effect, the combination therapy with SUs and DPP-4 inhibitors seems to be most effective for glycemic control compared to that

with other OADs and DPP-4 inhibitors. On the other hand, the main pathophysiology of Caucasian type 2 diabetes is insulin resistance compared to that of Japanese type 2 diabetes [25, 26]. Dosage of basal insulin in BOT in Caucasian patients is greater than that in Japanese patients. For example, in 4-T study, the mean dosage of basal insulin was 86 U (1.03 U/kg) [8], while 8.5 U (0.15 U/kg) in Japanese type 2 diabetes [10], and 15 U (0.24 U/kg) in our study. Therefore, it is not sure if basal insulin could be replaced with DPP-4 inhibitors even in subjects treated with high dosage of basal insulin. However, there is still possibility that in Caucasian subjects whose BMI is less than 25 kg/m<sup>2</sup> and CPI is over 1.3, basal insulin could be replaced with DPP-4 inhibitors. Or, if the combination therapy with high dosage of MET and DPP-4 inhibitors is more effective for glycemic control compared to other combinations in Caucasian type 2 diabetes, basal insulin with MET could be replaced with DPP-4 inhibitors and metformin.

During the course of the disease, type 2 diabetes patients are treated with several OHAs [27, 28]. However, if the HbA1c level does not reach less than 7%, insulin treatment is considered the next step [1, 2]. BOT is often selected for outpatients because once daily injection is acceptable and the glycemic control is superior, with fewer hypoglycemic episodes and less weight gain compared to biphasic insulin [8]. In Japan, the commonly used SUs are combined with basal insulin in BOT [10]. One of the biggest problems of

combination therapy with basal insulin and SUs is the high level of postprandial blood glucose while fasting blood glucose is within normal range. An increase in dosage of SUs or basal insulin does not resolve this problem, and sometimes leads to increased hypoglycemia. However, our results show that better glycemic control and lower frequency of hypoglycemia is obtained when switching from basal insulin to sitagliptin in subjects with sufficiently preserved insulin secretion capacity.

The advantages of discontinuation of basal insulin are 1) patients become free from daily injections; 2) they do not need to regularly perform self-monitoring of blood glucose (SMBG); and 3) oral therapy costs less than insulin therapy.

In summary, basal insulin in BOT can be switched to sitagliptin if CPI and/or SUIIT are equal to or higher than 1.19 or 36.4, respectively. On the other hand, sitagliptin can be added to insulin therapy if insulin secretion capacity is not sufficient for switching to sitagliptin. However, the effectiveness of combination therapy with basal insulin and sitagliptin on glycemic control in type 2 patients with CPI and/or SUIIT less than 1.19 or 36.4, respectively, is unknown. Further studies are required to determine the optimum insulin secretion capacities for switching BOT therapy to sitagliptin combined with SUs or combination therapy with sitagliptin and basal insulin or GLP-1 receptor analogues.

## **Acknowledgement**

None of the authors have any conflicts of interest to declare.

The study conception and protocol were by SH and NI. Patient examinations were by SH, TF, TK, and MA. Collection of data was by YW and MO. The statistical analysis was by DT, SY, and YF. The manuscript development was by Dr. SH and NI.

## References

- 1 *Nathan DM, Buse JB, Davidson MB, Ferrannini E, Holman RR, Sherwin R, Zinman B; American Diabetes Association; European Association for Study of Diabetes.*  
Medical management of hyperglycemia in type 2 diabetes: a consensus algorithm for the initiation and adjustment of therapy: a consensus statement of the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care* 2009; 32: 193-203.
- 2 *Nathan DM, Buse JB, Davidson MB, Ferrannini E, Holman RR, Sherwin R, Zinman B; American Diabetes Association; European Association for Study of Diabetes.*  
Medical management of hyperglycemia in type 2 diabetes mellitus: a consensus algorithm for the initiation and adjustment of therapy: a consensus statement from the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetologia* 2009; 52: 17-30.
- 3 *Heinemann L, Linkeschova R, Rave K, Hompesch B, Sedlak M, Heise T.*  
Time-action profile of the long-acting insulin analog insulin glargine (HOE901) in comparison with those of NPH insulin and placebo. *Diabetes Care* 2000; 23:644-649.
- 4 *Plank J, Bodenlenz M, Sinner F, Magnes C, Görzer E, Regittnig W, Endahl LA, Draeger E, Zdravkovic M, Pieber TR.* A double-blind, randomized, dose-response

- study investigating the pharmacodynamic and pharmacokinetic properties of the long-acting insulin analog detemir. *Diabetes Care* 2005; 28: 1107-1112.
- 5 *Swinnen SG, Simon AC, Holleman F, Hoekstra JB, Devries JH.* Insulin detemir versus insulin glargine for type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2011; 6: CD006383.
  - 6 *Blicklé JF, Hancu N, Piletic M, Profozic V, Shestakova M, Dain MP, Jacqueminet S, Grimaldi A.* Insulin glargine provides greater improvements in glycaemic control vs. intensifying lifestyle management for people with type 2 diabetes treated with OADs and 7-8% A1c levels. The TULIP study. *Diabetes Obes Metab* 2009; 11: 379-386.
  - 7 *Schreiber SA, Ferlinz K, Haak T.* The long-term efficacy of insulin glargine plus oral antidiabetic agents in a 32-month observational study of everyday clinical practice. *Diabetes Technol Ther* 2008; 10: 121-127.
  - 8 *Holman RR, Farmer AJ, Davies MJ, Levy JC, Darbyshire JL, Keenan JF, Paul SK; 4-T Study Group.* Three-year efficacy of complex insulin regimens in type 2 diabetes. *N Engl J Med* 2009; 361: 1736-1747.
  - 9 *Pfohl M, Dippel FW, Kostev K, Fuchs S, Kotowa W.* Different persistence on initial basal supported oral therapy in Type 2 diabetics is associated with unequal



- distributions of insulin treatment regimens under real-life conditions in Germany. *Int J Clin Pharmacol Ther* 2010; 48: 761-766.
- 10 *Kadowaki T*. Analysis of ALOHA (add-on Lantus® to OHA) study: Basal supported oral therapy (BOT) with insulin glargine resulted in reduction of HbA1c, FPG and PPG with nearly 2% incidence of adverse drug reactions. Dubai: World Diabetes Congress, 2011.
- 11 *Seino Y, Fukushima M, Yabe D*. GIP and GLP-1, two incretin hormones: Similarities and differences. *J Diabetes Invest* 2010; 1: 8-23.
- 12 *Tajima N, Kadowaki T, Odawara M, Nishi M, Taniguchi T, Arjona Ferreira JC*. Addition of sitagliptin to ongoing glimepiride therapy in Japanese patients with type 2 diabetes over 52 weeks leads to improved glycemic control. *Diabetol Int* 2011; 2: 32-44.
- 13 *Hermansen K, Kipnes M, Luo E, Fanurik D, Khatami H, Stein P*. Efficacy and safety of the dipeptidyl peptidase-4 inhibitor, sitagliptin, in patients with type 2 diabetes mellitus inadequately controlled on glimepiride alone or on glimepiride and metformin. *Diabetes, Obes, Metab* 2007; 9: 733-745.
- 14 *Harashima SI, Ogura M, Tanaka D, Fukushima T, Wang Y, Koizumi T, Aono M, Murata Y, Seike M, Inagaki N*. Sitagliptin add-on to low dosage sulfonylureas:

efficacy and safety of combination therapy on glycemic control and insulin secretion capacity in type 2 diabetes. *Int J Clin Prac* 2012; 66: 465-476.

15 *Kubota A, Matsuba I, Saito T, Nabe K, Seino Y.* Secretory units of islets in transplantation index is a useful clinical marker to evaluate the efficacy of sitagliptin in treatment of type 2 diabetes mellitus. *J Diabetes Invest*, doi:10.1111/j.2040-1124.2011.00109.x, 2011.

16 *Fukushima M, Usami M, Ikeda M, Nakai Y, Taniguchi A, Matsuura T, Suzuki H, Kurose T, Yamada Y, Seino Y.* Insulin secretion and insulin sensitivity at different stages of glucose tolerance: a cross-sectional study of Japanese type 2 diabetes. *Metabolism* 2004; 53: 831-835.

17 *Fukushima M, Suzuki H, Seino Y.* Insulin secretion capacity in the development from normal glucose tolerance to type 2 diabetes. *Diabetes Res Clin Pract* 2004; 66: S37-S47.

18 *Abdul-Ghani MA, Matsuda M, Jani R, Jenkinson CP, Coletta DK, Kaku K, DeFronzo RA.* The relationship between fasting hyperglycemia and insulin secretion subjects with normal or impaired glucose tolerance. *Am J Physiol Endocrinol Metab* 2008; 295: E401-406.

- 19 *Funakoshi S, Fujimoto S, Hamasaki A, Fujiwara H, Fujita Y, Ikeda K, Hamamoto Y, Hosokawa M, Seino Y, Inagaki N.* Analysis of factors influencing pancreatic beta-cell function in Japanese patients with type 2 diabetes: Association with body mass index and duration of diabetic exposure. *Diabetes Res Clin Pract* 2008; 82: 353-358.
- 20 *Yamada Y, Fukuda K, Fujimoto S, Hosokawa M, Tsukiyama K, Nagashima K, Fukushima M, Suzuki H, Toyoda K, Sassa M, Funakoshi S, Inagaki N, Taniguchi A, Sato TS, Matsumoto S, Tanaka K, Seino Y.* SUIT, secretory units of islets in transplantation: An index for therapeutic management of islet transplanted patients and its application to type 2 diabetes. *Diabetes Res Clin Pract* 2006; 74: 222-226.
- 21 *Esposito K, Cozzolino D, Bellastella G, Maiorino MI, Chiodini P, Ceriello A, Giugliano D.* Dipeptidyl peptidase-4 inhibitors and HbA1c target of <7% in type 2 diabetes: meta-analysis of randomized controlled trials. *Diabetes Obes Metab* 2011; 13: 594-603.
- 22 *Gallwitz B, Häring HU.* Future perspectives for insulinotropic agents in the treatment of type 2 diabetes-DPP-4 inhibitors and sulphonylureas. *Diabetes Obes Metab* 2010; 12: 1-11.

- 23 Zhang CL, Katoh M, Shibasaki T, Minami K, Sunaga Y, Takahashi T, Yokoi N, Iwasaki M, Miki T, Seino S. The cAMP sensor Rpac2 is a direct target of antidiabetic sulfonylurea drugs. *Science* 2009; 25: 607-610.
- 24 Arai K, Matoba K, Hirao K, Matsuba I, Takai M, Takeda H, Kanamori A, Yamauchi M, Mori H, Terauchi Y. Present status of sulfonylurea treatment for type 2 diabetes in Japan: second report of a cross-sectional survey of 15,652 patients. *Endocr J* 2010; 57: 499-507.
- 25 Welch S, Gebhart SS, Bergman RN, Phillips LS. Minimal model analysis of intravenous glucose tolerance test-derived insulin sensitivity in diabetic subjects. *J Clin Endocrinol Metab* 1990; 71: 1508-1518.
- 26 Taniguchi A, Nakai Y, Fukushima M, Kawamura H, Imura H, Nagata I, Tokuyama K. Pathogenic factors responsible for glucose intolerance in patients with NIDDM. *Diabetes* 1992; 41: 1540-1546.
- 27 Tahrani AA, Bailey CJ, Del Prato S, Barnett AH. Management of type 2 diabetes: new and future developments in treatment. *Lancet* 2011; 378: 182-197.
- 28 DeFronzo RA. Current issues in the treatment of type 2 diabetes. Overview of newer agents: where treatment is going. *Am J Med* 2010; 123: S38-48.

## Figure legends

**Fig. 1** Cut-off values and receiver-operator characteristic curves of (a) diabetes duration, (b) fasting plasma glucose, (c) HbA1c, (d) 0-min CPR, (e) 6-min CPR, (f) delta CPR, (g) CPI, and (h) SUIIT at baseline. CPR, C-peptide reaction; CPI, C-peptide index; SUIIT, the secretory unit of islet in transplantation.

**Fig. 2** Relationship between change in HbA1c in 8-week and results of glucagon loading test, CPI, and SUIIT at baseline. Change in HbA1c in 8-week and 0-min CPR (a), 6-min CPR (b), delta CPR (c), CPI (d), and SUIIT index (e) at baseline. CPR, C-peptide reaction; CPI, C-peptide index; SUIIT, the secretory unit of islet in transplantation.

\* $P < 0.05$

Figure 1

	Diabetes Duration (Years)	FPG (mM)	HbA1c (%)	0-min CPR (ng/mL)	6-min CPR (ng/mL)	Delta CPR (ng/mL)	CPI	SUIT
Cut-off value	16.5	8.2	7.7	1.25	2.80	1.60	1.34	37.5
AUC	0.468	0.507	0.252	0.530	0.570	0.541	0.509	0.482
Specificity	0.600	0.600	0.800	0.800	0.800	0.800	0.600	0.600
Sensitivity	0.545	0.568	0.159	0.432	0.455	0.364	0.545	0.432




Figure 2 Correlation between change in A1c in 8-week and insulin secretion capacity




Table 1 Demographic and clinical features of subjects participating in the study

Subjects (n)	49	Basal insulin	15.0 ± 8.4 units
Age (years)	70.0 ± 10.2	Medications	SU 100 %
			Glimepiride 34.7%
			1.67 ± 1.47mg
Male	60.8%		Gliclazide 65.3%
			33.8 ± 12.0mg
Diabetes duration (year)	14.3 ± 8.2		Metformin 22.4%
			636 ± 131mg
Complications	Nephropathy		Thiazolidinedione 16.3%
	61.2%		10.3 ± 3.9mg
	Retinopathy		α-glucosidase inhibitors
	69.4%		8.1%
	Neuropathy	Glucagon test	
	42.8%	0-min CPR (ng/mL)	1.65 ± 1.02
	Cardiovascular diseases	6-min CPR (ng/mL)	3.37 ± 1.98
	34.7%	Delta CPR (ng/mL)	1.72 ± 1.23
Weight (kg)	62.3 ± 10.4	CPI	1.19 ± 0.64
BMI (kg/m <sup>2</sup> )	24.3±3.8	SUIT	36.5 ± 22.1
HbA1c (%)	7.97±0.81		



Table 2 Change in HbA1c and dosage of SUs and sitagliptin in final subjects

Subjects (n)	Dropout rate (%) (n)	HbA1c level baseline (%)	HbA1c level 52-week (%)	Change in HbA1c (%) (95%CI)	Original dosage of SUs (mg)	Initial dosage of SUs (mg)	Final dosage of SUs (mg)	Final dosage of sitagliptin (mg)
Final 33	32.6% 16	7.75 ± 0.70	7.35 ± 0.68**	-0.40%* (-0.50, -0.32%)	-	-	-	74.2 ± 25.4
Glimepiride 12	29.4% 5	7.90 ± 0.85	7.16 ± 0.77**	-0.74%* (-1.03, -0.45%)	1.58 ± 0.93	0.96 ± 0.40*	1.42 ± 0.57*	70.8 ± 25.7
Gliclazide 21	34.4% 11	7.66 ± 0.60	7.46 ± 0.61*	-0.20%* (-0.35, -0.04%)	38.2 ± 14.0	24.8 ± 8.7*	31.4 ± 12.0*	77.3 ± 25.5
SUs 23	30.3% 10	7.63 ± 0.67	7.26 ± 0.68*	-0.38%* (-0.58, -0.19%)	-	-	-	67.4 ± 24.3
SUs + Met 7	30.0% 3	7.99 ± 0.80	7.43 ± 0.72	-0.54%* (-0.90, -0.20%)	-	-	-	87.5 ± 23.1
SUs + TZD 3	50.0% 3	8.10 ± 0.56	7.87 ± 0.40	-0.23%* (-0.43, -0.04%)	-	-	-	100 ± 0.0

\**P*<0.05

\*\**P*<0.01

Table 3 Change in weight, BMI, and frequency in hypoglycemia

	Weight (kg)	BMI (kg/m <sup>2</sup> )	Hypoglycemia (times/month)
0-week	64.2 ± 9.5	24.8 ± 3.6	1.21 ± 1.05
52-week	63.5 ± 8.7	24.5 ± 3.4	0.06 ± 0.24***
Change	-0.71*	-0.27	-1.21*
(95%CI)	(-1.42, -0.004)	(-0.54, 0.004)	(-1.5, -0.80)

\* $P < 0.05$

\*\*\* $P < 0.001$

Table 4 Change in HbA1c and background of final and dropped subjects

	Final subjects 33	Dropped subjects 16		Final subjects 33	Dropped subjects 16
0 Wk HbA1c (%)	7.75 ± 0.70	8.42 ± 0.87	Original dosage of SUs (mg)	Glimepiride 1.58 ± 0.93 Gliclazide 36.2 ± 10.2	Glimepiride 2.70 ± 2.05 Gliclazide 38.2 ± 14.1
8 Wk HbA1c (%)	7.45 ± 0.61***	8.86 ± 0.99**	Basal insulin (Units)	14.8 ± 9.3	15.2 ± 6.4
Delta HbA1c (%) (95%CI)	-0.32* (-0.45, -0.20)	0.64* (0.29, 1.00)	FPG (mM)	7.4 ± 1.5	8.9 ± 2.9*
Age (years)	69.8 ± 10.7	70.5 ± 9.3	HbA1c (%)	7.75 ± 0.70	8.42 ± 0.87**
Male (%)	66.7	56.3	Glucagon test 0-min CPR (ng/mL)	1.95 ± 1.25	1.37 ± 0.64*
Diabetes duration (years)	12.1 ± 6.6	18.7 ± 9.5*	6-min CPR (ng/mL)	3.81 ± 2.13	2.42 ± 1.21*
Weight (kg)	64.2 ± 9.5	58.4 ± 11.5	Delta CPR (ng/mL)	1.98 ± 1.35	1.16 ± 0.69*
BMI (kg/m <sup>2</sup> )	24.8 ± 3.6	23.4 ± 4.0	CPI	1.35 ± 0.68	0.92 ± 0.51*
			SUIT	42.7±23.0	23.1±10.6**

\**P*<0.05 \*\**P*<0.01 \*\*\**P*<0.001

Table 5 Correlation between change in HbA1c and insulin secretion capacity

Change in HbA1c (%)	0-min CPR (ng/mL)	6-min CPR (ng/mL)	Delta CPR (ng/mL)	CPI	SUIT
0 .0	1.64	3.36	1.71	1.19	36.4
-0.5	1.86	3.83	1.98	1.36	41.3

Table 6 Cut-off values of baseline glycemic control and endogenous beta-cell function

	Diabetes Duration (Years)	FPG (mM)	HbA1c (%)	0-min CPR (ng/mL)	6-min CPR (ng/mL)	Delta CPR (ng/mL)	CPI	SUIT
Cut-off value	16.5	8.2	7.8	1.25	2.80	1.60	1.34	37.5