

Fluorescence distribution pattern allows to distinguish two species of *Eugymnanthea* (Leptomedusae: Eirenidae)

SHIN KUBOTA¹, PATRIZIA PAGLIARA² AND CINZIA GRAVILI²

¹Seto Marine Biological Laboratory, Field Science Education and Research Center, Kyoto University, Shirahama, Nishimuro, Wakayama 649-2211, Japan, ²Department of Biological and Environmental Science and Technology, University of Salento, Lecce, Via per Monteroni 73100 Lecce, Italy

The auto-fluorescence patterns of the medusae observed under a fluorescent microscope with blue light excitation allows to distinguish two species of Eugymnanthea, this even when they are still attached to the hydroid as small medusa buds despite the occurrence of a sex-dependant pattern in E. japonica. A total of four distribution patterns of green fluorescence, including non-fluorescence, could be found. Three of them are found in E. japonica, called 'subumbrellar fluorescence type' except for non-fluorescence, while another type is found in E. inquilina, called 'umbrellar margin fluorescence type'. During the short life of the medusa the latter type remained invariable for up to six days in E. inquilina, while the pattern observed for up to seven days in E. japonica changed sometimes, but it always remained distinguishable from the pattern found in E. inquilina. Therefore, the fluorescence pattern is a reliable taxonomic character. Fluorescence was not found in unfertilized eggs, planulae 2–8 days old, parthenogenetically produced larvae, or in the hydroids of the two species. The auto-fluorescent and possible bioluminescent tissues of these Eugymnanthea medusae could have some unknown biological significance.

Keywords: fluorescence distribution patterns, *Eugymnanthea* species, various developmental stages

Submitted 9 August 2007; accepted 24 June 2008; first published online 23 September 2008

INTRODUCTION

Eugymnanthea is a commensal hydroid associated mainly with *Mytilus galloprovincialis* and other bivalves inhabiting shallow waters. It occurs in the Mediterranean Sea (*E. inquilina*) and around the Japanese Sea (*E. japonica*) (Kubota, 1989, 1992, 2004; Piraino *et al.*, 1994; Rayyan *et al.*, 2002; Baba *et al.*, 2007). The life cycle of both species comprises a benthic hydroid and a planktonic medusa stage. Species of *Eugymnanthea* have a simplified medusa that is released with already formed gonads. They only spend a very short time in the plankton and they die after spawning. The two species of *Eugymnanthea*, which evolved as the most derived bivalve-inhabiting hydrozoans from an ancestral *Eirene* and/or *Eutima*-like progenitor, resemble each other both in the hydroid and medusa stages. The similarity is likely due to a parallel, progenetic evolution, which also explains some of the subtle morphological differences (Kubota, 2000). Individuals from the Mediterranean and Japan are very similar at any developmental stage, but by examining a large number of specimens of several populations, Kubota (1991, 2004) found some morphological differences of the medusae that permit to discriminate the two species. These are: presence of a manubrium and a smaller number of statoliths per statocyst (usually 1) in the Japanese form, versus the absence of a manubrium and several statoliths per statocyst (usually 2 or 3) in the Mediterranean form.

Govindarajyan *et al.* (2005) demonstrated that they are two distinct species by using a cross-fertilization test, a mesogloaeal adhesion and spreading test, and by 16S rDNA sequence comparisons. In the present study, we describe a new morphological character that allows both species to be distinguished, i.e. different distribution pattern of green auto-fluorescence.

MATERIALS AND METHODS

A total of 77 medusae of both sexes of *Eugymnanthea inquilina* were examined, originating from 53 specimens of *Mytilus galloprovincialis* Lamarck from Taranto (39 specimens) and Lago Fusaro, the type locality (14 specimens), one specimen of *Mytilaster minimus* (Poli) from Taranto, and eight specimens of *Chylamys glabra* (Linnaeus) from Taranto. They were collected from October 1999 to January 2000 (Kubota, 2004). *Mytilus galloprovincialis* associated with hydroids were kept in the biological laboratory of Lecce (University of Salento) from mid-October 1999 to mid-March 2000 at 23°C and a 15hL:9hD photoperiod. They were kept in several thousand ml glass or plastic containers with natural seawater taken from rocky coasts near Porto Cesareo or from inside the port of Porto Cesareo and fed with newly hatched *Artemia* nauplii. The seawater (38–40 ppt) was changed nearly every day and almost all of the released mature medusae were examined before they spawned in early morning. A Zeiss Standard Axioplan microscope equipped with a halogen lamp (Hg 100) light and blue light excitation with a BP 485/20 excitation filter, an FT 510 chromatic beam splitter, and an LP 520 barrier filter were used to observe the slides prepared from

Corresponding author:

S. Kubota

Email: shkubota@medusanpolyp.mbox.media.kyoto-u.ac.jp

each sample. Hydroids with or without a medusa bud were removed from the bivalves, and reared and examined under the microscope as described above.

A total of 474 medusae of both sexes of *Eugymnanthea japonica* Kubota, 1979 were examined, originating from 47 specimens of *Mytilus galloprovincialis* from Shirahama (27 specimens) and Atami, near to the type locality Shimoda (20 specimens), Japan, which were reared and examined as described above (Kubota, 2004).

Medusae of both species (12 *E. japonica* and 4 *E. inquilina*) were reared until they died and the change of the fluorescence pattern was examined every day. Unfertilized eggs that were not older than a few hours and planulae obtained after conspecific crossings (see Govindarajan *et al.*, 2005) of both *Eugymnanthea* species were examined similarly together with parthenogenetically produced larvae if present.

RESULTS

The presence and type of green fluorescence is described in relation to the developmental stage (Tables 1 & 2). The changes of the fluorescence pattern observed during the life span of the medusa, maximally for seven days, are summarized in Table 3.

Medusa

The auto-fluorescence pattern observed in the medusae of *E. inquilina* (Figure 1A, B) and *E. japonica* (Figure 1C, D) was

constantly different, showing 'umbrellar margin fluorescence type' in the former and a 'subumbrellar fluorescent type' in the latter. In addition, in *E. inquilina* only the peripheral part of gonads showed an auto-fluorescence (Figure 1B).

Two subtypes of fluorescence distribution were found in *E. japonica*, namely the subumbrellar fluorescence type confined to individual auto-fluorescent photocytes (Figure 1D) and another type with a uniform overall auto-fluorescence (Table 2). No pattern-variation was found for *E. inquilina* (Tables 1 & 2; Figure 1B). Only a small fraction of *E. japonica* medusae (2.7% in 474) showed no fluorescence at all (Table 2). Summarizing, we observed four different patterns in the medusae of *Eugymnanthea*.

The type of subpattern found in *E. japonica* seems to depend somewhat on the sex as females had no auto-fluorescent particles, thus the subumbrella has a brilliant, uniform fluorescence (92.6% in 394 examined females), while the auto-fluorescence in males tends to be concentrated into auto-fluorescent particles (63.6% in 33 males examined).

During the short life span of the medusa of *E. japonica*, maximally a week, a change of fluorescence pattern was detected in three out of four examined males (from the subumbrellar type to absence type or vice versa), and the changes were all irreversible (Table 3). The changes did not correlate with the age of the medusa or with senescence. Females of *E. japonica* did not change their pattern (examined in eight individuals; Table 3). In *E. inquilina* the fluorescence pattern was constant in all of the examined individuals, i.e. in two males and two females.

Table 1. Presence of fluorescence in various developmental stages of two species of *Eugymnanthea*.

Developmental stage	Appearance of fluorescent type in	
	<i>E. inquilina</i>	<i>E. japonica</i>
Medusa (see Table 2)	1 type (77, 3*, 2**)	2 types + absent (474, 1, 2)
Unfertilized eggs within several hours after spawning	Absent (82 from 9 females, 1, 1)	Absent (203 from 26 females, 1, 2)
Planula	Absent (8 3–8 days old, 1 pair, 1, 1)	Absent (2 2–3 days old, 1 pair, 1, 1)
Parthenogenetic larva	Not appeared	Absent (2 1-day-old, 1 pair, 1, 1)
Hydroid	Absent (137, 3*, 2**)	Absent (116, 1, 2)
Medusa bud	Same as medusa (97, 3*, 1)	Same as medusa (103, 1, 2)

*, *Mytilus galloprovincialis* (53 specimens), *Mytilaster minimus* (1 specimen), and *Chlamys glabra* (8 specimens); **, Taranto and Lago Fusaro. N, number of host sp. examined, number of localities surveyed.

Table 2. Fluorescent subtype of medusa and its frequency in the number of individuals examined in the two species of *Eugymnanthea*.

Distribution of fluorescence	<i>E. inquilina</i> collected from 3 host spp. from 2 localities	<i>E. japonica</i> collected from 1 host sp. from 2 localities
Subumbrella with photocytes	0	21 males + 26 females + 12*
Subumbrella without photocytes	0	4 males + 365 females + 33*
Umbrellar margin (marginal bulbs with photocytes) and peripheral part of gonads	34 males + 16 females + 10* + 17 hermaphrodites	0
Non-fluorescent	0	8 males + 3 females + 2*

*, sex undetermined.

Table 3. Change of fluorescent type in the life span of the medusa of the two species of *Eugymnanthea*.

Fluorescence	Number of medusae examined	Age of change in days and/or state continuation
<i>E. japonica</i> from <i>Mytilus galloprovincialis</i> collected from two localities		
Changes		
from subumbrella without photocytes to non-fluorescent	1 male	Change on the 4th day, then continued until the 5th day
from non-fluorescent to subumbrella without photocytes	2 males	Change on the 2nd day, then continued until the 3rd day
No changes		
subumbrella with photocytes	1 male	For up to 3 days
subumbrella without photocytes	8 females	For up to 4–7 days
<i>E. inquilina</i> from <i>Mytilus galloprovincialis</i> and <i>Chlamys glabra</i> collected from Taranto		
No changes		
umbrellar margin	2 females	For up to 3 days
umbrellar margin	2 males	For up to 6 days




Fig. 1. Photomicrographs of medusa of (A, B) *Eugymnanthea inquilina* and (C, D) *E. japonica*. Auto-fluorescence image under blue light excitation (B, D) and transmitted light image (A, C) of same individuals. Scale bars: 0.5 mm except for inset in (B) that shows a marginal bulb.

Medusa bud

In *E. inquilina* medusa buds we could detect auto-fluorescence in the distal part of the bud in every one of the 97 examined individuals (Figure 2A, B), while in *E. japonica* the whole subumbrella showed a uniform fluorescence in all 103 examined medusa buds (Figure 2C, D; Table 1). The auto-fluorescence of the gonads in *E. inquilina* appeared only later, when the gonads were well developed.

Eggs, planulae and hydroids

The auto-fluorescence was completely absent in 285 (203 in *E. japonica* + 82 in *E. inquilina*) unfertilized eggs, 10 (2 + 8) 2–8 days old planulae, even in two parthenogenetically produced larva of *E. japonica*, and 253 (103 + 97) hydroids in the two *Eugymnanthea* species (Table 1).

DISCUSSION

We found that the auto-fluorescent distribution pattern of the medusa bud and the free medusa is a reliable taxonomic




Fig. 2. Photomicrographs of medusa buds of (A, B) *Eugymnanthea inquilina* and (C, D) *E. japonica*. Auto-fluorescence image under blue light excitation (B, D) and transmitted light image (A, C) of same individuals of respective species. Scale bars: 0.1 mm.

character, which permits to distinguish the two *Eugymnanthea* species, despite the occurrence of a sex-dependant pattern in *E. japonica*. Such an auto-fluorescence pattern might also be useful for distinguishing other similar hydrozoan species, particularly for species evolved convergently or sibling species. As was pointed out by Morin & Reynolds (1969, 1974), fluorescence microscopy is easily applied to living material without complicated, time-consuming techniques. It is noteworthy that the fluorescent pattern is different in the two *Anemonia* species common to the Mediterranean Sea (Leutenegger *et al.*, 2007).

Several studies have been carried out on natural bioluminescence produced by Cnidaria and Ctenophora due to the presence of natural bioluminescent compounds (Haddock & Case, 1999; Haddock *et al.*, 2001). Some photoproteins have been isolated from Hydrozoa: aequorin (Harvey, 1935; Davenport & Nicol, 1955; Nicol, 1962; Shimomura, 2005) and obelin (Campbell, 1974; Vysotskii *et al.*, 1990, 1993, 1995). Morin & Reynolds (1974) found that in *Obelia*

hydroids the distribution of the fluorescent regions correlates with the ability to produce bioluminescence and that it may have some functional significance. One eminent ecological example of fluorescent structures serving as lures is reported in an *Erenna* siphonophore that preys upon fish in deep waters (Haddock *et al.*, 2005). The auto-fluorescent and possible bioluminescent tissues of present *Eugymnanthea* medusae could likewise have some unknown biological significance as pointed out by Morin (1983). It is noteworthy that in contrast to complete absence of auto-fluorescence in the eggs and planula larvae of the two species of *Eugymnanthea*, the eggs and embryos of many kinds of sea urchin display auto-fluorescence and the pluteus larvae emit green fluorescence (Nakamura *et al.*, 2005) though their function is unknown.

ACKNOWLEDGEMENTS

We thank Drs Ferdinando Boero, Stefano Piraino, Christina Maria Buia, Genuario Belmonte, Antonio Terlizzi, Maria Pia Miglietta, Akio Murakami, and also Roberto Pagliara and Cataldo Pierri for their kind help with various ways of support during the study. Thanks are due to anonymous referees, and editors of *JMBA* for critical reading of the manuscript.

REFERENCES

- Baba K., Miyazono A., Matsuyama K., Kohno S. and Kubota S. (2007) Occurrence and detrimental effects of the bivalve-inhabiting hydroid *Eutima japonica* on juvenile of the Japanese scallop *Mizuhopecten yessoensis* in Funka Bay, Japan: relationship to juvenile massive mortality in 2003. *Marine Biology (Berlin)* 151, 1977–1987.
- Campbell A.K. (1974) Extraction, partial-purification and properties of obelin, calcium activated luminescent protein from hydroid *Obelia geniculata*. *Biochemical Journal* 143, 411–418.
- Davenport D. and Nicol J.A.G. (1955) Luminescence in Hydromedusae. *Proceedings of the Royal Society, Biological Sciences Series B* 144, 399–411.
- Govindarajan A.F., Piraino S., Gravili C. and Kubota S. (2005) Species identification of bivalve-inhabiting marine hydrozoans of the genus *Eugymnanthea*. *Invertebrate Biology* 124, 1–10.
- Haddock S.H.D. and Case J.F. (1999) Bioluminescence spectra of shallow and deep-sea gelatinous zooplankton: ctenophores, medusae and siphonophores. *Marine Biology (Berlin)* 133, 571–582.
- Haddock S.H.D., Rivers T.J., Robison B.H. (2001) Can coelenterates make coelenterazine? Dietary requirement for luciferin in cnidarian bioluminescence. *Proceedings of the National Academy of Sciences of the United States of America* 98, 11148–11151.
- Haddock S.H.D., Dunn C.W., Pugh P.R. and Schnitzler C.E. (2005) Bioluminescent and red-fluorescent lures in a deep-sea siphonophore. *Science* 309, 263.
- Harvey E.N. (1935) Studies on bioluminescence. XIII. Luminescence in the coelenterates. *Biological Bulletin. Marine Biological Laboratory, Woods Hole* 41, 280–287.
- Kubota S. (1989) Systematic study of a paedomorphic derivative hydrozoan *Eugymnanthea* (Thecata–Leptomedusae). *Zoological Science* 6, 147–154.
- Kubota S. (1991) Crossing-experiments between Japanese populations of three hydrozoans symbiotic with bivalves. *Hydrobiologia* 216/217, 429–436.
- Kubota S. (1992) Four bivalve-inhabiting hydrozoans in Japan differing in range and host preference. *Scientia Marina* 56, 149–159.
- Kubota S. (2000) Parallel, paedomorphic evolutionary processes of the bivalve-inhabiting hydrozoans (Leptomedusae, Eirenidae) deduced from the morphology, life cycle and biogeography, with special reference to taxonomic treatment of *Eugymnanthea*. *Scientia Marina* 64, Supplement 1, 241–247.
- Kubota S. (2004) Some new and reconfirmed biological observations in two species of *Eugymnanthea* (Hydrozoa, Leptomedusae, Eirenidae) associated with bivalves. *Biogeography* 6, 1–5.
- Leutenegger A., Kredel S., Gundel S., D'Angelo C., Salih A. and Wiedenmann J. (2007) Analysis of fluorescent and non-fluorescent sea anemones from the Mediterranean Sea during a bleaching event. *Journal of Experimental Marine Biology and Ecology* 353, 221–234.
- Morin J.G. (1983) Coastal bioluminescence: patterns and functions. *Bulletin of Marine Science* 33, 787–817.
- Morin J.G. and Reynolds G.T. (1969) Fluorescence and time distribution of photon emission of bioluminescent photocytes in *Obelia geniculata*. *Biological Bulletin. Marine Biological Laboratory, Woods Hole* 137, 410.
- Morin J.G. and Reynolds G.T. (1974) The cellular origin of bioluminescence in the colonial hydroid *Obelia*. *Biological Bulletin. Marine Biological Laboratory, Woods Hole* 147, 397–410.
- Nakamura S., Mikamori M., Hiramatsu M., Eura S., Takamoto H. and Watanabe M. (2001) Spectacular fluorescence emission in sea urchin larvae. *Zoological Science* 18, 807–810.
- Nicol J.A.C. (1962) Animal luminescence. *Advances in Comparative Physiology and Biochemistry* 1, 217–273.
- Piraino S., Todaro C., Geraci S. and Boero F. (1994) Ecology of the bivalve-inhabiting hydroid *Eugymnanthea inquilina* in the coastal sounds of Taranto (Ionian Sea, SE Italy). *Marine Biology (Berlin)* 118, 695–703.
- Rayyan A., Christidis J. and Chintiroglou C.C. (2002) First record of the bivalve-inhabiting hydroid *Eugymnanthea inquilina* in the eastern Mediterranean Sea (Gulf of Thessaloniki, north Aegean Sea, Greece). *Journal of the Marine Biological Association of the United Kingdom* 82, 851–853.
- Shimomura O. (2005) The discovery of aequorin and green fluorescent protein. *Journal of Microscopy* 217, 3–15.
- Vysotskii E.S., Bondar V.S., Gitelson I., Petrunyaka V.V., Gamalei I.A. and Kaulin A.B. (1990) Extraction, some properties and application of obelin, calcium activated photoprotein. In Jezowska-Trzebiatowska B., Kochel B., Slawinski J. and Sterk W. (eds) *Biological luminescence*. Singapore and New Jersey: World Scientific, pp. 386–395.
- Vysotskii E.S., Trofimov K.P., Bondar V.S. and Gitelson J.I. (1993) Luminescence of Ca²⁺ activated photoprotein obelin initiated by NaOCl and MnCl₂. *Journal of Bioluminescence and Chemiluminescence* 8, 301–305.
- and
- Vysotskii E.S., Trofimov C.P., Bondar V.S., Frank L.A., Markova S.V. and Illarionov B.A. (1995) Mn⁻²⁺ activated luminescence of the photoprotein obelin. *Archives of Biochemistry and Biophysics* 316, 92–99.

Correspondence should be addressed to:

Shin Kubota
Seto Marine Biological Laboratory, Field Science Education and Research Center,
Kyoto University, Shirahama, Nishimuro, Wakayama 649-2211, Japan
email: shkubota@medusanpolyp.mbox.media.kyoto-u.ac.jp