TITLE:

Effects of a 4-week static stretch training program on passive stiffness of human gastrocnemius muscle-tendon unit in vivo(Dissertation_全文)

AUTHOR(S):

Nakamura, Masatoshi

CITATION:

Nakamura, Masatoshi. Effects of a 4-week static stretch training program on passive stiffness of human gastrocnemius muscle-tendon unit in vivo. 京都大学, 2014, 博士(人間健康科学)

ISSUE DATE:

2014-03-24

URL:

<https://doi.org/10.14989/doctor.k18200>

RIGHT:

Effects of a four-week static stretch training program on passive stiffness of human gastrocnemius muscle-tendon unit in vivo.

Masatoshi Nakamura^{1, 2*}, Tome Ikezoe¹, Yohei Takeno¹, Noriaki Ichihashi¹

1) Human Health Sciences, Graduate School of Medicine, Kyoto University

2) Research Fellow of the Japan Society for the Promotion of Science, Japan.

Abstract

Static stretch is commonly used to prevent contracture and to improve joint mobility. However, it is unclear whether the components of the muscle-tendon unit are affected by a static stretch training program. This study investigated the effect of a four-week static stretch training program on the viscoelastic properties of the muscle-tendon unit and muscle. The subjects comprised eighteen male participants (mean age: 21.4 years $\pm$ 1.7). The range of motion (ROM), passive torque, muscle-tendon junction (MTJ) displacement and muscle fascicle length of the gastrocnemius muscle were assessed using both ultrasonography and a dynamometer whilst the ankle was passively dorsiflexed. After the initial test, the participants were assigned either to a group that stretched for four-week (N = 9) or to a control group (N = 9). The tests were repeated after the static stretch training program.

ROM and MTJ displacement significantly increased, and the passive torque at 30 deg significantly decreased, in the stretching group after the study period. However, there was no significant increase in muscle fascicle length. These results suggest that a four-week static stretch training program changes the flexibility of the overall MTU without causing concomitant changes in muscle fascicle length.

Key words:

Static stretch, Long-term effects, Ultrasonography, Muscle tendon unit, Gastrocnemius

Introduction

Static stretch (SS) is commonly employed to prevent contracture and to improve joint mobility. There have been many studies about the effects of SS training programs (Chan et al. 2001; Covert et al. 2010; Gajdosik 1991, 2001; Marques et al. 2009; Reid and McNair 2004; Santonja Medina et al. 2007), and in these studies SS training programs were shown to have increased the maximum range of motion (ROM). However, the measurement of maximum ROM has a number of limitations. For example, the method is influenced by factors such as pain, stretch tolerance and reflex activation of the agonist muscle (Magnusson et al. 1996a; McHugh et al. 1998). It has been pointed out that an alternative approach to assess the passive torque would be useful for determining the stiffness of overall muscle-tendon unit (MTU) (Toft et al. 1989).

Many studies have indicated that SS training decreased MTU stiffness, and changed MTU viscosity and elasticity (Feland et al. 2001; Kubo et al. 2002; Mahieu et al. 2007; Willy et al. 2001). On the other hand, it has also been reported that changes occurred in maximum ROM, but not in MTU stiffness, after 3- and 8-week SS training programs (Ben and Harvey 2010; Bjorklund et al. 2001; Folpp et al. 2006; Magnusson et al. 1996; Ylinen et al. 2009). Weppeler and Magnusson (2010) pointed out that the improvement in maximum ROM at the conclusion of the SS training program was predominantly due to a modification in stretch tolerance (i.e. subjects' sensation). Thus, it is still unclear whether any decrease in MTU stiffness after stretching can be attributed to alterations in the muscle, or alterations in another component of the viscoelastic MTU.

Morse et al. (2008) reported that muscle and tendon stiffness could be estimated using ultrasonography to measure the movement of the myotendinous junction (MTJ) during passive stretching. In addition, they reported that there was a decrease in the stiffness of the overall MTU and muscle (no change was apparent in the tendon) immediately after a five-minute SS session. However, it is unclear whether long-term SS training program affects muscle stiffness or other viscoelastic properties of MTU during passive stretching.

The purpose of this study was to determine the effect of a four-week SS training program on the human gastrocnemius MTU, i.e. the properties of the muscle and other structures in vivo.

Methods

Participants

Eighteen healthy males volunteered for this study (mean age: 21.4 years $\pm$ 1.7, mean height: 172.6 cm $\pm$ 6.6, mean weight: 63.0 kg $\pm$ 5.2). Subjects with a history of neuromuscular disease or musculoskeletal injury involving their lower limbs were excluded. This study was approved by the ethics committee of the Kyoto University Graduate School and by the Faculty of Medicine (E-816), and it conformed to the principles set out in the Declaration of Helsinki.

After successfully completing the initial test session, participants were randomly assigned to either a stretching group (N = 9) or a control group (N = 9). The

characteristics of the subjects and the ROM of dorsiflexion for each group are presented in Table 1. Unpaired t-test results indicated no significant differences for these characteristics between the 2 groups.

SS training program

The participants in the stretching group were required to hold the SS position for 60 s and complete 2 repetitions, for a total of 120 s of stretch during each session. Each session was performed on a daily basis over a 4-week period. The stretch manoeuvre was self-administered by participants. The participants stood with arms supported on the wall anterior to the body. Both legs were straight, the hip in neutral rotation, with only the forefoot resting on the platform. The ankle joint was dorsiflexed progressively by leaning towards the wall until they felt the largest stretch that they were willing to tolerate. The participants filled out a form on a daily basis to register compliance. They were instructed not to initiate any new form of training. They did not perform any stretch training on the day of the test.

Measurements

All measurements were performed prior to the SS training program (PRE) and after the SS training program (POST). The participants were familiarised with the procedure and were instructed to remain relaxed during measurement. To take into account the influence of the intervention, POST measurements in the stretching group were performed at least 24 hours after the last SS session. The person taking the measurements was not blinded to the intervention.

ROM of dorsiflexion

The participants were secured at the hip in a prone position on the dynamometer (MYORET RZ-450; Kawasaki Heavy Industries, Kobe, Japan), with their knees in full extension and the foot of their dominant leg attached securely to the footplate of the dynamometer. The footplate of the dynamometer was moved at a constant velocity of 1 degree/s by the motor control of the dynamometer, starting from 0 degrees to the dorsiflexion angle at which the participants felt discomfort or pain. This dorsiflexion angle was defined as the ROM.

Passive torque

Passive plantar flexion torque was measured using the dynamometer; participants held a position similar to that used during the measurement of ROM. The footplate of the dynamometer was moved at a constant velocity of 1 degree/s from 0 to 30 degrees of dorsiflexion, which was achieved by all participants without pain. Passive plantar flexion torque was measured at 0 and 30 degrees during passive ankle dorsiflexion.

Ultrasound measurements

B-mode ultrasonography (Famio Cube SSA-520A; Toshiba Medical Systems Corporation, Tochigi, Japan) was used to determine the displacement of MTJ of the medial head of the gastrocnemius muscle (GM) during passive ankle dorsiflexion. MTJ was identified as described Maganaris and Paul (1999) and visualised as a continuous sagittal plane ultrasound image using an 8 MHz linear-array probe. An acoustically reflective marker made from softened vinyl film, was placed between the skin and the probe to confirm that the probe did not move during measurements (Morse et al. 2008). We defined MTJ displacement as the distance between MTJ and the acoustically reflective marker secured to the probe. A custom-made fixation device was used to secure the probe to the skin. Ultrasound images of the MTJ were quantified using the open source digital measurement software (Image J, NIH, USA). The MTJ was identified at the inner-most edges of the fascia surrounding the muscle where it fuses with the tendon, to accurately measure MTJ. MTJ displacement was measured between 0 and 30 degrees of ankle dorsiflexion (Fig. 1).

Another B-mode ultrasonography (LOGIQe; GE Healthcare Japan, Tokyo, Japan) was used to obtain fascicle length, which was calculated from the muscle thickness and pennation angle. GM thickness was measured halfway along the lower leg; i.e. at a point equidistant from the lateral malleolus of the fibula and the lateral condyle of the tibia. The pennation angle of GM was determined from the angle of fascicle insertion into the deep aponeurosis. Muscle thickness and the pennation angle were measured at 0 and 30 degrees of ankle dorsiflexion. Movement of dynamometer was stopped at 0 and 30 degrees by the motor, and the ultrasonography and dynamometer measurements were performed synchronously. The fascicle length was calculated using the following formula (Kumagai et al. 2000):

$$\text{Fascicle length} = \text{muscle thickness} / \sin(\text{pennation angle})$$

Morse et al. (2008) reported that the contribution of the fascicle to MTU length is proportional to the cosine of the angle of pennation; i.e. the fascicle length resolved along the axis of the muscle (resolved fascicle length). The pennation angle decreases with ankle dorsiflexion while the fascicle length and resolved fascicle length increase. The change in factors other than muscle fascicle length was estimated from the change in the resolved fascicle length (Δ resolved fascicle length) and MTJ displacement (Δ MTJ) from 0 to 30 degrees of ankle dorsiflexion, elaborated as follows (Morse et al. 2008):

Change in factors other than fascicle length = Δ MTJ at 30 degrees – Δ resolved fascicle length (Fig. 2)

EMG

We monitored the electromyographic (EMG) activity in the GM during the test procedure (for about 40 sec) to confirm that the subject was relaxed and to ensure the absence of high EMG activity. EMG activity was recorded using an EMG system (TeleMyo2400; Noraxon USA, Inc., Scottsdale, AZ, USA) with surface electrodes (11mm; Blue Sensor N, Ambu, Denmark). EMG activity was calculated using the root mean square (RMS), and full wave rectification was performed using a RMS smoothing algorithm at a window interval of 50 ms. The EMG activity within 3 sec was calculated

during isometric maximum voluntary contraction (MVC) of the ankle plantar flexors with the ankle kept at 0 deg. The EMG activity recorded during the tests was expressed as a percentage of MVC. The EMG sampling rate was 1500 Hz.

Reliability of the ultrasound measurements of MTJ

The ultrasonographic measurement of the MTJ displacement from 0 deg to 30 deg was repeated twice on different days to assess test-retest reliability (n=7). Tests were performed with at least 1 week interval, but not longer than 2 weeks, between the two tests.

Statistics

SPSS (version 17.0; SPSS Japan INC., Tokyo, Japan) was used for statistical analyses.

For all variables, significant differences between PRE and POST were determined in both the stretching and control groups using a paired t-test. Significance of differences between the stretching and control groups at PRE was assessed using an unpaired t-test. Significant differences were determined between within-group changes, defined as POST minus PRE, in both the stretching and control group using an unpaired t-test. In addition, a two-way ANOVA [(groups) $\times$ (test time)] was used to analyse the interaction effects. Differences were considered statistically significant at an alpha level of $P < 0.05$.

The reliability of MTJ displacement measurements was examined using the intraclass correlation coefficient (ICC). Descriptive data are shown as means $\pm$ S.E.M and 95% confidence intervals (CI) for the differences in changes score.

Results

Reliability and validity of ultrasound measurements

ICC for MTJ displacement measurements was 0.985 (trial 1: 1.21 ± 0.08 cm, trial 2: 1.22 ± 0.09 cm). During all tests, the EMG activity of GM was $<2\%$ MVC, which confirmed the lack of a contractile component contribution to the passive torque, MTJ displacement, muscle thickness and pennation angle.

Change in the ROM

There was no significant difference in the ROM between the stretching and control groups at PRE. The ROM increased significantly after SS training in the stretching group, whereas no significant difference was noted between PRE and POST in the control group (Table 2). A two-way ANOVA showed a significant interaction effect ($F = 10.9$, $P < 0.01$).

Changes in the passive torque and MTJ displacement

There were no significant differences in the passive torque at 0 and 30 degrees dorsiflexion between the stretching and control groups at PRE. The passive torque at 0

degrees showed no significant differences between PRE and POST in either the stretching or the control group. However, the passive torque value at 30 degrees decreased significantly after SS training, whereas no significant difference was noted between PRE and POST in the control group (Table 2). A significant interaction effect was observed ($F = 15.1$, $P < 0.01$).

There was no significant difference in MTJ displacement between the stretching and the control groups at PRE. MTJ displacement increased significantly at 30 degrees dorsiflexion after SS training, whereas no significant difference was noted between PRE and POST in the control group. There was a significant interaction effect ($F = 43.6$, $P < 0.01$).

Change in the fascicle length and resolved fascicle length

Table 2 shows changes in the fascicle length and resolved fascicle length. The fascicle length and resolved fascicle length at 0 and 30 degrees showed no significant differences between PRE and POST in either stretching or control group. The fascicle length at 0 and 30 degrees showed no significant interaction effects ($F = 0.09$; $P = 0.76$, $F = 0.03$ and $P = 0.88$, respectively) and the resolved fascicle length at 0 and 30 degrees also showed no significant interaction effects ($F = 0.10$; $P = 0.76$ and $F = 0.02$; $P = 0.89$, respectively).

Change in factors other than muscle fascicle length

Regarding the change in resolved fascicle length (Δ resolved fascicle length) from 0 to 30 degrees of ankle dorsiflexion, no significant difference was observed between PRE and POST in either stretching or control group (Table 2). No significant interaction effect was observed ($F = 0.85$, $P = 0.37$).

With regard to the value for factors other than muscle fascicle length, calculated by subtracting Δ resolved fascicle length from Δ MTJ from 0 to 30 degrees, a significant increase was observed after SS training in the stretching group. However, no significant difference was noted between PRE and POST in the control group. A significant interaction effect was observed ($F = 84.4$, $P < 0.01$).

Discussion

Reliability and validity of ultrasound measurements

We assessed the test-retest reliability of our ultrasound measurement of MTJ displacement using the ICC. The measured MTJ displacement value demonstrated substantial agreement because the ICC score was 0.985.

During measurement, the EMG activity of GM was very low ($<2\%$). It thus seems rational that no voluntary or reflex contraction occurred during measurement, which indicates that the measured passive torque and MTJ displacement values reflect passive properties of MTU.

The change in MTU after SS training program

In this study, the ROM increased significantly in the stretching group, although no significant change was found in the control group. This result suggests that a four-week SS training program is effective for increasing ROM, which is consistent with previous studies (Chan et al. 2001; Covert et al. 2010; Gajdosik 1991, 2001; Marques et al. 2009; Reid and McNair 2004; Santonja Medina et al. 2007). Animal studies showed that the number of sarcomeres in a series of muscles can be changed by prolonged immobilisation in extreme positions (Goldspin.G et al. 1974; Tabary et al. 1972; Williams and Goldspink 1978), leading to speculation that increases of maximum ROM may be related to increases in the number of sarcomeres in series, and a concurrent increase in length of the stretched muscles (Chan et al. 2001; Gajdosik 1991, 2001; Reid and McNair 2004). However, in our study, the fascicle length (calculated from muscle thickness and pennation angle) did not change after SS training program, which suggests that the muscle length was not increased by SS training in vivo.

Morse et al. (2008) investigated the immediate effects of five-minute SS sessions on not only the flexibility of overall MTU but also on muscle and tendon stiffness. However, the effects of a long-term SS training program on MTU properties remain unclear. In this study, the passive torque at 30 degrees decreased significantly from PRE to POST in the stretching group, whereas no significant changes were evident over time in the control group. These results suggested that the SS training program decreased the stiffness of the overall MTU. In addition, MTJ displacement increased significantly in the stretching group, whereas no significant change was noted in the control group. These results suggest that the stiffness of the gastrocnemius muscle decreased after the four-week SS training program. Our findings show that the SS training program is effective in decreasing MTU stiffness, in particular muscle stiffness, which is consistent with the immediate effect of SS (Morse et al. 2008). In this study, POST measurements in the stretching group were performed at least 24 hours after the last SS session to exclude the influence of the intervention. Therefore, we consider that these results depend not on the immediate effects of stretching, but on the long-term effects of the four week intervention.

With regard to the mechanism of decrease in muscle stiffness, Morse et al. (2008) reported that MTJ displacement and the change in resolved fascicle length were virtually identical before passive stretching, whereas MTJ displacement was 0.19 cm greater than the change in resolved fascicle length after 5 min of SS training. They therefore concluded that the additional displacement of MTJ was, at least in part, the result of changes in structures other than the muscle fascicle. In this study, MTJ displacement in the stretching group PRE was almost completely explained by the change in fascicle length from 0 to 30 degrees of ankle dorsiflexion, since the change in factors other than muscle fascicle length before SS training was negligible (0.01 cm). However, a significantly larger change in factors other than muscle fascicle length was observed after SS training (PRE: 0.01 cm, POST: 0.24 cm). On the other hand, there were no significant changes in fascicle length and resolved fascicle length after SS training. This suggests that the increased MTJ displacement observed after SS training might be associated with factors other than muscle fibre length; i.e. components of MTU proximal to MTJ.

Muscle fibres are surrounded by a complex connective tissue network, which includes the endomysium, perimysium and epimysium. Perimysium bundles contain several muscle fibres, while the endomysium surrounds individual muscle fibres. Furthermore, the epimysium covers the entire muscle. Gajdosik et al. (2001) suggested that the cytoskeleton of the sarcomere and the intramuscular connective tissue are made up of parallel elastic components; for example, the endomysium, perimysium and epimysium; which cause passive tension and therefore modification of these tissues could lead to a change in MTU stiffness. Furthermore, Purslow (1989) reported that the connective tissue, particularly the perimysium, is a major extracellular contributor to passive stiffness. Thus, changes in properties of the intramuscular connective tissue that cause passive tension are considered to be related to a decrease in muscle stiffness. In addition, Kubo et al. (2002) reported that three-week SS training program decreased MTU stiffness but not tendon stiffness, which implied that SS training program affected the connective tissue elements in parallel with the muscle fibres. Taken together, changes in properties of the intramuscular connective tissue such as the endomysium, perimysium and epimysium instead of lengthening of muscle fiber, may also contribute to the decrease in muscle stiffness found in our study.

There are some limitations of this study. First, the person taking measurements was not blinded to the intervention. Second, the MTJ of the medial head of the gastrocnemius muscle is not sharply delineated. In performing the MTJ measurements, we identified the MTJ at the inner-most edges of the fascia surrounding the muscle where it joins the tendon. The test-retest reliability of the measurement of MTJ displacement in this study was very high. Therefore, we consider that the error in measuring MTJ displacement was negligible. The third limitation of this study is that passive torque is influenced not only by the gastrocnemius muscle but also by joint capsules, ligaments and other muscles such as the soleus muscle. Further work is needed to clarify the effect of long-term SS training program focusing on these factors.

In conclusion, the results of this study suggests that a four-week SS training program decreases passive torque and increases the MTJ displacement of the gastrocnemius at 30 degrees dorsiflexion, without causing changes in muscle fascicle length.

Conflict of Interest Statement

The authors have no conflicts of interest to declare.

References

- Ben M, Harvey LA (2010) Regular stretch does not increase muscle extensibility: a randomized controlled trial. *Scand J Med Sci Sports* 20:136–144
- Bjorklund M, Hamberg J, Crenshaw AG (2001) Sensory adaptation after a 2-week stretching regimen of the rectus femoris muscle. *Arch Phys Med Rehabil* 82:1245–1250
- Chan SP, Hong Y, Robinson PD (2001) Flexibility and passive resistance of the hamstrings of young adults using two different static stretching protocols. *Scand J Med Sci Sports* 11:81–86
- Covert CA, Alexander MP, Petronis JJ, Davis DS. (2010) Comparison of ballistic and static stretching on hamstring muscle length using an equal stretching dose. *J Strength Cond Res* 24:3008–3014
- Feland JB, Myrer JW, Schulthies SS, Fellingham GW, Measom GW (2001) The effect of duration of stretching of the hamstring muscle group for increasing range of motion in people aged 65 years or older. *Phys Ther* 81:1110–1117
- Folpp H, Deall S, Harvey LA, Gwinn T (2006) Can apparent increases in muscle extensibility with regular stretch be explained by changes in tolerance to stretch? *Aust J Physiother* 52:45–50
- Gajdosik RL (1991) Effects of static stretching on the maximal length and resistance to passive stretch of short hamstring muscles. *J Orthop Sports Phys Ther* 14:250–255
- Gajdosik RL (2001) Passive extensibility of skeletal muscle: review of the literature with clinical implications. *Clin Biomech* 16:87–101
- Goldspin G, Tabary C, Tabary JC, Tardieu C, Tardieu G (1974) Effect of Denervation on Adaptation of Sarcomere Number and Muscle Extensibility to Functional Length of Muscle. *J Physiol* 236:733–742
- Kubo K, Kanehisa H, Fukunaga T (2002) Effect of stretching training on the viscoelastic properties of human tendon structures *in vivo*. *J Appl Physiol* 92:595–601
- Kumagai K, Abe T, Brechue WF, Ryushi T, Takano S, Mizuno M (2000) Sprint performance is related to muscle fascicle length in male 100-m sprinters. *J Appl Physiol* 88:811–816
- Maganaris CN, Paul JP (1999) *In vivo* human tendon mechanical properties. *J Physiol* 521:307–313
- Magnusson SP, Simonsen EB, Aagaard P, Kjaer M (1996a) Biomechanical responses to repeated stretches in human hamstring muscle *in vivo*. *Am J Sports Med* 24:622–628
- Magnusson SP, Simonsen EB, Aagaard P, Sorensen H, Kjaer M (1996b) A mechanism for altered flexibility in human skeletal muscle. *J Physiol* 497:291–298
- Mahieu NN, McNair P, De Muynck M, Stevens V, Blanckaert I, Smits N, Witvrouw E (2007) Effect of static and ballistic stretching on the muscle-tendon tissue properties. *Med Sci Sports Exerc* 39:494–501
- Marques AP, Vasconcelos AA, Cabral CM, Sacco IC (2009) Effect of frequency of static stretching on flexibility, hamstring tightness and electromyographic activity. *Braz J Med Biol Res* 42:949–953

- 1 McHugh MP, Kremenec II, Fox MB, Gleim GW (1998) The role of mechanical and
2 neural restraints to joint range of motion during passive stretch. *Med Sci Sports*
3 *Exerc* 30:928–932
- 4 Morse CI, Degens H, Seynnes OR, Maganaris CN, Jones DA (2008) The acute effect of
5 stretching on the passive stiffness of the human gastrocnemius muscle tendon unit.
6 *J Physiol* 586:97–106
- 7 Purslow PP (1989) Strain-induced reorientation of an intramuscular connective tissue
8 network: implications for passive muscle elasticity. *J Biomech* 22:21–31
- 9 Reid DA, McNair PJ (2004) Passive force, angle, and stiffness changes after stretching
10 of hamstring muscles. *Med Sci Sports Exerc* 36:1944–1948
- 11 Tabary JC, Tabary C, Tardieu G, Tardieu C, Goldspin G (1972) Physiological and
12 Structural Changes in Cats Soleus Muscle Due to Immobilization at Different
13 Lengths by Plaster Casts. *J Physiol* 224:231–244
- 14 Toft E, Espersen GT, Kalund S, et al. Passive tension of the ankle before and after
15 stretching. *Am J Sports Med.* 1989; 17:489-94.
- 16 Santonja Medina FM, Sainz De Baranda Andujar P, Rodriguez Garcia PL, Lopez
17 Minarro PA, Canteras Jordana M (2007) Effects of frequency of static stretching
18 on straight-leg raise in elementary school children. *J Sports Med Phys Fitness*
19 47:304–308
- 20 Webright WG, Randolph BL, Perrin DH (1997) Comparison of nonballistic active knee
21 extension in neural slump position and static stretch techniques on hamstring
22 flexibility. *J Orthop Sport Phys* 26:7–13
- 23 Weppler CH, Magnusson SP (2010) Increasing muscle extensibility: a matter of
24 increasing length or modifying sensation? *Phys Ther* 90:438–449
- 25 Wessling KC, Devane DA, Hylton CR (1987) Effects of Static Stretch Versus Static
26 Stretch and Ultrasound Combined on Triceps Surae Muscle Extensibility in
27 Healthy Women. *Phys Ther* 67:674–679
- 28 Williams PE, Goldspink G (1978) Changes in Sarcomere Length and Physiological
29 Properties in Immobilized Muscle. *J Anat* 127:459–468
- 30 Willy RW, Kyle BA, Moore SA, Chleboun GS (2001) Effect of cessation and
31 resumption of static hamstring muscle stretching on joint range of motion. *J Orthop*
32 *Sports Phys Ther* 31:138–144
- 33 Ylinen J, Kankainen T, Kautiainen H, Rezasoltani A, Kuukkanen T, Hakkinen A (2009)
34 Effect of stretching on hamstring muscle compliance. *J Rehabil Med* 41:80–84

Figure 1.

Ultrasound imaging showing the measurements taken to determine MTJ displacement

An acoustically reflective marker (X) was placed between the skin and the ultrasonic probe to confirm that the probe did not move during measurements. The distance between X and the myotendinous junction (MTJ) of the medial head of the gastrocnemius muscle (GM) was measured at 0 and 30 degrees, and the difference between 0 and 30 degrees was defined as “MTJ displacement”.

Figure 2.

Schematic diagram showing measurements taken to determine fascicle length, resolved fascicle length and changes in factors other than muscle fascicle length

d: muscle thickness, θ : pennation angle, FL: fascicle length, RFL: resolved fascicle length, MTJ: myotendinous junction

$$FL = d/\sin\theta, RFL = FL \times \cos\theta$$

Changes in factors other than fascicle length = MTJ displacement – (RFL' – RFL)

Table 1.

Characteristics of subjects in the stretching group and those in the control group.

	stretching group (N=9)	control group (N=9)
age (years)	21.1±2.3 (19-26)	21.8±0.8 (20-23)
height (cm)	173.8±6.8 (163-184)	171.4±6.6 (165-186)
weight (kg)	64.4±4.9 (54-70)	61.6±5.4 (53-72)
ROM (deg) ^a	32.9±2.8 (30-36)	32.0±3.0 (30-37)

Data are means ± S.E.M. (95% confidence interval)

^aROM=Range of Motion

Table 2.

Changes in variables between PRE and POST in both the stretching and control groups.

	Stretching Group			Control Group			P-value	
	PRE	Change ^f	95%CI ^g	PRE	Change ^f	95%CI ^g	of change ^h	Interaction effect ⁱ
ROM (deg) ^a	32.9±2.8	6.7±1.7**	2.8 to 10.5	32.0±3.0	-0.7±0.7	-2.2 to 0.9	P < 0.01	P < 0.01
Passive torque at 0deg (Nm)	3.6±0.6	1.0±0.7	-0.6 to 2.5	4.0±0.4	0.0±0.3	-0.8 to 0.8	P=0.21	P = 0.21
at 30 deg (Nm)	46.2±5.6	-6.2±1.1**	-8.8 to -3.6	37.4±2.8	2.2±1.8	-2.0 to 6.4	P < 0.01	P < 0.01
MTJ displacement at 30 deg (cm) ^b	0.93±0.06	0.40±0.06**	0.27 to 0.53	1.03±0.09	0.01±0.01	-0.02 to 0.04	P < 0.01	P < 0.01
fascicle length at 0 deg (cm)	5.9±0.5	-0.3±0.4	-1.3 to 0.8	5.8±0.3	0.0±0.1	-0.2 to 0.2	P=0.60	P = 0.76
at 30 deg (cm)	6.8±0.5	-0.1±0.4	-1.2 to 1.0	6.7±0.3	0.0±0.0	0.0 to 0.1	P=0.79	P = 0.88
difference of fascicle length (cm) ^c	0.89±0.05	0.15±0.04	-0.04 to 0.33	0.98±0.08	0.03±0.01	-0.22 to 0.28	P=0.38	P = 0.43
resolved fascicle length at 0 deg (cm)	5.7±0.5	-0.3±0.4	-1.3 to 0.8	5.5±0.3	0.1±0.3	-0.1 to 0.2	P=0.60	P = 0.76
at 30 deg (cm)	6.6±0.5	-0.1±0.4	-1.2 to 0.4	6.6±0.3	0.0±0.3	0.0 to 0.1	P=0.81	P = 0.89
Δ resolved fascicle length (cm) ^d	0.92±0.06	0.17±0.07	-0.02 to 0.36	1.00±0.08	0.03±0.02	-0.01 to 0.08	P=0.14	P = 0.37
factors other than fascicle length (cm) ^e	0.01±0.01	0.22±0.02**	0.17 to 0.27	0.03±0.01	-0.01±0.01	-0.05 to 0.02	P < 0.01	P < 0.01

** : P < 0.01 significant difference in change between PRE and POST.

Data are means ± S.E.M.

^aROM = Range of Motion

^bMTJ = Myotendinous junction displacement at 30 deg

^c difference of fascicle length = fascicle length at 30 deg – fascicle length at 0 deg

^dΔ resolved fascicle length = resolved fascicle length at 30 deg – resolved fascicle length at 0 deg

^e factors other than fascicle length = MTJ displacement – Δ resolved fascicle length

^fChange = POST value – PRE value

^g 95 % CI: 95 % confidence intervals

^h Difference of change was determined in both stretching and control group using an unpaired t-test.

ⁱ Interaction effect was determined using a two-way ANOVA [(groups) × (test time)].