TITLE:

Development and validation of a short scale to measure how social relationships support the continuous and conscious endeavour to lose weight.

AUTHOR(S):

Takada, Akemi; Takahashi, Yoshimitsu; Nishimura, Shuzo; Kosugi, Shinji

CITATION:

Takada, Akemi ...[et al]. Development and validation of a short scale to measure how social relationships support the continuous and conscious endeavour to lose weight.. Journal of biosocial science 2014, 46(5): 561-579

ISSUE DATE:

2014-09

URL:

<http://hdl.handle.net/2433/189433>

RIGHT:

© Cambridge University Press 2014; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

Development and validation of a short scale to measure how social relationships support
the continuous and conscious endeavour to lose weight

AKEMI TAKADA*, YOSHIMITSU TAKAHASHI**, SHUZO NISHIMURA†, SHINJI KOSUGI‡

*Department of Cognitive and Behavioral Sciences, Graduate School of Human and Environmental Studies, Kyoto University, Kyoto, Japan

**Department of Health Informatics, Health Administration, Kyoto University School of Public Health, Kyoto, Japan

†National Institute of Population and Social Security Research, Tokyo, Japan

‡Department of Medical Ethics, Health Administration, Kyoto University School of Public Health, Kyoto, Japan

Running Heading: Social Relationships to Prevent Obesity

Summary. This paper reports the development of a short scale (10 items) entitled Social Relationships to Prevent Obesity (SRPO), which examines how social relationships support the continuous and conscious endeavour to lose weight. The construct and criterion validity of this scale were ascertained in this study. Factor structure and reliability were examined using data from a randomized controlled trial. A confirmatory factor analysis of the SRPO revealed three relevant factors. The results suggest that the SRPO has both validity and clinical utility and can thus be used as a screening tool in weight-loss interventions and to assess the degree of and trends in self-control for weight loss in individuals. The scale can also be used to examine the environmental and self-

control problems faced by obese people—factors that should be considered when conducting weight-loss interventions.

Introduction

Obesity is typically prevented through dieting and exercise regimens, and these two measures are related to the individual's surroundings (Sallis *et al.*, 1987). Moreover, on theoretical grounds, lifestyle-related obesity (henceforth, obesity)—like most social and personal problems (smoking, substance abuse, etc.)—seems to involve a substantial component of deficient self-control (Tangney *et al.*, 2004). In the context of the present study, 'self-control' means a continuous and conscious endeavour to lose weight that leads to a modification of the daily habits that are responsible for current obesity and the failure of previous weight-loss efforts (Kan, 2007). This term is synonymous to what Frank (1988) termed 'commitment'. Effective weight-loss interventions should enable participants to acquire social support through a social network (Sallis *et al.*, 1987). Such support should promote the participants' self-control and commitment to weight loss, and aid in the elimination of habits that lead to obesity.

Factors influencing the maintenance of a healthy weight and lifestyle

Obesity researchers recognize the importance of not only biological but also environmental and behavioural factors in sustaining a healthful lifestyle. Thus, future policies aimed at addressing obesity should involve interventions that mitigate unhealthy behavioural and environmental influences. (Cutler & Glaeser, 2005; Christakis & Fowler, 2007; Eid *et al.*, 2008; Hill *et al.*, 2008; Li *et al.*, 2009; Yakusheva *et al.*, 2011). Specifically, people with deficient self-control require support from others and the environment in their continuous and conscious efforts to adopt and maintain the

behaviours that prevent obesity. These behaviours include consuming an optimal amount of food; paying attention to the amount of fat and salt in one's diet, particularly while preparing one's own meals; and avoiding a sedentary life style (Sallis *et al.*, 1992; Komlos *et al.* 2004; Cutler & Glaeser, 2005). In order for the participants of a weight-loss program to develop strong self-control, the intervention needs to be targeted accurately and demonstrate clearly to recipients the factors involved in successful weight loss.

Studies using behavioural modification theories have demonstrated the considerable influence of social relationships and social support—for example, from one's spouse and friends—on dieting and exercising (Sallis *et al.*, 1987; Prochaska *et al.*, 1992; Sallis *et al.*, 1992; Marcus & Simkin, 1994; Unger & Johnson, 1995; Deforche & De Bourdeaudhuij, 2000). The peer effect, in terms of social interactions with friends, family, and acquaintances within a social network, has been shown to influence weight problems (Wallston *et al.*, 1978; Cutler & Glaeser, 2005; Christakis & Fowler, 2007; Cohen-Cole & Fletcher, 2008; Fowler & Christakis, 2008; Renna *et al.*, 2008; Trogon *et al.*, 2008; Ali *et al.*, 2011; Fortin & Yazbeck, 2011; Yakusheva *et al.*, 2011; Dale, Williams, & Bowyer, 2012).

In recent years, the importance of peer support through expansive social networks for weight loss has been enthusiastically examined in light of the obesity pandemic. At least three dimensions of support have been identified: the existence and quantity of social relationships, the structure of those relationships, and their functional content. The first two dimensions are more correctly conceived of as aspects of a social network, while the last dimension captures the behaviours by which one person actually supports another (Sallis *et al.*, 1987; Manski, 1993; Steptoe & Ayers, 2005; McDowell & Newell, 1996).

Social relationships and support have a long-standing association with health. Instrumental, emotional, and ongoing support have been shown to be important to sustained behaviour change and health in research involving people living with chronic condition, such as diabetes, cancer, cardiovascular diseases, mental illness, and HIV and AIDS (Boothroyd and Fisher, 2010). Conversely, social isolation has been shown to predict mortality and morbidity (House, Landis, & Umberson, 1988; Brummett *et al.*, 2001; House, 2001). In particular, peer support can provide assistance with daily management tasks, provide social and emotional support to stay motivated and deal with the stress chronic disease often brings, and help people stay connected to clinical care and improve outcomes in self-management (Boothroyd and Fisher, 2010).

An important mediator in the relationship between self-control and individual behaviour is locus of control (LOC). LOC refers to generalized expectations about the determinants of one's circumstances. On the basis of their experiences and learning history, individuals come to expect that future outcomes will be determined by either internal factors such as their own actions or characteristics (i.e. internal locus of control) or by external factors and opportunities not dependent on their own efforts or abilities (i.e. external locus of control) (Rotter, 1954, 1966; Contrada & Goyal, 2005). This theory forms the basis for an important idea in the field of behaviour modification: an individual's health behaviour might successfully be modified by promoting an internal locus of control (Nir & Neumann, 1995). Researchers have found strong and consistent correlation between an external locus of control and failure to comply with healthful behaviours (Wallston *et al.*, 1978; Macgregor *et al.*, 1997; Fujita & Noguchi, 2009).

The need for a short scale to measure social support in the weight-loss context

Many interventions have been conducted using these basic social support and LOC theories, but research findings have not always been consistent (Sallis *et al.*, 1987; Gorin *et al.*, 2008; Bahr *et al.*, 2009; Finnerty *et al.*, 2010). With regard to social relationships, for instance, some people lose weight if supported in terms of diet and exercise by cooperative spouses, while others, naturally, are influenced by a shared environment of obesity-promoting habits and traditions. Dieters may be unaware of these influences and may not understand why they fail to lose weight despite their best efforts. This inconsistency is primarily due to the shortcomings of the instruments used to measure the influences of social relationships on obesity (Brownell & Stunkard, 1981; Black & Lantz, 1984; McLean *et al.*, 2003; Yakusheva *et al.*, 2011). The instruments used in these previous studies were not designed to directly address the self-control problems of people with excess weight. Further, these instruments typically have an excessively large number of questions (Funch *et al.*, 1986; Sallis *et al.*, 1987; Karlsson *et al.*, 1995; Yata *et al.*, 2003; Gruber, 2008; Sherrill-Mittleman *et al.*, 2009). Consequently, respondents might have submitted unreliable or inappropriate responses to questions on self-control, because people with poorly regulated self-control often show impatience and destructive patterns of persistence when confronted with a large number of questions (Stunkard & Messick, 1985; Baumeister & Heatherton, 1996; Kan, 2004; Tangney *et al.*, 2004; Wills *et al.*, 2007). An additional shortcoming of these instruments is that they fail to adequately measure psychological adjustment to and compliance with healthy behaviour, both of which need to be examined in a weight-loss study to assess participants' degree of self-control (Tangney *et al.*, 2004; Carver, 2005).

In order to overcome these shortcomings, a short scale titled Social Relationships to Prevent Obesity (SRPO) was developed in this study. Through items that place as

little burden on respondents as possible, the scale examines the social relationships that support self-control with regard to weight loss, taking into consideration family- and community-based physical activities and dietary habits and the social-communication environment. This short scale is one of the first scales developed to measure the degree of and trends in social support for self-control by using a minimum number of items related to weight control interventions.

The objective of the present study was to develop and evaluate the validity and reliability of the SRPO. For this purpose, it uses secondary data from Takada *et al.*, 2011. The SRPO is a refined and extended version of a previous scale titled Social Support to Counter Obesity (SSCO; Takada *et al.*, 2010); the SRPO differs from the SSCO in that the latter does not explicitly measure social support that promotes self-control.

Methods

Source of secondary data: the Takada *et al.* (2011) study

In a previous study, a tele-care intervention for weight-loss was assessed through a randomized controlled trial. The participants were registered members of a community health club in Kyoto, Japan, recruited through a public advertisement. They were all obese but otherwise healthy men and women between 20 and 70 years old who met the study's strict eligibility requirements (for more details, see Takada *et al.*, 2011). There were 118 participants at baseline, and 21 dropped out before the intervention started. The participants were administered a questionnaire in person at a health check-up conducted as a part of the study. The questionnaire included the SRPO and questions on dietary habits, such as (1) number of meals, snacks, instances of eating out, and alcoholic drinks consumed; (2) regular eating of breakfast; and (3) late-night meals. The questionnaire

also requested information about descriptive variables, including gender, age, body mass index, education, job status, marital status, income, and property owned (see Table 1). The remaining participants were randomized into two groups—a tele-care group and a self-help group—matched by age, sex, and body mass index. Of the 97 participants, only 66 completed the SRPO because some participants who were retired could not respond to some questions about level of support from employers (response rate: 55.9% [66/118]). The data from the 66 participants who completed the SRPO were used in the present study. The reliability and validity of the SRPO were examined using the participants' baseline data before the tele-care or self-help interventions. The research was approved by the ethics committee of the Graduate School of Medicine, Kyoto University. Analyses were conducted using SPSS v. 15.0 (SPSS Inc., Chicago, IL, USA).

Composition of the SRPO

To develop this short scale, existing instruments were reviewed and items were chosen that addressed peer support for weight loss and social interaction that promotes constructive behaviour and adjustment. Items were chosen from the following instruments, all of which have been shown to have good reliability and validity.

Social Adjustment Scale: Two questions were taken from the 54-item Social Adjustment Scale (Weissman & Bothwell, 1976), which is one of the few scales designed to measure adjustment to community living among both psychotherapy patients and healthy individuals (McDowell & Newell, 1996). The first question was 'How many times in the last two weeks have you gone out socially? For example, visited friends, gone to movies, bowling, church, restaurants, etc.?' The available response options ranged from 1 (None) to 5 (More than three times). The second question was 'How much time have you spent on hobbies or spare time interests during the last two weeks? For

example, bowling, sewing, gardening, sports, reading?’ The response options ranged from 1 (I did not spend any time on hobbies or watching TV) to 5 (I spend a lot of time on hobbies almost every day).

Rand Social Health Battery: Two questions were adapted from the 11-item Rand Social Health Battery (Donald & Ware, 1984), which is one of the few social health scales not designed for use with patients. This scale records social interactions and resources for social support but does not evaluate the subjective experience of support (McDowell & Newell, 1996). The first question was ‘To how many volunteer groups or organisations do you belong (e.g. church, temple, or shrine groups; clubs in the community; or parent groups)?’ The response options ranged from 1 (None) to 5 (More than three groups or organizations). For the second question—‘How active are you in the affairs of the groups or clubs to which you belong?’—the response options ranged from 1 (Do not belong to any groups or attend any meetings) to 4 (Very active, attend most meetings).

Social Support and Exercise Survey and Social Support and Eating Habits Survey: Two questions were taken from the Social Support and Exercise Survey and two from the Social Support and Eating Habits Survey. These instruments are two of four separate scales (with 43 total items) designed to assess social support for diet and exercise (Sallis *et al.*, 1987). These scales were developed using a behavioural modification theory known as the ecological model (an approach characterized by its focus on levels of influence from the individual to the community) (Sallis *et al.*, 1987; Ståhl *et al.*, 2001; Uechi, 2006). Two originally separate items—‘My family or friends exercised with me’ and ‘gave me helpful reminders to exercise’—were combined in the SRPO. Three other items were adopted verbatim: (1) ‘My family or friends helped plan

activities around my exercise’, (2) ‘My family or friends reminded me not to eat high fat, high salt foods’, and (3) ‘discussed my eating habit changes with me’. The response options ranged from 1 (Strongly disagree/Not at all) to 5 (Strongly agree/Very often).

Medical Outcomes Study Social Support Survey: Two questions were taken from the 12-item Medical Outcomes Study Social Support Survey (Sherbourne & Stewart, 1991), which, though designed for use in chronically ill patients, is also universally applicable owing to its sound validity and reliability, despite being relatively short (McDowell & Newell, 1996). This instrument attempts to determine how often various kinds of support are available to the respondent. The following two items were chosen: ‘[How often do you have] Someone to get together with for relaxation?’ and ‘[How often do you have] Someone to prepare your meals if you were unable to make them yourself?’ The response options ranged from 1 (None of the time) to 5 (All of the time).

Since this was an exploratory study, a simple retranslation of the above items was employed, taking Japanese culture into consideration. It was explained to participants that ‘family’ was defined as ‘relatives who were also members of the household’. ‘Environment’ was defined to include both people and physical surroundings—siblings, family members, colleagues, co-workers, neighbours, and the shared environment (Rotter, 1966; Manski, 1993; Christakis & Fowler, 2007, Cohen-Cole & Fletcher, 2008; Yakusheva *et al.*, 2011).

Further, to ensure that the participants faced no inconvenience in giving their responses, the questions and response options were modified to be as brief as possible. The researchers attempted to use as few items as possible in the scale by combining related items and eliminating redundancies. Furthermore, the original scale used response options ranging from 1 (None) to 8 (Does not apply); we reduced the number of

options to make the scale easier for participants to complete. The responses of all items were summed. We hypothesized that a higher score on the SRPO indicates a greater amount of social support and social interaction that promotes self-control.

The exploratory version of the SRPO originally contained 14 questions. Four questions concerned the number of athletic facilities within a convenient distance and the level of health support from the participant's employer. These items were prepared by referring to the Social Functioning Schedule (Remington & Tyrer, 1979), which is intended to assess the problems experienced in normal social functioning (such as work problems and problems in relationships with others at work, home, and elsewhere), and the SLOTH model (Pratt *et al.*, 2004), which is intended to enhance health through public announcements, health promotion programs, worksite interventions, and the like. However, since these items had poor response rates, they were omitted from the scale.

The final version of the SRPO contained 10 items that were carefully chosen to effectively capture the key aspects of the scales from which they were taken (for more details, see Table 2: the SRPO questions and responses).

Evaluation procedures, analysis, and results

Factor analysis

Confirmatory factor analysis was used to examine whether the data fit the model previously hypothesized by the researchers (French *et al.*, 2005). To confirm construct validity, factor loading was calculated. A value of .40 or greater is generally considered acceptable for this purpose. Confirmatory factor analysis revealed that the 10 items of the SRPO can be clustered into three factors. Factor 1 (covering the four questions from the Social Support and Exercise Survey and the Social Support and Eating Habits Survey)

was called ‘family and environmental support’. Factor 2 (covering the two questions from the Rand Social Health Battery and the two questions from the Medical Outcomes Study Social Support Survey) was called ‘social interaction’. Factor 3 (covering the two questions from the Social Adjustment Scale) was called ‘social adjustment’. The items, final factor loadings, explained variances, and eigenvalues are presented in Table 3.

High factor loading values indicate high consistency among the items of the scale. The Kaiser–Meyer–Olkin measure of sampling adequacy is used to test whether the partial correlations among variables are small. The results of this test can vary between 0 and 1, and values closer to 1 indicate that factor analysis is appropriate. The Kaiser–Meyer–Olkin value was .66 so factor analysis was undertaken. Bartlett’s test of sphericity indicated that the factor model was appropriate ($p < .01$) (Table 3).

Reliability

Reliability was assessed by examining the scores of the tele-care and self-help groups using Cronbach’s alpha, the generalized formula used to express the internal consistency of a test. Higher internal consistency can also mean higher test–retest reliability (McDowell & Newell, 1996), and a value of .70 or above is generally considered adequate. The Cronbach’s alpha values for the SRPO were .77 and .73 for the tele-care and self-help groups, respectively. In terms of internal consistency, Cronbach’s alpha coefficients were high for the tele-care and self-help groups (Table 4).

Validity

One of the ways of establishing a scale’s validity is to determine whether its scores are positively related to scores obtained on other scales that measure related or similar constructs. As the tele-care intervention utilized in the previous study was based on the Transtheoretical Model (which is effective in designing behaviour modification

interventions), various original scales and theories related to this model were utilized to assess the validity of the SRPO: the Stages of Change Theory, the level of Motivation for Exercise (Prochaska & DiClemente, 1983; Marcus, Rakowski & Rossi, 1992), Decisional Balance for Exercise Scale (Marcus & Simkin, 1994), and the Self-Efficacy for Exercise Scale (Prochaska *et al.*, 1992). In the previous study, the participants were administered these scales along with the main questionnaire at the health check-up (see Takada *et al.*, 2011, for details).

Construct validity: Marital status and improvements on the “Motivation for Exercise Scale” were used to confirm construct validity. The stages of change and the level of motivation for exercise theories are significant core factors of the Transtheoretical Model, which describes stages along a continuum of behavioural change and the participant’s motivation at each stage (1: not intending to exercise, 2: intending to exercise within 6 months, 3: intending to exercise next month, 4: sustaining exercise for 6 months or less, 5: sustaining exercise for over 6 months) (Marcus, Rakowski, & Rossi, 1992). These levels express the degree to which one is prepared for lifestyle modifications related to exercise. The level of motivation for exercise is strongly correlated with the level of social support (Prochaska, Norcross & Diclemente, 1994). Thus, these two variables—marital status and improvements in the level of motivation for exercise—could be used to represent social-relationship elements that prevent obesity by promoting exercise (Ruggiero & Prochaska, 1993; Glanz *et al.*, 1994; Vallis *et al.*, 2003). First, the model was constructed with already collected data, to estimate whatever outcomes possible considering the flexibility of these data. Then, to evaluate the model’s validity, split sample validation was performed. Although split sample validation is an accepted method, researchers should ideally collect new data to confirm model fit (Katz,

1999). In split sample validation, the full sample is split into two groups of approximately equal size and results of each group are compared. Therefore, in the current study, to confirm the validity of the scale and suitability of the analytical methods used, the sample was split into randomly selected groups and verified whether the previously obtained findings were robust. For further confirmation, analyses were conducted with randomly selected subgroups of 33 participants each.

A high degree of association was found between the SRPO and scores obtained on the Decisional Balance for Exercise Scale (Table 5). This instrument measures what people think the pros and cons of exercise are. The SRPO score was significantly and negatively correlated with the scores for the Cons subscale of this instrument, which measures perceived disadvantages of exercise, and factor 1 of the SRPO positively associated with the marital status. The SRPO score was also found to be positively and significantly correlated with the level of motivation for exercise.

The correlation between the scores on the Decisional Balance for Exercise Scale and the SRPO confirm the SRPO's validity. Although Factor 3 by itself had little correlation with the score for the Decisional Balance for Exercise Scale, a higher total score on the SRPO, including Factor 3, was found to correlate with a significantly higher score on the Decisional Balance for Exercise Scale (Table 5). Moreover, the level of motivation for exercise was significantly correlated with family and environmental support, social interaction, and adjustment (Factors 1, 2, and 3) and with the SRPO total score. Further analysis revealed, the tele-care and self-help groups also indicated to have similar correlations as with the full sample. Another randomly selected groups was indicated the reliability was similar to that reported above (Group 1 $\alpha = .81$, Group 2 $\alpha = .67$; results not shown in Table 5). Thus, split-sample validation suggests that the

scale is well calibrated (Table 5). Thus, the SRPO's construct validity is substantiated by the high correlations between the SRPO and the level of motivation for exercise and between the SRPO and the Decisional Balance for Exercise Scale (Yata *et al.*, 2003).

Concurrent criterion-related validity: To determine concurrent criterion-related validity, using two-tailed Pearson correlations, correlations between the SRPO and data obtained from the questions about eating habits were examined: (1) number of meals, breakfast daily, regular mealtimes, the hours between the last meal of the day and bedtime, (2) the number of snacks and alcoholic drinks consumed, and (3) instances of eating out. The responses to the questions pertaining to eating habits are shown in Table 6, and the correlations between eating habits and SRPO items are shown in Table 7. Items 3 and 4 of the SRPO, both of which concern eating habits, were significantly correlated with the number of times snacks and alcohol was consumed. Item 8 and 10 also were significantly correlated with consumption of snacks. Items 5 and 9 were significantly correlated with the number of meals. Item 5 was significantly correlated with instances of eating out. Item 10 was significantly correlated with consumption of breakfast and snacks. No items significantly correlated with regular mealtime and the hours between the last meal of the day and bedtime.

The association between eating habits and the SRPO is proved by the following correlations: (1) The presence of cordial relations with family members was inversely related to drinking and positively correlated to eating snacks. (2) The number of social activities and participation in social affairs were inversely related to the number of meals and instances of eating out and directly related to eating snacks. (3) The time spent on hobbies was directly related to the number of meals, consumption of snacks and inadequate breakfasts. The correlations between items for social adjustment or

communication in the SRPO and items on dietary habits suggest that the quality and quantity of a person's food intake may be affected by the person's relationships with his or her supporters. Moreover, the response to each question on dietary habits was significantly correlated with Factors 1, 2, and 3 of the SRPO (Table 7).

Discussion

Advantages of the SRPO

Many instruments have been developed in studies on weight control, but these contain a bewildering number of items on many different aspects, including the physical and psychosocial (Stunkard & Messick, 1985; McDowell & Newell, 1996). Because the SRPO contains fewer items with high factor-loading values, validity, and reliability, it is a more useful and convenient instrument for study participants and researchers than previous scales. The response rate in the Takada *et al.* (2011) study (55.9%) shows that the participants faced no inconvenience in providing responses, except for the excluded questions described above. Health information and knowledge helps people to choose a healthier lifestyle by improving their understanding of the relationships between health behaviour and health outcomes (Kenkel, 1991). The protective health effects of social relationships may be as important as the negative effects of established risk factor, such as smoking, obesity, and high blood pressure (House, Landis, & Umberson, 1988; Boothroyd & Fisher, 2010). Each item in the SRPO provides information about key health behaviours in social relationship. To convince the clinical practitioner in weight control of the SRPO's usefulness, it would be helpful to show that the SRPO directly correlates with the decisional balance for exercise in the Transtheoretical Model.

This study finds that the quantity and quality of social relationships are positively related to marital status. It seems that the SRPO score might be influenced, in implicit or explicit ways, by a spouse, family members, or other important people outside the family through a peer effect (Wallston *et al.*, 1978). In terms of LOC, it was assumed that a higher SRPO score would indicate greater environmental influence or the presence of a large number of high-quality social relationships. This effect could result in a dependency-related tendency, perhaps caused by the influence of strong relationships with others, such as a spouse, siblings, family members, colleagues, co-workers, and neighbours, and the effect of the shared environment (Wallston *et al.*, 1978; Macgregor *et al.*, 1997; Renna *et al.*, 2008; Fujita & Noguchi, 2009; Fortin & Yazbeck, 2011; Yakusheva *et al.*, 2011). Although Nir and Neumann (1995) reported no significant differences in weight loss between those with internal and external LOC during the post-intervention period of their study, the internal group gained less weight than the external group did. This evidence leads to the conclusion that internal LOC has a long-term effect and that modification of an external LOC is required to bring about meaningful change.

The importance of peer support has policy significance. Group-level interventions might be more cost-effective, successful, and open to variation than individual interventions (Christakis & Fowler, 2007; Fowler & Christakis, 2008; Renna *et al.*, 2008; Trogon *et al.*, 2008; Bahr *et al.*, 2009; Cobb *et al.*, 2011; Fortin & Yazbeck, 2011; Yakusheva *et al.*, 2011). Several studies have examined policy interventions targeted at altering the environment in such a way as to increase people's physical activity levels (Sallis & Owen, 1998; Ståhl *et al.*, 2001; Pratt *et al.*, 2004; Roux *et al.*, 2008; Li *et al.*, 2009; Cobb *et al.*, 2011; Montes *et al.*, 2012). Results from the current study suggest that individuals who have high SRPO scores at baseline are more influenced by their spouses,

family members and friends. When conducting a weight-loss intervention, special attention should be paid to the possibility that a self-control problem might be interfering with weight-loss for participants who express the characteristic of external LOC and who have spouses, close family members, or friends who are obese or have untreated obesity-related disease. Moreover, it is possible that participants' attitudes towards future obesity risks (i.e. obesity-related behaviour, sedentary lifestyle, impatience, indifferent attitude toward risk aversion, obesity-related family eating traditions, and food choices) can be measured with the SRPO in future weight control studies (Yakusheva *et al.*, 2010; Pachucki, Jacques & Christakis, 2011; Takada *et al.*, 2011). Effective weight-loss interventions that incorporate the acquisition of social support through a reliable social network should be used as an aid for self-control and a strong commitment to weight loss. Additionally, participants should be encouraged to develop a reliable social network that helps them maintain healthier habits.

Limitations

This study had some limitations. First, the items on the number of convenient athletic facilities and employer support were deleted because of their extremely poor response rates. Many participants could not respond to questions about employer direction and support because they were retired or owned their own businesses. However, the literature shows that these two factors play an important role in preventing obesity (Sallis *et al.*, 1992). Second, the participants in the Takada *et al.* (2011) study were recruited through a public advertisement, so there may have been a self-selection bias, although participants were subsequently randomized into the tele-care and self-help groups. If the SRPO is to be employed in a particular population, demographic characteristics, in particular, potential confounding factors such as age and job status,

should be taken into consideration. Third, as this scale was constructed by selecting items from other scales, the direct correlation between weight loss and scores on this scale may not strongly reflect the relationship between weight loss and self-control in social relationships, because we did not examine correlations between weight loss and other variables that have been previously shown to be related to weight loss. Fourth, it is necessary to determine the intrinsic differences between individuals who can and cannot develop and maintain strong self-control for healthy behaviours and devise a means to measure these differences.

Applications

The SRPO has moderate validity, reliability, and clinical utility in examining how social relationships support self-control with regard to weight loss or obesity prevention. Thus, it can be used as a screening tool in weight-loss interventions. The SRPO can also be used to examine the social environment and self-control problems in obese people, factors that should be considered when conducting a weight-loss intervention since obese people may have self-control problems that interfere with weight-loss plans (Kan, 2007).

References

- Ali, M. M., Amialchuk, A. & Heiland, F. W.** (2011) Weight-related behavior among adolescents: the role of peer effects. *PLoS One* **6**, e21179.
- Bahr, D. B., Browning, R. C., Wyatt, H. R. & Hill, J. O.** (2009) Exploiting social networks to mitigate the obesity epidemic. *Obesity* **17**, 723–728.
- Baumeister, R. F. & Heatherton, T. F.** (1996) Self-regulation failure: An overview. *Psychological Inquiry* **7**, 1-15.

- Black, D. R. & Lantz, C. E.** (1984) Spouse involvement and a possible long-term follow-up trap in weight loss. *Behaviour Research and Therapy* **22**, 557–562.
- Boothroyd, R. I. & Fisher, E. B.** (2010) Peers for progress: promoting peer support for health around the world. *Family Practice* **27** Suppl 1, i62–68.
- Brownell, K. D. & Stunkard, A. J.** (1981) Couples training, pharmacotherapy, and behavior therapy in the treatment of obesity. *Archives of General Psychiatry* **38**, 1224–1229.
- Brummett, B. H., Barefoot, J. C., Siegler, I. C., Clapp-Channing, N. E., Lytle, B. L., Bosworth, H. B., Williams, R. B., Jr. & Mark, D. B.** (2001) Characteristics of socially isolated patients with coronary artery disease who are at elevated risk for mortality. *Psychosomatic Medicine* **63**, 267–272.
- Carver, C. S.** (2005) Impulse and constraint: perspectives from personality psychology, convergence with theory in other areas, and potential for integration. *Personality and Social Psychology Review*, **9**, 312–333.
- Christakis, N. A. & Fowler, J. H.** (2007) The spread of obesity in a large social network over 32 years. *The New England Journal of Medicine* **357**, 370–379.
- Cobb, N. K., Graham, A. L., Byron, M. J., Niaura, R. S. & Abrams, D. B.** (2011) Online social networks and smoking cessation: a scientific research agenda. *Journal of Medical Internet Research* **13**, e119.
- Cohen-Cole, E. & Fletcher, J. M.** (2008) Is obesity contagious? Social networks vs. environmental factors in the obesity epidemic. *Journal of Health Economics* **27**, 1382–1387.
- Contrada, R. J. & Goyal, T. M.** (2005) Individual differences, health and illness: the role of emotional traits and generalized expectancies. In Sutton, S., Baum, A. &

- Johnston, M. (eds) *The SAGE Handbook of Health Psychology*. SAGE Publications, London, pp. 143–168.
- Cutler, D. M. & Glaeser, E. L.** (2005) What explains differences in smoking, drinking and other health-related behaviors? *American Economic Review* **95**, 238–242.
- Dale, J. R., Williams, S. M. & Bowyer, V.** (2012) What is the effect of peer support on diabetes outcomes in adults? A systematic review. *Diabetic Medicine* **29**, 1361–1377.
- Deforche, B. & De Bourdeaudhuij, I.** (2000) Differences in psychosocial determinants of physical activity in older adults participating in organised versus non-organised activities. *Journal of Sports Medicine and Physical Fitness* **40**, 362–372.
- Donald, C. A. & Ware, J. E. Jr.** (1984) The measurement of social support. *Research in Community and Mental Health* **4**, 325–370.
- Eid, J., Overman, H. G., Puga, D. & Turner, M. A.** (2008) Fat city: questioning the relationship between urban sprawl and obesity. *Journal of Urban Economics* **63**, 385–404.
- Finnerty, T., Reeves, S., Dabinett, J., Jeanes, Y. M. & Vögele, C.** (2010) Effects of peer influence on dietary intake and physical activity in schoolchildren. *Public Health Nutrition* **13**, 376–383.
- Fortin, B. & Yazbeck, M.** (2011) Peer effects, fast food consumption and adolescent weight gain. *CIRANO Scientific Series* 2011s-20.
- Fowler, J. H. & Christakis, N. A.** (2008) Estimating peer effects on health in social networks: a response to Cohen-Cole and Fletcher; and Trogon, Nonnemaker, and Pais. *Journal of Health Economics* **27**, 1400–1405. DOI: 10.1016/j.jhealeco.2008.07.001.

- Frank, R. H.** (1988) *Passion Within Reason: The Strategic Role of the Emotions*. W. W. Norton, New York.
- French, D. P., Yardley, L. & Sutton, S.** (2005) Research methods in health psychology. In Sutton, S., Baum, A. & Johnston, M. (eds) *The SAGE Handbook of Health Psychology*. SAGE Publications, London, pp. 326–359.
- Fujita, T. & Noguchi, A.** (2009) A study of the relationship between self-control and academic procrastination behavior in college students (in Japanese). *Bulletin of the Center for Educational Research and Development* **18**, 101–106.
- Funch, D. P., Marshall, J. R. & Gebhardt, G. P.** (1986) Assessment of a short scale to measure social support. *Social Science & Medicine* **23**, 337–344.
- Glanz, K., Patterson, R. E., Kristal, A. R., DiClemente, C. C., Heimendinger, J., Linnan, L. & McLerran, D. F.** (1994) Stages of change in adopting healthy diets: fat, fiber, and correlates of nutrient intake [Erratum in *Health Education Quarterly* **22**, 261, 1995]. *Health Education Quarterly* **21**, 499–519.
- Gorin, A. A., Wing, R. R., Fava, J. L., Jakicic, J. M., Jeffrey, R., West, D. S., Breije, K. DiLillo, V. G. Look AHEAD Home Environment Research Group** (2008) Weight loss treatment influences untreated spouses and the home environment: evidence of a ripple effect. *International Journal of Obesity* **32**, 1678–1684.
- Gruber, K. J.** (2008) Social support for exercise and dietary habits among college students. *Adolescence* **43**, 557–575.
- Hill, J. O., Peters, J. C., Catenacci, V. A. & Wyatt, H. R.** (2008) International strategies to address obesity. *Obesity Reviews* **9**(Suppl 1), 41–47.
- House, J. S.** (2001) Social isolation kills, but how and why? *Psychosomatic Medicine* **63**, 273–274.

- House, J. S., Landis, K. R. & Umberson, D.** (1988) Social relationships and health. *Science* **241**, 540–545.
- Kan, K.** (2007) Cigarette smoking and self-control. *Journal of Health Economics*, **26**, 61–81.
- Karlsson, J., Sjostrom, L. & Sullivan, M.** (1995) Swedish obese subjects (SOS)—an intervention study of obesity. Measuring psychosocial factors and health by means of short-form questionnaires. Results from a method study. *Journal of Clinical Epidemiology* **48**, 817–823.
- Katz, M. H.** (2006) *Multivariable analysis -A practical guide for clinicians, 2nd edition*. Cambridge University Press, Cambridge.
- Kenkel, D. S.** (1991) Health behavior, health knowledge, and schooling. *Journal of Political Economy* **2**, 313–325.
- Komlos, J., Smith, P. K. & Bogin, B.** (2004) Obesity and the rate of time preference: is there a connection? *Journal of Biosocial Science* **36**, 209–219.
- Li, F., Harmer, P., Cardinal, B. J., Bosworth, M., Johnson-Shelton, D., Moore, J. M., Acock, A. & Vongjaturapat, N.** (2009) Built environment and 1-year change in weight and waist circumference in middle-aged and older adults: Portland Neighborhood Environment and Health Study. *American Journal of Epidemiology* **169**, 401–408.
- Manski, C. F.** (1993) Identification of endogenous social effects: the reflection problem. *Review of Economic Studies* **60**, 531–542.
- Marcus, B. H. & Simkin, L. R.** (1994) The transtheoretical model: applications to exercise behavior. *Medical and Science in Sports and Exercise* **26**, 1400–1404.

- Marcus, B. H., Rakowski, W. & Rossi, J. S.** (1992) Assessing motivational readiness and decision making for exercise. *Health Psychology* **11**, 257–261.
- McDowell, I. & Newell, C.** (1996) *Measuring Health: A Guide to Rating Scales and Questionnaires, 2nd Edition*. Oxford University Press, London.
- McLean, N., Griffin, S., Toney, K. & Hardeman, W.** (2003) Family involvement in weight control, weight maintenance and weight-loss interventions: a systematic review of randomised trials. *International Journal of Obesity and Related Metabolic Disorders* **27**(9), 987–1005.
- Macgregor, I. D. M., Regis, D. & Balding, J.** (1997) Self-concept and dental health behaviors in adolescents. *Journal of Clinical Periodontology* **24**, 335–339.
- Montes, F., Sarmiento, O. L., Zarama, R., Pratt, M., Wang, G., Jacoby, E., Schmid, T. L., Ramos, M., Ruiz, O., Vargas, O., Michel, G., Zieff, S. G., Valdivia, J. A., Cavill, N. & Kahlmeier, S.** (2012) Do health benefits outweigh the costs of mass recreational programs? An economic analysis of four ciclovia programs. *Journal of Urban Health* **89**, 153–170.
- Nir, Z. & Neumann, L.** (1995) Relationship among self-esteem, internal-external locus of control, and weight change after participation in a weight reduction program. *Journal of Clinical Psychology* **51**, 482–490.
- Pachucki, M. A., Jacques, P. F. & Christakis, N. A.** (2011) Social network concordance in food choice among spouses, friends, and siblings. *American Journal of Public Health* **101**, 2170–2177.
- Pratt, M., Macera, C. A., Sallis, J. F., O'donnell, M. & Frank, L. D.** (2004) Economic intervention to promote physical activity application of the SLOTH model. *American Journal of Preventive Medicine* **27**, 136–145.

- Prochaska, J. O. & Diclemente, C. C.** (1983) Stages and processes of self-change of smoking: toward an integrative model of change. *Journal of Consulting and Clinical Psychology* **51**, 390–395.
- Prochaska, J. O., DiClemente, C. C. & Norcross, J. C.** (1992) In search of how people change. Applications to addictive behaviors. *American Psychologist* **47**, 1102–1114.
- Prochaska, J. O., Norcross, J. C. & Diclemente, C. C.** (1994) *Changing for good: The Revolutionary Program That Explains the Six Stages of Change and Teaches You How to Free Yourself from Bad Habits*. Harper Collins Publishers, New York.
- Remington, M., Tyrer, P.** (1979) The social functioning schedule—a brief semi-structured interview. *Social Psychiatry* **14**, 151–7.
- Renna, F., Grafova, I. B. & Thakur, N.** (2008) The effect of friends on adolescent body weight. *Economics and Human Biology* **6**, 377–387. DOI: 10.1016/j.ehb.2008.06.005.
- Rotter, J. B.** (1954) *Social Learning and Clinical Psychology*. Prentice-Hall, Englewood Cliffs.
- Rotter, J. B.** (1966) Generalized expectancies for internal versus external control of reinforcement. *Psychological Monographs* **80**, 1–28.
- Roux, L., Pratt, M., Yore, M. M., Yanagawa, T. L., Van Den Bos, J., Brownson, R. C., Powell, K. E., Hea, H. W. R., Teusch, S., Cawley, J., Lee, I. M., West, L. & Buchner, D. M.** (2008) Cost effectiveness of community-based physical activity interventions. *American Journal of Preventive Medicine* **35**, 578–588.
- Ruggiero, L. & Prochaska, J. O.** (1993) Readiness for change: application of the transtheoretical model to diabetes. *Diabetes Spectrum* **6**, 22–60.

Sallis, J. F., Grosman, R. M., Pinski, R. B., Patterson, T. L. & Nader, P. R. (1987)

The development of scales to measure social support for diet and exercise behaviors.

Preventive Medicine **16**, 825–836.

Sallis, J. F., Hovell, M. F. & Hofstetter, C. R. (1992) Predictors of adoption and

maintenance of vigorous physical activity in men and women. *Preventive Medicine*

21, 237–251.

Sallis, J. F. & Owen, N. (1998) *Physical activity and behavioral medicine*. SAGE

Publications, London.

Sherbourne, C. D. & Stewart, A. L. (1991) The MOS Social Support Survey. *Social*

Science & Medicine **32**, 705–714.

Sherrill-Mittleman, D. A., Klesges, L. M., Lanctot, J. Q., Stockton, M. B. & Klesges,

R. C. (2009) Measurement characteristics of dietary psychosocial scales in a

weight-gain prevention study with 8- to-10-year-old African-American girls.

Health Education Research **24**, 586–595.

Ståhl T., Rütten, A., Nutbeam, D., Bauman, A., Kannas, L., Abel, T., Lüschen, G.,

Rodriguez Diaz J. A., Vinck, J. & Van der Zee, J. (2001) The importance of the

social environment for physically active lifestyle—results from an international

study. *Social Science & Medicine* **52**, 1–10.

Steptoe, A. & Ayers, S. (2005) Stress, health and illness. In Sutton, S., Baum, A. &

Johnston, M. (eds) *The SAGE Handbook of Health Psychology*. SAGE Publications,

London, pp. 169–196.

Stunkard, A. J. & Messick, S. (1985) The three-factor eating questionnaire to measure

dietary restraint, disinhibition and hunger. *Journal of Psychosomatic Research*, **29**,

71–83.

- Takada, A., Nakamura, R., Furukawa, M., Takahashi, Y., Nishimura, S. & Kosugi, S.** (2011) The relationship between weight loss and time and risk preference parameters: a randomized controlled trial [Erratum in: *Journal of Biosocial Science* **43**, 505–508, 2011]. *Journal of Biosocial Science* **43**, 481–503
- Takada, A., Takahashi, Y. & Kosugi, S.** (2010). The development of scale for social support to counter obesity (SSCO). *The Joint Scientific Meeting of the International Epidemiological Association Western Pacific Region and the Japan Epidemiological Association, Program and Abstracts*, S123.
- Tangney, J. P., Baumeister, R. F. & Boone, A. L.** (2004) High self-control predicts good adjustment, less pathology, better grades, and interpersonal success. *Journal of Personality* **72**, 271–324.
- Trogon, J. G., Nonnemaker, J. & Pais, J.** (2008) Peer effects in adolescent overweight. *Journal of Health Economics* **27**, 1388–1399. DOI: 10.1016/j.jhealeco.2008.05.003.
- Uechi, H.** (2006) The perspectives of behavioral modification theories, models, and techniques (in Japanese). *L'esprit d'aujourd'hui* **2**, 95–108.
- Unger, J. B. & Johnson, C. A.** (1995) Social relationships and physical activity in health club members. *American Journal of Health Promotion* **9**, 340–343.
- Vallis, M., Ruggiero, L., Greene, G., Jones, H., Zinman, B., Rossi, S., Edwards, L., Rossi, J. S. & Prochaska, J. O.** (2003) Stages of change for healthy eating in diabetes: relation to demographic, eating-related, health care utilization, and psychosocial factors. *Diabetes Care* **26**, 1468–1474.
- Wallston, K. A., Wallston, B. S. & DeVellis, R.** (1978) Development of the Multidimensional Health Locus of Control (MHLC) Scales. *Health Education Monographs* **6**, 160–170.

- Weissman, M. M. & Bothwell, S.** (1976) Assessment of social adjustment by patient self-report. *Archives of General Psychiatry* **33**, 1111–1115
- Wills, T. A., Isasi, C. R., Mendoza, D. & Ainett, M. G.** (2007) Self-control constructs related to measures of dietary intake and physical activity in adolescents. *Journal of Adolescent Health* **31**, 551–558.
- Yakusheva, O., Kapinos, K. A. & Weiss, M.** (2011) Peer effects and the freshman 15: evidence from a natural experiment. *Economics and Human Biology* **9**, 119–132.
- Yata, K., Yokoda, K. & Takama, S.** (2003) Development of an instrument to measure family support of patients with diabetes (in Japanese). *The Journal of the Nursing Society of the Toyama Medical and Pharmaceutical University* **5**, 97–104.

Table 1. Summary statistics

Variables	n	%
Gender	(n = 66)	
Male	23	(34.8)
Female	43	(65.2)
Age	(n = 66)	
< 40	24	(36.4)
> = 40	42	(63.6)
BMI	(n = 66)	
< 25	28	(42.4)
> = 25	38	(57.6)
Education	(n = 65)	
Junior High School	1	(1.5)
High School	19	(29.2)
Vocational School	7	(10.8)
Junior Collage	11	(16.9)
University(literature)	20	(30.8)
University(science)	3	(4.6)
Graduate School	4	(6.2)
Job status	(n = 65)	
None	15	(23.1)
Current(with wages)	50	(76.3)
Marital Status	(n = 65)	
Single, divorces, widowed	19	(29.2)
Married	46	(70.8)
Income	(n = 59)	
None	10	(16.9)
< 100 million	10	(16.9)
101- 200 million	14	(23.8)
201- 400 million	10	(16.9)
> 400 million	15	(25.5)
Property	(n =53)	
None	10	(18.9)
< 500 million	14	(26.4)
501-1000 million	7	(13.2)
1001-1500 million	4	(7.5)
> 1500 million	18	(34.0)

Table 2. Results of factor analysis of the SRPO

Items	Factor loading
Factor I. Family and environmental support	
1. My family or friends exercised with me and gave me helpful reminders to exercise	.76
2. My family or friends helped plan activities around me to ensure more time for exercise	.81
3. My family or friends reminded me not to eat high-salt, high-fat foods	.87
4. My family or friends discussed my eating habit changes with me	.82
Factor II. Social interaction	
5. To how many volunteer groups or organizations do you belong (e.g. church, temple, shrine groups, clubs in the community, or parent groups)?	.91
6. How active are you in the affairs of the groups or clubs to which you belong?	.89
7. [How often do you have] Someone to get together with for relaxation?	.43
8. [How often do you have] Someone to prepare your meals if you were unable to make them yourself?	.38
Factor III. Social adjustment	
9. How many times in the last two weeks have you gone out socially (visited friends, gone to movies, churches, restaurants, etc.)?	.83
10. How much time have you spent on hobbies or items of interest in the last two weeks?	.75
	*Eigenvalue
Factor I	3.4
Factor II	2.2
Factor III	1.2
	% variance explained
Factor I	32.6
Factor II	20.1
Factor III	13.4
	Cronbach's alpha
Factor I	.85
Factor II	.70
Factor III	.51

Note. SRPO: Social Relationships to Prevent Obesity Scale

A higher score on the SRPO indicates a higher degree of social support

Kaiser-Meyer-Olkin measure of sampling adequacy = .66

Bartlett's test of sphericity = 264.03, $p < .01$

*Eigenvalues are the variances of the factors; a value over 1 indicates that factor analysis can be performed

Table 3. Means, standard deviations, and coefficients of the SRPO scores across groups

	Total (N = 66)			Tele-care (n = 36) group			Self-help (n = 30) group		
	Mean	SD	Cronbach's alpha	Mean	SD	Cronbach's alpha	Mean	SD	Cronbach's alpha
Baseline	27.9	6.5	.75	27.6	6.7	.77	28.3	6.2	.73

Table 4. Correlations between the factors of the SRPO and other indices

	SRPO	Factor I	Factor II	Factor III
	Total	Family and environment	Social support [†] interaction	Social adjustment
Decisional Balance for Exercise Scale (n = 65)	.38**	.32*	.27*	.18
Pros (n = 65) [‡]	.12	.06	.12	.02
Cons (n = 65)	-.48**	-.42**	-.34**	-.22
Motivation for exercise (n = 63)	.47**	.28*	.35**	.45**
Marital status (n = 65)	.18	.34*	.02	-.24

Tele-care group (n=36)

	SRPO	Factor I	Factor II	Factor III
	Total	Family and environment	Social support [†] interaction	Social adjustment
Decisional Balance for Exercise Scale (n = 35)	.27	.26	.20	.05
Pros (n = 35) [‡]	.14	.09	.13	-.07
Cons (n = 35)	-.31	-.34*	-.22	-.15
Motivation for exercise (n = 33)	.49**	.31	.41*	.46**
Marital status (n = 36)	.18	.28	.03	-.15

Self-help group (n=30)

	SRPO	Factor I	Factor II	Factor III
	Total	Family and environment	Social support [†] interaction	Social adjustment
Decisional Balance for Exercise Scale (n = 30)	.47**	.38*	.31	.33
Pros (n = 30) [‡]	.09	.00	.08	.13
Cons (n = 30)	-.62*	-.55**	-.42*	-.34
Motivation for exercise (n = 29)	.40*	.13	.33	.38*
Marital status (n = 28)	.16	.41**	-.02	-.33

Note. SRPO: Social Relationships to Prevent Obesity Scale

*: $p < .05$; **: $p < .01$

[†] : Support from family and surroundings in monitoring calorie intake and exercising

[‡] : Pros/Cons: Perceived advantages/disadvantages of exercise

Values are scores at baseline

Table 5. Descriptive statistics: Dietary habits of the participants

Variables	n	%
How many meals do you have in a day?	(n = 63)	
Two	4	(6.3)
Three	58	(92.1)
Four	1	(1.6)
Do you have breakfast daily?	(n = 64)	
Yes	60	(93.8)
No	4	(6.2)
Do you have regular mealtimes?	(n = 63)	
Yes	51	(81.0)
No	12	(19.0)
How many hours do you usually keep between your last meal of the day and bedtime?	(n = 63)	
1 hour	4	(6.3)
1–2 hours	7	(11.1)
Over 2 hours	52	(82.6)
How many times a day do you have a snack between meals?	(n = 60)	
Never	11	(18.3)
Every few days	1	(1.7)
Once	30	(50.0)
Twice	14	(23.3)
Three times	4	(6.7)
How many alcoholic drinks do you have in a week?	(n = 64)	
None	31	(48.4)
One drink a week a few drinks a month	19	(29.7)
One almost every day	14	(21.9)
How many times in a month do you eat out at a restaurant or such?	(n = 61)	
Never	31	(50.8)
Less than 4 times	14	(23.0)
8 times or less	2	(3.3)
9 times or more	14	(23.0)

Table 6. Significant correlations between eating habits and SRPO items

Factors and items	Number of meals	Breakfast	Snacks	Drinks	Eating out	Meal time	Hours
Factor I. Family and environmental support							
1. My family or friends exercised with me and gave me Helpful reminders to exercise	-.05	-.06	.11	-.17	.07	.09	.09
2. My family or friends helped plan activities around my exercise	.02	-.08	.21	-.24	.07	.02	.09
3. My family or friends reminded me not to eat high fat/salt foods	-.17	.03	.30*	.25*	.06	-.08	.07
4. My family or friends discussed my eating habit changes with me	-.19	.00	.37**	.33**	-.03	-.12	.22
Factor II. Social interaction							
5. How many volunteer groups or organizations do you belong to (e.g. church, temple, shrine groups, clubs in the community, or parent groups)?	-.25*	.12	-.05	.06	-.27*	-.04	-.03
6. How active are you in the affairs of the groups or clubs to which you belongs?	-.24	.10	-.08	-.02	-.23	-.04	-.11
7. [How often do you have] Someone to get together with for relaxation?	-.24	.09	.08	-.12	-.25	-.00	-.06
8. [How often do you have] Someone to prepare your meals if you were unable to make them yourself?	-.16	-.11	.36**	-.13	-.01	-.09	-.03
Factor III. Social adjustment							
9. How many times in the last two weeks have you gone out socially (visited friends, gone to movies, churches, restaurants, etc.)?	.26*	-.19	-.04	-.02	-.07	.19	-.12
10. How much time have you spent on hobbies or items of interests during the last two weeks?	.19	-.27*	.26*	-.10	-.05	.11	-.17

Note. SRPO: Social Relationships to Prevent Obesity Scale

Meal time: Regular mealtimes Hours: the hours between the last meal of the day and bedtime

*: $p < .05$

** : $p < .01$

Appendix. Items and response options for the Social Relationships to Prevent Obesity Scale (SRPO; N = 66)

Items	Response options	n	%
Factor I. Family and environmental support			
1. My family or friends exercised with me and gave me helpful reminders to exercise			
	a. Strongly disagree/Not at all	17	25.8
	b. Disagree	12	18.2
	c. Undecided	14	21.2
	d. Agree	19	28.8
	e. Strongly agree/Very often	4	6.0
2. My family or friends helped plan activities around my exercise			
	a. Strongly disagree/Not at all	17	25.8
	b. Disagree	23	34.8
	c. Undecided	18	27.3
	d. Agree	6	9.1
	e. Strongly agree/Very often	2	3.0
3. My family or friends reminded me not to eat high-fat, high-salt foods			
	a. Strongly disagree/Not at all	11	16.6
	b. Disagree	14	21.2
	c. Undecided	17	25.8
	d. Agree	17	25.8
	e. Strongly agree/Very often	7	10.6
4. My family or friends discussed my eating habit changes with me			
	a. Strongly disagree/Not at all	16	24.2
	b. Disagree	13	19.7
	c. Undecided	18	27.3
	d. Agree	13	19.7
	e. Strongly agree/Very often	6	9.1
Factor II. Social interaction			
5. To how many volunteer groups or organizations do you belong to, like church, temple, shrine groups, clubs in the community, or parent groups, etc.?			
	a. None	31	47.0
	b. One	16	24.2
	c. Two	12	18.2
	d. Three	4	6.0
	e. More than three groups or organizations	3	4.6

continued

6. How active are you in the affairs of the groups or clubs to which you belong?			
a.	Do not belong to any groups or attend any meetings	32	48.5
b.	Not active, belong but hardly ever go	0	0.0
c.	Fairly active, attend fairly often	15	22.7
d.	Very active, attend most meetings	19	28.8
7. [How often do you have] Someone to get together with for relaxation?			
a.	None of the time	3	4.6
b.	A little of the time	19	28.8
c.	Some of the time	28	42.4
d.	Most of the time	14	21.2
e.	All of the time	2	3.0
8. [How often do you have] Someone to prepare your meals if you were unable to make them yourself?			
a.	None of the time	13	19.7
b.	A little of the time	25	37.9
c.	Some of the time	11	16.7
d.	Most of the time	15	22.7
e.	All of the time	2	3.0
Factor III. Social adjustment			
9. How many times in the last two weeks have you gone out socially (visited friends, gone to movies, churches, restaurants, etc.)?			
a.	None	1	1.5
b.	Once	6	9.1
c.	Twice	8	12.1
d.	Three times	9	13.7
e.	More than three times	42	63.6
10. How much time have you spent on hobbies or items of interests during the last two weeks?			
a.	I did not spend any time on hobbies or watching TV	2	3.0
b.	I usually did not spend any time on hobbies but did watch TV	12	18.2
c.	I spent a little time on hobbies	26	39.4
d.	I spent some time on hobbies on most days	17	25.8
e.	I spent a lot of time on hobbies almost every day	9	13.6