TITLE:

Protistan grazing and viral lysis losses of bacterial carbon production in a large mesotrophic lake (Lake Biwa)

AUTHOR(S):

Takasu, Hiroyuki; Kunihiro, Tadao; Nakano, Shin-ichi

CITATION:

Takasu, Hiroyuki ...[et al]. Protistan grazing and viral lysis losses of bacterial carbon production in a large mesotrophic lake (Lake Biwa). *Limnology* 2014, 15(3): 257-270

ISSUE DATE:

2014-08

URL:

<http://hdl.handle.net/2433/190411>

RIGHT:

The final publication is available at Springer via <http://dx.doi.org/10.1007/s10201-014-0431-6>; 許諾条件により本文は2015-04-19に公開.; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

Protistan grazing and viral lysis losses of bacterial carbon production in a large mesotrophic lake (Lake Biwa)

Hiroyuki Takasu^{1,2}, Tadao Kunihiro^{3,4} and Shin-ichi Nakano^{1*}

¹ *Center for Ecological Research (CER), Kyoto University, Hirano-cho 2 chome, 509-3, Otsu, Shiga 520-2113, Japan*

² *Present address: Atmosphere and Ocean Research Institute (AORI), The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, Chiba 277-8564, Japan*

³ *Center for Marine Environmental Studies (CMES), Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Ehime, Japan*

⁴ *Present address: Department of Ecosystems Studies / Department of Marine Microbiology, Royal Netherlands Institute for Sea Research (NIOZ), Korrिंगaweg 7, 4400 AC Yerseke, The Netherlands*

*Correspondence to: Shin-ichi Nakano: Mailing address: Center for Ecological Research (CER), Kyoto University, Hirano-cho 2 chome, 509-3, Otsu, Shiga 520-2113, Japan.

Phone and Fax: +81-77-549-8239, +81-77-549-8201

E-mail: nakano@ecology.kyoto-u.ac.jp

Running title: Losses of bacterial carbon production in Lake Biwa

Key words: Bacterial growth, Grazing mortality, Viral-mediated mortality

24 **Abstract**

25 The grazing and lysis mortality of planktonic bacteria were estimated using the modified
26 dilution method and respiratory quinone (RQ) analysis in mesotrophic Lake Biwa, Japan. The
27 planktonic bacterial assemblages in the lake consisted of various RQs subgroups with
28 different growth and mortality rates. The sum of total bacterial mortalities due to protistan
29 grazing and viral lysis accounted for 96.6% (range; 89.0 - 107.2%) of daily total bacterial
30 production. This is the first report which successfully demonstrates a balanced relationship
31 between bacterial production and losses using the modified dilution method in a lake. The
32 growth rates of ubiquinone (UQ)-containing bacteria were faster than those of
33 menaquinone-containing bacteria. Especially, the dominant and fastest-growing bacterial
34 groups in the present study were the bacterial groups containing UQ-8 or UQ-10. The sum of
35 their production and loss accounted for 60% of carbon fluxes within the microbial loop. Thus,
36 a large portion of the carbon cycling through the bacterial community in Lake Biwa can be
37 explained by the carbon fluxes through dominant bacterial groups.

38

39

Introduction

In the pelagic environments of freshwater and marine systems, a significant fraction of primary production is consumed by heterotrophic bacteria via dissolved organic matter (DOM) released from phytoplankton (Cole et al. 1988; Ducklow 2000). Protistan grazing and viral lysis are two important determinants of the fate of bacterial biomass (Azam et al. 1983; Proctor and Fuhrman 1991), with different ecological and biogeochemical implications. Protistan grazing transfers bacterial biomass to higher trophic-level organisms via the microbial loop (Azam et al. 1983), whereas viral lysis leads to the recycling of carbon and nutrients, both of which are derived from lysed bacterial biomass and are re-supplied to bacteria (Bratbak et al. 1990; Gobler et al. 1997; Proctor and Fuhrman 1991).

Natural bacterial assemblages consist of various subgroups in terms of ecological and biogeochemical features. The relationship between dynamics of bacterial community structure and variations in carbon cycling within the microbial loop are not well understood in natural aquatic systems, though the results of some ecological models suggest that changes in bacterial community structure can affect the carbon fluxes through bacterial communities (Miki et al. 2008). Only limited attempts have so far been conducted to make simultaneous estimation of grazer-induced and virus-induced bacterial mortality for distinct bacterial groups (Šimek et al. 2001; 2007). Obviously, different mortalities among distinct bacterial groups can influence bacterial community composition. The ‘size-selective mortality’ for flagellates changes the size distribution of bacterial community (Pernthaler 2005), whereas the ‘host-specificity mortality’ for viruses changes the bacterial community structure (Thingstad 2000). It is, therefore, important to simultaneously estimate grazing-induced mortality and virus-induced mortality for better understanding on carbon fluxes through bacterial community.

Specific growth and mortality rates of microbial populations can be simultaneously estimated from observed differences in their rates of population growth in a series of

incubated diluted and undiluted natural water samples (Landry and Hassett 1982; Landry et al. 1984). The dilution technique, originally developed for the estimation of grazing pressure by micro-zooplankton on phytoplankton (Landry and Hassett 1982), is valid to estimate protistan bacterivory (Landry et al. 1984; Trremaïne and Mills 1987). Evans et al. (2003) proposed a modified dilution technique to estimate the impacts of protistan grazing and viral lysis on the picoeukaryote *Micromonas* spp. Their technique was successfully applied to estimate the impact of viral lysis on planktonic bacteria in freshwater lake (Tijdens et al. 2008) and ocean (Taira et al. 2009).

Fluorescence *in situ* hybridization (FISH) technique as one of the most powerful quantitative molecular approaches has been used for quantifying and visualizing bacterial cells in freshwater and seawater (Pernthaler et al. 1998; Glöckner et al. 1999). The FISH technique is suitable for targeting at specific phylogenetic group levels but less suitable for analysis of the full bacterial community, because quantitative application for analysis of all bacterial groups requires the use of many target-specific probes and also need to optimize its protocol for each target groups (Bouvier and del Giorgio 2003). Despite the superiority of FISH technique-based approaches in terms of phylogenetic identification, respiratory quinone (RQ) analysis has been successfully used to quantify bacterial biomass and to overview bacterial community composition in freshwater (Takasu et al. 2013). Respiratory quinone (RQ), including ubiquinone (UQ) and menaquinone (MK), are electron-transporting compounds in bacterial plasma membranes. Different types of RQ differ in their preference of electron accepters for energy metabolism (Hedrick and White 1986). A bacterial phylum has generally only one dominant molecular species of respiratory quinone (Collins and Jones 1981; Hedrick and White 1986). The RQ analysis provides a less laborious and accurate method for simultaneously determining bacterial carbon biomass and community composition because of the chemical analytical-based method with a standardized quantitative extraction protocol (Hu et al. 1999; Takasu et al., 2013). The modified dilution technique combined with

RQ analysis provides us quantitative information about carbon fluxes through distinct bacterial groups in complex microbial food web.

Lake Biwa is a large (surface area, 674 km²; water volume, 27.3 km³; watershed area, 3848 km²), deep (maximum depth, 104 m), mesotrophic and monomictic lake in Japan. Large lakes in the world are generally important freshwater resources (Herdendorf 1990). In addition, large lakes are a precious food production site (Constanza et al. 1997). Net bacterial production in Lake Biwa was estimated as 30% of primary production (Nagata et al. 1990), and their total consumption (net bacterial production plus respiration) was accounted for 50 to 100% of the primary production (Nagata et al. 2012). It has been regarded that carbon fluxes through bacterial community are highly active, and that bacteria are a potentially important basis of the pelagic food web in Lake Biwa (Nagata 1990). Thus, elucidating the carbon fluxes through bacterial subgroups is important for our better understanding about carbon cycling in ecosystem of Lake Biwa.

In the present study, we quantified the carbon fluxes through the bacterial community by estimating carbon production and losses calculated from growth and mortality rates of planktonic bacteria in Lake Biwa. We hypothesized that each bacterial group would have different ecological roles within microbial loop. To test this hypothesis, we determined the grazing and lysis mortalities of each bacterial group using the modified dilution technique and RQ analysis. Our data demonstrated that a large portion of the carbon cycling through the bacterial community in Lake Biwa can be explained by the carbon fluxes through dominant bacterial groups.

Materials and Methods

Dilution experiments

The dilution experiments were conducted in June and October 2011 and May, June, and July 2012, using lake water collected from St. Ie-1 (35°12'58"N, 135°59'55"E; maximum water

depth 73 m) in the pelagic area of the north basin of Lake Biwa, Japan. Approximately 100 L of lake water was collected from 5 m depth using a 10-L acrylic water sampler. Enumeration of microbes was performed by placing 200 mL of the water sample in a polypropylene bottle and fixing immediately with glutaraldehyde at a final concentration of 1%. Water samples for the dilution experiments were poured into acid-washed 10-L polyethylene bags or 20-L polyethylene tanks. The lake water was gently filtered through 20 μm mesh to remove mesozooplankton, and 50 L of the filtrate was gravity filtered through 0.2 μm filter cartridges (PALL Acropak Supor membrane capsules) and collected into tanks. After the filtration, half of the 0.2 μm filtrate was passed through a 30 kDa tangential flow filtration system (Millipore PES membrane) to prepare a virus-free diluent. The 20 μm filtrate was diluted in 0.2 μm or 30 kDa diluents to dilution levels of 1.0, 0.8, 0.4, and 0.2, in 5-L polycarbonate bottles washed with 1.2M HCl before use. The bottles were then incubated for 36 to 48 hours at *in situ* temperatures, in dark conditions. Subsamples for the measurement of RQs were collected into clean polyethylene bags at the beginning (0 hours) and end of the incubations.

Enumeration of microbes

For the enumeration of bacteria, 1 or 2 mL of the fixed water sample was used. Bacterial cells were counted using an epifluorescence microscope (BX60, Olympus) under ultraviolet excitation by the DAPI (4,6-diamidino-2-phenylindole) method (Porter and Feig 1980) using 0.2 μm pore-size black polycarbonate filters (Advantec). At least 300 bacterial cells were counted, and a minimum of 20 fields were randomly selected. Fifteen milliliters of the fixed water sample were used for the enumeration of nanoflagellates, and 0.1 mL (1 mL from the 10x diluted samples with 0.02 μm filtered distilled water) was used for the enumeration of viral-like particles (VLP). Heterotrophic nanoflagellates (HNF) and pigmented nanoflagellates (PNF) were counted using epifluorescence microscopy under ultraviolet and green excitation respectively, using the primulin method (Caron 1983), using 0.8 μm pore-size black polycarbonate filters (Corning). Cells of PNF were enumerated by

autofluorescence using an epifluorescence microscope under green excitation. For HNF and PNF counting, a minimum of 100 fields were randomly inspected. VLP were counted using epifluorescence microscopy under blue excitation by the SYBR Green I method (Noble and Fuhrman 1998; Patel et al. 2008), using 0.02 μm pore-size Anodisc filters (Whatman). More than 300 VLP were counted and then a minimum of 20 fields were randomly examined.

Chemical variables

Samples for dissolved organic carbon (DOC) measurements were filtered through 0.2 μm polycarbonate filters (Whatman) washed with 1.2M HCl before use. DOC concentrations were determined using a total organic carbon analyzer (TOC-5000A; Shimadzu).

To determine chlorophyll *a* (chl. *a*) concentrations, 100 mL water samples were filtered through 0.2- μm polycarbonate filters (Whatman) and analyzed using the *N'*-*N*-dimethylformamide method (Moran and Porath 1980) with a fluorescence spectrometer (RF-5300PC; Shimadzu).

Quinone analysis

For RQ analysis, bacteria-sized particles in 5-L of the water samples were collected using 0.2 μm pore-size Teflon filters (Advantec) after passing through 2.0 μm nominal pore-sized GMF-2UM glass fiber filters (Whatman) to remove large particles such as phytoplankton and zooplankton. About 97.2–99.9% of bacterial cells passed through the GMF2UM filter (Takasu et al. 2013). In the experiment in June 2011, the GMF-2UM glass fiber filter was not used.

The RQ concentrations were determined using a modified method previously described by Hu et al. (1999). Briefly, quinones were extracted from the filters with a chloroform-methanol mixture (2:1, v/v) and re-extracted into hexane. UQs and MKs contained in the crude extract were separated and purified using Sep-Pak[®] Plus Silica (Waters). The molecular species and concentrations of quinones were determined using a high performance liquid chromatography (HPLC) system equipped with an ODS column (pore size,

3.5 μm ; Eclipse Plus C18, $3.0 \times 150 \text{ mm}$; Agilent) and a photodiode array detector (SPD-M20A; Shimadzu). Details on the analytical conditions have been described by Takasu *et al.* (2013). The type of quinones was identified according to the UV spectrum of each peak observed in the photodiode array detector. The quinone species were identified by the linear relationship between the logarithm of the retention times of quinones and the number of their isoprene units, using the identification-supporting sheet (made by T. Kunihiro) based on the equivalent number of isoprene units (ENIU) of quinone components as described by Tamaoka *et al.* (1983).

In the present study, we express each RQ type as follows: ubiquinone, UQ- n ; menaquinone, MK- n . The number (n) indicates the number of isoprene units in the side chain of the quinone. For example, UQ-10 represents a ubiquinone with 10 isoprenoid units, and MK-9(H_8) represents a menaquinone with 9 isoprenoid units where one of the nine units is hydrogenated with eight hydrogen atoms.

Calculations

The synthesis and destruction rates of each RQ are proxies of the growth and mortality of bacterial groups with different types of RQ. Thus, the apparent growth rates (μ_{app} , d^{-1}) of bacterial subgroups with different types of RQ were calculated from the concentrations of each RQ at the beginning and end of the incubation experiment, with the assumption that bacterial growth would follow an exponential model (Landry and Hassett 1982)

$$\mu_{\text{app}} = (1/t) \ln (N_t/N_0)$$

where t is the duration of the incubation (days), and N_0 and N_t are RQ concentrations (pmol L^{-1}) at the beginning and end of the incubation, respectively. Two dilution series were prepared: a 30 kDa dilution series to estimate the combined effects of protistan grazing and viral lysis rate ($M_{\text{g+v}}$, d^{-1}) and a 0.2 μm dilution series to determine the protistan grazing rate (M_{g} , d^{-1}) on bacteria. The slope of the regression lines from the 0.2 μm dilution series represents the grazing rate. The difference between the slopes of the regression lines

represents the bacterial mortality rate due to viral lysis (M_v , d^{-1}), and this difference was tested using analysis of covariance (ANCOVA). The intercept of the 30 kDa dilution series gives the instantaneous growth rate (μ , d^{-1}) of bacteria when neither grazing nor viral lysis occurs (Evans et al. 2003).

Carbon fluxes through bacterial subgroups with different RQ types were determined by combining the carbon conversion factor from RQ (Takasu et al. 2013) and data from the dilution experiments. For each specific bacterial subgroup, the carbon production (CP , $\mu g C L^{-1} d^{-1}$), losses to grazing (GL , $\mu g C L^{-1} d^{-1}$), and losses to viruses (VL , $\mu g C L^{-1} d^{-1}$) were calculated using the formulas of Baudoux et al. (2008):

$$CP = \mu \times P_m$$

$$GL = M_g \times P_m$$

$$VL = M_v \times P_m$$

$$P_m = P_0 \times [e^{(\mu - M_{g+v})t} - 1] / (\mu - M_{g+v})t$$

where μ (d^{-1}) is the dilution-based specific growth (y-intercept of the 30 kDa regression), M_g and M_v are the dilution-based grazing and viral lysis rates (in d^{-1}), respectively, P_0 (in $\mu g C L^{-1}$) is the initial carbon biomass of bacteria, P_m (in $\mu g C L^{-1}$) is the geometric mean carbon biomass of bacteria during the incubation, and t (in d) is the time of incubation.

Statistical analysis

Analyses with Student's t-test and ANCOVA were conducted using the free statistical environment R (R Development Core Team 2011).

Results

Physicochemical and biological conditions of the sampling site

Water samples used for the modified dilution technique, covered wide ranges of physicochemical properties: water temperatures ranged from 16.0 to 27.1°C; DOC concentrations from 1.21 to 1.56 $mg C L^{-1}$; and chl. *a* concentrations from 3.09 to 31.4 $\mu g L^{-1}$

(Table 1). The HNF number (0.5×10^6 to 2.7×10^6 cells L⁻¹) was in most cases higher than the PNF number (0.3×10^6 to 0.6×10^6 cells L⁻¹). The bacteria:Total nanoflagellates (TNF) abundance ratios in Lake Biwa ranged between 725 and 2,417 (Table 1). The VLP number varied from 2.4×10^{10} to 4.1×10^{10} VLP L⁻¹ (Table 1). The range of viruses to bacteria abundance ratios (range: 12.3 to 36.2) falls within the average reported for freshwater systems (Maranger and Bird 1995).

Estimation of growth and mortality rates

The RQ concentrations at a dilution level of 1.0 varied in the range of 32.4 to 90.9 pmol L⁻¹ at the beginning of the dilution experiments (Fig. 1A). A total of 12 types of RQ were detected, and UQ-8, UQ-10, MK-8, MK-9, and MK-9(H₈) were detected as major RQs (Fig. 1B). In general, negative relationships between the dilution factor and apparent growth rate (RQ synthesis rate) were found in the <0.2 μm diluent (Table 2, Fig. S1). We regarded relationships with $r^2 > 0.8$ and significance levels of $p < 0.1$ as statistically significant. We discussed growth and mortality rates using only the statistically significant data based on the criteria, though previous studies have used not only significant but insignificant growth and mortality values (e.g. Tijdens et al. 2008). Out of the 39 cases, 23 and 14 met these criteria for significance using the <0.2 μm diluents and the <30 kDa diluents, respectively (Table 2). In 5 out of the 39 experiments, both the <0.2 μm and <30 kDa dilution series were statistically significant, and the differences in the slopes of the two dilution series were also statistically significant (Table 2). Estimates of the growth and mortality due to grazing and lysis of UQ-8-containing bacteria in June 2011; total bacterial community (expressed as RQ in Table 2), UQ-, and UQ-8-containing bacteria in October 2011; and UQ-containing bacteria in May 2012 were statistically significant. However, most of the linear relationships between growth and the dilution factor were statistically insignificant (Table 2, Fig. S1). Rates could not be determined for UQ-10- and MK-containing bacteria in some experiments using the <30 kDa diluents because of the positive relationship (against theory) between the dilution factor and

apparent growth rate (Table 2, Fig. S1).

The growth rate of the total bacterial community (expressed as RQ in Table 3) varied from 1.05 to 2.42 d⁻¹ (Table 3), and the grazing and lysis rates of total bacteria varied from 0.46 to 0.74 d⁻¹ and 0.69 d⁻¹, respectively (Table 3). Growth and grazing rates varied among individual bacterial groups, and those of UQ-containing bacteria tended to be higher than those of MK-containing bacteria (Table 3). The sums of the grazing and lysis rates of individual groups had values close to their growth rates.

The changing patterns in the mortality due to grazing and lysis (M_{g+v}) of total bacteria and UQ-8- and UQ-10-containing bacteria were similar to the growth rates of those bacteria, even though grazing rates (M_g) on those bacteria were constant (Fig. 2).

Bacterial carbon production and losses

Daily carbon production (CP) in the present study ranged between 16.3 and 52.5 $\mu\text{g C L}^{-1} \text{ d}^{-1}$ (average: 37.4 $\mu\text{g C L}^{-1} \text{ d}^{-1}$) (Table 4), close to the estimates from previous studies in Lake Biwa using tritiated thymidine uptake (5 to 59 $\mu\text{g C L}^{-1} \text{ d}^{-1}$) (Nagata 1987) and the frequency of dividing cells (4.1 to 33 $\mu\text{g C L}^{-1} \text{ d}^{-1}$) (Nagata 1987). Grazing loss (GL) was two times higher (average: 18.5 $\mu\text{g C L}^{-1} \text{ d}^{-1}$) than lysis loss (VL) (average: 9.2 $\mu\text{g C L}^{-1} \text{ d}^{-1}$) (Table 4). GL showed a pattern of change similar to that of initial carbon biomass (Fig. 3). Contributions of grazing loss and lysis loss to bacterial production in Lake Biwa fell into the ranges of previous estimates in other lakes (Table 5). Total losses (TL) of bacteria were almost the same (average: 36.2 $\mu\text{g C L}^{-1} \text{ d}^{-1}$) as CP , and this tendency was found in individual bacterial groups (Table 4). The net carbon production of bacteria (CP_{net} : $CP - TL$) was calculated to estimate the remaining bacterial carbon biomass (Table 4). Estimates of total bacterial CP_{net} were positive on average (average: $1.2 \pm 4.4 \mu\text{g C L}^{-1} \text{ d}^{-1}$), varying from negative ($-3.1 \mu\text{g C L}^{-1} \text{ d}^{-1}$) to positive ($5.8 \mu\text{g C L}^{-1} \text{ d}^{-1}$) values. Positive estimates of CP_{net} were mostly found in UQ-8-containing bacteria (average: 0.8 $\mu\text{g C L}^{-1} \text{ d}^{-1}$, range: 0.21 to 1.78 $\mu\text{g C L}^{-1} \text{ d}^{-1}$) (Table 4). Overall, the CP_{net} values of total bacteria and individual bacterial groups were less than

274 10% of each *CP* value.

275

276 **Discussion**

277 *Grazing and lysis mortality of bacteria*

278 The dilution technique is based on a critical assumption, which assumes that the plankton
279 mortality rates resulting from predation are proportional to the dilution effect on predator
280 (grazer and/or virus) densities (Landry & Hassett 1982). This assumption may not always
281 meet because non-linear relationships between the dilution factor and apparent growth rate
282 have been frequently reported in previous studies (e.g. Tijdens et al. 2008, Personnic et al.
283 2009). In the present study, the positive relationship between the dilution factor and apparent
284 growth rate was found from some experiments using the <30 kDa diluent (Table 2, Fig. S1). It
285 is known that viral lysis of bacterial cells leads to the recycling of carbon and nutrients, both
286 of which are derived from lysed bacterial biomass and are re-supplied to bacteria (Bratbak et
287 al. 1990; Gobler et al. 1997; Proctor and Fuhrman 1991). Thus, the possible interpretation
288 could be that growth of bacteria was stimulated by the lysed bacterial cells in some of the <30
289 kDa diluent series.

290 In limnetic and oceanic systems, concentrations of organic substrates and inorganic
291 nutrients are generally low, causing death of planktonic bacteria to starvation (Amy and
292 Morita 1983). So, planktonic bacteria are required to have some physiological adaptation for
293 efficient uptake and utilization of substrates as well as for long-term survival under carbon
294 and energy limitation (del Giorgio and Gasol 2008). In Lake Biwa, however, the sum of total
295 bacterial mortalities due to protistan grazing and viral lysis accounted for 96.6% (range; 89.0 -
296 107.2%) of daily total bacterial production in the present study. Thus, almost all bacterial
297 production is efficiently consumed by protists and viruses before bacterial death due to other
298 environmental factors in Lake Biwa within a day. In other words, fate of bacterial production
299 in the epilimnion of Lake Biwa may be mainly determined through microbial interactions.

This is the first study which demonstrates a balanced relationship between bacterial production and losses in a freshwater lake using the direct estimation method. The modified dilution method was rarely applied to the estimation of protistan grazing and virus lysis of freshwater bacteria, and only two previous studies are so far available in freshwater systems (Personnic et al. 2009, Tijdens et al. 2008) (Table 5). In those studies, however, estimates of grazing losses frequently exceeded the sum of grazing plus lysis loss estimates. Personnic et al. (2009) suggested that some complex interactions among microbes such as synergistic and antagonistic effects of viral lysis and protistan grazing on bacterial production occurred in their incubation experiments. By contrast, the sum of the grazing and lysis losses of bacteria (*TL*) was almost equal to *CP* in the present study (Table 4). Thus, complex interactions among microbes might be negligible in our experiments, and almost all bacterial production is efficiently consumed by protists and viruses in Lake Biwa.

Bacterial abundances in the epilimnion of Lake Biwa were relatively constant and changed within the order of 10^9 cells L^{-1} (Nagata 1987; Nakano 1992; Nishimura et al. 2005; Takasu et al. 2013), whereas other microbes such as picophytoplankton, nanoflagellates, and viruses respectively showed large variations from 10^5 to 10^8 cells L^{-1} (Nagata 1988), from 10^2 to 10^5 cells L^{-1} (Nagata 1988), and from 10^9 to 10^{11} VLP L^{-1} (Nishimura and Nagata 2007; Pradeep Ram et al. 2010). In addition, bacterial abundance is generally less variable than bacterial production (Nagata 1987; Nakano 1992). In the present study, the percentages of remaining bacterial carbon production ($\%CP_{\text{net}} = (CP_{\text{net}} / CP) \times 100$) ranged from -7.2 to 11.0% (average $\pm$ SD, 3.4 ± 9.4). These estimates suggest that the less than 11% of daily bacterial production remains or excessively loses, and those small increase or decrease in net bacterial production does not change the whole bacterial abundance so much. This may be the major reason why the bacterial abundance in the epilimnion of Lake Biwa is relatively constant.

Owing to food limitation, nanoflagellates in lakes show clearance rates independent of

bacterial density (Bird and Kalff 1993). The bacteria:TNF ratios in Lake Biwa were low (average: 1346, Table 1) and similar to those of oligotrophic lakes and oceans (Sanders et al. 1992). Thus, nanoflagellates in Lake Biwa are also under food limitation. When we calculated the specific grazing loss ($SGL = GL / \text{initial TNF density}$), SGL estimates were relatively constant ($1.24 \pm 0.29 \times 10^{-5} \mu\text{g C TNF cell}^{-1} \text{d}^{-1}$) despite variations in the bacteria:TNF ratio (Table 1). The previous study suggested that nanoflagellates are clearing as much water as is physically possible independent of bacterial density when they are under food limitation (Bird and Kalff 1993). The results of the present study also indicate that individual TNF in Lake Biwa ingest as much bacteria as possible due to the chronically food limitation. The elimination mechanisms of bacteria should be different between protistan grazing and viral lysis, though bacterial production and the sum of grazing and lysis losses were almost equal in Lake Biwa (Table 4). In the present study, most of the viral lysis rates (M_v) were not statistically significant (Table 2), similar to those in previous studies (e.g. Tijdens et al. 2008; Personic et al. 2009). The sum of grazing and lysis mortality rates (M_{g+v}) seasonally changed, though grazing rates (M_g) were relatively constant during the study period (Fig. 2). Thus, the changing pattern of M_{g+v} is due to that of M_v . So, the changing pattern of M_{g+v} was similar to that of growth rate, suggesting that viral mortality is coupled with bacterial growth rate (Fig. 2). Similar results have been reported in previous studies (e.g., Weinbauer 2003). In Lake Biwa, the loss processes of bacterial production can be explained by a combination of protistan grazing pressure, which is independent of bacterial abundance, and viral lysis, which is dependent on bacterial growth (Fig. 2, RQ).

Carbon fluxes through major bacterial groups and differentiating the ecological traits of bacterial groups

The present study elucidated that planktonic bacterial assemblages in Lake Biwa consisted of various RQs subgroups with different growth and mortality rates (Table 3). The growth rates of UQ-containing bacteria were higher than those of MK-containing bacteria in Lake Biwa

(Table 3). Because of the large midpoint potential of UQs, UQs are thermodynamically favorable compounds to use oxygen as an electron acceptor compared to MKs, and this makes energy gains by UQs higher than those of MKs (Søballe and Pool 1999). The coupling between bacterial production and oxygen consumption (respiration) has been reported in aquatic systems (reviewed by del Girogio and Cole 1998; Robinson 2008). Thus, UQ-containing bacteria may have an advantage over other bacteria in the epilimnion of Lake Biwa because of the lake's oxygenated condition. UQ-containing bacteria showed higher GL than MK-containing bacteria (Table 3), probably due to higher abundance which would have higher encounter with nanoflagellates. In addition, there may be another reason why UQ-containing bacteria had higher GL. In freshwater planktonic bacterial community, UQ-8- and UQ-10-containing bacteria are generally gram-negative Proteobacteria, whereas MK-8, MK-9, and MK-9(H₈)-containing bacteria are gram-positive Actinobacteria (e.g. Hiraishi and Kato 1999). The consumption of gram-positive freshwater planktonic bacteria (mainly Actinobacteria) is selectively avoided by nanoflagellates due to various protection mechanisms of gram-positive bacteria such as cell surface charge and cell size reduction (reviewed by Pernthaler 2005). Thus, it is possible that UQ-containing bacteria are preferentially grazed by nanoflagellates.

UQ-8- and UQ-10-containing bacteria were the most dominant and fastest-growing bacterial groups during the study period (Fig. 1, Table 3). The average *CPs* of UQ-8- and UQ-10-containing bacteria were respectively estimated at 12.2 and 13.8 $\mu\text{g C L}^{-1} \text{d}^{-1}$, accounting for 31.7% (range: 28.4 to 35.2%) and 28.5% (range: 25.0 to 32.1%) of total bacterial *CP* (Table 4). The sum of the average *TL* of UQ-8- and UQ-10-containing bacteria (26.0 $\mu\text{g C L}^{-1} \text{d}^{-1}$) also accounted for 60% of total bacterial *TL* (Table 4). Hence, the fate of those two bacterial groups explained a large portion of the carbon fluxes within the microbial loop of Lake Biwa. Our estimation thus implies that the magnitude of carbon fluxes within the microbial loop can be regulated by the production of major bacterial groups.

R-BT065 cluster bacteria (a subcluster of Beta-proteobacteria) have been reported to be the most abundant (up to 50% of total bacteria) (Zwart et al. 2002; 2003) and have UQ-8 as their major RQ (e.g., Hahn et al. 2010a; 2010b; Kasalický et al. 2010). R-BT065 cluster bacteria have been reported to be the fastest-growing segment of bacterial communities in European freshwater lakes (Šimek et al. 2006; Salcher et al. 2008). These bacteria preferentially inhabit the oxygenated layer of oligo-mesotrophic lakes (Piburger See, Austria; Lake Zurich, Switzerland) (Salcher et al. 2008; 2011). The environmental conditions of the epilimnetic water in the north basin of Lake Biwa may be similar to those of the aforementioned European lakes (Nishimura et al. 2005; Kim et al. 2006; Takasu et al. 2013). UQ-10-containing bacteria have been classified as Alpha-proteobacteria (Hiraishi 1999). In the freshwater Alpha-proteobacteria, members of the LD12 clade are one of the most abundant ubiquitous lineages (Salcher et al. 2011). Previous study suggested that LD12 bacteria generally prefer the upper epilimnetic water layers during nutrient limited summer season (Salcher et al. 2011). This feature is well consistent with UQ-10-containing bacteria in Lake Biwa (Takasu et al. 2013).

The CP_{net} of UQ-8-containing bacteria showed positive estimates, whereas the CP_{net} of UQ-10 containing bacteria showed negative estimates in the present study (Table 4). In our previous study, UQ-8-containing bacteria dominated in Lake Biwa throughout a year, whereas biomass of UQ-10-containing bacteria was relatively low (Takasu et al. 2013). In addition, biomass of those bacterial groups showed different seasonal changing patterns (Takasu et al. 2013). Thus, it is likely that relatively low biomass of UQ-10-containing bacteria was due to higher loss of the bacteria by protistan grazing and/or viral lysis than that of UQ-8-containing bacteria in Lake Biwa.

Conclusion

Our study is the first to demonstrate a balanced relationship between bacterial production and losses in a freshwater lake using the modified dilution method. UQ-8- and UQ-10-containing

bacteria were the two dominant groups, and the sum of their production and losses explained 60% of the carbon fluxes within the microbial loop. Thus, a large portion of carbon fluxes through the bacterial community may be explained by the carbon fluxes of dominant bacterial groups.

Acknowledgments

We are grateful to Dr. Masayuki Ushio for his invaluable comments as well as his encouragement throughout this study. We also thank Mr. Tadatoshi Koitabashi and Dr. Yukiko Goda for their assistance during field sampling, and thank other colleagues at the Center for Ecological Research, Kyoto University, for their valuable comments on this study. This study was partly supported by Japan Society for the Promotion of Science (JSPS) KAKENHI Grant Numbers 23370010 and Japan Science and Technology Strategic International Research Cooperative Program (Japan-China) on Science and Technology for Environmental Conservation and Construction of a Society with Less Environmental Burden "Fate of dissolved organic matter in lakes with special reference to loading and pollution" to S.N. and 21710081, 23710010 to T.K. H.T. was supported by JSPS KAKENHI Grant Number 11J00658.

422

423 **References**

- 424 Amy PS, Morita RY (1983) Starvation-survival patterns of sixteen freshly isolated
425 open-ocean bacteria. *Appl Environ Microbiol* 45:1109–1115
- 426 Azam F, Fenchel T, Field JG, Gray JS, Meyer-Reil LA, Thingstad F (1983) The
427 ecological role of water-column microbes in the sea. *Mar Ecol Prog Ser* 10:257–263
- 428 Baudoux AC, Veldhuis MJW, Noordeloos AAM, van Noort G, Brussaard CPD (2008)
429 Estimates of virus- vs. grazing induced mortality of picophytoplankton in the North
430 Sea during summer. *Aquat Microb Ecol* 52:69–82
- 431 Bettarel Y, Amblard C, Sime-Ngando T, Carrias J-F, Sargos D, Garabétian F, Lavandier
432 P (2003) Viral lysis, flagellate grazing potential, and bacterial production in Lake
433 Pavin. *Microb Ecol* 45:119–127
- 434 Bird D, Kalff J (1993) Protozoan grazing and the size–activity structure of limnetic
435 bacterial communities. *Can J Fish aquat Sci* 50:370–380
- 436 Bouvier T, del Giorgio PA (2003) Factors influencing the detection of bacterial cells
437 using fluorescence in situ hybridization (FISH): A quantitative review of published
438 reports. *FEMS Microbiol Ecol* 44:3–15
- 439 Bratbak G, Heldal M, Norland S, Thingstad TF (1990) Viruses as partners in spring
440 bloom microbial trophodynamics. *Appl Environ Microbiol* 56:1400–1405
- 441 Caron DA (1983) Technique for enumeration of heterotrophic and phototrophic
442 nanoplankton, using epifluorescence microscopy, and comparison with other
443 procedures. *Appl Environ Microbiol* 46: 491–498
- 444 Cole JJ, Findlay SS, Pace ML (1988) Bacterial production in fresh and saltwater
445 ecosystems: a cross-system overview. *Mar Ecol Prog Ser* 43:1–10

- 446 Collins MD, Jones D (1981) Distribution of isoprenoid quinone structural types in
447 bacteria and their taxonomic implications. *Microbiol Rev* 45: 316–354
- 448 Constanza R, d’Arge R, de Groot R, Farber S, Grasso M, Hannon B, Limburg K,
449 Naeem S, O’Neill RV, Paruelo J, Raskin RG, Sutton P, van den Belt M (1997) The
450 value of the world’s ecosystem services and natural capital. *Nature* 387: 253–260
- 451 del Girogio PA, Cole JJ (1998) Bacterial growth efficiency in natural aquatic systems.
452 *Ann Rev Ecol Systematics* 29: 503–541
- 453 del Giorgio PA, Gasol JM (2008) Physiological structure and single-cell activity in
454 marine bacterioplankton, In Kirchman DL (ed) *Microbial ecology of the oceans*.
455 (2nd edition) Wiley, New York, pp. 243–298
- 456 Ducklow HW (2000) Bacterial production and biomass in the oceans. In Kirchman DL
457 (ed) *Microbial ecology of the oceans*. Wiley, New York, pp 85–120
- 458 Evans C, Archer SD, Jacquet S, Wilson WH (2003) Direct estimates of the contribution
459 of viral lysis and microzooplankton grazing to the decline of a *Micromonas* spp.
460 population. *Aquat Microb Ecol* 30:207–219
- 461 Glöckner FO, Fuchs BM, Amann R (1999) Bacterioplankton compositions of lakes and
462 oceans: a first comparison based on fluorescence in situ hybridization. *Appl Environ*
463 *Microbiol* 65:3721–3726
- 464 Gobler CJ, Hutchins DA, Fisher NS, Coper EM, Sanudo-Wilhelmy SA (1997) Release
465 and bioavailability of C, N, P, Se, and Fe following viral lysis of a marine
466 chrysophyte. *Limnol Oceanogr* 42:1492–1504
- 467 Gobler CJ, Davis TW, Deonarine SN, Saxton MA, Lavrentyev PJ, Jochem FJ, Wilhelm
468 SW. (2008) Grazing and virus-induced mortality of microbial populations before
469 and during the onset of annual hypoxia in Lake Erie. *Aquat Microb Ecol* 51: 117–

- 470 128
- 471 Hahn MW, Kasalický V, Jezbera J, Brandt U, Jezberová J, Šimek K (2010a)
- 472 *Limnohabitans curvus* gen. nov., sp. nov., a planktonic bacterium isolated from a
- 473 freshwater lake. Int J Syst Evol Microbiol 60:1358–1365
- 474 Hahn MW, Kasalický V, Jezbera J, Brandt U, Šimek K (2010b) *Limnohabitans*
- 475 *australis* sp. nov., isolated from a freshwater pond, and emended description of the
- 476 genus *Limnohabitans*. Int J Syst Evol Microbiol 60:2946–2950
- 477 Hedrick DB, White DC (1986) Microbial respiratory quinones in the environment. J
- 478 Microbiol Meth 5:243–254
- 479 Ducklow HW (2000) Bacterial production and biomass in the oceans. In Kirchman DL
- 480 (ed) Microbial ecology of the oceans. Wiley, New York, pp 85–120
- 481 Herdendorf CE (1990) Distribution of the world's large lakes. In Tilzer MM, Serruya C
- 482 (ed) Large lakes: ecological structure and function. Springer Verlag, Berlin. pp. 3–
- 483 38
- 484 Hiraishi A (1999) Isoprenoid quinones as biomarkers of microbial populations in the
- 485 environment. J Biosci Bioeng 88:449–460
- 486 Hiraishi A, Kato K (1999) Quinone profiles in lake sediments: Implications for
- 487 microbial diversity and community structures. J Gen Appl Microbiol 45: 221–227
- 488 Hu H-Y, Fujie K, Urano K (1999) Development of a novel solid phase extraction
- 489 method for the analysis of bacterial quinones in activated sludge with a higher
- 490 reliability. J. Biosci. Bioeng. 87: 378–382
- 491 Jacquet S, Domaizon I, Personnic S, Pradeep Ram AS, Hedal M, Duhamel S,
- 492 Sime-Ngando T (2005) Estimates of protozoan- and viral-mediated mortality of
- 493 bacterioplankton in Lake Bourget (France). Freshw Biol 50:627–645

- 494 Kasalický V, Jezbera J, Šimek K, Hahn MW (2010) *Limnohabitans planktonicus* sp.
495 nov. and *Limnohabitans parvus* sp. nov., planktonic betaproteobacteria isolated
496 from a freshwater reservoir, and emended description of the genus *Limnohabitans*.
497 Int J Syst Evol Microbiol 60:2710–2714
- 498 Kim C, Nishimura Y, Nagata T (2006) Role of dissolved organic matter in hypolimnetic
499 mineralization of carbon and nitrogen in a large, monomictic lake. Limnol Oceanogr
500 51:70–78
- 501 Kunihiro T, Takasu H, Miyazaki T, Uramoto Y, Kinoshita K, Yodnarasri S, Hama D,
502 Wada M, Kogure K, Ohwada K, Tsutsumi H (2011) Increase in
503 *Alphaproteobacteria* in association with a polychaete, *Capitella* sp. I, in the
504 organically enriched sediment. The ISME J 5:1818–1831
- 505 Landry MR, Hassett RP (1982) Estimating the grazing impact of marine
506 micro-zooplankton. Mar Biol 67:283–288
- 507 Landry MR, Haas LW, Fagerness VL (1984) Dynamics of microbial plankton
508 communities: experiments in Kaneohe Bay, Hawaii. Mar Ecol Prog Ser 16:127–133
- 509 Maranger R, Bird DF (1995) Viral abundance in aquatic systems: a comparison between
510 marine and fresh waters. Mar Ecol Prog Ser 121:217–226
- 511 Miki T, Yokokawa T, Nagata T, Yamamura N (2008) Immigration of prokaryotes to
512 local environments enhances remineralization efficiency of sinking particles: a
513 metacommunity model. Mar Ecol Prog Ser 366:1–14
- 514 Moran R, Porath D (1980) Chlorophyll determination in intact tissues using
515 *N,N*-Dimethylformamide. Plant Physiol 65:478–479
- 516 Nagata T (1987) Production rate of planktonic bacteria in the north basin of Lake Biwa,
517 Japan. Appl Environ Microbiol 53:2872–2882

- 518 Nagata T (1988) The microflagellate–picoplankton food linkage in the water column of
519 Lake Biwa. *Limnol Oceanogr* 33:504–517
- 520 Nagata T (1990) Contribution of picoplankton to the grazer food chain of Lake Biwa. In
521 Tilzer MM, Serruya C (ed) Large lakes: ecological structure and function. Springer
522 Verlag, Berlin. pp. 526–539
- 523 Nagata T, Kumagai M, Yoshiyama K (2012) *Ondanka no kosho-gaku*. Kyoto
524 University press (in Japanese)
- 525 Nakano S-I (1992) Changes in bacterioplankton production and dominant algal species
526 in the north basin of Lake Biwa. *Jpn J Limnol* 53:145–149
- 527 Nishimura Y, Kim C, Nagata T (2005) Vertical and Seasonal Variations of
528 Bacterioplankton subgroups with different nucleic acid contents: possible regulation
529 by phosphorus. *Appl Microbiol* 71:5828–5836
- 530 Nishimura Y, Nagata T (2007) Alphaproteobacterial dominance in a large mesotrophic
531 lake (Lake Biwa, Japan). *Aquat Microb Ecol* 48:231–240
- 532 Noble RT, Fuhrman JA (1998) Use of SYBR Green I for rapid epifluorescence counts
533 of marine viruses and bacteria. *Aquat Microb Ecol* 14:113–118
- 534 Patel A, Nobel RT, Steele JA, Schwalbach MS, Hewson I, Fuhrman JA (2008) Virus
535 and prokaryote enumeration from planktonic aquatic environments by
536 epifluorescence microscopy with SYBR Green I. *Nat Protoc* 2:269–276
- 537 Pernthaler J, Glöckner FO, Unterholzner S, Alfreider A, Psenner R, Amann R (1998)
538 Seasonal community and population dynamics of pelagic Bacteria and Archaea in a
539 high mountain lake. *Appl Environ Microbiol* 64:4299–4306
- 540 Pernthaler J (2005) Predation on prokaryotes in the water column and its ecological
541 implications. *Nature Rev Microbiol* 3:537–546

- Personnic S, Domaizon I., Sime-Ngando T, Jacquet S (2009) Seasonal variations of microbial abundances and virus- versus flagellate-induced mortality of picoplankton in three peri-alpine lakes. *J Plankton Res* 31: 1161–1177
- Pirlot S, Unrein F, Descy JP, Servais P (2007) Fate of heterotrophic bacteria in Lake Tanganyika (East Africa). *FEMS Microbial Ecol* 62:354–364
- Porter KG, Feig YS (1980) The use of DAPI for identifying and counting aquatic microflora. *Limnol Oceanogr* 25:943–948
- Pradeep Ram AS, Nishimura Y, Tomaru Y, Nagasaki K, Nagata T (2010) Seasonal variation in viral-induced mortality of bacterioplankton in the water column of a large mesotrophic lake (Lake Biwa, Japan). *Aquat Microb Ecol* 58: 249–259
- Proctor LM, Fuhrman JA (1991) Roles of viral infection in organic particulate flux. *Mar Ecol Prog Ser* 69:133–142
- R Development Core Team (2011) R: A language and environment for statistical computing. Vienna, Austria
- Robinson C (2008) Heterotrophic bacterial respiration. In Kirchman DL (ed) *Microbial ecology of the oceans*. Wiley, New York, pp 299–334
- Salcher MM, Pernthaler J, Zeder M, Psenner R, Posch T (2008) Spatio-temporal niche separation of planktonic *Betaproteobacteria* in an oligo-mesotrophic lake. *Environ Microbiol* 10:2074–2086
- Salcher MM, Pernthaler J, Posch T (2011) Seasonal bloom dynamics and ecophysiology of the freshwater sister clade of SAR11 bacteria ‘that rule the waves’ (LD12). *The ISME J* 5:1242–1252
- Salcher MM, Pernthaler J, Frater N, Posch T (2011) Vertical and longitudinal distribution patterns of different bacterioplankton populations in a canyon-shaped,

- 566 deep prealpine lake. *Limnol Oceanogr* 56:2027–2039
- 567 Sanders RW, Caron DA, Berninger U-G (1992) Relationships between bacteria and
568 heterotrophic nanoplankton in marine and fresh waters: an inter-ecosystem
569 comparison. *Mar Ecol Prog Ser* 86:1–14
- 570 Šimek K, Pernthaler J, Weinbauer MG, Horňák K, Dolan JR, Nedoma J, Mašín M,
571 Amann R (2001) Changes in bacterial community composition and dynamics and
572 viral mortality rates associated with enhanced flagellate grazing in a mesotrophic
573 reservoir. *Appl Environ Microbiol* 67: 2723–2733
- 574 Šimek K, Horňák K, Jezbera J, Nedoma J, Vrba J, Straškrábová V, Macek N, Dolan JR,
575 Hahn MW (2006) Maximum growth rates and possible life strategies of different
576 bacterioplankton groups in relation to phosphorus availability in a freshwater
577 reservoir. *Environ Microbiol* 8:1613–1624
- 578 Šimek K, Weinbauer G, Horňák K, Jezbera J, Nedoma J, Dolan JR (2007) Grazer and
579 virus-induced mortality of bacterioplankton accelerates development of
580 *Flectobacillus* populations in a freshwater community. *Environ Microbiol* 9: 789–
581 800
- 582 Søballe B, Pool KP (1999) Microbial ubiquinones: multiple roles in respiration, gene
583 regulation and oxidative stress management. *Microbiology* 145: 1817–1830
- 584 Taira Y, Uchimiya M, Kudo I (2009) Simultaneous estimation of viral lysis and
585 protozoan grazing on bacterial mortality using a modified virus-dilution method.
586 *Mar Ecol Prog Ser* 379: 23–32
- 587 Takasu H, Kunihiro T, Nakano S-I (2013) Estimation of carbon biomass and community
588 structure of planktonic bacteria in Lake Biwa using respiratory quinone analysis.
589 *Limnology* 14: 247–256

- 590 Tamaoka J, Katayama-Fujimura H, Kuraishi H (1983) Analysis of bacterial
591 menaquinone mixtures by high performance liquid chromatography. *J Appl*
592 *Microbiol* 54: 31–36
- 593 Thingstad TF (2000) Elements of a theory for the mechanisms controlling abundance,
594 diversity, and biogeochemical role of lytic bacterial viruses in aquatic systems.
595 *Limnol Oceanogr* 45:1320–1328
- 596 Tjeldens M, Van de Wall BD, Slovackova H, Hoogveld HL, Gons HJ (2008) Estimates
597 of bacterial and phytoplankton mortality caused by viral lysis and microzooplankton
598 grazing in a shallow eutrophic lake. *Freshw Biol* 53:1126–1141
- 599 Tremaine SC, Mills AL (1987) Tests of the critical assumptions of the dilution method
600 for estimating bacterivory by microeucaryotes. *Appl Environ Microbiol* 53:2914–
601 2921
- 602 Weinbauer MG, Höfle, MG (1998) Significance of viral lysis and flagellate grazing as
603 factors controlling bacterioplankton production in a eutrophic lake. *Appl Environ*
604 *Microbiol* 64: 431–438
- 605 Weinbauer MG, Christaki U, Nedoma J, Šimek K. (2003) Comparing the effects of
606 resource enrichment and grazing on viral production in a meso-eutrophic reservoir.
607 *Aquat Microb Ecol* 31:137–144
- 608 Zwart G, Crump BC, Kamst-van Agterveld MP, Hagen F, Han S-K (2002) Typical
609 freshwater bacteria: an analysis of available 16S rRNA gene sequences from
610 plankton of lakes and rivers. *Aquat Microb Ecol* 28:141–155
- 611 Zwart G, van Hannen EJ, Kamst-van Agterveld MP, der Gucht KV, Lindström ES,
612 Wichelen JV, Lauridsen T, Crump BC, Han S-K, Declerck S (2003) Rapid
613 screening for freshwater bacterial groups by using reverse line blot hybridization.

614 Appl Environ Microbiol 69: 5875–5883

615

Table 1. Physico-chemical and biological variabes

Date	WT (°C)	DOC (mg C L ⁻¹)	Chl. <i>a</i> (µg L ⁻¹)	Bacteria (10 ⁹ cells L ⁻¹)	TNF (10 ⁶ cells L ⁻¹)	HNF (10 ⁶ cells L ⁻¹)	PNF (10 ⁶ cells L ⁻¹)	VLP (10 ¹⁰ VLP L ⁻¹)	Bacteria:TNF	VBR
22 June 2011	22.0	ND	9.07	2.0	3.0	2.7	0.3	2.4	725	12.3
19 October 2011	20.0	ND	3.09*	1.2	1.1	0.5	0.6	3.1	2417	25.3
16 May 2012	16.0	ND	31.1*	0.9	1.4	1.1	0.3	3.1	759	36.2
25 June 2012	21.5	1.21	31.4	2.5	2.4	1.8	0.6	3.4	1389	13.3
24 July 2012	27.1	1.56	6.16	2.6	2.3	1.8	0.5	4.1	1441	15.7

WT, Water Temperature

ND, Not Determined

VBR, Viruses to bacterium ratio

*Water samples were filtered using different type of filter (GF/F glass fiber filter, Whatman)

(I. Mukherjee and S.D. Thottathil, personal communication).

Table 2. Summary of growth (μ), grazing mortality (M_g), lysis mortality (M_v), and total mortality (M_{g+v}) from results of the dilution experiments.

Date	Bacterial RQ type	Diluent	Dilution level	Linear fit		Regression slopes	μ	M_g	M_v	M_{g+v}
				r^2	p limit	p limit				
22 June 2011	RQ	0.2 μ m	4	0.993	<0.01	NS	1.128	0.697	0.123	0.820
		30 kDa	4	0.659	NS					
	UQ	0.2 μ m	4	0.996	<0.01	NS	1.206	0.777	0.133	0.910
		30 kDa	4	0.727	NS					
	MK	0.2 μ m	4	0.688	NS	NS	0.714	0.42	-0.082	0.338
		30 kDa	4	0.175	NS					
	UQ-8	0.2 μ m	4	0.957	<0.05	<0.05	1.943	0.578	1.095	1.673
		30 kDa	4	0.965	<0.05					
	UQ-10	0.2 μ m	4	0.989	<0.01	NS	0	0.812		
		30 kDa	4	0.004	NS					
	MK-8	0.2 μ m	4	0.127	NS	NS	0.360	0.070	0.178	0.248
		30 kDa	4	0.304	NS					
19 October 2011	RQ	0.2 μ m	3*	0.865	<0.1	NS	0	-0.248		
		30 kDa	4	0.464	NS					
	RQ	0.2 μ m	4	0.980	<0.01	<0.01	1.225	0.459	0.689	1.148
		30 kDa	4	0.996	<0.01					
	UQ	0.2 μ m	4	0.944	<0.05	<0.01	1.767	0.268	1.471	1.733
		30 kDa	4	0.978	<0.05					
	MK	0.2 μ m	4	0.968	<0.05	<0.05	0	0.602		
		30 kDa	4	0.436	NS					
	UQ-8	0.2 μ m	4	0.926	<0.05	<0.01	2.188	0.543	1.554	2.097
		30 kDa	4	0.998	<0.01					
	UQ-10	0.2 μ m	4	0.440	NS	NS	0.192			0.161
		30 kDa	4	0.063	NS					
	MK-7	0.2 μ m	4	0.986	<0.01	<0.1	0	0.968		
		30 kDa	4	0.069	NS					
	MK-8	0.2 μ m	4	0.819	<0.1	<0.05	0	0.533		
		30 kDa	4	0.768	NS					
	MK-9	0.2 μ m	4	0.873	<0.1	<0.1	0	0.589		
		30 kDa	3*	0.651	NS					
	MK-9(H ₈)	0.2 μ m	4	0.882	<0.1	<0.05	0	0.351		
		30 kDa	4	0.601	NS					
16 May 2012	RQ	0.2 μ m	4	0.612	NS	<0.05	2.424	0.646	1.511	2.157
		30 kDa	4	0.948	<0.05					
	UQ	0.2 μ m	4	0.829	<0.1	<0.05	1.382	2.066	-0.936	1.13
		30 kDa	4	0.999	<0.001					
	MK	0.2 μ m	4	0.591	NS	NS	0.995	1.136	-0.859	0.277
		30 kDa	4	0.427	NS					
	UQ-8	0.2 μ m	4	0.648	NS	<0.05	2.962	0.721	2.052	2.773
		30 kDa	4	0.952	<0.05					
	UQ-10	0.2 μ m	4	0.710	NS	<0.05	2.524	0.552	2.012	2.564
		30 kDa	4	0.944	<0.05					
	MK-8	0.2 μ m	4	0.559	NS	NS	0.481	1.027	-0.718	0.309
		30 kDa	4	0.198	NS					
	MK-9(H ₈)	0.2 μ m	4	0.098	NS	NS	0.253	0.290	-0.188	0.102
		30 kDa	3*	0.101	NS					

Table 2. Continued

Date	Bacterial RQ type	Diluent	Dilution level	Linear fit		Regression slopes	μ	M_g	M_v	M_{g+v}
				r^2	p limit	p limit				
25 June 2012	RQ	0.2 μ m	4	0.968	<0.05					
		30 kDa	3**	0.958	NS	<0.1	1.448	0.644	0.720	1.364
	UQ	0.2 μ m	4	0.976	<0.05					
		30 kDa	3**	0.954	NS	NS	1.567	1.368	0.080	1.448
	MK	0.2 μ m	4	0.831	<0.1					
		30 kDa	3**	0.963	NS	<0.05	1.349	0.113	1.183	1.296
	UQ-8	0.2 μ m	4	0.985	<0.01					
		30 kDa	4	0.961	NS	NS	1.245	1.589	-0.381	1.208
	UQ-10	0.2 μ m	4	0.958	<0.05					
		30 kDa	3**	0.937	NS	NS	1.816	1.077	0.522	1.599
	MK-8	0.2 μ m	4	0.019	NS					
		30 kDa	3**	0.911	NS	NS	0	0.088		
	MK-9	0.2 μ m	4	0.595	NS					
		30 kDa	3**	0.259	NS	NS	0.220	0.388	-0.126	0.262
	MK-9(H ₈)	0.2 μ m	4	0.003	NS					
		30 kDa	3**	0.983	<0.1	NS	0.242	0.046	0.277	0.323
24 July 2012	RQ	0.2 μ m	4	0.862	<0.1					
		30 kDa	4	0.870	<0.1	NS	1.048	0.744	0.379	1.123
	UQ	0.2 μ m	4	0.92	<0.05					
		30 kDa	4	0.897	<0.1	NS	1.446	1.007	0.492	1.499
	MK	0.2 μ m	4	0.76	NS					
		30 kDa	4	0.655	NS	NS	0.416	0.644	-0.24	0.404
	UQ-8	0.2 μ m	4	0.930	<0.05					
		30 kDa	4	0.891	<0.1	NS	1.202	0.926	0.256	1.182
	UQ-10	0.2 μ m	4	0.911	<0.05					
		30 kDa	4	0.913	<0.05	NS	1.636	1.120	0.647	1.767
	MK-8	0.2 μ m	4	0.393	NS					
		30 kDa	4	0.222	NS	NS	0.026	0.392	-0.269	0.123
	MK-9	0.2 μ m	4	0.535	NS					
		30 kDa	4	0.841	<0.1	NS	1.398	0.409	0.893	1.302
	MK-9(H ₈)	0.2 μ m	4	0.923	<0.05					
		30 kDa	4	0.455	NS	<0.1	0.056	0.933	-0.684	0.249

Statistically meaningful values are shown in bold.

NS, Not significant

*20% Initial sample was under the detection limit.

**40% Initial sample could not be measured due to laboratory accident.

Table 3. Ranges and averages of daily growth (μ), grazing mortality (M_g), lysis mortality (M_v),
and total mortality (M_{g+v})

Bacterial RQ type	μ (d ⁻¹)		M_g (d ⁻¹)		M_v (d ⁻¹)		M_{g+v} (d ⁻¹)	
	Range	Average $\pm$ SD	Range	Average $\pm$ SD	Range	Average $\pm$ SD	Range	Average $\pm$ SD
	(Min - Max)		(Min - Max)		(Min - Max)		(Min - Max)	
RQ	1.05 – 2.42	1.57 $\pm$ 0.75	0.46 – 0.74	0.64 $\pm$ 0.13	–	0.69	1.12 – 2.16	1.48 $\pm$ 0.59
UQ	1.38 – 1.77	1.53 $\pm$ 0.21	0.27 – 2.07	1.10 $\pm$ 0.67	-0.94 – 1.47	0.27 $\pm$ 1.70	1.13 – 1.73	1.45 $\pm$ 0.30
MK	–	–	0.11 – 0.60	0.36 $\pm$ 0.36	–	–	–	–
UQ-8	1.20 – 2.96	2.07 $\pm$ 0.73	0.54 – 1.59	0.91 $\pm$ 0.49	1.10 – 1.55	1.33 $\pm$ 0.33	1.18 – 2.77	1.93 $\pm$ 0.67
UQ-10	1.64 – 2.52	2.08 $\pm$ 0.63	0.81 – 1.12	1.00 $\pm$ 0.17	–	–	1.77 – 2.56	2.17 $\pm$ 0.56
MK-8	–	–	–	0.53	–	–	–	–
MK-9	–	1.40	–	0.59	–	–	–	1.30
MK-9(H ₈)	–	0.24	-0.25 – 0.93	0.35 $\pm$ 0.59	–	–	–	0.32

SD, Standard Deviation

Table 4. Ranges and averages of daily carbon production (CP), grazing loss (GL), lysis loss (LL), total losses (TL) and net carbon production (CP_{net})

Bacterial RQ type	CP		GL		LL		TL		CP_{net}	
	$(\mu\text{g C L}^{-1} \text{ d}^{-1})$		$(\mu\text{g C L}^{-1} \text{ d}^{-1})$		$(\mu\text{g C L}^{-1} \text{ d}^{-1})$		$(\mu\text{g C L}^{-1} \text{ d}^{-1})$		$(\mu\text{g C L}^{-1} \text{ d}^{-1})$	
	Range (Min - Max)	Average $\pm$ SD	Range (Min - Max)	Average $\pm$ SD	Range (Min - Max)	Average $\pm$ SD	Range (Min - Max)	Average $\pm$ SD	Range (Min - Max)	Average $\pm$ SD
RQ	16.3 – 52.5	37.4 $\pm$ 18.8	6.1 – 30.9	18.5 $\pm$ 17.5	–	9.2	15.3 – 46.7	36.2 $\pm$ 18.1	-3.1 – 5.8	1.2 $\pm$ 4.4
UQ	8.0 – 27.0	19.7 $\pm$ 18.8	1.3 – 40.3	19.4 $\pm$ 19.7	-18.3 – 6.9	-5.7 $\pm$ 17.8	8.1 – 24.7	18.3 $\pm$ 8.9	-0.9 – 4.9	1.4 $\pm$ 3.1
MK	–	–	–	–	–	–	–	–	–	–
UQ-8	5.1 – 18.5	12.2 $\pm$ 5.5	1.3 – 9.5	4.9 $\pm$ 4.2	3.6 – 7.2	5.4 $\pm$ 2.5	4.9 – 17.3	11.3 $\pm$ 5.1	0.2 – 1.8	0.8 $\pm$ 0.8
UQ-10	10.9 – 16.8	13.8 $\pm$ 4.2	–	7.4	–	–	11.7 – 17.1	14.4 $\pm$ 3.8	-0.3 – -0.9	-0.6 $\pm$ 0.4
MK-8	–	–	–	–	–	–	–	–	–	–
MK-9	–	5.8	–	–	–	–	–	5.4	–	0.4
MK-9(H ₈)	–	0.6	–	–	–	–	–	0.8	–	-0.2

SD, Standard Deviation

Table 5. Studies of simultaneous determination of grazing and viral lysis rates in freshwater systems

Method	Grazing % potential production	Lysis % potential production	Location and water layer	Trophic status	Reference
TC & FVIC	81.8 - 108.0	7.7 - 27.8	Lake Plußsee, Epilimnion	Eutrophic	Weinbauer & Hofle 1998
	2.9 - 27.6	19.6 - 46.8	Lake Plußsee, Metalimnion	–	Weinbauer & Hofle 1998
	5.0 - 8.9	38.4 - 97.3	Lake Plußsee, Hypolimnion	–	Weinbauer & Hofle 1998
FLB & FVIC	50	25	Řimov Reservoir, Surface water	Meso-Eutrophic	Šimek et al. 2006
TC & FVIC	10.3	6.4	Lake Pavin, Epilimnion	Oligomesotrophic	Bettarel et al. 2003
	8.4	15.6	Lake Pavin, Epilimnion	–	Bettarel et al. 2003
Tritiated thymidine labeled bacteria	58.2	26.3	Lake Tanganyika, Upper water (Wet season)	Oligotrophic	Pirlot et al. 2007
	88.5	39.6	Lake Tanganyika, Upper water (Dry season)	–	Pirlot et al. 2007
CD & FVIC	78*	7.7*	Lake Erie, Surface and Deep water	Eutrophic	Gobler et al. 2008
FLMB & viral dilution & FVIC	18 - 63*	35 - 60*	Lake Bourget, Surface layer	Mesotrophic	Jacquet et al. 2005
MD	45.9	100.5	Lake Loosdrecht, Surface layer	Eutrophic	Tijdens et al. 2008
MD	37.3-76.5	ND	Lake Geneva, Surface layer	Mesotrophic	Personnic et al. 2009
	18.2 - 56.8	ND	Lake Bourget, Surface layer	Mesotrophic	Personnic et al. 2009
	5.3	ND	Lake Annecy, Surface layer	Oligotrophic	Personnic et al. 2009
MD	37.5 – 71.0	56.2	Lake Biwa, Epilimnion	Mesotrophic	This study

TC, theoretical calculation; FVIC, frequency of visible infected cells; FLB, fluorescence labeled bacteria; CD, conventional dilution method; FLMB, fluorescence labeled micro-beads; MD, modified dilution method

*Grazing or lysis % standing stock d⁻¹

ND, not determined

1

2 **Figure legends**

3 **Fig. 1.** RQs concentration (A) and composition (B) of 100% samples at the beginning
4 experiments.

5 **Fig. 2.** Changes in growth (μ), grazing mortality (M_g), lysis mortality (M_v), and total
6 mortality (M_{g+v}) rates of total bacteria (RQ) and major bacterial groups. An asterisk
7 indicates that the rate is statistically meaningful (see Table 2).

8 **Fig. 3.** Changes in carbon biomass (CB), production (CP), grazing loss (GL), lysis loss
9 (LL) and total losses (TL) of total bacteria (RQ) and major bacterial groups. An asterisk
10 indicates that the rate is statistically meaningful (see Table 2).

11

12 **Appendix**

13 **Fig. S1.** Regression analyses of dilution experiments to estimate growth and mortality
14 raty

Takasu et al. Fig. 1

Takasu et al. Fig. 2

Takasu et al. Fig. 3

Takasu et al. Fig. S1

Jun. 2012

Jul. 2012

Takasu et al. Fig. S1 continued.