TITLE:

Seed dormancy and germination traits of an endangered aquatic plant species, *Euryale ferox* Salisb. (Nymphaeaceae)

AUTHOR(S):

Imanishi, Ayumi; Imanishi, Junichi

CITATION:

Imanishi, Ayumi ...[et al]. Seed dormancy and germination traits of an endangered aquatic plant species, *Euryale ferox* Salisb. (Nymphaeaceae). *Aquatic Botany* 2014, 119: 80-83

ISSUE DATE:

2014-10

URL:

<http://hdl.handle.net/2433/191026>

RIGHT:

© 2014 Elsevier B.V.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

1 Seed dormancy and germination traits of an endangered aquatic plant species, *Euryale ferox*

2 Salisb. (Nymphaeaceae)

3

4 Ayumi Imanishi^a and Junichi Imanishi^b

5

6 ^aFaculty of Applied Sociology, Kinki University, 228-3 Shinkamikosaka, Higashiosaka, Osaka

7 577-0813, Japan

8 ^bGraduate School of Global Environmental Studies, Kyoto University, Oiwake-cho, Kitashirakawa,

9 Sakyo-ku, Kyoto 606-8502, Japan

10

11 E-mail addresses: AI, ayumi.imanishi@a.email.ne.jp; JI, imanishi.junichi.6c@kyoto-u.ac.jp

12

13 Corresponding author: Ayumi Imanishi

14 Phone number: +81-6-4307-4168

15

16 **ABSTRACT**

17 Populations of *Euryale ferox* Salisb. have declined throughout its global range because of habitat

18 loss and degradation. The present study aimed to evaluate the influence of storage temperature (4

and 20 °C), storage period (0, 90, 180, 270, and 360 days), light condition (light and darkness), germination temperature (constant 10, 15, 20, 25, and 30 °C), and seed size (two classes from 0.5 cm to less than 1.2 cm, and from 1.2 cm to less than 1.6 cm) on germination of *E. ferox* seeds. Our results indicated that seeds were dormant when shed and 4 °C stratification promoted germination more effectively than 20 °C. After stratification at 4 °C, the germination frequency of the larger seeds peaked at 90 days' stratification, whereas the smaller seeds maintained high germination frequencies up to 180 days' stratification. The different germination responses between smaller and larger seeds may reflect a difference in the rate of induction of dormancy in the annual dormancy cycle. Seeds germinated in both light and darkness, which demonstrated that light is not involved in the regulation of *E. ferox* seed germination. The optimal temperature for germination was 25 °C. Light condition × germination temperature interaction caused significantly higher germination frequency at 30 °C in light than in darkness, and the opposite trend at 15 °C.

Keywords: Nymphaeaceae, vulnerable, cold stratification, induction of dormancy, interaction of light and temperature

1. Introduction

Euryale ferox Salisb. (Nymphaeaceae) is a prickly annual aquatic herb with gigantic floating leaves 0.3–1.5 m in diameter. It is distributed from northwestern India to Japan and inhabits meso- and eutrophic water bodies such as lakes, ponds, reservoirs and rivers. In Japan, the species' distribution ranges from Kyushu to the northern part of Honshu, but habitat loss caused by drainage or land reclamation and water pollution has led to population decline and the species is classified as “vulnerable” in the Red List of Threatened Plants of Japan (Ministry of the Environment of Japan, 2012). Moreover, its populations have declined throughout its global range (Schneider et al., 2003).

It is well known that natural populations of *E. ferox* are subject to considerable annual variation in number of individuals (Miyashita, 1983; Kume, 1987). The species' seeds are believed capable of remaining dormant, even over several decades, when the external environment is unsuitable for germination (Wakita, 1959; Kadono, 1983; Ohtaki, 1987). This trait may influence annual variation in population sizes. However, seed dormancy and the germination characteristics of this species are poorly understood.

Seed dormancy is an adaptive mechanism to ensure that germination takes place in a suitable location and in suitable conditions (e.g., Baskin and Baskin, 2001; Fenner and Thompson, 2005).

Temperature is regarded as the most important factor influencing seed dormancy (Bouwmeester and

Karssen, 1992, 1993). Kumaki and Minami (1973) reported that about 30% of *E. ferox* seeds germinated after prechilling at 2–3 °C for 1 month, whereas at constant room temperature no seeds germinated. This result indicated that cold stratification breaks dormancy of the seeds. However, since the optimum period of stratification varies among species (Baskin and Baskin, 2001), and thus it is necessary to determine the optimal period of cold stratification for maximum germination and the influence of extended seed preservation on germination.

After release of seed dormancy, environmental factors such as temperature and light promote germination (Benvenuti et al., 2001; Penfield et al., 2005; Jha et al., 2010). A systematic quantitative study has not been conducted on the effect of temperature and light on germination of *E. ferox* seeds, but from field observations it is estimated that the seeds germinate between 20 and 25 °C and do not require light for germination (Okada, 1935; Wakita, 1959).

In addition to temperature and light, seed size may play an important role in seed germination for some species (Cideciyan and Malloch, 1982; Zammit and Zedler, 1990; Leverett and Jolls, 2013). Seeds of *E. ferox* are about 1 cm in diameter and their size varies among populations (Okada, 1928; Miyashita, 1983), among individuals in a population (Miyashita 1983; Hashimoto 1986), and even within the same individual (Hagiwara 1993). The phenomenon that small seeds of *E. ferox* require a shorter after-ripening period than large seeds has been observed (Okada 1935; Wakita 1959). However, a quantitative study of the relationship between seed size and germination has not been

conducted.

The aim of the present research was to quantitatively verify seed dormancy and germination traits of *E. ferox* focusing on the influence of seed storage period, storage temperature, light condition, germination temperature, and seed size.

2. Materials and methods

2.1 Seed collection and storage

Fresh fruits and floating fresh seeds with arils were collected from late October to early November 2009 in the Hiranosawa pond, Kameoka, western Japan (35°04'03" N, 135°33'55" E). Hiranosawa pond consists of three small ponds with a total area of about 10 ha. It has been used as an agricultural reservoir. Genetic diversity of *E. ferox* is low within and among populations in Japan and the genotype of *E. ferox* in the Hiranosawa pond is the most widely distributed one in western Japan (the authors' unpublished data). Thus we considered that our samples are representative of the genetic diversity in this region. Seeds were separated from the fruits and the arils were removed in the laboratory. Small green- or skin-colored immature seeds were excluded.

The maximum diameter of collected seeds was measured with a digital caliper and seeds were divided into five size groups: from 0.5 cm to less than 1.0 cm; from 1.0 cm to less than 1.2 cm; from

1 1.2 cm to less than 1.4 cm; from 1.4 cm to less than 1.6 cm; and greater than 1.6 cm. The number of
2 seeds in each size group was about 1,800 seeds, about 6,500 seeds, about 4,400 seeds, about 1,000
3 seeds, and 19 seeds, respectively.

4 Seeds of each size group were stored in separate plastic containers with four replicates wrapped
5 in aluminum foil in order to avoid seed deterioration caused by fungal infection. The containers were
6 filled with water and placed at either a constant low temperature (4 °C) to simulate the winter water
7 temperature or a constant warm temperature (20 °C) to simulate the early summer water temperature
8 in a small pond (Shimomura et al., 2010). Water was replaced regularly in complete darkness under a
9 green safelight.

10

11 2.2 Germination tests

12 Germination tests were performed in light and darkness at constant temperatures of 10, 15, 20,
13 25, and 30 °C after 0, 90, 180, 270, and 360 days of stratification at 4 and 20 °C. Four replicates of
14 30 seeds each were placed in plastic containers and the seeds were submerged with water. The
15 containers were placed in an incubator (TG-280CCFL-5LD, NKsystem, Osaka, Japan). On the basis
16 of the number of collected seeds in each size group, four (from 0.5 cm to less than 1.0 cm), 14 (from
17 1.0 cm to less than 1.2 cm), 10 (from 1.2 cm to less than 1.4 cm) and two seeds (from 1.4 cm to less
18 than 1.6 cm) per container were sown, respectively. Seeds of greater than 1.6 cm diameter were not

1 used in germination tests because of the limited number of seeds available. We defined seeds from
2 0.5 to less than 1.2 cm diameter as small seeds and those from 1.2 to less than 1.6 cm diameter as
3 large seeds.

4 In the light treatments, seeds were exposed to warm white fluorescent light providing
5 photosynthetically active radiation of $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ for 12 h day^{-1} . For the darkness treatments, all
6 handling of containers was conducted in complete darkness under a green safelight and the
7 containers were wrapped with aluminum foil after seeding.

8 For the light treatments, germination counts were made at intervals of 2–3 days for 60 days. For
9 the darkness treatments, the containers were unwrapped at the end of the germination period (60
10 days), and the number of germinated seeds was counted. Seeds were recorded as germinated when
11 the protruding radicle was ≥ 1 mm in length as defined by Okada (1935).

12

13 2.3 Data analysis

14 To examine the effects of light condition, germination temperature, and seed size on
15 germination, we analyzed germination of seeds stratified at 4°C for 90 days by constructing a
16 generalized linear model with a binomial distribution with number of germinated seeds as the
17 response variable and three factors (light condition, germination temperature excluding 10°C , and
18 seed size), three interactions of combinations of two factors (light condition $\times$ germination

temperature excluding 10 °C, light condition × seed size, and germination temperature excluding 10 °C × seed size), and one interaction of the three factors (light condition × germination temperature excluding 10 °C × seed size) as explanation variables. We adopted the best model by stepwise selection. For the statistical analysis, we used the glm and stepAIC functions in R ver. 3.1.0 (R Core Team, 2014) and set the significance level as $p < 0.05$.

3. Results

Few seeds germinated immediately after harvest (no stratification) (Fig. 1). Regarding the seeds stratified at 4 °C, small seeds maintained high germination frequencies until at least 180 days (Fig. 1A and B). The germination frequency of large seeds was highest at 90 days and decreased at 180 days regardless of light condition (Fig. 1C and D).

With regard to the seeds stratified at 4 °C that germinated in light, the median germination time (i.e., the number of days until at least 50% germination frequency was attained in all of the four replicates within 60 days) was evaluated for the seven cases listed in Table 1. The shortest median germination time was 5.5 ± 0.5 days (mean ± SD) for small seeds stratified for 180 days and germinated at 25 °C.

Regarding the seeds stratified at 20 °C for 90 days, mean germination frequencies in light were

0, 0, 1.7 ± 1.9 , 28.3 ± 7.9 , and 29.2 ± 10.7 % (mean $\pm$ SD) at 10, 15, 20, 25, and 30 °C, respectively, and in darkness were 0, 0, 12.5 ± 7.4 , 47.5 ± 18.9 , and 18.3 ± 8.8 % at 10, 15, 20, 25, and 30 °C, respectively. After 180, 270 or 360 days of stratification at 20 °C, few or no seeds germinated.

The generalized linear model focusing on 4 °C stratification for 90 days revealed significant effects of light condition, germination temperature, seed size, light condition $\times$ germination temperature interaction, and germination temperature $\times$ seed size interaction (Table 2). The light condition $\times$ germination temperature interaction was significant owing to higher germination frequency at 30 °C in light (mean $\pm$ SD; 44.2 ± 6.9 %) than that in darkness (18.3 ± 12.3 %), and higher germination frequency at 15 °C in darkness (53.3 ± 9.8 %) than that in light (31.7 ± 9.8 %). The germination temperature $\times$ seed size interaction was significant owing to higher germination frequency of small seeds at 30 °C (37.5 ± 8.0 %) than that of large seeds (21.9 ± 12.3 %), and higher germination frequency of large seeds at 15 °C (49.0 ± 7.4 %) than that of small seeds (38.2 ± 4.6 %).

4. Discussion

Our results indicated that seeds of *E. ferox* are in a dormant state when shed and stratification for several months can break the dormancy (Fig. 1). Dormancy release was possible with 4 and 20 °C stratification, but was more strongly promoted by 4 °C stratification. With regard to seeds

1 stratified at 4 °C, the germination frequency of large seeds peaked at 90 days (Fig. 1C and D),
2 whereas small seeds maintained high germination frequencies until at least 180 days (Fig. 1A and B).
3 It was hypothesized that non-germinated seeds did not lose viability but dormancy was re-induced
4 because the embryos were white and hard when dissected after completion of the germination tests,
5 although the induction of dormancy was not investigated. The different germination responses
6 between small and large seeds may reflect a difference in the rate of induction of dormancy in annual
7 dormancy cycle.

8 Most flowers of *E. ferox* are cleistogamous, but chasmogamous flowers are also produced
9 (Okada and Otaya, 1930). In many species, germination responses differ between seeds from
10 cleistogamous and chasmogamous flowers (e.g. Weiss, 1980; Trapp and Hendrix, 1988; Ferreira and
11 Reinhardt, 1999). Kadono and Schneider (1987) reported that seeds of *E. ferox* from chasmogamous
12 flowers are slightly larger than those from cleistogamous flowers because fruits from chasmogamous
13 flowers contain significantly fewer seeds than cleistogamous flowers and seed size is inversely
14 correlated to the total number of seeds in each fruit. Although more research is required, it is
15 hypothesized that *E. ferox* may produce many small seeds, in which the rate of induction of
16 dormancy is relatively slow, from cleistogamous flowers and fewer large seeds that show relatively
17 rapid induction of dormancy from chasmogamous flowers.

18 Seeds of *E. ferox* germinated in both light and darkness (Fig. 1), which indicated that light is

1 unimportant for regulation of germination in this species. It is believed that *E. ferox* can form a
2 persistent seed bank (Wakita, 1959; Kadono, 1983; Otaki, 1987). A light requirement for germination
3 is a common feature among species that form persistent seed banks (Pons 2000). However,
4 germination in large-seeded species might be expected to be insensitive to light because seedlings
5 from large seeds can emerge successfully from a much greater depth that light cannot penetrate
6 (Pons, 2000; Milberg et al., 2000; Pearson et al., 2002; Fenner and Thompson, 2005). Seeds of *E.*
7 *ferox* are large (about 1 cm in diameter), which may be a factor in their not requiring light for
8 germination.

9 The present study quantitatively verified that the optimal temperature for germination was
10 25 °C because the median germination times at 25 °C were shorter than those at 20 °C (Table 1), and
11 the germination frequency was high at both 20 and 25 °C (Fig. 1). In some species, light interacts
12 with temperature to regulate seed germination (Baskin and Baskin, 2001). In the present study, the
13 generalized linear model focusing on 4 °C stratification of 90 days revealed that light condition ×
14 germination temperature interaction significantly affected germination (Table 2). The significant
15 interaction was due to higher germination frequencies at 30 °C in light than darkness and higher
16 germination frequencies at 15 °C in darkness than in light. The interaction between light and
17 temperature is considered to be beneficial for germination of *E. ferox*, as the seeds often sink to the
18 bottom of water bodies. If the temperature around the seed is low even though light reaches the

- 1 bottom of the water body, germination will be inhibited, which enables avoidance of an unsuitable
- 2 external environment for growth. On the other hand, germination could occur at a low temperature
- 3 when the seeds are buried at the bottom of a water body and do not receive light, which permits
- 4 increased opportunity for germination close to the level that would be observed if the seeds were not
- 5 covered by soil.

- 1 **Acknowledgments:** We are grateful to Mr. Ryuzo Hayashi in Kameoka city for collecting seeds.
- 2 This work was supported in part by the Osaka University Program for the Support of Networking
- 3 among Present and Future Researchers.

1 **References**

2

3 Baskin, C.C., Baskin, J.M., 2001. Seeds: Ecology, Biogeography, and Evolution of Dormancy and
4 Germination. Academic Press, San Diego.

5 Benvenuti, S., Macchia, M., Miele, S., 2001. Light, temperature and burial depth effects on *Rumex*
6 *obtusifolius* seed germination and emergence. Weed Res. 41, 177-186.

7 Bouwmeester, H.J., Karssen, C.M., 1992. The dual role of temperature in the regulation of the
8 seasonal changes in dormancy and germination of seeds of *Polygonum persicaria* L. Oecologia
9 90, 88-94.

10 Bouwmeester, H.J., Karssen, C.M., 1993. Annual changes in dormancy and germination in seeds of
11 *Sisymbrium officinale* (L.) Scop. New Phytol. 124, 179-191.

12 Cideciyan, M.A., Malloch, A.J.C., 1982. Effects of seed size on the germination, growth and
13 competitive ability of *Rumex crispus* and *Rumex obtusifolius*. Jour. Ecol. 70, 227-232.

14 Fenner, M., Thompson, K., 2005. The Ecology of Seeds. Cambridge University Press, Cambridge.

15 Ferreira, M.I., Reinhardt, C.F., 1999. The role of temperature in the germination of subterranean and
16 aerial seeds of *Commelina benghalensis* L. S. Afr. J. Plant Soil 16, 165-168.

17 Hagiwara, T., 1993. *Euryale ferox* Salisb. Bull. Water Plant Soc., Jpn 51, 32-34 (in Japanese).

18 Hashimoto, T., 1986. *Euryale ferox* Salisb. of Chizuka-pond in Fukuyama city. Bull. Water Plant

- 1 Soc., Jpn 26, 6-11 (in Japanese).
- 2 Jha, P., Norsworthy, J.K., Riley, M.B., Bridges Jr, W., 2010. Annual changes in temperature and light
- 3 requirements for germination of Palmer amaranth (*Amaranthus palmeri*) seeds retrieved from
- 4 soil. Weed Sci. 58, 426-432.
- 5 Kadono, Y., 1983. Natural history of *Euryale ferox* Salisb. Nat. Stud. 29, 63-66 (in Japanese).
- 6 Kadono, Y., Schneider, E.L., 1987. The life history of *Euryale ferox* Salisb. in south-western Japan
- 7 with special reference to reproductive ecology. Plant Sp. Biol. 2, 109-115.
- 8 Kumaki, Y., Minami, Y., 1973. Seed germination of “Onibasus” *Euryale ferox* Salisb. (II). Bull. Fac.
- 9 Educ., Kanazawa Univ., Nat. Sci. 22, 71-78.
- 10 Kume, O., 1987. Growth situation of *Euryale ferox* Salisb. in Kagawa prefecture I. Bull. Water Plant
- 11 Soc., Jpn 27, 16-19 (in Japanese).
- 12 Leverett, L.D., Jolls, C.L., 2013. Cryptic seed heteromorphism in *Packera tomentosa* (Asteraceae):
- 13 differences in mass and germination. Plant Spec. Biol.. doi: 10.1111/1442-1984.12011
- 14 Maeda, O., Nakamichi, T., Takahashi, G., 1983. *Euryale ferox* Salisb. in Shishitsuka-Oike. Tsukuba
- 15 no kankyo kenkyu 7, 80-85 (in Japanese).
- 16 Milberg, P., Andersson, L., Thompson, K., 2000. Large-seeded species are less dependent on light
- 17 for germination than small-seeded ones. Seed Sci. Res. 10, 99-104.
- 18 Ministry of the Environment of Japan, 2012. Red List of Threatened Plants of Japan,

- 1 http://www.biodic.go.jp/english/rdb/rdb_f.html [accessed 22 July 2013].
- 2 Miyashita, Y., 1983. *Euryale ferox* Salisb. of Sakata lagoon in Niigata prefecture. Bull. Water Plant
- 3 Soc., Jpn 11, 4-6 (in Japanese).
- 4 Otaki, S., 1987. *Euryale ferox* Salisb. in Japan. Nihon no seibutsu 1(4), 48-55. (in Japanese)
- 5 Okada, Y., 1928. Study of *Euryale ferox* Salisb. I. On the size of leaves, fruits, etc., with some
- 6 remarks on the mode of expansion of the leaf blade. Sci. Rep. Tohoku Imp. Univ., Ser. 4, 3,
- 7 271-278.
- 8 Okada, Y., 1935. Long-term dormancy of *Euryale ferox* Salisb. seeds. Seitaigakuteki kenkyu 1,
- 9 14-22 (in Japanese).
- 10 Okada, Y., Otake, T., 1930. Study of *Euryale ferox* Salisb. VI. Cleistogamous versus chasmogamous
- 11 flowers. Bot. Mag. (Tokyo) 44, 369-373.
- 12 Pearson, T.R.H., Burslem, D.F.R.P., Mullins, C.E., Dalling, J.W., 2002. Germination ecology of
- 13 neotropical pioneers: interacting effects of environmental conditions and seed size. Ecology 83,
- 14 2798-2807.
- 15 Penfield, S., Josse, E.-M., Kannangara, R., Gilday, A.D., Halliday, K.J., Graham, I.A., 2005. Cold
- 16 and light control seed germination through the bHLH transcription factor SPATULA. Curr. Biol.
- 17 15, 1998-2006.
- 18 Pons, T.L., 2000. Seed responses to light. In: Fenner, M. (Eds.), Seeds: The Ecology of Regeneration

- 1 in Plant Communities (2nd edition). CAB International, Wallingford, pp. 237-260.
- 2 R Core Team, 2014. R: A language and environment for statistical computing. R Foundation for
- 3 Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- 4 Schneider, E.L., Tucker, S.C., Williamson, P.S., 2003. Floral development in the Nymphaeales. Int. J.
- 5 Plant Sci. 164, S279-S292.
- 6 Shimomura, H., Kamakura, T., Masuda, H., 2010. Seasonal variations of water quality and the
- 7 sediment quality of Benten pond in Joetsu University of Education: as one of the basic data to
- 8 utilize a small reservoir for an environmental education. Bull. Joetsu Univ. Educ. 29, 249-255
- 9 (in Japanese with English abstract).
- 10 Trapp, E.J., Hendrix, S.D., 1988. Consequences of a mixed reproductive system in the hog peanut,
- 11 *Amphicarpaea bracteata*, (Fabaceae). Oecologia 75, 285-290.
- 12 Wakita, H., 1959. Study on the fresh water plants in Nagoya and north-eastern part of Owari
- 13 province; including ecological study of *Euryale ferox* Salisbury. Chubu nihon shizen kagaku
- 14 chosadan hokoku 3, 5-7 (in Japanese).
- 15 Weiss, P.W., 1980. Germination, reproduction and interference in the amphicarpic annual *Emex*
- 16 *spinosa* (L.) Campd. Oecologia 45, 244-251.
- 17 Zammit, C., Zedler, P., 1990. Seed yield, seed size and germination behavior in the annual *Pogogyne*
- 18 *abramsii*. Oecologia 84, 24-28.
- 19

1 **Figure captions**

2

3 **Fig. 1.** Germination frequency of *E. ferox* seeds stratified at 4 °C and germinated at five constant
4 temperatures ranging from 10 to 30 °C. (A) Small seeds germinated in light, (B) small seeds
5 germinated in darkness, (C) large seeds germinated in light, and (D) large seeds germinated in
6 darkness. Symbols and error bars represent the mean $\pm$ SD ($n = 4$).

Table 1 Median germination time for *E. ferox* seeds, which is the days required for the germination frequency of all four replicates to reach at least 50% within the 60-day experimental period.

Germination temperature	Storage periods			
	90 days		180 days	270 days
	small seeds	large seeds	small seeds	small seeds
20 °C	20.8 ± 9.5	–	11.5 ± 6.1	–
25 °C	10.8 ± 2.6	8.5 ± 1.5	5.5 ± 0.5	14.5 ± 1.7
30 °C	–	–	8.0 ± 1.6	–

Values are the mean ± SD

"-" indicates that germination frequency of all of the four replicates did not reach 50% within the 60-day experimental period.

Table 2 Effects of germination temperature, light condition, and seed size on germination of *E. ferox* seeds

	Estimate	SE	z-value	P
Germination temperature	-0.12	0.02	-5.087	< 0.001
Light condition (in light)	-2.83	0.56	-5.081	< 0.001
Seed size (small)	-1.17	0.57	-2.048	0.04
Germination temperature × Light condition (in light)	0.12	0.02	5.19	< 0.001
Germination temperature × Seed size (small)	0.05	0.02	2.14	0.03
Intercept	2.86	0.54	5.29	< 0.001

AIC (selected model/null model) : 437.9 / 441.1

Models that have the highest power of explanation are shown.