TITLE:

Notes on the Parthenopidae (Crustacea: Decapoda: Brachyura) collection of the Natural History Museum, Florence University, Italy, with the quotation of the world's largest specimen of parthenopid crab

AUTHOR(S):

NAGAI, Shinji; INNOCENTI, Gianna

CITATION:

NAGAI, Shinji ...[et al]. Notes on the Parthenopidae (Crustacea: Decapoda: Brachyura) collection of the Natural History Museum, Florence University, Italy, with the quotation of the world's largest specimen of parthenopid crab. Publications of the Seto Marine Biological Laboratory 2015, 43: 30-38

ISSUE DATE:

2015-09-15

URL:

<http://hdl.handle.net/2433/193237>

RIGHT:

Notes on the Parthenopidae (Crustacea: Decapoda: Brachyura) collection of the Natural History Museum, Florence University, Italy, with the quotation of the world's largest specimen of parthenopid crab

SHINJI NAGAI¹ and GIANNA INNOCENTI^{2*}

¹5045-1 Fukushima, Kiso-Machi, Kiso-Gun, Nagano, 397-0001 Japan

²Sezione di Zoologia “La Specola”, Museo di Storia Naturale dell’Università degli Studi di Firenze,
via Romana 17, 50125 Firenze, Italy

*Corresponding author: E-mail: gianna.innocenti@unifi.it

Abstract New information on the geographical distribution of seven species of Parthenopidae is given. The fifth record of *Daldorfia spinosissima* (A. Milne-Edwards, 1862) is reported, presently exhibited in the Crustacean room of the Natural History Museum of Florence University, Italy.

Key words: Parthenopidae, systematics, museum collection

Introduction

The senior author visited the Natural History Museum of the University of Florence (MZUF), Italy, in June, 1991 for a research on the Coleoptera Lucanidae type material. During his visit, he noticed a very large parthenopid crab in the Crustacean exhibition room. Subsequently, in 2010 he contacted the MZUF Crustacean curator, in order to get information and photographs of the gigantic specimen. Recently, the senior author received several papers on Parthenopidae from Dr Tan Swee Hee, which helped with the identification of this large crab.

The following paper gives information on the parthenopid collection in the Museum from the Indo-Pacific Ocean, and adds new data on Parthenopidae distribution. An asterisk (*) marks new localities for the species. The Parthenopidae crabs listed in the present paper are preserved in the Natural History Museum, Florence, Italy and the number after the acronym MZUF indicates their collection number.

Systematics

Family **Parthenopidae** MacLeay, 1838

Subfamily **Parthenopinae** MacLeay, 1838

Genus ***Cryptopodia*** H. Milne-Edwards, 1834

This genus is distributed from the Indian Ocean to the central area of the Pacific Ocean, and thirteen species are known. The carapace is flatten at the edges, fan shaped at the posterior half, wide and triangular shaped at the anterior half. Legs are slender, short, and hidden under the carapace.

Cryptopodia pan Laurie, 1906
(Figure 1A)

Specimens examined: 1 male (MZUF 4287), carapace width: 22.5 mm, Sri Lanka, Trincomalee, -18-27 m, II.1889, leg. K. Fristedt; 1 male (MZUF 4286), carapace width: 15.4 mm, Sri Lanka, Trincomalee, -18-27 m, II.1889, leg. K. Fristedt.

Distribution: from Seychelles to New Caledonia.

Remarks: This species is close to *C. collifer* Flipse, 1930, recorded from Thailand, Philippines and Fiji, but it is distinguishable by the following differences: frontal projection strongly produced distinctly forward and frontal corners somehow angulated at the epibranchial margins. These two male specimens are probably the oldest collected ones amongst all parthenopid specimens.

Genus *Rhinolambrus* A. Milne-Edwards, 1878

There are thirteen species in this genus, and they are distributed from the Indian Ocean to the central area of the Pacific Ocean.

Rhinolambrus pelagicus (Rüppell, 1830)
(Figure 1B)

Specimens examined: 1 male (MZUF 4280), carapace width: 14.4 mm, Somalia, Gesira, sheltered pools, X.1986, leg. M. Vannini; 1 male (MZUF 4282), carapace width: 12.0 mm, Kenya, N of Mombasa, Twiga, XI.1991, leg. M. Vannini; 1 male (MZUF 4283), carapace width: 15.2 mm, Kenya, N of Mombasa, Malindi, XI.1999, leg. S. Cianfanelli; 1 male (MZUF 4281), carapace width: 15.3 mm, Tanzania, S of Dar es Salaam, Ras Dege, X.2005, leg. S. Cannicci; 1 female (MZUF 4285), carapace width: 22.0 mm, India, Paumben, -2-5 m, IV.1889, leg. K. Fristedt; 4 juvenile males, carapace width, respectively: 8.3 mm, 7.4 mm, 5.9 mm, 5.7 mm, 1 juvenile female, carapace width: 6.8 mm (MZUF 4291), Sri Lanka, Trincomalee, -7-11 m, II.1889, leg. K. Fristedt; 1 male (MZUF 4284), carapace width: 31.3 mm, Samoa, Upola, IV.1889, purchased by Pohl.

Distribution: *Somalia, *Kenya, *Tanzania, Madagascar, Red Sea, Sudan, Djibouti, India, Sri Lanka, Andaman Is., Thailand, Indonesia, Malaysia, Singapore, Philippines, Taiwan, Japan, Korea, Caroline Is., Palau, Australia, New Caledonia, Fiji, Samoa.

Remarks: The lectotype of this species was designated on a specimen collected in the Red Sea, and the species is distributed from East off Madagascar through the central area of the Pacific Ocean. With this report, the following new localities are added: Somalia, Kenya and Tanzania.

Rhinolambrus turriger (White, 1847)
(Figure 1C)

Specimen examined: 1 male (MZUF 4289), carapace width: 11.9 mm, Sri Lanka, Trincomalee, -18-27 m, II.1889, leg. K. Fristedt.

Distribution: Mayotte, Seychelles, Maldives, Laccadives, Sri Lanka, Andaman Islands, Philippine, China, Indonesia, Australia and Palau.

Remarks: This specimen is very similar to *Rhinolambrus sisimanensis* (Sérene & Umali, 1972), but differently from this species, the mesobranchial spines are very short (Tan, 2004).

Genus *Aulacolambrus* Paul'son, 1875

This genus consists of eight species widely distributed from the Indian Ocean to the western area of the Pacific Ocean.

Aulacolambrus hoplonotus (Adams & White, 1849)
(Figure 1D)

Specimen examined: 1 female (MZUF 4288), carapace width: 19.1 mm, Sri Lanka, Trincomalee, -7-11 m, II.1889, leg. K. Fristedt.

Distribution: Seychelles, Sri Lanka, Japan, Taiwan, Guam, Hawaiian Island, New Caledonia, Loyalty Island and Fiji.

Genus *Enoplolambrus* A. Milne- Edwards, 1878

This genus consists of seven species distributed from the Indian Ocean to the western Pacific Ocean.

Enoplolambrus cf. *carenatus* (H. Milne Edwards, 1834)
(Figure 1E)

Specimen examined: 1 male (MZUF 4290), carapace width: 16.0 mm, Sri Lanka, Trincomalee, -7-11 m, II.1889, leg. K. Fristedt.

Distribution: Sri Lanka.

Remarks: This specimen is very similar to *Lambrus holdswarthi* Miers, 1879 and *Lambrus* (*Platylambrus*) *carenatus* var. *alcocki* Laurie, 1906, and both the species were originally collected from Sri Lanka. Those two species are presently synonyms of *Enoplolambrus carenatus* (H. Milne-Edwards, 1834).

Subfamily **Daldorfiinae** Ng & Rodríguez, 1986
Genus *Daldorfia* Rathbun, 1904

This genus consists of twelve species, most of which are distributed in the Indian Ocean through the western area of the Pacific Ocean. One species, *D. bouvieri* (A. Milne Edwards 1869), is known from the eastern part of the Atlantic Ocean and another one, *D. glasselli* (Garth, 1958), is distributed in the western part of the Atlantic Ocean. Most species are very robust and the carapace of some species are very irregular, rugose or spinose.

This genus includes the largest species of the family Parthenopidae.

Daldorfia spinosissima (A. Milne-Edwards, 1862)
(Figure 2)

Specimen examined: 1 male (MZUF 4292), carapace width: 200.0 mm, Mauritius Is., IV.1875, purchased by L. De Greaux; both fourth legs are missing.

Distribution: Yemen (Socotra Is.), Réunion Is., *Mauritius Is.

Remarks: This specimen was identified by Adolfo Targioni Tozzetti as *Parthenope horrida* and subsequently updated in 1997 by the junior author as *Daldorfia horrida* (Linnaeus, 1758), but, after a more careful examination of the specimen, we identified it as *Daldorfia spinosissima* (A. Milne-Edwards, 1862), according to the revision of Daldorfiinae by Tan & Ng (2007). The type species is a female from Réunion (formerly Bourbon Is.). After that, a pair of specimens has been reported from the Bengal bay by Alcock

(1895), and two male specimens reported respectively from Socotra Is. and Réunion Is. by Tan & Ng (2007). At date, only three males (ranging from 151.3 and 167.2 mm) and a female (165.9 mm) are known. This species was reported from Japan (Sakai, 1976), Bay of Bengal (Alcock, 1895), Philippines (Serène & Umali, 1972) and Indonesia, Ambon (Flipse, 1930), but all these reports are to be considered as large specimens of *D. triangularis* Sakai 1974, as discussed by Tan & Ng (2007).

This specimen is the oldest and largest record for males (and the largest specimen in the Parthenopidae family), moreover it is the first report from Mauritius Is.

Acknowledgments

We wish to thank Dr. Luca Bartolozzi (MZUF) and Anna Kraczyna for their collaboration. A special thank to Dr Tan Swee Hee, NUS, Singapore, who helped in giving his papers on Parthenopidae and for his useful remarks in a previous version of the manuscript. Saulo Bambi (MZUF) is the author of the photographs of the specimens.

Literature cited

- Adams, A. and White, A. 1849. Crustacea. In, Adams, A. (ed.) The Zoology of the voyage of H.M.S. *Samarang*; under the command of Captain Sir Edward Belcher, C.B., F.R.A.S., F.G.S., during the years 1843–1846. London: Reeve, Bentham and Reeve, pp. 33–66.
- Alcock, A. 1895. Materials for a carcinological fauna of India. No. 1. The Brachyura Oxyrhyncha. Journal of the Asiatic Society of Bengal 64, 157–291.
- Flipse, H.J. 1930. Decapoda F. (Decapoda Brachyura continued) Oxyrhyncha: Parthenopidae. Siboga Expédition 39(C2), 1–96.
- Garth, J.S. 1958. Brachyura of the Pacific coast of America: Oxyrhyncha. Allan Hancock Pacific Expeditions 21(1), 1–499.
- Laurie, R.D. 1906. Report on the Brachyura collected by Professor Herdman, at Ceylon, in 1902. In, Herdman, W.A. (ed.) Report to the Government of Ceylon on the Pearl Oyster Fisheries of the Gulf of Manaar with Supplementary Reports Upon the Marine Biology of Ceylon by Other Naturalists, Part 5, Suppl. Rep. 40, pp. 349–432.
- Linnaeus, C. 1758. Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis. (10 Edition) Vol. 1. Holmiae: Laurentii Salvii. iii, 824 pp.
- MacLeay, W.S. 1838. On the brachyurous decapod Crustacea brought from the Cape by Dr. Smith. In, Smith, A. (ed.) Illustrations of the Annulosa of South Africa; being a portion of the objects of natural history chiefly collected during an expedition into the interior of South Africa, under the direction of Dr. Andrew Smith, in the years 1834, 1835 and 1836; fitted out by “The Cape of Good Hope Association for Exploring Central Africa”. London: Smith, Elder, and Co., pp. 53–71.
- Miers, E.J. 1879. On the classification of the Maioid Crustacea or Oxyrhyncha, with a synopsis of the families, sub-families and genera. Journal of the Linnean Society of London, Zoology, 14, 634–673.
- Milne Edwards, H. 1834. Histoire naturelle des Crustacés, comprenant l’anatomie, la physiologie et la classification de ces animaux. Librairie Encyclopédique de Roret. Vol. 1, pp. 468.
- Milne-Edwards, A. 1862. Faune carcinologique de l’île de la Réunion. In, Maillard, L. (ed.) Notes sur l’île de la Réunion, Vol. Annexe F, Paris, pp. 1–16.
- Milne-Edwards, A. 1869. Description de quelques espèces nouvelles de Crustacés provenant du voyage de M. A. Bouvier aux îles du Cap Vert. Revue et Magasin de Zoologie, 1-13.

- Milne-Edwards, A. 1878. Études sur les xiphosures et les crustacés podophthalmes de la région mexicaine. In, Mission scientifique au Mexique et dans l'Amérique centrale. Recherches zoologiques pour servir à l'histoire de la faune de l'Amérique centrale et du Mexique, Cinquième partie, Paris, Imprimerie Nationale, Livraison 4, 121-184.
- Ng, P.K.L. and Rodríguez G. 1986. New records of *Mimilambrus wileyi* Williams, 1979 (Crustacea: Decapoda: Brachyura), with notes on the systematics of the Mimilambridae Williams, 1979 and Parthenopoidea MacLeay, 1838 sensu Guinot, 1978. Proceedings of the Biological Society of Washington, 99, 88-99.
- Paul'son, O. 1875. Studies on Crustacea of the Red Sea with notes regarding other seas. Part I. Podophthalmata and Edriophthalmata (Cumacea). [In Russian, English translation by the Israel Program for Scientific Translations, Jerusalem, 1961, Kiev, S.V. Kul'zhenko, pp. 144.
- Rathbun, M.J. 1904. Some changes in crustacean nomenclature. Proceedings of the Biological Society of Washington, 17, 169-172.
- Rathbun, M.J. 1916. New species of crabs of the families Inachidae and Parthenopidae. Proceedings of the United States National Museum, 50, 527-559.
- Rüppell, E. 1830. Beschreibung und Abbildung von 24 Arten kurzschwänzigen Krabben, also Beitrag zur Naturgeschichte des Rothen Meeres, Frankfurt am Main, H. L. Brönnner, 28 pp.
- Sakai, T. 1974. Notes from the carcinological fauna of Japan (V). Researches on Crustacea, 6, 86-102.
- Sakai, T. 1976. Crabs of Japan and the Adjacent Seas, Tokyo, Kodansha Ltd, 773 pp.
- Serène, R. and Umali A.F. 1970. The family Raninidae and other new and rare species of brachyuran decapods from the Philippines and adjacent regions. Philippine Journal of Science, 99(1-2), 21-105.
- Tan, S.H. 2004. A systematic revision of the Parthenopidae (Crustacea: Decapoda: Brachyura): 1-730. Unpublished Ph.D. Thesis, Department of Biological Sciences, National University of Singapore, available at: <http://scholarbank.nus.edu.sg/handle/10635/27686>
- Tan, S.H. and Ng P.K.L. 2007. Review of the subfamily Daldorfiinae Ng & Rodríguez, 1986 (Crustacea: Decapoda: Brachyura: Parthenopidae). The Raffles Bulletin of Zoology, Supplement 16, 121-167.
- White, A. (1847) Descriptions of new Crustacea from the Eastern Seas. Proceedings of the Zoological Society of London, 1847(XV): 56-58.

Received: 9 Dec 2014

Accepted: 25 Dec 2014

Published online: 20 Jan 2015

Figure 1. **A**, *Cryptopodia pan*, 1 male (MZUF 4287), carapace width: 22.5 mm, Sri Lanka, Trincomalee, -18-27 m, II.1889, leg. K. Fristedt. **B**, *Rhinolambrus pelagicus*, 1 male (MZUF 4282), carapace width: 12.0 mm, Kenya, N of Mombasa, Twiga, XI.1991, leg. M. Vannini.

Figure 1 (continue). **C**, *Rhinolambrus turriger*, 1 male (MZUF 4289), carapace width: 11.9 mm, Sri Lanka, Trincomalee, -18-27 m, II.1889, leg. K. Fristedt. **D**, *Aulacolambrus hoplonotus*, 1 female (MZUF 4288), carapace width: 19.1 mm, Sri Lanka, Trincomalee, -7-11 m, II.1889, leg. K. Fristedt

Figure 1 (continue). **E**, *Enoplolambrus* cf. *carenatus*, 1 male (MZUF 4290), carapace width: 16.0 mm, Sri Lanka, Trincomalee, -7-11 m, II.1889, leg. K. Fristedt.

Figure 2. *Daldorfia spinosissima*, 1 male (MZUF 4292), carapace width: 200.0 mm, Mauritius Is., IV.1875, purchased by L. De Greaux.