TITLE:

Ontogenetic habitat shifts of green turtles
(*Chelonia mydas*) suggested by the size
modality in foraging aggregations along the
coasts of the western Japanese main islands

AUTHOR(S):

Hamabata, Tomoko; Hikida, Tsutomu; Okamoto,
Kei; Watanabe, Saya; Kamezaki, Naoki

CITATION:

Hamabata, Tomoko ...[et al]. Ontogenetic habitat shifts of green turtles (*Chelonia mydas*) suggested by the size modality in foraging aggregations along the coasts of the western Japanese main islands. *Journal of Experimental Marine Biology and Ecology* 2015, 463: 181-188

ISSUE DATE:

2015-02

URL:

<http://hdl.handle.net/2433/193271>

RIGHT:

© 2014 Elsevier B.V.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

Ontogenetic habitat shifts of green turtles (*Chelonia mydas*) suggested by the size modality
in foraging aggregations along the coasts of the western Japanese main islands

Tomoko Hamabata^a, Tsutomu Hikida^a, Kei Okamoto^b, Saya Watanabe^c and Naoki
Kamezaki^{d, e}

^a Graduate School of Science, Kyoto University, Kitashirakawa-Oiwakecho, Sakyo, Kyoto
606-8502, Japan

^b Orido, Shimizu, Shizuoka, Shizuoka 424-0902, Japan

^c Sea Turtle Association of Japan, 5-17-18 Nagaomotomachi Hirakata, Osaka 573-0163,
Japan

^d Suma Aqualife Park Kobe, 1-3-5 Wakamiyacho Suma, Kobe, Hyogo 654-0049, Japan

^e Faculty of Biosphere-Geosphere Science, Okayama University of Science, 1-1 Ridai-mae
Kita, Okayama, Okayama 700-0005, Japan

Corresponding author: T. Hamabata

E mail: hamabata@zoo.zool.kyoto-u.ac.jp

Phone: +81-75-753-4091; Fax: +81-75-753-4114

25 **Abstract**

26 To understand the life histories and ontogenetic habitat utilization of green turtles along the
27 coasts of the western Japanese main islands, we collected size frequency and genetic data of
28 green turtles captured by pound nets in three foraging grounds (FG): Nomaike (n = 38),
29 Muroto (n = 93), and Kumano-nada (n = 31), and compared their natal origins among
30 different size classes. Population genetic analyses based on an 820-bp fragment of
31 mitochondrial DNA showed that the three FG were part of a single multiple-coast FG.
32 Although turtles from all size classes originated mainly from rookeries in the Ogasawara
33 Group, the size distributions clearly exhibited bimodality, with low occurrences of turtles in
34 the 50–70-cm straight carapace length (SCL) range. The bimodal size distributions could
35 not be attributed to demographic shifts in rookeries, because the number of female green
36 turtles in Ogasawara has exhibited an increasing trend since 1979. We also examined
37 whether factors such as seasonality and predation risk could have caused the size bimodality.
38 There were, however, no strong relationships between sea-surface temperatures when
39 turtles were captured and the sizes of the turtles ($r^2 < 0.2$), and it appeared that predation risk
40 could not result in the size modality observed in the FG. Our results strongly suggest that
41 after switching from a pelagic to a neritic lifestyle, the green turtles in the neritic FG along
42 the western Japanese main islands undergo another ontogenetic habitat shift upon reaching
43 ~50-cm SCL. Here, we explore the possibility that developmental growth might stimulate a

44 habitat shift, resulting in habitat differentiations by size and growth phase in the long-lived
45 green turtle.

46

47 **Keywords**

48 Size distribution, Foraging ground, Ontogenetic habitat shift, Green turtle, Developmental
49 change

50

51 **Abbreviations**

52 FG: Foraging ground

53 MSA: Mixed Stock Analysis

54 mtDNA: Mitochondrial DNA

55 SCL: Straight carapace length

56 SST: Sea surface temperature

57 RMU : Regional Management Unit

58

59 **1. Introduction**

60 The green turtle (*Chelonia mydas*) has a circumglobal distribution and is highly migratory
61 and long-lived. It is believed that in its typical life cycle, after hatching on beaches the green
62 turtle spends several years in the oceanic environment in its pelagic stage before moving
63 back into the neritic zone (neritic stage), where its major food sources, seagrass and sea
64 algal beds, are distributed (Bolten, 2003). Although this typical life cycle suits some
65 populations, several studies have revealed that green turtles have non-homogenous

66 lifestyles, and may exhibit diversity or plasticity in behavior, food resources, and timing of
67 ontogenetic dietary and habitat shifts among populations, regions, and individuals (e.g.,
68 Burkholder et al., 2011; Cardona et al., 2009; González Carman et al., 2012; Hatase et al.,
69 2006; Hays et al., 2002; Parker et al., 2011). This lifestyle diversity suggests that the
70 ecologies and life histories of green turtles may vary depending on their local environments,
71 the availability of resources, and their biological requirements.

72 Foraging grounds (FG), where green turtles aggregate and spend the vast majority
73 of their life spans, are some of the most important areas for understanding their ecology,
74 migration, and life history. Previous studies have constructed hypothetical scenarios of the
75 foraging aggregation processes of marine turtles by using mixed-stock analysis (MSA),
76 estimating the contributions of various source rookeries to FG based on the haplotype
77 compositions of mitochondrial DNA (mtDNA) in rookeries and FG (reviewed by Jensen et
78 al., 2013). Many studies have focused on the early passive recruitment to neritic FG after the
79 pelagic stage and proposed factors that might influence foraging aggregation, such as
80 distance from rookeries (Bass and Witzell, 2000), relative sizes of rookeries (Lahanas et al.,
81 1998), and oceanic currents (Bass et al., 2006). Yet in some FG, immature turtles were
82 thought to have migrated from their initial FG to suitable FG closer to their natal rookeries
83 (Luke et al., 2004).

84 As suggested by several mechanisms proposed above, green turtle size distribution

85 data in FG have also demonstrated that the timing of habitat shifts and patterns of habitat use
86 are varied and complex across regions, even after the shift from a pelagic to a neritic
87 lifestyle. For example, in Atlantic sites many green turtle FG are dominated by immature
88 turtles, and some FG are shared seasonally with migratory adults (Meylan et al., 2011). Also,
89 evidence from the direct tracking of the developmental migrations of large immature green
90 turtles in the Atlantic has shown that they can actively swim among different FG (Godley et
91 al., 2003). In contrast, the FG in Pacific sites have well mixed compositions of adult and
92 immature turtles year-round (Balazs, 1980; Limpus et al., 2005; Sterling et al., 2013). These
93 mixed-size compositions were believed to be a result of their long-term strong fidelity to
94 their foraging areas based on mark and recapture studies that showed limited movement
95 between FG (Limpus et al., 1992). One study conducted MSA in the green turtle FG of the
96 Torres Strait, in the southwestern Pacific, however, demonstrated that the contributions of
97 source rookeries varied between the juveniles and the adults, suggesting either
98 developmental migration by juveniles, or demographic shifts due to reduced hatching
99 success at source rookeries (Jensen, 2010). Moreover, at Palmyra Atoll, in the central
100 Pacific, discrepancies between the directions of oceanic currents and the distributions of
101 haplotype data suggested that biological processes may be important factors driving green
102 turtle foraging aggregation (Naro-Maciel et al., 2014). These results indicate that foraging
103 aggregations were not always governed by the passive early recruitments. Thus, detailed

investigations of the composition and dynamics of each foraging aggregation, as well as information on the source rookeries, are critical for revealing the main influences on the lifestyle, ecology, and migration of this species.

Distributions of foraging green turtles on the coasts of the Japanese main islands in the northwestern Pacific, were determined based on incidental catch by coastal net fisheries, direct observation, and coastal stranding data (Kameda and Ishihara, 2009; Kamezaki et al., 2007; Okamoto et al., 2011; Shimada, 2009). Previous studies using MSA have revealed that green turtles in foraging aggregations along the Japanese main islands originate primarily from rookeries in the Ogasawara Group, the largest nesting site in Japan (Nishizawa et al., 2014, 2013). Foraging aggregations, however, have been described without regard to population dynamics, because available data on the size compositions of foraging green turtles around the Japanese main islands were limited (e.g., Okamoto et al., 2011). Here we describe the size compositions of green turtle foraging aggregations around the western Japanese main islands, discuss how these foraging aggregations were formed, and provide new insights into foraging aggregation dynamics and the factors affecting the ontogenetic habitat shifts of green turtles.

2. Materials and methods

2.1. Descriptions of the study sites and sample collection

The Japanese main islands are all situated beyond the northern limit of the female green turtles' nesting sites in the North Pacific, and green turtles distributed along these coasts are therefore considered to be aggregated for the purpose of foraging. Our data set consists of the aggregations from three FG located in the coastal areas of the western Japanese main islands: Nomaike FG (31°25'N, 130°08'E), Muroto FG (33°15'N, 134°11'E), and Kumano-nada FG (34°07'N, 136°27'E) (Fig. 1A). The Nomaike FG was located on the southwestern coast of Kyushu Island. The pound net used at Nomaike was set at a depth of 27 m in the inner bay. The Muroto FG was located on the southeastern coast of Shikoku Island. Our samples from the Muroto FG were taken from individuals captured in three pound nets located within an 8-km area near the tip of the Cape. These pound nets were set at depths ranging from 35–78 m, near the edge of the narrow continental shelf. The Kumano-nada FG was located at the west side of the Kii Peninsula, in the central part of the Pacific coast of the Japanese main island. These pound nets were set at depths of ~60 m in the ria coasts.

Tissue samples were collected from green turtles captured from the three FG from 2004 to 2012 (Table 1, n = 162): 38 turtles from the Nomaike FG, 93 turtles from the Muroto FG, and 31 turtles from the Kumano-nada FG. All turtles from the Nomaike FG and 59 turtles from the Muroto FG were previously examined by Hamabata et al. (2009). Twelve turtles previously examined by Hamabata et al. (2009) that were listed as samples from

Owase were regarded as samples from the Kumano-nada FG in this study, as Owase is a part of Kumano-nada. Two turtles from the Muroto FG and two from the Kumano-nada FG were the same individuals as used in a morphological study by Okamoto and Kamezaki (2014); these four turtles correspond to ID nos. 8–11 in Okamoto and Kamezaki (2014; Appendix 1). Living turtles were released after the attachment of plastic tags, Inconel tags, or both, and multiple samples from the same individual were avoided. Size frequency structures were constructed based on the straight carapace length (SCL). The relationship between water temperature and the sizes of turtles aggregating in FG were examined by regression analysis. Sea surface temperature (SST) data were used for water temperature. We obtained the SST data from the Japan Oceanographic Data Center website (http://www.jodc.go.jp/index_j.html), the Kochi Prefectural Fisheries Experiment Station (http://www.suisan.tosa.pref.kochi.lg.jp/kaikyo_inf/show), and the Mie Prefectural Fisheries Research Institute (<http://www.mpstpc.pref.mie.lg.jp/SUI/kaikyo/index.htm>). For size analyses, the turtles from each FG were grouped into three size classes: i) $SCL < 50$ cm; ii) $50 \text{ cm} \leq SCL < 70 \text{ cm}$; and iii) $SCL \geq 70 \text{ cm}$. These size groupings were arbitrarily defined, with consideration of the shapes of the size distributions.

2.2. Molecular techniques and haplotype determination

Skin or muscle samples were preserved in 99% ethanol until laboratory analysis. We

161 isolated DNA from skin or muscle samples by phenol/chloroform extraction or DNeasy
162 Blood and Tissue Kits (QIAGEN). Polymerase chain reaction (PCR) was performed using
163 the primers LCM15382 and H950 (Abreu-Grobois et al., 2006), designed to target an
164 820-bp fragment containing partial sequence of the tRNA^{Pro} gene and the 5' end of the
165 control region of the mtDNA genome. For each PCR, 1–2 µl of template DNA was used in a
166 12.5- or 15.0-µl reaction volume under the following conditions: hot start at 94°C for 3 min;
167 35–40 cycles of 94°C for 30 s, 55°C for 30 s, and 72°C for 60 s; a final extension at 72°C for
168 3 min; and then storage at 4°C. Sequences were obtained using an ABI model 3130xl
169 (Applied Biosystems Inc.) sequencer, with all variable positions confirmed by comparing
170 sequences from the forward and reverse strands. Sequences were assembled using DNA
171 BASER (Heracle Biosoft S.R.L), and aligned with Muscle in MEGA 6 (Tamura et al., 2013).
172 We sequenced all 162 samples, because short fragments of haplotypes (500-bp) were used
173 in the previous study by Hamabata et al. (2009). Sequences were compared to previously
174 described haplotypes based on 380-bp, 500-bp, and >700-bp fragments of the mtDNA
175 control region reported by Chassin-Noria et al. (2004), Cheng et al. (2008), Dethmers et al.
176 (2006, 2010), Dutton et al. (2008), Hamabata et al. (2009, 2014), Nishizawa et al. (2010,
177 2011, 2013, 2014), and Norman et al. (1994). The standardized haplotype names for the
178 Indo-Pacific region were assigned to new sequences.

179

2.3. Population genetic analyses

Haplotype diversity (h) and nucleotide diversity (π) were estimated for each FG and each size class of the FG using ALREQUIN V.3.5. (Excoffier and Lischer, 2010) based on 380-bp sequences, to compare our results with FG from previous studies. Annual variations were examined in the Muroto and Kumano-nada FG (Muroto FG: $n = 37$ in 2005, $n = 9$ in 2006, $n = 25$ in 2007, and $n = 17$ in 2008; Kumano-nada FG, $n = 14$ in 2004, $n = 3$ in 2005, $n = 6$ in 2008, and $n = 8$ in 2009). Five turtles in the Muroto FG ($n = 1$ in 2004, $n = 2$ in 2009, and $n = 2$ in 2011) were excluded from this analysis due to the small sample sizes for annual comparisons. Annual variations were not examined in the Nomaike FG because all samples were collected in 2004. To examine seasonal variations within the FG, turtles were grouped based on two different SST as follows: i) $SST < 20^{\circ}\text{C}$ ($n = 15$ in the Muroto FG, $n = 9$ in the Kumano-nada FG) and $SST \geq 20^{\circ}\text{C}$ ($n = 78$ in the Muroto FG, $n = 22$ in the Kumano-nada FG), and ii) $SST < 25^{\circ}\text{C}$ ($n = 6$ in Nomaike FG, $n = 56$ in Muroto FG, $n = 16$ in Kumano-nada FG) and $SST \geq 25^{\circ}\text{C}$ ($n = 32$ in Nomaike FG, $n = 37$ in Muroto FG, $n = 15$ in Kumano-nada FG). In the Nomaike FG, only one turtle was captured at a $SST < 20^{\circ}\text{C}$. Therefore, the Nomaike FG was excluded from the former analysis. Annual and seasonal variations were examined by exact tests of population differentiation (Raymond and Rousset, 1995) using a Markov chain length of 500,000 steps with 10,000 dememorization steps, implemented in ARLEQUIN.

199 Genetic data from the three FG examined in this study were also compared by
200 exact tests with data from four other FG around Japan: Yaeyama and Ginoza in the Ryukyus,
201 and Kanto and Sanriku in the eastern Japanese main islands (Nishizawa et al., 2013, 2014).
202 The comparisons were conducted using shorter sequences, truncated to ~380 bp. The
203 p-values of the multiple comparisons were corrected using the B-Y method (Benjamini and
204 Yekutieli, 2001).

205

206 2.4. *Mixed-stock analyses*

207 We performed MSA on pooled size group data from the Nomaie, Muroto, and
208 Kumano-nada FG, because no significant differences were observed between the same size
209 classes among the three FG. Following Naro-Maciel et al. (2014), potential source rookeries
210 outside of the northwestern Pacific were classified under the following regional
211 management units (RMU) developed by Wallace et al. (2010): Hawaii, Eastern Pacific,
212 Western and South Central Pacific, Southwestern Pacific, Eastern Indian, and Southeast
213 Asia. The Northwestern Pacific RMU was divided into the following four regions for a
214 more detailed investigation: the Taiwan and Hong Kong, Yaeyama, Central Ryukyu, and
215 Ogasawara regional groups of rookeries (Fig. 1B). Haplotype data for the above rookeries
216 were derived from Dethmers et al. (2006), Dutton et al. (2008), Chassin-Noria et al. (2004),
217 Cheng et al. (2008), Hamabata et al. (2014), Nishizawa et al. (2011, 2013), Ng et al. (2014),

218 and Naro-Maciel et al. (2014). Additional data from four female turtles nested on the
219 northwestern Amami Oshima Island, Central Ryukyus, in 2013, were also included for
220 analysis (all four females possessed haplotype CmP39.1). The rookery size data for the
221 MSA were obtained from Amorocho et al. (2012), Cheng et al. (2008), Dethmers et al.
222 (2006), Maison et al. (2010), and Hamabata et al. (2014). Following Nishizawa et al. (2013,
223 2014), MSA estimations were conducted in two ways using Bayesian methods:
224 many-to-one (M2O) analysis using the program BAYES, which examines each size class of
225 the combined FG independently (Pella and Masuda, 2001), and many-to-many (M2M)
226 analysis using the software package R which enables the estimation of multiple FG
227 simultaneously (Bolker et al., 2007). In the M2M analyses we included the data from four
228 other FG around Japan: the Yaeyama, Ginoza, Kanto, and Sanriku FG (Nishizawa et al.,
229 2013, 2014). Both methods were carried out under two priors: (1) uninformative priors
230 assumed that each rookery had the same likelihood of contributing individuals to the
231 foraging aggregations (M2O₁ and M2M₁), and (2) informative priors incorporated the
232 relative size of each rookery (M2O₂ and M2M₂). For the M2O analyses, six chains,
233 corresponding to potentially contributing sources, were run with 20,000 Markov chain
234 Monte Carlo (MCMC) steps and a burn-in of 10,000 runs to calculate the posterior
235 distribution. For the M2M analyses, six chains were run with 50,000 MCMC steps and a
236 burn-in of 25,000 runs. The Gelman and Rubin shrink factor diagnostic was calculated to

test that the posterior probability distribution of all chains had converged (shrink factor < 1.2). Orphan haplotypes, defined as haplotypes observed only in the foraging grounds and not in any of the nesting rookeries, were removed from the analyses.

3. Results

3.1. Size compositions

The sizes of turtles in the Nomaie, Muroto, and Kumano-nada FG ranged from 40.6 to 96.7, 37.2 to 105.2, and 37.3 to 95.4 cm SCL, respectively. All size distributions (three individual distributions and one pooled distribution), plotted in 5-cm increments, did not exhibit bell-shaped curves, but bimodal size distributions, with peaks in the 45–49.9-cm range, and either the 70–74.9- or 75–79.9-cm range (Fig. 2). There was no strong relationship between SST and SCL ($r^2 < 0.2$, Fig. 3).

3.2. Haplotype composition, genetic diversity, and differentiation

Twenty-seven 820-bp haplotypes were identified from a total of 162 samples from green turtle FG around the Japanese main islands (Table 1). Twenty-three haplotypes matched previously reported shorter haplotypes (380- or 500-bp), and 16 matched previously identified longer haplotypes (>700-bp). Two haplotypes did not match any previously identified sequences, and were assigned standardized haplotype designations. One new

haplotype, found in the Kumano-nada FG, differed from CmP50.1 by one base pair, and was assigned the name CmP210.1 (GenBank accession no. **AB896707**). The other new haplotype, found in the Muroto FG, was characterized by a 10-bp insertion difference from CmP39.1, and has been assigned the name CmP208.1 (GenBank accession no. **AB896708**). The Hawaiian and Eastern Pacific haplotypes (CmP4.1, CmP6.1, and CmP15.1) were observed in the Muroto and Kumano-nada FG (Table 1). Orphan haplotypes (CmP51.1, CmP79.1, CmP93.1, CmP122.1, CmP131.1, CmP208.1.1, CmP210.1.1, CmP213.1) made up 6.8% ($n = 11$) of the total sample population. Both the haplotype and nucleotide diversity in the Muroto FG and the haplotype diversity in the Kumano-nada FG were highest in the 50–70-cm SCL class despite the smaller sample sizes than the other size classes; however, the nucleotide diversities of the 50–70 cm SCL classes in the Nomaie and Kumano-nada FG were lower than those of the other size classes, showing no consistent pattern in diversity indices (Table 2). The diversity of the total sample population was similar to that of the Sanriku FG, and lower than the other FG in the Ryukyus and Kanto (Table 2).

Neither annual nor seasonal variations were observed in any of the FG (annual: $p > 0.055$, and seasonal $p > 0.482$). Exact tests revealed that significant population differentiation occurred between the FG in the Ryukyus (Yaeyama and Ginoza) and the Muroto and Kumano-nada FG, but no significant differences were observed among the Nomaie, Muroto, and Kumano-nada FG (Table 3). In addition, significant population

differentiations were supported between the Kumano-nada and Kanto FG by the exact tests, even after the correction for multiple comparisons (Table 3), but no significance was observed among the three FG of the present study and the Sanriku FG, which was more distant than the Kanto FG. Although a significant difference was observed between the 50–70-cm SCL class of the Muroto FG and the >70-cm SCL class of the Kumano-nada FG, the difference was absent after correction for multiple comparisons (Table 4).

3.3. *Mixed-stock analyses*

All estimations by MSA indicated that in the FG of the western Japanese main islands, many turtles in all size classes originated from rookeries in the Ogasawara Group, although the proportion decreased in M2M analyses, especially in M2M of the 50–70-cm SCL classes (Fig. 4). While the contributions from the Northwestern Pacific rookeries (the Taiwan, Hong Kong, Yaeyama, and central Ryukyu rookeries) were very small in the M2O analyses (the lower limits of 95% probability intervals included zero), in M2M the 95% probability intervals of these rookeries were larger (Fig. 4). The probability intervals were broader in the M2M of the 50–70-cm SCL classes (Fig. 4).

4. Discussion

4.1. *Genetic structure of the foraging aggregations along the Japanese coasts*

294 The non-differentiation of genetic compositions among the three FG examined in the
295 present study suggests that turtles migrate among the coasts of the western Japanese main
296 islands as if they constitute a single foraging site. This is supported by a report that one
297 turtle tagged at the Kumano-nada FG on 3 November 2005 was recaptured at the Muroto
298 FG eleven days later (Okamoto and Kamezaki, 2014). Our additional population genetic
299 analyses highlighted significant differences between the Yaeyama FG and all FG along the
300 Japanese main islands. Yet, the extent of the multiple-coast foraging site was unclear,
301 because the significant differences did not show a clear pattern. For example, the
302 Kumano-nada FG was not significantly differentiated from the Sanriku FG, but was
303 significantly differentiated from the closer Kanto FG. These results suggested that the
304 boundaries among the FG were complex or that more samples are needed to reveal the
305 boundaries conclusively.

306

307 *4.2. Size compositions in the FG and population trends in the natal region*

308 The range of SCL in the present study indicated that green turtles in various growing stages
309 aggregate around the Japanese main islands. The mixed-size compositions of all three FG in
310 the present study were consistent with size ranges from other FG in the Pacific. The smallest
311 turtles in the three FG were ~40 cm SCL. This size is similar to that at which pelagic
312 juveniles in the Pacific appear to switch to a neritic lifestyle (e.g., Balazs, 1980; Limpus et

al., 2005). The size distributions of the three FG, however, clearly demonstrated a characteristic bimodality with a low frequency of turtles of 50–70 cm SCL, similar to reports from FG in Shoalwater Bay in eastern Australia (Limpus et al., 2005), and in eastern Taiwan (Cheng and Chen, 1997). Bresette et al. (2010) also reported a bimodal size distribution of pooled green turtles from Mooney Harbor and the eastern Quicksands, west of Key West, Florida, USA, although the bimodality was not noted in their study. Interestingly, the peak sizes of the present size distributions were consistent among all three FG. Some FG in the Atlantic are seasonally shared by immature and adult green turtles, and their size compositions could change temporarily to be similar to a bimodal distribution (Meylan et al., 2011). The bimodality of the present study, however, was not a result of seasonal sharing among turtles in different life stages, as both smaller and larger turtles were captured in various sea surface temperatures.

One of the most important factors affecting the population demographics was the number of births in the rookeries. The estimations by MSA indicated that in the three FG of the present study, the main natal origin of turtles of all size groups was the Ogasawara Group, although some turtles from the central Ryukyu and Yaeyama rookeries were also observed in the FG. Therefore, variations in the numbers of births in Japan, especially Ogasawara, the largest nesting site in Japan and the predominant source for the present FG, would substantially affect the demographics of these foraging aggregations. The number of

332 nests in the Ogasawara Group has been monitored since 1979 and demonstrated an
333 increasing trend up to 2005 (Chaloupka et al., 2008; Yamaguchi et al., 2005). The precise
334 age to reach ~50 cm SCL has not been estimated for wild green turtles born in the
335 Ogasawara Group; however, based on estimates from other Pacific regions that females
336 reach sexual maturity at around 20–40 years (Zug et al., 2002), an increasing number of
337 nests over the past 25 years would not result in such SCL bimodality. In addition, although
338 mortality during the pelagic life stage or in the FG could influence the size compositions of
339 foraging aggregations, in the past decades no specific factors that could have increased the
340 mortality of green turtles in the pelagic stage or in the FG are known. Therefore, there is no
341 reason to believe that the bimodal size compositions of the three FG reflects skewed
342 population demographics in which green turtles of 50–70 cm SCL were less abundant in the
343 wild. Presumably, the size modality is attributable to location shifts by turtles of ~50-cm
344 SCL into habitats that were not sampled, as indicated by data from the Shoalwater Bay,
345 where over 18 years of sampling adults were captured at a higher frequency than late-stage
346 juveniles (Limpus et al., 2005).

348 *4.3. Possible factors contributing to the bimodal size distributions*

349 What factors could cause green turtles to choose their habitat locations according to their
350 sizes? In several FG, green turtles were known to demonstrate size-partitioning of habitats

as a result juveniles inhabiting shallower waters, and larger turtles inhabiting deeper water (Balazs, 1980; Bresette et al., 2010; Koch et al., 2007; Limpus et al., 2005; López-Mendilaharsu et al., 2005). Such size-partitioning of habitats has been explained by the minimization of predation risk. Yet, all pound nets in our study sites were set at depths deep enough (> 27 m) for large predators such as sharks to approach turtles of all sizes. Thus, the differences in predation risk among sizes probably did not cause the bimodality.

We speculate that a developmental change, which commonly occurs in green turtles, could have stimulated habitat shifts in some FG based on the evidence that low occurrences of turtles in the 50–70-cm SCL range were common in all of the FG exhibiting bimodal size distributions. The size range of 50–70 cm SCL corresponds to an accelerated somatic growth phase that occurs before the sub-adult stage (Chaloupka et al., 2004). The requirement for food resources at this stage is probably increased compared to younger stages, resulting in increased competition within the habitat. Meylan et al. (2011) surmised that in the Atlantic, FG dominated by immature turtles formed because immature green turtles were avoiding intraspecific competition with adults, as green turtles showed density-dependent growth, indicating that intraspecific competition can limit the growth rate (Bjorndal et al., 2000). Similarly, in some FG, the developmental change that is characteristically observed in green turtles upon reaching ~50-cm SCL may drive them to depart to other FG, where they can maintain their growth rate by avoiding intraspecific

competition with turtles of other size classes. The possibility of habitat shifts corresponding to developmental growth has been reported in Kemp's ridley turtles, *Lepidochelys kempii* (Schmid et al. 2003). At present, there is no evidence to support this hypothesis or data showing conditional differences, such as higher intraspecific competition in the three FG from the present study than in other FG with unimodal size distributions. Future studies examining to which location the 50–70-cm SCL turtles move and evaluating differences in the growth rates between unimodal and bimodal FG are needed to verify size-specific habitat preferences and ontogenetic habitat shifts. Nevertheless, our results strongly suggest that developmental growth in green turtles can cause shifts in habitat selection. It is likely that the green turtle foraging aggregations along the coasts of the western Japanese main islands are not maintained by long-term residents, but by periodic and continually dynamic populations resulting from ontogenetic habitat shifts.

Acknowledgments

We thank Takaoka, Shiina, Mitsu, Shirose, and Haida pound net fishery associations, Shimakatsu pound net association Co., Ltd., S. Yamashita, N. Yamashita, and K. Miyauchi for their cooperation in collecting green turtles. We thank Dr. Nishizawa for his help in the MSA of this study. We also thank Dr. Jensen to help us to confirm new haplotypes and assign the standardized names of longer haplotypes. We wish to acknowledge use of the

389 Maptool program of SEATURTLE.ORG for maps in this paper. This research was
390 financially supported in part by grants for Excellent Graduate Schools, MEXT, Japan, from
391 the Kuroshio Biological Research Foundation, and the Sasagawa Scientific Research Grant
392 from The Japan Science Society.

393

394 **References**

- 395 Abreu-Grobois, F.A., Horrocks, J.A., Krueger, B., Formia, A., Beggs, J., 2006. New
396 mtDNA dloop primers which work for a variety of marine turtle species may increase
397 the resolution of mixed stock analyses, in: Book of Abstracts from the 26th Annual
398 Symposium on Sea Turtle Biology and Conservation. International Sea Turtle Society.
399 p. 179. ISBN: 9608792614.
- 400 Amoroch, D.F., Abreu-Grobois, F.A., Dutton, P.H., Reina, R.D., 2012. Multiple distant
401 origins for green sea turtles aggregating off Gorgona Island in the Colombian eastern
402 Pacific. PLoS One 7, e31486. doi:10.1371/journal.pone.0031486
- 403 Balazs, G.H., 1980. Synopsis of biological data on the green turtle in the Hawaiian Island.
404 National Oceanic and Atmospheric Administration, Southwest Fisheries Center
405 Administrative Report H-79-24-C.
- 406 Bass, A.L., Epperly, S.P., Braun-McNeill, J., 2006. Green turtle (*Chelonia mydas*) foraging
407 and nesting aggregations in the Caribbean and Atlantic: impact of currents and
408 behavior on dispersal. J. Hered. 97, 346–54. doi:10.1093/jhered/esl004
- 409 Bass, A.L., Witzell, W.N., 2000. Demographic composition of immature green turtles
410 (*Chelonia mydas*) from the east central Florida coast: evidence from mtDNA makers.
411 Herpetologica 56, 357–367.
- 412 Benjamini, Y., Yekutieli, D., 2001. The control of the false discovery rate in multiple
413 testing under dependency. Ann. Stat. 29, 1165–1188.
- 414 Bjørndal, K.A., Bolten, A.B., Chaloupka, M.Y., 2000. Green turtle somatic growth model:
415 evidence for density dependence. Ecol. Appl. 10, 269–282.

- 416 Bolker, B.M., Okuyama, T., Bjorndal, K.A., Bolten, A.B., 2007. Incorporating multiple
417 mixed stocks in mixed stock analysis: “many-to-many” analyses. *Mol. Ecol.* 16,
418 685–695. doi:10.1111/j.1365-294X.2006.03161.x
- 419 Bolten, A.B., 2003. Chapter 9: Variation in sea turtle life history patterns: neritic vs. oceanic
420 developmental stages, in: Lutz, P.L., Musick, J.A., Wyneken, J. (Eds.), *The Biology of*
421 *Sea Turtles Volume II*. CRC Press, pp. 243–257.
- 422 Bresette, M., Witherington, B., Herren, R., Bagley, D., Gorham, J., Traxler, S., Crady, C.,
423 Hardy, R., 2010. Size-class partitioning and herding in a foraging group of green
424 turtles *Chelonia mydas*. *Endanger. Species Res.* 9, 105–116. doi:10.3354/esr00245
- 425 Burkholder, D., Heithaus, M., Thomson, J., Fourqurean, J., 2011. Diversity in trophic
426 interactions of green sea turtles *Chelonia mydas* on a relatively pristine coastal
427 foraging ground. *Mar. Ecol. Prog. Ser.* 439, 277–293. doi:10.3354/meps09313
- 428 Cardona, L., Aguilar, A., Pazos, L., 2009. Delayed ontogenic dietary shift and high levels of
429 omnivory in green turtles (*Chelonia mydas*) from the NW coast of Africa. *Mar. Biol.*
430 156, 1487–1495. doi:10.1007/s00227-009-1188-z
- 431 Chaloupka, M., Bjorndal, K.A., Balazs, G.H., Bolten, A.B., Ehrhart, L.M., Limpus, C.J.,
432 Suganuma, H., Troëng, S., Yamaguchi, M., 2008. Encouraging outlook for recovery of
433 a once severely exploited marine megaherbivore. *Glob. Ecol. Biogeogr.* 17, 297–304.
434 doi:10.1111/j.1466-8238.2007.00367.x
- 435 Chaloupka, M., Limpus, C.J., Miller, J., 2004. Green turtle somatic growth dynamics in a
436 spatially disjunct Great Barrier Reef metapopulation. *Coral Reefs* 23, 325–335.
437 doi:10.1007/s00338-004-0387-9
- 438 Chassin-Noria, O., Abreu-Grobois, F.A., Dutton, P.H., Oyama, K., 2004. Conservation
439 genetics of the east Pacific green turtle (*Chelonia mydas*) in Michoacan, Mexico.
440 *Genetica* 121, 195–206.
- 441 Cheng, I.-J., Chen, T.-H., 1997. Incidental Capture of five species of sea turtles by coastal
442 setnet fisheries in the eastern waters of Taiwan. *Biol. Conserv.* 82, 235–239.
- 443 Cheng, I.-J., Dutton, P.H., Chen, C.-L., Chen, H.-C., Chen, Y.-H., Shea, J.-W., 2008.
444 Comparison of the genetics and nesting ecology of two green turtle rookeries. *J. Zool.*
445 276, 375–384. doi:10.1111/j.1469-7998.2008.00501.x
- 446 Dethmers, K.E.M., Broderick, D., Moritz, C., Fitzsimmons, N.N., Limpus, C.J., Lavery, S.,
447 Whiting, S., Guinea, M., Prince, R.I.T., Kennett, R., 2006. The genetic structure of

- 448 Australasian green turtles (*Chelonia mydas*): exploring the geographical scale of
449 genetic exchange. *Mol. Ecol.* 15, 3931–3946. doi:10.1111/j.1365-294X.2006.03070.x
- 450 Dethmers, K.E.M., Jensen, M.P., FitzSimmons, N.N., Broderick, D., Limpus, C.J., Moritz,
451 C., 2010. Migration of green turtles (*Chelonia mydas*) from Australasian feeding
452 grounds inferred from genetic analyses. *Mar. Freshw. Res.* 61, 1376.
453 doi:10.1071/MF10084
- 454 Dutton, P.H., Balazs, G.H., LeRoux, R.A., Murakawa, Sh.K.K., Zarate, P., Martines, L.S.,
455 2008. Composition of Hawaiian green turtle foraging aggregations: mtDNA evidence
456 for a distinct regional population. *Endanger. Species Res.* 5, 37–44.
457 doi:10.3354/esr00101
- 458 Excoffier, L., Lischer, H.E.L., 2010. Arlequin suite ver 3.5: a new series of programs to
459 perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.*
460 10, 564–567. doi:10.1111/j.1755-0998.2010.02847.x
- 461 Godley, B., Lima, E., Åkesson, S., Broderick, A.C., Glen, F., Godfrey, M.H., Luschi, P.,
462 Hays, G., 2003. Movement patterns of green turtles in Brazilian coastal waters
463 described by satellite tracking and flipper tagging. *Mar. Ecol. Prog. Ser.* 253, 279–288.
464 doi:10.3354/meps253279
- 465 González Carman, V., Falabella, V., Maxwell, S., Albareda, D., Campagna, C., Mianzan, H.,
466 2012. Revisiting the ontogenetic shift paradigm: The case of juvenile green turtles in
467 the SW Atlantic. *J. Exp. Mar. Bio. Ecol.* 429, 64–72. doi:10.1016/j.jembe.2012.06.007
- 468 Hamabata, T., Kamezaki, N., Hikida, T., 2014. Genetic structure of green turtle (*Chelonia*
469 *mydas*) peripheral populations nesting in the northwestern Pacific rookeries: evidence
470 for northern refugia and postglacial colonization. *Mar. Biol.* 161, 495–507.
471 doi:10.1007/s00227-013-2352-z
- 472 Hamabata, T., Nishida, S., Kamezaki, N., Koike, H., 2009. Genetic structure of populations
473 of the green turtle (*Chelonia mydas*) in Japan using mtDNA control region sequences.
474 *Bull. Grad. Sch. Soc. Cult. Stud. Kyushu Univ.* 15, 35–50.
- 475 Hatase, H., Sato, K., Yamaguchi, M., Takahashi, K., Tsukamoto, K., 2006. Individual
476 variation in feeding habitat use by adult female green sea turtles (*Chelonia mydas*): are
477 they obligately neritic herbivores? *Oecologia* 149, 52–64.
478 doi:10.1007/s00442-006-0431-2
- 479 Hays, G.C., Glen, F., Broderick, A.C., Godley, B.J., Metcalfe, J.D., 2002. Behavioural
480 plasticity in a large marine herbivore: contrasting patterns of depth utilisation between

- 481 two green turtle (*Chelonia mydas*) populations. Mar. Biol. 141, 985–990.
482 doi:10.1007/s00227-002-0885-7
- 483 Jensen, M.P., 2010. Assessing the composition of green turtle (*Chelonia mydas*) foraging
484 grounds in Australasia using mixed Stock Analyses. PhD dissertation, University of
485 Canberra, Australia.
- 486 Jensen, M.P., FitzSimmons, N.N., Dutton, P.H., 2013. Chapter 6: Molecular genetics of sea
487 turtles, in: Wyneken, J., Lohmann, K.J., Musick, J.A. (Eds.), The Biology of Sea
488 Turtles, Volume III. CRC Press, pp. 135–161.
- 489 Kameda, K., Ishihara, T., 2009. Gut contents analysis of green turtles (*Chelonia mydas*) in
490 Japan. Umigame News Lett. 17–23 (in Japanese with English summary).
- 491 Kamezaki, N., Matsuzawa, Y., Mizuno, K., Shima, T., 2007. Abstract of the 45th annual
492 meeting of Herpetological Society of Japan, Distribution of marin turtles in coastal
493 waters of Japan based on their washed-up carcasses. Bull. Herpetol. Soc. Japan 1, 78
494 (in Japanese).
- 495 Koch, V., Brooks, L.B., Nichols, W.J., 2007. Population ecology of the green/black turtle
496 (*Chelonia mydas*) in Bahía Magdalena, Mexico. Mar. Biol. 153, 35–46.
497 doi:10.1007/s00227-007-0782-1
- 498 Lahanas, P.N., Bjorndal, K.A., Bolten, A.B., Encalada, S.E., Miyamoto, M.M., Valverde,
499 R.A., Bowen, B.W., 1998. Genetic composition of a green turtle (*Chelonia mydas*)
500 feeding ground population: evidence for multiple origins. Mar. Biol. 130, 345–352.
501 doi:10.1007/s002270050254
- 502 Limpus, C.J., Limpus, D.J., Arther, K.E., Parmenter, C.J., 2005. Monitoring green turtle
503 population dynamics in Shoalwater Bay : 2000 - 2004. Queensland Environmental
504 Protection Agency and the Great Barrier Reef Marine Park Authority.
- 505 Limpus, C.J., Miller, J.D., Paramenter, C., Reimer, D., McLachlan, N., Webb, R., 1992.
506 Migration of green (*Chelonia mydas*) and loggerhead (*Caretta caretta*) turtles to and
507 from eastern Australian rookeries. Wildl. Res. 19, 347. doi:10.1071/WR9920347
- 508 López-Mendilaharsu, M., Gardner, S.C., Seminoff, J.A., Riosmena-Rodriguez, R., 2005.
509 Identifying critical foraging habitats of the green turtle (*Chelonia mydas*) along the
510 Pacific coast of the Baja California peninsula, Mexico. Aquat. Conserv. Mar. Freshw.
511 Ecosyst. 15, 259–269. doi:10.1002/aqc.676

- 512 Luke, K., Horrocks, J.A., LeRoux, R.A., Dutton, P.H., 2004. Origins of green turtle
513 (*Chelonia mydas*) feeding aggregations around Barbados, West Indies. Mar. Biol. 144,
514 799–805. doi:10.1007/s00227-003-1241-2
- 515 Maison, K.A., Kelly, I.K., Frutchey, K.P., 2010. Green turtle nesting sites and sea turtle
516 legislation throughout Oceania: NOAA Tech. Memo. NMFS-F/SPO-110.
- 517 Meylan, P.A., Meylan, A.B., Grey, J.A., 2011. The ecology and migrations of sea turtles 8.
518 Tests of the developmental habitat hypothesis, in: Bulletin of the American Museum of
519 Natural History.
- 520 Naro-Maciel, E., Gaughran, S.J., Putman, N.F., Amato, G., Arengo, F., Dutton, P.H.,
521 McFadden, K.W., Vintinner, E.C., Sterling, E.J., 2014. Predicting connectivity of
522 green turtles at Palmyra Atoll, central Pacific: a focus on mtDNA and dispersal
523 modelling. J. R. Soc. Interface 11, 20130888. doi:10.1098/rsif.2013.0888
- 524 Ng, C.K., Dutton, P.H., Chan, S.K., Cheung, K., Qiu, J., Sun, Y., 2014. Characterization
525 and Conservation Concerns of Green Turtles (*Chelonia mydas*) Nesting in Hong Kong,
526 China. Pacific Sci. 68, 231–243. doi:10.2984/68.2.5
- 527 Nishizawa, H., Abe, O., Okuyama, J., Kobayashi, M., Arai, N., 2011. Population genetic
528 structure and implications for natal philopatry of nesting green turtles *Chelonia mydas*
529 in the Yaeyama Islands, Japan. Endanger. Species Res. 14, 141–148.
530 doi:10.3354/esr00355
- 531 Nishizawa, H., Naito, Y., Suganuma, H., Abe, O., Okuyama, J., Hirate, K., Tanaka, S.,
532 Inoguchi, E., Narushima, K., Kobayashi, K., Ishii, H., Tanizaki, S., Kobayashi, M.,
533 Goto, A., Arai, N., 2013. Composition of green turtle feeding aggregations along the
534 Japanese archipelago: implications for changes in composition with current flow. Mar.
535 Biol. 160, 2671–2685. doi:10.1007/s00227-013-2261-1
- 536 Nishizawa, H., Narazaki, T., Fukuoka, T., Sato, K., Hamabata, T., Kinoshita, M., Arai, N.,
537 2014. Juvenile green turtles on the northern edge of their range: mtDNA evidence of
538 long-distance westward dispersals in the northern Pacific Ocean. Endanger. Species
539 Res. 24, 171–179. doi:10.3354/esr00592
- 540 Nishizawa, H., Okuyama, J., Kobayashi, M., Abe, O., Arai, N., 2010. Comparative
541 phylogeny and historical perspectives on population genetics of the Pacific hawksbill
542 (*Eretmochelys imbricata*) and green turtles (*Chelonia mydas*), inferred from feeding
543 populations in the Yaeyama Islands, Japan. Zoolog. Sci. 27, 14–18.
544 doi:10.2108/zsj.27.14

- 545 Norman, J.A., Moritz, C., Limpus, C.J., 1994. Mitochondrial DNA control region
546 polymorphisms: genetic markers for ecological studies of marine turtles. *Mol. Ecol.* 3,
547 363–373.
- 548 Okamoto, K., Ishihara, T., Taniguchi, M., Yamashita, N., Kamezaki, N., 2011. Occurrence
549 of the sea turtles at the coastal water of Kumanonada. *Umigame News Lett.* 88, 13–17
550 (in Japanese).
- 551 Okamoto, K., Kamezaki, N., 2014. Morphological variation in *Chelonia mydas* (Linnaeus,
552 1758) from the coastal waters of Japan, with special reference to the turtles allied to
553 *Chelonia mydas agassizii* Bocourt, 1868. *Curr. Herpetol.* 33, 46–56.
554 doi:10.5358/hsj.33.46
- 555 Parker, D.M., Dutton, P.H., Balazs, G.H., 2011. Oceanic diet and distribution of haplotypes
556 for the green turtle, *Chelonia mydas*, in the Central North Pacific. *Pacific Sci.* 65,
557 419–431. doi:10.2984/65.4.419
- 558 Pella, J., Masuda, M., 2001. Bayesian methods for analysis of stock mixtures from genetic
559 characters. *Fish. Bull. Natl. Mar. Fish. Serv. Seattle* 99, 151–167.
- 560 Raymond, M., Rousset, F., 1995. An exact test for population. *Evolution* (N. Y.) 49,
561 1280–1283.
- 562 Schmid, J.R., Alan, B.B., Kalen, A.B., Lindberg, W.J., Percival, H.F., Zwick, P.D., 2003.
563 Home range and habitat use by Kemp's ridley turtles in west-central Florida. *J. Wildl.*
564 *Manage.* 67, 196–206.
- 565 Shimada, T., 2009. Report of preliminary research of sea turtles in Hachijo Island.
566 *Umigame News Lett.* 7–8 (in Japanese with English summary).
- 567 Sterling, E.J., Mcfadden, K.W., Holmes, K.E., Vintinner, E.C., Arengo, F., Naro-Maciel, E.,
568 2013. Ecology and conservation of marine turtles in a central Pacific foraging ground.
569 *Chelonian Conserv. Biol.* 12, 2–16.
- 570 Tachikawa, H., 1991. Carapace length and body weight of adult green turtle in Ogasawara.
571 *Umigame News Lett.* 8, 7–10 (in Japanese).
- 572 Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S., 2013. MEGA6: Molecular
573 Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.* 30, 2725–9.
574 doi:10.1093/molbev/mst197

- 575 Wallace, B.P., DiMatteo, A.D., Hurley, B.J., Finkbeiner, E.M., Bolten, A.B., Chaloupka,
576 M.Y., Hutchinson, B.J., Abreu-Grobois, F.A., Amorocho, D., Bjorndal, K.A., Bourjea,
577 J., Bowen, B.W., Dueñas, R.B., Casale, P., Choudhury, B.C., Costa, A., Dutton, P.H.,
578 Fallabrino, A., Girard, A., Girondot, M., Godfrey, M.H., Hamann, M.,
579 López-Mendilaharsu, M., Marcovaldi, M.A., Mortimer, J.A., Musick, J.A., Nel, R.,
580 Pilcher, N.J., Seminoff, J.A., Troëng, S., Witherington, B., Mast, R.B., 2010. Regional
581 management units for marine turtles: a novel framework for prioritizing conservation
582 and research across multiple scales. PLoS One 5, e15465.
583 doi:10.1371/journal.pone.0015465
- 584 Yamaguchi, M., Suganuma, H., Narushima, K., 2005. Nesting status of green turtles
585 (*Chelonia mydas*) in Chichijima Islands, Ogasawara in 2005 and a nesting trend over
586 the last 27 years. Umigame News Lett. 2–6 (in Japanese with English summary).
- 587 Zug, G.R., Balazs, G.H., Wetherall, J.A., Parker, D.M., Murakawa, S.K.K., 2002. Age and
588 growth of Hawaiian green seaturtles (*Chelonia mydas*): an analysis based on
589 skeletochronology. Fish. Bull. 100, 117–127.
- 590
- 591

Table 1. Frequencies of 820-bp mtDNA haplotypes for each size class in the FG. The size

classes are denoted as follows: i, $SCL < 50$ cm; ii, $50 \text{ cm} \leq SCL < 70$ cm; and iii, $SCL \geq 70$

cm.

Haplotype name 820-bp	Nomaike FG			Muroto FG			Kumano-nada FG			GenBank Accession no.
	i	ii	iii	i	ii	iii	i	ii	iii	
CmP4.1									1	<u>KC306666</u>
CmP6.1					1					<u>KC306657</u>
CmP15.1						1				<u>KC306649</u>
CmP18.1				1						<u>AB896713</u>
CmP20.1			1							<u>AB819806</u>
CmP20.3						1				<u>KF311745</u>
CmP32.1			2							<u>KF311749</u>
CmP39.1	3	6	12	19	3	27	7	3	7	<u>AB819807</u>
CmP39.2						1				<u>AB896709</u>
CmP49.1		1	1	1						<u>AB819808</u>
CmP50.1	1		2	5	4	9	3			<u>AB819809</u>
CmP51.1				1						<u>AB896706</u>
CmP53.1			1				1		1	<u>AB819810</u>
CmP54.1			2	1	1	3				<u>AB819811</u>
CmP79.1		1				2				<u>AB896712</u>
CmP93.1								1		<u>FJ917194</u>
CmP95.1						2		1		<u>FJ917196</u>
CmP121.1						2	1		1	<u>AB819813</u>
CmP122.1						1	1		1	<u>AB896710</u>
CmP126.1				1						<u>AB819815</u>
CmP127.1	1		1			1		1		<u>AB856321</u>
CmP128.1		1		1						<u>AB896711</u>
CmP130.1	1			1						<u>AB973567</u>
CmP131.1				1						<u>AB973568</u>
CmP208.1						2				<u>AB896708</u>
CmP210.1									1	<u>AB896707</u>
CmP213.1			1							<u>AB973569</u>
Total	6	9	23	32	9	52	13	6	12	

Table 2. Haplotype (h) and nucleotide (π) diversities of green turtles in FG along the Japanese coasts. Values were calculated for both the total samples from each FG and the three size groups from each FG and regional FG based on 380-bp haplotypes. Data for the Yaeyama, Ginoza, and Kanto FG were from Nishizawa et al. (2013), and data for the Sanriku FG were from Nishizawa et al. (2014).

Foraging ground		N	h	π
Nomaike	total	38	0.6913 ± 0.0823	0.02363 ± 0.01236
	< 50 cm	6	0.8000 ± 0.1721	0.02721 ± 0.01672
	50–70 cm	9	0.5833 ± 0.1833	0.01073 ± 0.00668
	> 70 cm	23	0.7273 ± 0.0971	0.02717 ± 0.01434
Muroto	total	93	0.6746 ± 0.0477	0.02320 ± 0.01193
	< 50 cm	32	0.6351 ± 0.0915	0.02106 ± 0.01116
	50–70 cm	9	0.7500 ± 0.1121	0.03248 ± 0.01838
	> 70 cm	52	0.6825 ± 0.0642	0.02189 ± 0.01140
Kumano-nada	total	31	0.6946 ± 0.0888	0.02450 ± 0.01284
	< 50 cm	13	0.6923 ± 0.1187	0.02747 ± 0.01505
	50–70 cm	6	0.8000 ± 0.1721	0.01839 ± 0.01160
	> 70 cm	12	0.6818 ± 0.1482	0.02722 ± 0.01504
Combined	total	162	0.6785 ± 0.0385	0.02324 ± 0.01189
	< 50 cm	51	0.6525 ± 0.0691	0.02257 ± 0.01175
	50–70 cm	24	0.7391 ± 0.0891	0.02391 ± 0.01268
	> 70 cm	87	0.6855 ± 0.0528	0.02352 ± 0.01209
Yaeyama	-	142	0.8355 ± 0.0215	0.03343 ± 0.01675
Ginoza	-	20	0.8789 ± 0.0432	0.03473 ± 0.01819
Kanto	-	47	0.7438 ± 0.0448	0.03054 ± 0.01563
Sanriku	-	39	0.6478 ± 0.0745	0.02313 ± 0.01210

Table 3. P-values from exact tests based on the 380-bp haplotypes identified in FGs around Japan. Data for the Yaeyama, Ginoza, and Kanto FGs were from Nishizawa et al. (2013), and data for the Sanriku FG were from Nishizawa et al. (2014).

	Yaeyama	Ginoza	Nomaike	Muroto	Kumano-nada	Kanto
Ginoza	0.3725					
Nomaike	< 0.001**	0.0469*				
Muroto	< 0.001**	0.0110**	0.2683			
Kumano-nada	< 0.001**	0.0063**	0.3074	0.1422		
Kanto	< 0.001**	0.1433	0.0166*	0.1981	0.0090**	
Sanriku	< 0.001**	0.0059**	0.0869	0.3910	0.1623	0.0547

*P < 0.05, **P < 0.0137 in B-Y method for 21 simultaneous tests

Table 4. P-values from exact tests of comparisons of FG size classes based on the 380-bp haplotypes.

Size class	Nomaike FG			Muroto FG			Kumano-nada FG	
	< 50 cm	50–70 cm	> 70 cm	< 50 cm	50–70 cm	> 70 cm	< 50 cm	50–70 cm
Nomaike								
50–70 cm	0.474							
> 70 cm	0.716	0.702						
Muroto								
< 50 cm	0.532	0.537	0.485					
50–70 cm	0.472	0.053	0.373	0.368				
> 70 cm	0.461	0.320	0.216	0.340	0.497			
Kumano-nada								
< 50 cm	0.633	0.278	0.672	0.611	0.405	0.771		
50–70 cm	1.000	0.474	0.468	0.225	0.118	0.267	0.177	
> 70 cm	0.566	1.000	0.426	0.201	0.032*	0.157	0.647	0.568

Significant differences were absent after correction for multiple comparisons ($P < 0.01198$ in B-Y method for 36 simultaneous tests).

* $P < 0.05$

Figure Legends

Fig. 1. Locations of the FG (A), and rookeries (black dots) and regional groups of rookeries (dashed circles) used in this study (B). Stars indicate the FG analyzed in this study. Circles indicate the referenced Sanriku, Kanto, Ginoza, and Yaeyama FG. Rookery location data were from Chassin-Noria et al. (2004), Dethmers et al. (2006), Cheng et al. (2008), Dutton et al. (2008), Naro-Maciel et al. (2014), Nishizawa et al. (2011 and 2013), and Hamabata et al. (2014).

Fig. 2. Size frequency distributions of straight carapace lengths (SCL) in the FG: Nomaie (A, $n = 38$), Muroto (B, $n = 93$), Kumano-nada (C, $n = 31$), and the combined data of the three FG (D, $n = 162$). The minimum sizes considered to be adults in male and female of the Ogasawara Group are 79.4 and 82.1 cm SCL, respectively (Tachikawa, 1991).

Fig. 3. Sea surface temperatures (SST) and sizes (SCL) of green turtles captured at each FG: Nomaie (A), Muroto (B), and Kumano-nada (C). Each point represents an individual.

Fig. 4. Estimated mixed-stock analysis (MSA) of green turtle foraging aggregations along the coasts of the western Japanese main islands. Circles and triangles represent M2O and M2M analyses, respectively. Bars indicate 95% probability intervals. Uninformative prior

637 estimations ($M2O_1$, $M2M_1$) are indicated in black, and informative prior estimations ($M2O_2$,
638 $M2M_2$) are indicated in white. Abbreviations of location and RMU are as follows: HK =
639 Hong Kong, SW Pacific = Southwestern Pacific, W & SC Pacific = Western and South
640 Central Pacific, and SE Asia = Southeast Asia. Size classes are as follows: A, $SCL < 50$ cm;
641 B, $50 \text{ cm} \leq SCL < 70$; C, $SCL \geq 70$ cm.

642