

The strong continuity in weakly o-minimal structures

近畿大学・工学部 田中 広志

Hiroshi Tanaka

Faculty of Engineering, Kinki University

Abstract

In this paper, we study the strong continuity of definable functions in weakly o-minimal structures with the strong cell decomposition property.

Throughout this paper, “definable” means “definable possibly with parameters” and we assume that a structure $\mathcal{M} = (M, <, \dots)$ is a dense linear ordering $<$ without endpoints.

A subset A of M is said to be *convex* if $a, b \in A$ and $c \in M$ with $a < c < b$ then $c \in A$. Moreover if $A = \emptyset$ or $\inf A, \sup A \in M \cup \{-\infty, +\infty\}$, then A is called an *interval* in M . We say that \mathcal{M} is *o-minimal* (*weakly o-minimal*) if every definable subset of M is a finite union of intervals (convex sets), respectively. A theory T is said to be *weakly o-minimal* if every model of T is weakly o-minimal. The reader is assumed to be familiar with fundamental results of o-minimality and weak o-minimality; see, for example, [1], [2], [3], or [4].

For any subsets C, D of M , we write $C < D$ if $c < d$ whenever $c \in C$ and $d \in D$. A pair $\langle C, D \rangle$ of non-empty subsets of M is called a *cut* in M if $C < D, C \cup D = M$ and D has no lowest element. A cut $\langle C, D \rangle$ is said

2010 *Mathematics Subject Classification.* 03C64.

Key Words and Phrases. Weakly o-minimal, cell decomposition and strongly continuous.

to be *definable* in \mathcal{M} if the sets C, D are definable in \mathcal{M} . The set of all cuts definable in \mathcal{M} will be denoted by \overline{M} . Note that we have $M = \overline{M}$ if \mathcal{M} is o-minimal. We define a linear ordering on \overline{M} by $\langle C_1, D_1 \rangle < \langle C_2, D_2 \rangle$ if and only if $C_1 \subsetneq C_2$. Then we may treat $(M, <)$ as a substructure of $(\overline{M}, <)$ by identifying an element $a \in M$ with the definable cut $\langle (-\infty, a], (a, +\infty) \rangle$.

We equip M (\overline{M}) with the *interval topology* (the open intervals form a base), and each product M^n (\overline{M}^n) with the corresponding product topology, respectively. For each positive integer n the topological closure in M^n of a set $A \subseteq M^n$ is denoted by $\text{cl}(A)$. We also write $\text{CL}(A)$ for the closure of a set $A \subseteq \overline{M}^n$ in \overline{M}^n .

Recall the notion of definable functions from [4]. Let n be a positive integer and $A \subseteq M^n$ definable. A function $f : A \rightarrow \overline{M}$ is said to be *definable* if the set $\{\langle x, y \rangle \in M^{n+1} : x \in A, y < f(x)\}$ is definable. A function $f : A \rightarrow \overline{M} \cup \{-\infty, +\infty\}$ is said to be *definable* if f is a definable function from A to \overline{M} , $f(x) = -\infty$ for all $x \in A$, or $f(x) = +\infty$ for all $x \in A$.

Recall the notion of strong cells from [5].

Definition 1. Suppose that $\mathcal{M} = (M, <, \dots)$ is a weakly o-minimal structure. For each positive integer n , we inductively define *strong cells* in M^n and their completions in \overline{M}^n .

- (1) A one-element subset of M is called a *strong 0-cell* in M . If $C \subseteq M$ is a strong 0-cell, then its completion $\overline{C} := C$.
- (2) A non-empty definable convex open subset of M is called a *strong 1-cell* in M . If $C \subseteq M$ is a strong 1-cell, then its completion $\overline{C} := \{x \in \overline{M} : (\exists a, b \in C)(a < x < b)\}$.

Assume that k is a non-negative integer, and strong k -cells in M^n and their completions in \overline{M}^n are already defined.

- (3) Let $C \subseteq M^n$ be a strong k -cell in M^n and $f : C \rightarrow M$ is a definable continuous function which has a continuous extension $\overline{f} : \overline{C} \rightarrow \overline{M}$. Then the graph $\Gamma(f)$ is called a *strong k -cell* in M^{n+1} and its completion $\overline{\Gamma(f)} := \Gamma(\overline{f})$.

- (4) Let $C \subseteq M^n$ be a strong k -cell in M^n and $g, h : C \rightarrow \overline{M} \cup \{-\infty, +\infty\}$ are definable continuous functions which have continuous extensions $\overline{g}, \overline{h} : \overline{C} \rightarrow \overline{M} \cup \{-\infty, +\infty\}$ such that
- (a) each of the functions g, h assumes all its values in one of the sets $M, \overline{M} \setminus M, \{\infty\}, \{-\infty\}$,
 - (b) $\overline{g}(x) < \overline{h}(x)$ for all $x \in \overline{C}$.

Then the set

$$(g, h)_C := \{\langle a, b \rangle \in M^{n+1} : a \in C, g(a) < b < h(a)\}$$

is called a *strong $(k+1)$ -cell* in M^{n+1} . The completion of $(g, h)_C$ is defined as

$$\overline{(g, h)_C} := \{\langle a, b \rangle \in \overline{M}^{n+1} : a \in \overline{C}, \overline{g}(a) < b < \overline{h}(a)\}.$$

- (5) Let C be a subset of M^n . The set C is called a *strong cell* in M^n if there exists some non-negative integer k such that C is a strong k -cell in M^n .

Let C be a strong cell of M^n . A definable function $f : C \rightarrow \overline{M}$ is said to be *strongly continuous* if f has a continuous extension $\overline{f} : \overline{C} \rightarrow \overline{M}$. A function which is identically equal to $-\infty$ or $+\infty$, and whose domain is a strong cell is also said to be *strongly continuous*.

Definition 2. Let $\mathcal{M} = (M, <, \dots)$ be a weakly o-minimal structure. For each positive integer n , we inductively define a *strong cell decomposition* (or a *decomposition into strong cells* in M^n) of a non-empty definable set $A \subseteq M^n$.

- (1) If $A \subseteq M$ is a non-empty definable set and $\mathcal{D} = \{C_1, \dots, C_k\}$ is a partition of A into strong cells in M , then \mathcal{D} is called a *decomposition of A into strong cells* in M .
- (2) Suppose that $A \subseteq M^{n+1}$ is a non-empty definable set and $\mathcal{D} = \{C_1, \dots, C_k\}$ is a partition of A into strong cells in M^{n+1} . Then \mathcal{D} is called a *decomposition of A into strong cells* in M^{n+1} if $\{\pi(C_1), \dots, \pi(C_k)\}$ is a decomposition of $\pi(A)$ into strong cells in M^n , where $\pi : M^{n+1} \rightarrow M^n$ is the projection on the first n coordinates.

Definition 3. Let $\mathcal{M} = (M, <, \dots)$ be a weakly o-minimal structure and n a positive integer. Suppose that $A, B \subseteq M^n$ are definable sets, $A \neq \emptyset$ and \mathcal{D} is a decomposition of A into strong cells in M^n . We say that \mathcal{D} *partitions* B if for each strong cell $C \in \mathcal{D}$, we have either $C \subseteq B$ or $C \cap B = \emptyset$.

Definition 4. A weakly o-minimal structure $\mathcal{M} = (M, <, \dots)$ is said to have the *strong cell decomposition property* if for any positive integers k, n and any definable sets $A_1, \dots, A_k \subseteq M^n$, there exists a decomposition of M^n into strong cells partitioning each of the sets A_1, \dots, A_k .

Let $\mathcal{M} = (M, <, +, \dots)$ be a weakly o-minimal expansion of an ordered abelian group $(M, <, +)$. Then, the weakly o-minimal structure \mathcal{M} is said to be *non-valuational* if for any definable cut $\langle C, D \rangle$ we have $\inf\{d - c : c \in C, d \in D\} = 0$.

Then, the following facts hold.

Fact 5 ([4, Fact 2.5]). *Let $\mathcal{M} = (M, <, \dots)$ be a weakly o-minimal structure with the cell decomposition property. Suppose that $X \subseteq M^n$ is definable and $f : X \rightarrow \overline{M}$ is definable. Then, there is a decomposition \mathcal{D} of X into strong cells in M^n such that for every $D \in \mathcal{D}$,*

1. $f|_D$ assumes all its values in one of the sets $M, \overline{M} \setminus M$,
2. $f|_D$ is strongly continuous.

Fact 6 ([4, Corollary 2.16]). *Let $\mathcal{M} = (M, <, +, \dots)$ be a weakly o-minimal expansion of an ordered abelian group $(M, <, +)$. Then the following conditions are equivalent.*

1. \mathcal{M} is non-valuational.
2. \mathcal{M} has the strong cell decomposition property.

Let \mathcal{M} be a weakly o-minimal structure with the cell decomposition property. For any strong cell $C \subseteq M^m$, we denote by \overline{R}_C the m -ary relation determined by \overline{C} , i.e. if $a \in \overline{M}^m$, then $\overline{R}_C(a)$ holds iff $a \in \overline{C}$. We define the structure $\overline{\mathcal{M}} := (\overline{M}, <, (\overline{R}_C : C \text{ is a strong cell}))$. The following fact is known.

Fact 7 ([4]). *Let \mathcal{M} be a weakly o-minimal structure with the cell decomposition property. Then, $\overline{\mathcal{M}}$ is o-minimal, and every set $X \subseteq \overline{M}^m$ definable in $\overline{\mathcal{M}}$ is a finite Boolean combination of completions of strong cells in M^m .*

Proposition 8. *Let $\mathcal{M} = (M, <, +, \dots)$ be a weakly o-minimal expansion with the cell decomposition property of an ordered abelian group. Let $X \subseteq M^n$ be definable and $f : X \rightarrow \overline{M}$ definable. Suppose that there is a decomposition \mathcal{D} of X into strong cells in M^n such that for every $D \in \mathcal{D}$,*

1. $f|_D$ assumes all its values in one of the sets $M, \overline{M} \setminus M$,
2. $f|_D$ is strongly continuous,
3. $\overline{f|_D}(\overline{D})$ is bounded.

Then, there exists some continuous extension $\overline{f} : \text{CL}(X) \rightarrow \overline{M}$ of f .

Corollary 9. *Let $\mathcal{M} = (M, <, +, \dots)$ be a weakly o-minimal expansion with the cell decomposition property of an ordered abelian group. Let C be a strong cell and $f : C \rightarrow M$ or $f : C \rightarrow \overline{M} \setminus M$. Suppose that f is definable and strongly continuous, and $\overline{f}(\overline{C})$ is bounded. Then, for any strong cell $D \subseteq C$, $f|_D$ is strongly continuous.*

Remark 10. *Let $\mathcal{M} = (M, <, \dots)$ be a weakly o-minimal structure with the cell decomposition property. Then, the following hold.*

1. *There exist strong cells C, D_1, D_2 such that $C = D_1 \cup D_2$ but $\overline{C} \neq \overline{D_1} \cup \overline{D_2}$.*
2. *There exist strong cells C, D such that $C \subseteq D$ but $\overline{C} \not\subseteq \overline{D}$.*

References

- [1] M. Coste, An introduction to o-minimal geometry, Dottorato di Ricerca in Matematica, Dip. Mat. Univ. Pisa, Istituti Editoriali e Poligrafici Internazionali (2000).
- [2] L. van den Dries, Tame topology and o-minimal structures, Lecture notes series 248, London Math. Soc. Cambridge Univ. Press (1998).

- [3] D. Macpherson, D. Marker and C. Steinhorn, Weakly o-minimal structures and real closed fields, *Trans. Amer. Math. Soc.* 352 (2000), no. 12, 5435–5483.
- [4] R. Wencel, Weakly o-minimal non-valuational structures, *Ann. Pure Appl. Logic* 154 (2008), no. 3, 139–162.
- [5] On the strong cell decomposition property for weakly o-minimal structures, *MLQ Math. Log. Q.* 59 (2013), no. 6, 452–470.