

## ABSTRACTS (PH D THESIS)

Characterization of *O*-methyltransferases involved in lignan biosynthesis

(Graduate School of Agriculture, Department of Applied Life Sciences, Laboratory of Metabolic Science of Forest Plants and Microorganisms, RISH, Kyoto University)

Safendrri Komara Ragamustari

Lignans are a group of phenolic compounds where two phenylpropanoid units are coupled by the central carbon (C8) of their side chains. These compounds have been gaining much interest due to their important biological activities, such as antitumor property, as well as unique stereochemical mechanisms for their biosynthesis. Albeit several reports on cDNAs encoding enzymes involved in lignan biosynthesis, little is known about *O*-methyltransferases (OMTs) in lignan biosynthesis (lignan OMTs), and no cDNAs encoding lignan OMTs have been isolated. In addition, nothing has been known about enzymes involved in the biosynthesis of phenylpropanoid monomers which are dedicated to the biosynthesis of lignans, but not lignins. In the present study, the author isolated and characterized cDNAs that encode lignan OMTs in relation to regio- and enantiomeric selectivities for the first time. The author also examined cinnamyl alcohol dehydrogenases (CADs) which are possibly involved in lignan biosynthesis.




Figure. The cinnamate/monolignol and lignan biosynthetic pathways. DIR: dirigent protein, PLR: pinoreosin/lariciresinol reductase, SIRD: secoisolariciresinol dehydrogenase. Boxed: isolated/characterized in this study.

---

## ABSTRACTS (PH D THESIS)

---

First, a *Carthamus tinctorius* (safflower) cDNA encoding an OMT methylating a lignan was characterized for the first time. Its recombinant OMT catalyzed regioselective methylation of matairesinol to give rise to an antitumor lignan, arctigenin (4'-*O*-methylmatairesinol) (Figure), and designated as *C. tinctorius* matairesinol OMT (CtMROMT). Gene expression analysis and biochemical characterization of the OMT indicated that CtMROMT is responsible for arctigenin biosynthesis in *C. tinctorius*. In addition, two other cDNAs encoding matairesinol-methylating OMT, each from *Anthriscus sylvestris* (cow parsley) and *Forsythia koreana* (Korean forsythia), were isolated and designated as *AsMROMT* and *FkMROMT*, respectively. AsMROMT showed the same regioselectivity of matairesinol methylation as CtMROMT, whereas FkMROMT gave rise to only isoarctigenin, exhibiting the regioselectivity opposite to those of Ct and AsMROMTs (Figure).

Next, enantiomeric control of the three MROMTs was examined; all MROMTs exhibited strict enantioselectivity, giving rise to only (–)-enantiomers of their products. This is in sharp contrast to the previously proposed mechanism for matairesinol methylation by a *Forsythia intermedia* (golden bell) crude OMT preparation, which showed neither regioselectivity nor enantiomeric selectivity.

Third, a cDNA responsible for the first *O*-methylation step in the biosynthesis of podophyllotoxin (another antitumor lignan) from matairesinol (Figure) was isolated from *A. sylvestris*. The recombinant OMT catalyzed regioselective *O*-methylation of thujaplicatin to afford 5-*O*-methylthujaplicatin. Based on the gene-expression and biochemical analyses, it was concluded that this is the first example of a cDNA encoding an enzyme in the podophyllotoxin biosynthetic pathway from matairesinol, and was designated as thujaplicatin OMT (AsTJOMT).

In addition, characterization of CAD-encoding cDNAs was also conducted. CADs are responsible for the last reductive step in monolignol biosynthesis, which are precursors to lignin and lignans (Figure). Three *C. tinctorius* CADs were isolated and designated as *CtCAD1*, *CtCAD2* and *CtCAD3*. Gene expression analysis and biochemical characterization of plant and recombinant CADs suggested that CtCAD2 and CtCAD3 were involved in lignin and lignan biosynthesis, while CtCAD1 is most likely to be involved in plant defense mechanism.

### References

- [1] Umezawa, T., Ragamustari, S.K., Nakatsubo, T., Wada, S., Li, L., Yamamura, M., Sakakibara, N., Hattori, T., Suzuki, S., and Chiang, V.L., “A novel lignan *O*-methyltransferase catalyzing the regioselective methylation of matairesinol in *Carthamus tinctorius*,” *Plant Biotech*, vol. 30, 97-109, 2013.
- [2] Ragamustari, S.K., Shiraiwa, N., Hattori, T., Nakatsubo, T., Suzuki, S., and Umezawa, T., “Characterization of three cinnamyl alcohol dehydrogenases from *Carthamus tinctorius*,” *Plant Biotech*, vol. 30, 315-326, 2013.
- [3] Ragamustari, S.K., Nakatsubo, T., Hattori, T., Ono, E., Kitamura, Y., Suzuki, S., Yamamura, M., and Umezawa, T., “A novel *O*-methyltransferase involved in the first methylation step of yatein biosynthesis in *Anthriscus sylvestris*,” *Plant Biotech*, vol. 30, 375-384, 2013.
- [4] Ragamustari, S.K., Yamamura, M., Ono, E., Hattori, T., Suzuki, S., and Umezawa, T., “Substrate-enantiomer selectivity of matairesinol *O*-methyltransferases,” *Plant Biotech*, in press.