TITLE:

Immature male gibbons produce female-specific songs.

AUTHOR(S):

Koda, Hiroki; Oyakawa, Chisako; Kato, Akemi;
Shimizu, Daisuke; Rizaldi; Koyama, Yasuhiro;
Hasegawa, Satoshi

CITATION:

Koda, Hiroki ...[et al]. Immature male gibbons produce female-specific songs.. Primates; journal of primatology 2013, 55(1): 13-17

ISSUE DATE:

2013-10-25

URL:

<http://hdl.handle.net/2433/196935>

RIGHT:

The final publication is available at Springer via <http://dx.doi.org/10.1007/s10329-013-0390-2>; この論文は著者最終稿です。内容が印刷版と異なることがありますので、引用の際には出版社版をご確認ご利用ください。This is the Accepted Author Manuscript. Please cite only the published version.

1 News and Perspectives to *Primates*

2 **Immature male gibbons produce female-**
3 **specific songs**

4
5 **Hiroki Koda^{a*}, Chisako Oyakawa^{ab}, Akemi Kato^a, Daisuke Shimizu^{cd}, Rizaldi^e,**
6 **Yasuhiro Koyama^f, Satoshi Hasegawa^g**

7
8 ^aPrimate Research Institute, Kyoto University, Japan

9 ^bGraduate School of Agricultural Science, Tohoku University, Japan

10 ^cJapan Monkey Center, Japan

11 ^dGraduate School of Science, Kyoto University, Japan

12 ^eDepartment of Biology, Faculty of Science and Mathematics, Andalas University

13 ^fNHK Enterprise Co. Ltd., Japan

14 ^gInfinity Co. Ltd., Japan

15
16 **Keywords:** Gibbon songs; sexual differentiation; sexual maturation; song development

17
18 *Corresponding author. *E-mail address:* koda.hiroki.7a@kyoto-u.ac.jp

2/Koda et al. Video evidences for great calls by immature male gibbon

Abstract

Gibbons are apes that are well known to produce characteristic species-specific loud calls, referred to as “songs.” Of particular interest is the sex specificity observed in “great calls” of female songs. However, little is known about the development of this call. While great calls have been observed to be given by female gibbons of different ages, they have never been recorded from males. Here, we report 2 observations of immature male gibbons from 2 different species, wild *Hylobates agilis* and captive *H. lar*, spontaneously singing female-specific great calls. Based on the video clips, we conclude that immature males also have the potential of producing great calls. Our observations lead us to propose a new hypothesis for the development of sexual differentiation in the songs of gibbons and bear on the general issue of sex-specific behaviors in primates.

Key words: Gibbon songs; sex specificity; sexual maturation; sexual differentiation

3/Koda et al. Video evidences for great calls by immature male gibbon

Gibbons are apes that are well-known to produce characteristic species-specific loud calls, which are referred to as “songs” (Preuschoft et al. 1984; Marshall and Sugardjito 1986). Currently, 4 genera of gibbons (*Hylobates*, *Symphalangus*, *Nomascus*, and *Bunopithecus*) have been identified, with this taxon occupying a wide range of habitats in southeast Asia (Lappan and Whittaker 2009). Like bird song, gibbon songs exhibit sequential structuring of single vocal utterances (“notes”), with both species and sexual specificity (Haimoff 1984). Vocal characteristics have been used to assess the systematic relationships among gibbon species (Family Hylobatidae) and to reconstruct their phylogeny and evolution (Marshall and Marshall 1976; Brockelman and Schilling 1984; Marshall and Sugardjito 1986; Geissmann 2002). It has been proposed that the production of gibbon songs is inherited, rather than involving a learning process (Brockelman and Schilling 1984).

Of particular interest is the sex-specificity of gibbon songs. A part of female-specific songs, termed the “great call,” is an approximately 10-sec phase that contains a more complicated species-specific acoustic pattern compared to male songs (Geissmann 2002). Great calls are typically given during vocal duets between a mated pair and have been hypothesized to strengthen the pair bond. Sex-specificity in songs is particularly noticeable in *Hylobates* and *Nomascus* (Geissmann 2002), with males and females in these 2 genera producing overlapping vocal notes, elements, and songs. In *Hylobates*, the female-specific acoustic patterns of great calls are substantially different in form compared to male song, i.e., none of the elements in great calls are ever produced by males (Haimoff 1984; Geissmann 2002). Since newborn and infant female gibbons do not give great calls nor other song types, great calls are expected to be acquired later in life. Because strong sexual dimorphism like gibbon songs is rarely observed in primate

4/Koda et al. Video evidences for great calls by immature male gibbon

vocalizations (Geissmann 2002), the developmental process of sexual differentiation in vocal signals is of special interest for the issue of vocal development in nonhuman primates. However, little is known about the developmental process of primate species-specific songs. In one study of *Nomascus*, a genus that differs from *Hylobates*, the acoustic features of great calls given by immature females were reported to change gradually during the early stages of song development (Merker and Cox 1999). Great calls were hypothesized to be a kind of female-specific behavior that emerges at the appropriate developmental stage of female sexual maturation. In contrast, great calls have never been observed at any developmental stage of male maturation. Consequently, the sexual differentiation of gibbon songs is expected to arise through this type of acoustic developmental process.

However, some studies indicate that the sexual differentiation of species-specific songs is not a straightforward developmental process. Interestingly, one early report noted that subadult male gibbons atypically sing some parts of female-specific songs in the species *Nomascus concolor* (Schilling 1984). This observation might imply that immature male gibbons are potentially able to sing some parts of female-specific songs, and suggests that both males and females develop common acoustic characteristics during the developmental process of song sexual differentiation, including female-specific patterns, with such female patterns disappearing in males after sexual maturation. This developmental scenario would challenge traditional views of sexual differentiation in the vocal behavior of gibbons and primates in general. The anecdotal nature of this previous report makes it difficult to evaluate, and in this paper, we report two observations of immature male gibbons from two different species, *Hylobates agilis* and *Hylobates lar*, spontaneously singing female-specific great calls.

5/Koda et al. Video evidences for great calls by immature male gibbon

Specifically, we present sound files and video clips of male gibbons singing female-like-songs.

Observational information*Subjects*

We observed one immature male of agile gibbon (*H. agilis*) in the wild and one immature male white-handed gibbon (*H. lar*) in captivity.

Wild gibbon observations

We investigated the behavior of wild agile gibbons in a tropical rainforest at the field station of Andalas University in Limau Manis (0°54'S, 100°28'E), Sumatra, Indonesia, from August 23 to September 28, 2008. Field research was initiated at this site in 2004 (Koda et al. 2012; Oyakawa et al. 2007). The subject of this report was a 4- or 5-year-old immature male gibbon (named “Air”), belonging to B group, in the vicinity of the research station. The members of B group were habituated to human researchers and were individually identifiable. During the study period, we confirmed that B group contained an adult mated-pair, their presumed offspring (including the present subject) and a 1-year-old infant. The total time of field observations exceeded 227 h over 32 d.

Captive gibbon observations

6/Koda et al. Video evidences for great calls by immature male gibbon

Observations of a captive gibbon were made at the Japan Monkey Center, Inuyama, Aichi, Japan, on July 11, 2008. The subject was an immature male gibbon (named Bobby), who was born on August 5, 2003. At the time of observation, the gibbon was 5 years old. He was housed together with his family, i.e., his mother, 2-year-old brother and adult male.

Results

Observations of male great calls

Supplementary Video S1 is a video clip of Air co-singing with his mother (named “Gula”). Figure 1 shows a typical example sound spectrogram of a female-specific song, with great calls (Supplementary Sound S1). The great calls of agile gibbons are characterized by a sequential organization of multiple notes. Typically, at first, the great call is initiated with 2 or 3 notes of long duration (around 500 msec) and little frequency modulation. This is followed by a pitch-rising note with large frequency modulations (around 1200 Hz). Finally, 2 or 3 high-pitched and loud notes are produced (Haimoff 1984; Oyakawa et al. 2007). We found that the acoustic structure of Air’s song resemble female great calls. This observation provides direct evidence for the possibility that immature males also sing female-specific songs in agile gibbons. It would be interesting to determine whether this is a one-off observation, or if males routinely sing female songs. During 768 min of focused observation of the B group, we recorded the same immature male producing female-specific calls 21 times.

Similar to the observations of a male gibbon in the wild, we recorded the

7/Koda et al. Video evidences for great calls by immature male gibbon

vocalization of an immature male in captivity. Supplementary Video S2, S3, and S4 are video clips of an immature male (Bobby) co-singing with his mother. Each video clip contains singing events by Bobby. Based on the video clips, we confirmed that the acoustic characteristics of his singing were similar to female-specific great calls. In all 3 events, his mother always initiated the singing, and then Bobby also started to sing. The acoustic characteristics and sequential organization of white-handed gibbon songs are quite similar to those of agile gibbons. Typically, at first, the great call is initiated by 2 or 3 notes of long duration (around 500 msec), with little frequency modulations. Then, it is followed by a pitch-rising note with large frequency modulations (around 1200 Hz). Finally, 2 or 3 high-pitched and loud notes are produced (Brockelman and Schilling 1984; Haimoff 1984; Geissmann 2002). Notes of long duration like those initiating great calls have not been previously reported in any types of male songs sung by white-handed gibbons. Thus, it is usually easy to distinguish these sequential notes as the “introductory part” of great calls. Sound spectrograms of the 3 video clips (Figure 2) reveal that the immature male clearly emitted the long-duration notes, followed by the typical pitch-rising note. His movements during singing also mimicked those of females when giving great calls. Typically, females brachiate during and after producing great calls, and interestingly, the immature male also brachiated during the climax of his singing. Based on both the acoustic characteristics of his songs and his movements, it is reasonable to consider that this immature male sang great calls.

Discussion

Are our observations of male great calls “abnormal” behavior?

8/Koda et al. Video evidences for great calls by immature male gibbon

153

154

155

156

157

158

159

160

161

Our observations clearly challenge the standard model for the development of great calls i.e., females acquire the ability to sing great calls as they mature, while males do not. However, a counter argument is that our observations were simply “exceptional” or “abnormal” might be plausible. Our observations from the wild group were obtained from just a single subject. Moreover, despite several long-term studies of gibbons (for review of the literature, see Lappan and Whittaker 2009), it is surprising that there have been no reports of males giving great calls. Our report of the captive white-handed gibbon might also be unusual and not characteristic of the species.

162

163

164

165

166

167

168

169

170

171

172

Successful recordings of male great calls from the captive subject were made on only one day, July 11, 2008. After making these recordings, we continued observations for one month, to confirm that the male gave great calls again; however, we failed to record great calls from this male subsequently. Interestingly, the keeper said that he had never heard any type of singing by this male prior to our observation, indicating that the great calls we recorded were the first time that this male had sung. Indeed, our observations of both wild and captive males are just case reports and might be interpreted as exceptions, instead of reflecting the species-typical behavior of male gibbons. More study is clearly needed before concluding that male gibbons habitually produce female great calls.

173

174

Implications for the sexual differentiation of gibbon songs

175

176

So far, previous studies have reported that gibbon songs are sexually dimorphic; however, the developmental course of song acquisition remains unclear.

9/Koda et al. Video evidences for great calls by immature male gibbon

Despite the obvious sex specificity in male and female songs, few data exist regarding the development of song. Based on the observations presented here, we speculate that additional study of song development will reveal more immature males giving great calls. Alternatively, it may be premature to generalize from the findings presented here. For example, we cannot overlook the fact that there have been no previous reports of male gibbons producing great calls, even though many captive gibbons have been housed in zoos throughout the world and gibbons have been the subjects of considerable research in the wild. Therefore, our observations may be anomalous. Unfortunately, it is unclear why the two immature males observed in this study gave great calls.

One possibility might be related to the idea that the great calls of mothers are a kind of “trigger” for the emergence of songs by their offspring. Recently, we reported the overlapping duets between mothers and their subadult daughters, where daughters acquired the acoustic features of her mothers’ great calls and routinely co-sang with their them (Koda et al. 2013). Subadult daughters co-sang great calls with their mothers just before dispersing from the family group to form a new pair. Mother-daughter duets (interactions) are routinely observed, with daughters rarely singing without any overlap with the mother’s great call. Here, we hypothesize that the great calls of mothers trigger their offspring to start the song when offspring just starts to produce sounds like song, even if the offspring is a son. Regardless of gender, offspring would first start to vocalize his/her songs together with the mother’s great call. Great calls might serve as a trigger to motivate offspring to start the singing. If male androgen levels are not yet elevated in immature male offspring, they might start to give great calls. Once androgen levels begin to increase, the female-like features of great calls might diminish and switch to a male-specific song. This model is consistent with evidence from the video

10/Koda et al. Video evidences for great calls by immature male gibbon

201 footage that appear to suggest that maternal great calls trigger offspring “singing.”
202 Careful observation of supplementary video S2 shows a very immature offspring
203 hugging his mother rhythmically, while producing high-pitched screams that overlap
204 with his mother’s great call. The published literature also documents observations of an
205 immature baby emitting screams while the mother sings her great calls (Preuschoft et al.
206 1984). Female great calls might play a role in triggering the onset of songs by offspring.
207 Our observation of great calls by immature males might be a result of a maternal
208 influence on song acquisition.

209 In sum, gibbon songs present a promising model for examining the
210 developmental course of sexual dimorphism in behavior. Given that primates, including
211 humans, rarely show clear sex-specific behavior, gibbon song would provide an ideal
212 model system to investigate development and its endocrine basis. Future research
213 should track how song changes in relation to hormonal states of singers. Such
214 information could be used to study the development of sexual differentiation in primate
215 vocal behavior.

216

11/Koda et al. Video evidences for great calls by immature male gibbon

217 **Acknowledgements**

218 We thank Akira Kato, Jun Koike, Yuji Takenoshita, Koichiro Zamma, Takahisa
219 Matsusaka, Noriko Ito, and all staff of the Japan Monkey Center for permitting us to
220 conduct this study and for their assistance. We are grateful to Syamsuardi, Santi
221 Nurulkamilah, and the staff of Andalas University, and many friends in Indonesia for
222 their tremendous support and friendship; and to John C. Mitani and Esther Clarke for
223 their comments on our manuscripts. Our fieldwork was conducted under the
224 Memorandum of Universities between KUPRI and UNAND. This study was supported
225 by a Grant-in-Aid for overseas research (B) (20405016) to Hirohisa Hirai (HK joined as
226 a co-PI), and by the Young Researcher Overseas Visits Program for the Vitalizing Brain
227 Circulation from the Primate Research Institute, Kyoto University.

228

229

12/Koda et al. Video evidences for great calls by immature male gibbon

References

- Brockelman WY, Schilling D (1984) Inheritance of stereotyped gibbon calls. *Nature* 312:634-636
- Geissmann T (2002) Duet-splitting and the evolution of gibbon songs. *Biological Review* 77:57-76
- Haimoff E (1984) Acoustic and organizational features of gibbon songs. In: Preuschoft H, Chivers D, Brockelman WY, Creel N (eds) *The lesser apes: evolutionary and behavioural biology*. Edinburgh University Press, Edinburgh
- Koda H, Lemasson A, Oyakawa C, Rizaldi, Pamungkas J, Masataka N (2013) Possible role of mother-daughter vocal interactions on the development of species-specific song in gibbons. *PLoS ONE* 8: e71432
- Koda H, Oyakawa C, Nurulkamilah S, Rizaldi, Sugiura H, Bakar A, Masataka N. (2012) Male replacement and stability of territorial boundary in a group of agile gibbons (*Hylobates agilis agilis*) in west Sumatra, Indonesia. *Primates* 53:327-332.
- Lappan S, Whittaker D (2009) *The gibbons: new perspectives on small ape socioecology and population biology*. In. Springer-Verlag, New York
- Marshall J, Marshall E (1976) Gibbons and their territorial songs. *Science* 193:235-237
- Marshall J, Sugardjito J (1986) Gibbon systematics. In: Swindler DR, Erwin J (eds) *Comparative Primate Biology*, vol 1. Alan R. Liss., New York, pp 137-185
- Merker B, Cox C (1999) Development of the female great call in *Hylobates gabriellae*: A case study. *Folia Primatologica* 70:97-106
- Oyakawa C, Koda H, Sugiura H (2007) Acoustic features contributing to the individuality of wild agile gibbon (*Hylobates agilis agilis*) songs. *American Journal of Primatology* 69:777-790
- Preuschoft H, Chivers D, Brockelman WY, Creel N (1984) *The lesser apes: evolutionary and behavioural biology*. In. Edinburgh University Press, Edinburgh
- Schilling D (1984) Song bouts and duetting in the concolor gibbon. In: Preuschoft H, Chivers D, Brockelman WY, Creel N (eds) *The lesser apes: evolutionary and behavioural biology*. Edinburgh University Press, Edinburgh, pp 390-403

13/Koda et al. Video evidences for great calls by immature male gibbon

262 Figure legends

263

264 Figure 1

265 Sound spectrogram of male great calls by a wild agile immature male gibbon (Sound
266 S1). The spectrograms were generated using fast Fourier transform (sound sampling
267 frequency 8 kHz, frame length 30 ms, Gaussian window, time step 1ms, frequency step
268 20 Hz, 40 dB dynamic range), by Praat ver 5.3.19, developed by Paul Boersma and
269 David Weenick, which is freely available online from www.praat.org. The extracted
270 pitch contours colored in orange and blue represent the mother and her male offspring,
271 respectively.

272

273 Figure 2

274 Sound spectrogram of male great calls by a captive white-handed immature male gibbon
275 with that of his mother, which are presented in Video S2 (A), S3 (B), and S4 (C). The
276 extracted pitch contours colored in orange and blue represent the mother and her male
277 offspring, respectively.

278

14/Koda et al. Video evidences for great calls by immature male gibbon

279 Legends for Supplementary video clips

280

281 Video S1

282 A sample video clip of an immature male gibbon singing great calls in the wild. The
283 larger gibbon was the mother, and smaller one was her son (Air). The two of them sang
284 together in the canopy of a 20-meter-high tree. At the time of this video clip, the mother
285 had already started to sing great calls, and Air had not previously begun to sing with her.
286 After the mother had sung the great calls several times, Air also sang with her. As the
287 great calls reached a climax, the two of them brachiated through the canopy of the forest.
288 In this video, the mother moved first and was then followed by Air replicating her
289 movements.

290

291 Video S2

292 This is the video clip in which we successfully filmed the first case of male great calls
293 in captivity. At the time (11th July, 2008), we were filming gibbon singing for another
294 purpose, so the video camera on the tripod was placed in front of the home cage,
295 covering a wide view of the cage, not focusing on the immature male (Bobby). Bobby
296 had a middle-sized body with black fur. The mother was a large gibbon with a brown
297 and buff fur color, and the smallest gibbon with white fur was her 2-year-old son. The
298 mother's male partner was not filmed, but his songs were recorded in this video. First,
299 the mother sang the great calls alone. Just after the mother started to sing the great call,
300 Bobby approached her closely. The 2-year-old son also approached her after Bobby. As
301 the mother's great call reached its climax, Bobby also began to sing. The mother
302 jumped from the branch to right side when singing the climax. Bobby also jumped to
303 the left side while singing, following the mother's great call and jumping behavior. The
304 2-year-old son also jumped following the mother and Bobby.

305

306 Video S3

307 This video clip was made after Video S2. After watching Bobby's great call, we tried to
308 film him at a closer distance. As in S2, the mother first started to sing a great call. As the
309 mother's great call reached its climax, Bobby also joined in singing. In this case, the
310 mother did not brachiate as much, while Bobby brachiated while singing his climax.
311 Interestingly, the 2-year-old son approached the mother when she singing the climax
312 part, and repeatedly called with very high pitch screaming. At the end of this video, the
313 male partner (black fur) entered the film, but he did not join in the duet at that time.

314

15/Koda et al. Video evidences for great calls by immature male gibbon

315 Video S4

316 The final observation was made after Video S3. In this case, when the mother started to
317 sing, Bobby almost simultaneously started to sing with her. Bobby remained close to his
318 mother during the early part of the singing. During their singing, the screams of the 2-
319 year-old son were sometimes heard too. At the climax part, only Bobby jumped in the
320 cage. A short conversation in Japanese between the observer (HK) and the keeper is
321 included in the video clip.

322

323 Sound S1

324 The typical example of an immature male gibbon singing great calls in the wild. The
325 sound spectrogram was shown in Figure 1.

326

Figure 1

Figure 2