

FINITE GENERATION OF THE LOG CANONICAL RING IN DIMENSION FOUR

OSAMU FUJINO

Dedicated to the memory of Professor Masayoshi Nagata

ABSTRACT. We treat two different topics on the log minimal model program, especially for four-dimensional log canonical pairs.

- (a) Finite generation of the log canonical ring in dimension four.
- (b) Abundance theorem for irregular fourfolds.

We obtain (a) as a direct consequence of the existence of four-dimensional log minimal models by using Fukuda's theorem on the four-dimensional log abundance conjecture. We can prove (b) only by using traditional arguments. More precisely, we prove the abundance conjecture for irregular $(n+1)$ -folds on the assumption that the minimal model conjecture and the abundance conjecture hold in dimension $\leq n$.

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. Log canonical ring	5
3.1. Appendix	8
4. Abundance theorem for irregular varieties	9
4.1. Appendix	12
References	13

1. INTRODUCTION

In this paper, we treat two different topics on the log minimal model program, especially for four-dimensional log canonical pairs. We will

Date: 2009/9/23, version 5.01.

2000 Mathematics Subject Classification. Primary 14J35; Secondary 14E30.

Key words and phrases. log canonical ring, log abundance conjecture, log minimal model program.

freely use the results on the three-dimensional log minimal model program (cf. [FA], [KeMM], etc.). Sorry, we do not always refer to the original papers since the results are scattered in various places.

1 (Finite generation of the log canonical ring in dimension four). The following theorem is the main result of Section 3 (cf. [F4, Section 3.1]).

Theorem 1.1 (Finite generation of the log canonical ring in dimension four). *Let $\pi : X \rightarrow Z$ be a proper surjective morphism from a smooth fourfold X . Let B be a boundary \mathbb{Q} -divisor on X such that $\text{Supp} B$ is a simple normal crossing divisor on X . Then the relative log canonical ring*

$$R(X/Z, K_X + B) = \bigoplus_{m \geq 0} \pi_* \mathcal{O}_X(\lfloor m(K_X + B) \rfloor)$$

is a finitely generated \mathcal{O}_Z -algebra.

It is easy to see that Theorem 1.1 is equivalent to Theorem 1.2.

Theorem 1.2. *Let $\pi : X \rightarrow Z$ be a proper surjective morphism from a four-dimensional log canonical pair (X, B) such that B is an effective \mathbb{Q} -divisor. Then the relative log canonical ring*

$$R(X/Z, K_X + B) = \bigoplus_{m \geq 0} \pi_* \mathcal{O}_X(\lfloor m(K_X + B) \rfloor)$$

is a finitely generated \mathcal{O}_Z -algebra.

In Section 3, we give a proof of Theorem 1.1 by using the existence theorem of four-dimensional log minimal models (cf. [B] and [S2]) and Fukuda's result on the log abundance conjecture for fourfolds (cf. [Fk]). A key point of Fukuda's result is the abundance theorem for semi log canonical threefolds in [F1].

2 (Abundance theorem for irregular fourfolds). In Section 4, we prove the abundance theorem for irregular $(n+1)$ -folds on the assumption that the minimal model conjecture and the abundance conjecture hold in dimension $\leq n$ (see Theorem 4.5). By this result, we know that the abundance conjecture for irregular varieties is the problem for lower dimensional varieties. Since the minimal model conjecture and the abundance conjecture hold in dimension ≤ 3 , we obtain the next theorem (see Corollary 4.7).

Theorem 1.3 (Abundance theorem for irregular fourfolds). *Let X be a normal complete fourfold with only canonical singularities. Assume that K_X is nef and the irregularity $q(X)$ is not zero. Then K_X is semi-ample.*

We also prove that there exists a good minimal model for any smooth projective irregular fourfold (see Theorem 4.8).

Theorem 1.4 (Good minimal models of irregular fourfolds). *Let X be a smooth projective irregular fourfold. If X is not uni-ruled, then there exists a normal projective variety X' such that X' has only \mathbb{Q} -factorial terminal singularities, X' is birationally equivalent to X , and $K_{X'}$ is semi-ample.*

We note that Sections 3 and 4 can be read independently.

Acknowledgments. The author was partially supported by the Grant-in-Aid for Young Scientists (A) #20684001 from JSPS. He was also supported by the Inamori Foundation.

We will work over \mathbb{C} , the complex number field, throughout this paper. We will freely use the notation in [KMM] and [KM]. Note that we do not use \mathbb{R} -divisors.

2. PRELIMINARIES

In this section, we collect basic definitions.

Definition 2.1 (Divisors, \mathbb{Q} -divisors). Let X be a normal variety. For a \mathbb{Q} -Weil divisor $D = \sum_{j=1}^r d_j D_j$ on X such that D_j is a prime divisor for every j and $D_i \neq D_j$ for $i \neq j$, we define the *round-down* $\lfloor D \rfloor = \sum_{j=1}^r \lfloor d_j \rfloor D_j$, where for every rational number x , $\lfloor x \rfloor$ is the integer defined by $x - 1 < \lfloor x \rfloor \leq x$.

We call D a *boundary* \mathbb{Q} -divisor if $0 \leq d_j \leq 1$ for every j .

We note that $\sim_{\mathbb{Q}}$ denotes the \mathbb{Q} -linear equivalence of \mathbb{Q} -divisors.

We call X *\mathbb{Q} -factorial* if and only if every Weil divisor on X is \mathbb{Q} -Cartier.

Definition 2.2 (Exceptional locus). For a proper birational morphism $f : X \rightarrow Y$, the *exceptional locus* $\text{Exc}(f) \subset X$ is the locus where f is not an isomorphism.

Let us quickly recall the definitions of singularities of pairs.

Definition 2.3 (Singularities of pairs). Let X be a normal variety and B an effective \mathbb{Q} -divisor on X such that $K_X + B$ is \mathbb{Q} -Cartier. Let $f : Y \rightarrow X$ be a resolution such that $\text{Exc}(f) \cup f_*^{-1}B$ has a simple normal crossing support, where $f_*^{-1}B$ is the strict transform of B on Y . We write

$$K_Y = f^*(K_X + B) + \sum_i a_i E_i$$

and $a(E_i, X, B) = a_i$. We say that (X, B) is *lc* (resp. *klt*) if and only if $a_i \geq -1$ (resp. $a_i > -1$) for every i . We note that *lc* (resp. *klt*) is an abbreviation of *log canonical* (resp. *Kawamata log terminal*). We also note that the *discrepancy* $a(E, X, B) \in \mathbb{Q}$ can be defined for every prime divisor E over X .

In the above notation, if $B = 0$ and $a_i > 0$ (resp. $a_i \geq 0$) for every i , then we say that X has only *terminal* (resp. *canonical*) singularities.

Definition 2.4 (Divisorial log terminal pair). Let X be a normal variety and B a boundary \mathbb{Q} -divisor such that $K_X + B$ is \mathbb{Q} -Cartier. If there exists a resolution $f : Y \rightarrow X$ such that

- (i) both $\text{Exc}(f)$ and $\text{Exc}(f) \cup \text{Supp}(f_*^{-1}B)$ are simple normal crossing divisors on Y , and
- (ii) $a(E, X, B) > -1$ for every exceptional divisor $E \subset Y$,

then (X, B) is called *divisorial log terminal* (*dlt*, for short).

For the details of singularities of pairs, see, for example, [KM] and [F2].

Definition 2.5 (Center, lc center). Let E be a prime divisor over X . The closure of the image of E on X is denoted by $c_X(E)$ and called the *center* of E on X .

Let (X, B) be an *lc* pair. If $a(E, X, B) = -1$, $c_X(E)$ is called an *lc center* of (X, B) .

The following definitions are now classical.

Definition 2.6 (Iitaka's D -dimension and numerical D -dimension). Let X be a normal complete variety and D a \mathbb{Q} -Cartier \mathbb{Q} -divisor. Assume that mD is Cartier for a positive integer m . Let

$$\Phi_{|tmD|} : X \dashrightarrow \mathbb{P}^{\dim |tmD|}$$

be rational mappings given by linear systems $|tmD|$ for positive integers t . We define *Iitaka's D -dimension*

$$\kappa(X, D) = \begin{cases} \max_{t>0} \dim \Phi_{|tmD|}(X), & \text{if } |tmD| \neq \emptyset \text{ for some } t, \\ -\infty, & \text{otherwise.} \end{cases}$$

In case D is nef, we can also define the *numerical D -dimension*

$$\nu(X, D) = \max\{e \mid D^e \neq 0\},$$

where \equiv denotes *numerical equivalence*. We note that $\nu(X, D) \geq \kappa(X, D)$ always holds.

Definition 2.7 (Nef and abundant divisors). Let X be a normal complete variety and D a \mathbb{Q} -Cartier \mathbb{Q} -divisor on X . Assume that D is nef. The nef \mathbb{Q} -divisor D is said to be *abundant* if the equality $\kappa(X, D) = \nu(X, D)$ holds. Let $\pi : X \rightarrow Z$ be a proper surjective morphism of normal varieties and D a π -nef \mathbb{Q} -divisor on X . Then D is said to be *π -abundant* if D_η is abundant, where $D_\eta = D|_{X_\eta}$ and X_η is the generic fiber of π .

Definition 2.8 (Irregularity). Let X be a normal complete variety with only rational singularities. We put

$$q(X) = h^1(X, \mathcal{O}_X) = \dim H^1(X, \mathcal{O}_X) < \infty$$

and call it the *irregularity* of X .

Let X be as above. If $q(X) \neq 0$, then we call X *irregular*.

If X' is a normal complete variety with only rational singularities such that X' is birationally equivalent to X , then it is easy to see that $q(X) = q(X')$.

3. LOG CANONICAL RING

In this section, we prove the following theorem: Theorem 1.1.

Theorem 3.1 (Finite generation of the four-dimensional log canonical ring). *Let $\pi : X \rightarrow Z$ be a proper surjective morphism from a smooth fourfold X . Let B be a boundary \mathbb{Q} -divisor on X such that $\text{Supp} B$ is a simple normal crossing divisor on X . Then the relative log canonical ring*

$$R(X/Z, K_X + B) = \bigoplus_{m \geq 0} \pi_* \mathcal{O}_X(\lfloor m(K_X + B) \rfloor)$$

is a finitely generated \mathcal{O}_Z -algebra.

The next proposition is well known and a slight generalization of [K2, Theorem 7.3].

Proposition 3.2. *Let (X, B) be a proper log canonical fourfold such that $K_X + B$ is nef and $\kappa(X, K_X + B) > 0$. Then $K_X + B$ is abundant, that is, $\kappa(X, K_X + B) = \nu(X, K_X + B)$.*

Proof. See, for example, [Fk, Proposition 3.1]. We note that we need the three-dimensional log minimal model program and log abundance theorem here (see [FA], [KeMM], and [KeMM2]). \square

Let us recall Fukuda's result in [Fk]. We will generalize this in Theorem 3.10.

Theorem 3.3 (cf. [Fk, Theorem 1.5]). *Let (X, B) be a proper dlt four-fold. Assume that $K_X + B$ is nef and that $\kappa(X, K_X + B) > 0$. Then $K_X + B$ is semi-ample.*

Proof. By Proposition 3.2, $\kappa(X, K_X + B) = \nu(X, K_X + B)$. We put $S = \lfloor B \rfloor$ and $K_S + B_S = (K_X + B)|_S$. Then the pair (S, B_S) is semi-divisorial log terminal and $K_S + B_S$ is semi-ample by [F1, Theorem 0.1]. Finally, by [F3, Corollary 6.7], we obtain that $K_X + B$ is semi-ample. \square

Remark 3.4. The proof of [Fk, Proposition 3.3] depends on [K2, Theorem 5.1]. It requires [K2, Theorem 4.3] whose proof contains a non-trivial gap. See [F2, Remark 3.10.3] and [F5]. So, we adopted [F3, Corollary 6.7] in the proof of Theorem 3.3.

In this section, we adopt Birkar's definition of the log minimal model (see [B, Definition 2.4]), which is slightly different from [KM, Definition 3.50]. See Remark 3.6 and Example 3.7 below.

Definition 3.5 (cf. [B, Definition 2.4]). Let (X, B) be a log canonical pair over Z . A *log minimal model* $(Y/Z, B_Y + E)$ of $(X/Z, B)$ consists of a birational map $\phi : X \dashrightarrow Y/Z$, $B_Y = \phi_* B$, and $E = \sum_j E_j$, where E_j is a prime divisor on Y and ϕ^{-1} -exceptional for every j , and satisfies the following conditions:

- (1) Y is \mathbb{Q} -factorial and $(Y, B_Y + E)$ is dlt,
- (2) $K_Y + B_Y + E$ is nef over Z , and
- (3) for every prime divisor D on X which is exceptional over Y , we have

$$a(D, X, B) < a(D, Y, B_Y + E),$$

where $a(D, X, B)$ (resp. $a(D, Y, B_Y + E)$) denotes the discrepancy of D with respect to (X, B) (resp. $(Y, B_Y + E)$).

Remark 3.6. In [KM, Definition 3.50], it is required that ϕ^{-1} has no exceptional divisors.

Example 3.7. Let $X = \mathbb{P}^2$ and D_X the complement of the big torus. Then $K_X + D_X$ is dlt and $K_X + D_X \sim 0$. Let $Y = \mathbb{P}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-1))$ and D_Y the complement of the big torus. Then (Y, D_Y) is a log minimal model of (X, D_X) in the sense of Definition 3.5. Of course, $K_Y + D_Y$ is dlt and $K_Y + D_Y \sim 0$. On the other hand, (Y, D_Y) is not a log minimal model of (X, D_X) in the sense of [KM, Definition 3.50].

We prepare the following two easy lemmas.

Lemma 3.8. *We use the notation in Definition 3.5. Then we have*

$$a(\nu, X, B) \leq a(\nu, Y, B_Y + E)$$

for every divisor ν over X . Thus, we obtain

$$R(X/Z, K_X + B) \simeq R(Y/Z, K_Y + B_Y + E).$$

Proof. It is an easy consequence of the negativity lemma. See, for example, [KM, Proposition 3.51 and Theorem 3.52]. \square

Lemma 3.9. *Let $\pi : X \rightarrow Z$ be a projective surjective morphism between projective varieties. Assume that (X, B) is log canonical and H is an ample Cartier divisor on Z . Let R be a $(K_X + B)$ -negative extremal ray of $\overline{NE}(X)$ such that*

$$R \cdot (K_X + B + (2 \dim X + 1)\pi^*H) < 0.$$

Then R is a $(K_X + B)$ -negative extremal ray of

$$\overline{NE}(X/Z) = \{z \in \overline{NE}(X) \mid z \cdot \pi^*H = 0\} \subset \overline{NE}(X).$$

In particular, if $K_X + B$ is π -nef, then $K_X + B + (2 \dim X + 1)\pi^*H$ is nef.

Proof. If (X, B) is klt, then it is obvious by Kawamata's bound of the length of extremal rays (see [K4]). When (X, B) is lc, it is sufficient to use [F4, Subsection 3.1.3] or [F7, Section 18]. \square

Let us start the proof of Theorem 1.1.

Proof of Theorem 1.1. We can assume that the fiber of π is connected. First, if $\kappa(X_\eta, K_{X_\eta} + B_\eta) = -\infty$, where η is the generic point of Z , then the statement is trivial. We note that the statement is obvious when Z is a point and $\kappa(X, K_X + B) = 0$. So, we can assume that $\kappa(X_\eta, K_{X_\eta} + B_\eta) \geq 0$ and that $\kappa(X, K_X + B) \geq 1$ when Z is a point. Since the problem is local, we can assume that Z is affine. By compactifying Z and X and taking a resolution of X , we can assume that X and Z are projective and that $\text{Supp} B$ is a simple normal crossing divisor. By the assumption, we can find an effective \mathbb{Q} -divisor M on X such that $K_X + B \sim_{\mathbb{Q}, \pi} M$, that is, there exists a \mathbb{Q} -divisor N on Z such that $K_X + B \sim_{\mathbb{Q}} M + \pi^*N$. We take a log minimal model of (X, B) over Z by using the arguments in [B, Section 3]. Then we obtain a projective surjective morphism $\pi_Y : Y \rightarrow Z$ such that $(Y/Z, B_Y + \sum_j E_j)$ is a log minimal model of $(X/Z, B)$, where B_Y is the pushforward of B on Y by $\phi : X \dashrightarrow Y$ and E_j is exceptional over X and is a prime divisor on Y for every j . Let A be a sufficiently ample general Cartier divisor on Z . Then $(Y, B_Y + E + \pi_Y^*A)$, where $E = \sum_j E_j$, is a log minimal model of $(X, B + \pi^*A)$ by Lemma 3.9. Since $\kappa(Y, K_Y + B_Y + E + \pi_Y^*A) \geq 1$, $K_Y + B_Y + E + \pi_Y^*A$ is semi-ample by Theorem 3.3. In particular,

$$K_Y + B_Y + E = K_Y + B_Y + E + \pi_Y^*A - \pi_Y^*A$$

is π_Y -semi-ample. Thus,

$$R(Y/Z, K_Y + B_Y + E) = \bigoplus_{m \geq 0} \pi_{Y*} \mathcal{O}_Y(\lfloor m(K_Y + B_Y + E) \rfloor)$$

is a finitely generated \mathcal{O}_Z -algebra. Therefore,

$$R(X/Z, K_X + B) = \bigoplus_{m \geq 0} \pi_* \mathcal{O}_X(\lfloor m(K_X + B) \rfloor)$$

is a finitely generated \mathcal{O}_Z -algebra by Lemma 3.8. We finish the proof. \square

The final theorem in this section is a generalization of Fukuda's theorem (see Theorem 3.3).

Theorem 3.10 (A special case of the log abundance theorem). *Let $\pi : X \rightarrow Z$ be a proper surjective morphism from a four-dimensional log canonical pair (X, B) such that B is an effective \mathbb{Q} -divisor and that $K_X + B$ is π -nef. When Z is a point, we further assume that $\kappa(X, K_X + B) > 0$. Then $K_X + B$ is π -semi-ample.*

Proof. Without loss of generality, we can assume that π has connected fibers. By Proposition 3.2 and the log abundance theorem in dimension ≤ 3 , $K_{X_\eta} + B_\eta$ is nef and abundant, where η is the generic point of Z . By Theorem 1.2, $\bigoplus_{m \geq 0} \pi_* \mathcal{O}_X(\lfloor m(K_X + B) \rfloor)$ is a finitely generated \mathcal{O}_Z -algebra. Therefore, $K_X + B$ is π -semi-ample by Lemma 3.12 below. \square

The next lemma is well known. We leave the proof as an exercise for the reader.

Lemma 3.11 (cf. [L, Theorem 2.3.15]). *Let $\pi : X \rightarrow Z$ be a projective surjective morphism from a smooth variety X to a normal variety Z and M a π -nef and π -big Cartier divisor on X . Then $\bigoplus_{m \geq 0} \pi_* \mathcal{O}_X(mM)$ is a finitely generated \mathcal{O}_Z -algebra if and only if M is π -semi-ample.*

By [KMM, Proposition 6-1-3], we can reduce Lemma 3.12 to Lemma 3.11.

Lemma 3.12. *Let $\pi : X \rightarrow Z$ be a proper surjective morphism between normal varieties and M a π -nef and π -abundant Cartier divisor on X . Then $\bigoplus_{m \geq 0} \pi_* \mathcal{O}_X(mM)$ is a finitely generated \mathcal{O}_Z -algebra if and only if M is π -semi-ample.*

3.1. Appendix. In this appendix, we explicitly state the results in dimension ≤ 3 because we can find no good references for the relative statements (cf. [Ft], [KeMM], and [KeMM2]).

Theorem 3.13. *Let $\pi : X \rightarrow Z$ be a proper surjective morphism between normal varieties. Assume that (X, B) is log canonical with $\dim X \leq 3$ and that B is an effective \mathbb{Q} -divisor. Then*

$$\bigoplus_{m \geq 0} \pi_* \mathcal{O}_X(\lfloor m(K_X + B) \rfloor)$$

is a finitely generated \mathcal{O}_Z -algebra.

Proof. When Z is a point, this theorem is well known (cf. [Ft], [KeMM], and [KeMM2]). So, we assume that $\dim Z \geq 1$. By the arguments in the proof of Theorem 1.1, we can prove that $\bigoplus_{m \geq 0} \pi_* \mathcal{O}_X(\lfloor m(K_X + B) \rfloor)$ is a finitely generated \mathcal{O}_Z -algebra. \square

Theorem 3.14. *Let $\pi : X \rightarrow Z$ be a proper surjective morphism such that (X, B) is log canonical with $\dim X \leq 3$. Assume that $K_X + B$ is π -nef and B is an effective \mathbb{Q} -divisor. Then $K_X + B$ is π -semi-ample.*

Proof. When Z is a point, this theorem is well known (cf. [Ft], [KeMM], and [KeMM2]). So, we assume that $\dim Z \geq 1$. Without loss of generality, we can assume that π has connected fibers. It is well known that $K_X + B$ is π -nef and π -abundant by the log abundance theorem in dimension ≤ 2 . By Theorem 3.13 and Lemma 3.12, $K_X + B$ is π -semi-ample. \square

We close this appendix with a remark.

Remark 3.15. Let $\pi : X \rightarrow Z$ be a proper surjective morphism between normal varieties. Assume that (X, B) is klt and that B is an effective \mathbb{Q} -divisor. Then

$$\bigoplus_{m \geq 0} \pi_* \mathcal{O}_X(\lfloor m(K_X + B) \rfloor)$$

is a finitely generated \mathcal{O}_Z -algebra by [BCHM]. Therefore, by Lemma 3.12, $K_X + B$ is π -semi-ample if and only if $K_X + B$ is π -nef and π -abundant by Lemma 3.12.

Of course, we know that $K_X + B$ is π -semi-ample if and only if $K_X + B$ is π -nef and π -abundant without appealing [BCHM]. See, for example, [F5]. It is known as Kawamata's theorem (cf. [K2]).

4. ABUNDANCE THEOREM FOR IRREGULAR VARIETIES

In this section, we treat the abundance conjecture for irregular varieties. Let us recall the following minimal model conjecture.

Conjecture 4.1 (Minimal model conjecture). *Let X be a smooth projective variety. Assume that K_X is pseudo-effective. Then there exists a normal projective variety X' which satisfies the following conditions:*

- (i) X' is birationally equivalent to X .
- (ii) X' has only \mathbb{Q} -factorial terminal singularities.
- (iii) $K_{X'}$ is nef.

We call X' a minimal model of X .

In Conjecture 4.1, if $K_{X'}$ is semi-ample, X' is usually called a *good minimal model* of X .

Conjecture 4.2 (Abundance conjecture). *Let X be a projective variety with only canonical singularities. Assume that K_X is nef. Then K_X is semi-ample. In particular, $\kappa(X) = \kappa(X, K_X)$ is non-negative.*

We know that Conjectures 4.1 and 4.2 hold in dimension ≤ 3 (cf. [KMM], [FA], etc.).

Remark 4.3. In Conjecture 4.1, by [BCHM], we can replace (ii) with the following slightly weaker condition: (ii') X' has at most canonical singularities. Similarly, we can assume that X has only \mathbb{Q} -factorial terminal singularities in Conjecture 4.2.

Remark 4.4. Let X be a smooth projective variety. Then X is uniruled if and only if K_X is not pseudo-effective by [BDPP].

The next theorem is the main theorem of this section.

Theorem 4.5 (Abundance theorem for irregular $(n+1)$ -folds). *Assume that Conjectures 4.1 and 4.2 hold in dimension $\leq n$. Let X be a normal complete $(n+1)$ -fold with only canonical singularities. If K_X is nef and $q(X) \neq 0$, then K_X is semi-ample.*

Proof. Let $\pi : \bar{X} \rightarrow X$ be a resolution and $\alpha : \bar{X} \rightarrow A = \text{Alb}(\bar{X})$ the Albanese mapping. By the assumption, we have $\dim A \geq 1$. Since X has only rational singularities, $\beta = \alpha \circ \pi^{-1} : X \rightarrow A$ is a morphism (cf. [R, Proposition 2.3], [BS, Lemma 2.4.1])

Claim 1. *We have $\kappa(X, K_X) = \kappa(\bar{X}, K_{\bar{X}}) \geq 0$.*

Proof of Claim 1. Let $f : \bar{X} \rightarrow S$ be the Stein factorization of α and F a general fiber of f . Then, by [K3, Corollary 1.2], we have

$$\kappa(\bar{X}, K_{\bar{X}}) \geq \kappa(F, K_F) + \kappa(\bar{S}, K_{\bar{S}}),$$

where \bar{S} is a resolution of S . We note that $\kappa(\bar{S}, K_{\bar{S}}) \geq 0$ because $S \rightarrow \beta(X) \subset A$ is generically finite (see, for example, [U, Theorem 6.10, Lemma 10.1]). We also note that $\kappa(F, K_F) = \kappa(G, K_G) \geq 0$ since $\dim G \leq n$, G has only canonical singularities, and K_G is nef, where $G = \pi(F)$. Here, we used Conjectures 4.1 and 4.2 in dimension $\dim G = \dim F \leq n$. Therefore, we obtain $\kappa(\bar{X}, K_{\bar{X}}) \geq 0$. \square

Claim 2. *If $\kappa(X, K_X) = 0$, then $\nu(X, K_X) = 0$.*

Proof of Claim 2. By Kawamata's theorem (cf. [K1, Theorem 1]), β is surjective and $\beta_*\mathcal{O}_X \simeq \mathcal{O}_A$. Let G be a general fiber of β . Then $\kappa(G, K_G) = 0$ by

$$0 = \kappa(X, K_X) \geq \kappa(G, K_G) + \kappa(A, K_A) = \kappa(G, K_G)$$

as in Claim 1 and $\kappa(G, K_G) \geq 0$ by Conjecture 4.2 in $\dim G \leq n$ since K_G is nef. We note that X and G have only canonical singularities. By Remark 3.15 and the assumption: Conjecture 4.2 in dimension $\leq n$, K_X is β -semi-ample. Therefore, $\beta : X \rightarrow A$ can be written as follows:

$$\beta : X \xrightarrow{f} S \xrightarrow{g} A,$$

where $K_X \sim_{\mathbb{Q}} f^*D$ for some g -ample \mathbb{Q} -Cartier \mathbb{Q} -divisor D on S , $g : S \rightarrow A$ is a birational morphism, and S is a normal variety. Since $\kappa(X, K_X) = 0$, we obtain $\kappa(S, D) = 0$. So, it is sufficient to prove that $D \sim_{\mathbb{Q}} 0$. By [A, Theorem 0.2], we can write $D \sim_{\mathbb{Q}} K_S + \Delta_S$ such that (S, Δ_S) is klt. In particular, Δ_S is effective. By Lemma 4.6 below, we obtain that g is an isomorphism. Therefore, $D \sim_{\mathbb{Q}} 0$ since $\kappa(S, D) = 0$ and $S = A$ is an Abelian variety. \square

By Claim 1 and Claim 2, $\nu(X, K_X) > 0$ implies $\kappa(X, K_X) > 0$. In this case, we obtain $\kappa(X, K_X) = \nu(X, K_X)$ by Kawamata's argument and the assumption: Conjectures 4.1 and 4.2 in dimension $\leq n$ (see the proof of [K2, Theorem 7.3]). Therefore, K_X is semi-ample by Remark 3.15. \square

We already used the following lemma in the proof of Claim 2.

Lemma 4.6. *Let $g : S \rightarrow A$ be a projective birational morphism from a klt pair (S, Δ_S) to an Abelian variety A . Assume that $K_S + \Delta_S$ is g -nef and $\kappa(S, K_S + \Delta_S) = 0$. Then g is an isomorphism.*

Proof. By replacing S with its small projective \mathbb{Q} -factorialization (cf. [BCHM]), we can assume that S is \mathbb{Q} -factorial. We note that $K_S = E$, where E is effective and $\text{Supp}E = \text{Exc}(g)$ since A is an Abelian variety. If $B = g_*\Delta_S \neq 0$, then $g^*B \leq m(K_S + \Delta_S)$ for some $m > 0$. In this case,

$$1 \leq \kappa(A, B) = \kappa(S, g^*B) \leq \kappa(S, K_S + \Delta_S) = 0.$$

It is a contradiction. Therefore, $B = 0$. This means that Δ_S is g -exceptional. Thus, $K_S + \Delta_S$ is effective, g -exceptional, and $\text{Exc}(g) = \text{Supp}(K_S + \Delta_S)$. By the assumption, $K_S + \Delta_S$ is g -nef. So, g is an isomorphism by the negativity lemma. \square

As a special case of Theorem 4.5, we obtain the abundance theorem for irregular fourfolds.

Corollary 4.7 (Abundance theorem for irregular fourfolds). *Let X be a normal complete fourfold with only canonical singularities. Assume that K_X is nef and the irregularity $q(X)$ is not zero. Then K_X is semi-ample.*

Proof. It is obvious by Theorem 4.5 because Conjectures 4.1 and 4.2 hold in dimension ≤ 3 (cf. [FA], [KM], etc.). \square

We close this section with the following theorem.

Theorem 4.8 (Good minimal models of irregular fourfolds). *Let X be a smooth projective irregular fourfold. If X is not uni-ruled, then X has a good minimal model. More precisely, there exists a normal projective variety X' such that X' has only \mathbb{Q} -factorial terminal singularities, X' is birationally equivalent to X , and $K_{X'}$ is semi-ample.*

Proof. We run the minimal model program. Then we obtain a minimal model X' of X since K_X is pseudo-effective by the assumption. Here, we used the existence and the termination of four-dimensional terminal flips (cf. [KMM, Theorem 5-1-15], [S1], and [HM, Corollary 5.1.2]). We note that $q(X') = h^1(X', \mathcal{O}_{X'}) = q(X) \neq 0$. Therefore, by Theorem 4.5, we obtain that $K_{X'}$ is semi-ample. \square

4.1. Appendix. In this appendix, we give a remark on the abundance conjecture for fourfolds for the reader's convenience.

Conjecture 4.9 (Abundance conjecture for fourfolds). *Let X be a complete fourfold with only canonical singularities. If K_X is nef, then K_X is semi-ample.*

This conjecture is still open. By Corollary 4.7 and Kawamata's argument (cf. [K2, Theorem 7.3]), we can reduce Conjecture 4.9 to the following two problems.

Problem 4.10. Let X be a smooth projective fourfold. If X is not uni-ruled and $q(X) = 0$, then $\kappa(X) \geq 0$.

Problem 4.11. Let X be a projective fourfold with only \mathbb{Q} -factorial terminal singularities. If K_X is nef, $q(X) = 0$, and $\kappa(X, K_X) = 0$, then K_X is numerically trivial, equivalently, $K_X \sim_{\mathbb{Q}} 0$.

We explain the reduction argument closely. Let X be a complete fourfold with only canonical singularities such that K_X is nef. If $q(X) \neq 0$, then K_X is semi-ample by Corollary 4.7. So, from now on, we can assume that $q(X) = 0$. By taking a resolution of X and running the minimal model program (cf. [KMM, Theorem 5-1-15], [S1], and [HM,

Corollary 5.1.2]), there exists a projective variety X' such that $K_{X'}$ is nef and that X' has only \mathbb{Q} -factorial terminal singularities. Let

$$X \xleftarrow{f} W \xrightarrow{g} X'$$

be a common resolution. Then $f^*K_X = g^*K_{X'}$ by the negativity lemma. Therefore, we can replace X with X' in order to prove Conjecture 4.9. If we solve Problem 4.10, then we obtain $\kappa(X, K_X) \geq 0$ since X has only terminal singularities. Furthermore, if we solve Problem 4.11, then we can prove that $\nu(X, K_X) > 0$ implies $\kappa(X, K_X) > 0$. By the proof of [K2, Theorem 7.3], we obtain $\nu(X, K_X) = \kappa(X, K_X)$ (cf. Proposition 3.2). Thus, K_X is semi-ample (cf. Remark 3.15).

REFERENCES

- [A] F. Ambro, The moduli b -divisor of an lc-trivial fibration, *Compos. Math.* **41** (2005), no. 2, 385–403.
- [BS] M. Beltrametti, A. Sommese, *The adjunction theory of complex projective varieties*, de Gruyter Expositions in Mathematics, **16**. Walter de Gruyter & Co., Berlin, 1995.
- [B] C. Birkar, On existence of log minimal models, preprint 2007.
- [BCHM] C. Birkar, P. Cascini, C. Hacon, and J. McKernan, Existence of minimal models for varieties of log general type, preprint 2006.
- [BDPP] S. Boucksom, J.-P. Demailly, M. Paun, and T. Peternell, The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension, preprint 2004.
- [F1] O. Fujino, Abundance theorem for semi log canonical threefolds, *Duke Math. J.* **102** (2000), no. 3, 513–532.
- [F2] O. Fujino, What is log terminal?, 49–62 in *Flips for 3-folds and 4-folds*, Oxford University Press (2007).
- [F3] O. Fujino, Base point free theorems—saturation, b -divisors, and canonical bundle formula—, preprint 2007.
- [F4] O. Fujino, Introduction to the log minimal model program for log canonical pairs, preprint 2008.
- [F5] O. Fujino, On Kawamata’s theorem, preprint 2009.
- [F6] O. Fujino, On injectivity, vanishing and torsion-free theorems for algebraic varieties, to appear in *Proc. Japan Acad. Ser. A Math. Sci.* **85** (2009).
- [F7] O. Fujino, Fundamental theorems for the log minimal model program, preprint 2009.
- [Ft] T. Fujita, Fractionally logarithmic canonical rings of surfaces, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **30** (1984), no. 3, 685–696.
- [Fk] S. Fukuda, On numerically effective log canonical divisors, *Int. J. Math. Math. Sci.* **30** (2002), no. 9, 521–531.
- [HM] C. Hacon, J. McKernan, Extension theorems and the existence of flips, in *Flips for 3-folds and 4-folds*, 76–110, Oxford Lecture Ser. Math. Appl., **35**, Oxford Univ. Press, Oxford, 2007.

- [K1] Y. Kawamata, Characterization of abelian varieties, *Compositio Math.* **43** (1981), no. 2, 253–276.
- [K2] Y. Kawamata, Pluricanonical systems on minimal algebraic varieties, *Invent. Math.* **79** (1985), no. 3, 567–588.
- [K3] Y. Kawamata, Minimal models and the Kodaira dimension of algebraic fiber spaces, *J. Reine Angew. Math.* **363** (1985), 1–46.
- [K4] Y. Kawamata, On the length of an extremal rational curve, *Invent. Math.* **105** (1991), no. 3, 609–611.
- [KMM] Y. Kawamata, K. Matsuda, and K. Matsuki, Introduction to the minimal model problem, *Algebraic geometry, Sendai, 1985*, 283–360, *Adv. Stud. Pure Math.*, **10**, North-Holland, Amsterdam, 1987.
- [KeMM] S. Keel, K. Matsuki, and J. McKernan, Log abundance theorem for threefolds, *Duke Math. J.* **75** (1994), no. 1, 99–119.
- [KeMM2] S. Keel, K. Matsuki, and J. McKernan, Corrections to “Log abundance theorem for threefolds”, *Duke Math. J.* **122** (2004), no. 3, 625–630.
- [FA] J. Kollár et al., *Flips and Abundance for algebraic threefolds*, *Astérisque* **211**, (1992).
- [KM] J. Kollár, and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, Vol. **134**, 1998.
- [L] R. Lazarsfeld, *Positivity in algebraic geometry. I. Classical setting : line bundles and linear series*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, **48**. Springer-Verlag, Berlin, 2004.
- [R] M. Reid, Projective morphisms according to Kawamata, preprint, Univ. of Warwick 1983.
- [S1] V. V. Shokurov, Prelimiting flips, *Tr. Mat. Inst. Steklova* **240** (2003), *Biratsion. Geom. Linein. Sist. Konechno Porozhdennye Algebr*, 82–219; translation in *Proc. Steklov Inst. Math.* 2003, no. 1 (240), 75–213.
- [S2] V. V. Shokurov, Letters of a bi-rationalist VII. Ordered termination, preprint 2006.
- [U] K. Ueno, *Classification theory of algebraic varieties and compact complex spaces*, Notes written in collaboration with P. Cherenack. *Lecture Notes in Mathematics*, Vol. **439**. Springer-Verlag, Berlin-New York, 1975.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KYOTO UNIVERSITY,
KYOTO 606-8502 JAPAN

E-mail address: fujino@math.kyoto-u.ac.jp