

Caffeine activates preferentially α 1-isoform of 5'AMP-activated protein kinase in rat skeletal muscle

Tatsuro Egawa¹, Taku Hamada², Xiao Ma¹, Kouhei Karaike¹, Naoko Kameda¹, Shinya Masuda¹, Nobumasa Iwanaka¹, and Tatsuya Hayashi¹

1 Laboratory of Sports and Exercise Medicine, Graduate School of Human and Environmental Studies, Kyoto University, Kyoto, 606-8501, Japan

2 Department of Health and Sports Management, Osaka University of Health and Sport Sciences, Osaka, 590-0496, Japan

Running head: CAFFEINE AND ISOFORM-SPECIFIC AMPK ACTIVATION

Address correspondence to:

Tatsuya Hayashi, M.D., Ph.D.

Laboratory of Sports and Exercise Medicine, Graduate School of Human and Environmental Studies, Kyoto University, Yoshida-nihonmatsu-cho, Sakyo-ku, Kyoto, 606-8501, Japan

Telephone, Fax: +81-75-753-6640, E-mail: tatsuya@kuhp.kyoto-u.ac.jp

Abstract

Aim: Caffeine activates 5'AMP-activated protein kinase (AMPK), a signalling intermediary implicated in the regulation of glucose, lipid, and energy metabolism in skeletal muscle. Skeletal muscle expresses two catalytic α subunits of AMPK, $\alpha 1$ and $\alpha 2$, but the isoform specificity of caffeine-induced AMPK activation is unclear. The aim of this study was to determine which α isoform is preferentially activated by caffeine in vitro and in vivo using rat skeletal muscle.

Methods: Rat epitrochlearis muscle was isolated and incubated in vitro in the absence or presence of caffeine. In another experiment, the muscle was dissected after intravenous injection of caffeine. Isoform-specific AMPK activity, the phosphorylation status of AMPK α Thr¹⁷² and acetyl-CoA carboxylase (ACC) Ser⁷⁹, the concentrations of ATP, phosphocreatine (PCr), and glycogen, and 3-O-methyl-D-glucose (3MG) transport activity were estimated.

Results: Incubation of isolated epitrochlearis muscle with 1 mM of caffeine for 15 min increased AMPK $\alpha 1$ activity, but not AMPK $\alpha 2$ activity; concentrations of ATP, PCr and glycogen were not affected. Incubation with 3 mM of caffeine activated AMPK $\alpha 2$ and reduced PCr and glycogen concentrations. Incubation with 1 mM of caffeine increased the phosphorylation of AMPK and ACC and enhanced 3MG transport. Intravenous injection of caffeine (5 mg kg⁻¹) predominantly activated AMPK $\alpha 1$ and increased 3MG transport without affecting energy status.

Conclusion: Our results suggest that of the two α isoforms of AMPK, AMPK $\alpha 1$ is predominantly activated by caffeine via an energy-independent mechanism and that activation of AMPK $\alpha 1$ increases glucose transport and ACC phosphorylation in

skeletal muscle.

Keywords: acetyl-CoA carboxylase, glucose transport, phosphocreatine, glycogen,
Ca²⁺/calmodulin-dependent protein kinase

Introduction

5'AMP-activated protein kinase (AMPK) is emerging as a signalling intermediary that controls the use of glucose and fatty acids in skeletal muscle. AMPK has been identified as part of the mechanism responsible for exercise-stimulated insulin-independent glucose transport (reviewed in Kahn et al., 2005, Hardie et al. 2006, Fujii et al. 2006) and is also implicated in GLUT4 expression (Zheng et al. 2001, Nakano et al. 2006), glycogen regulation (Jørgensen et al. 2004a, Miyamoto et al. 2007), fatty acid oxidation (Winder and Hardie, 1996, Vavvas et al. 1997), mitochondrial biogenesis (Jager et al. 2007, Garcia-Roves et al. 2008), and enhanced insulin sensitivity (Fiedler et al. 2001, Nakano et al. 2006, Tanaka et al. 2007).

Caffeine (1,3,7-trimethylxanthine) is a xanthine alkaloid compound that acts as a stimulant of skeletal muscle metabolism. Caffeine increases the rate of glucose transport in the absence of insulin in rodent skeletal muscles (Wright et al. 2004, Cantó et al. 2006, Jensen et al. 2007a, Egawa et al. 2009), enhances GLUT4 mRNA and protein expression in cultured myotubes (Ojuka et al. 2002, Mukwevho et al. 2008), and promotes fatty acid metabolism in perfused rat skeletal muscles (Raney and Turcotte, 2008). AMPK is considered part of the mechanism leading to the metabolic activation by caffeine (Wright et al. 2004, Jensen et al. 2007a, Raney and Turcotte 2008, Egawa et al. 2009).

AMPK is a heterotrimeric kinase comprising a catalytic α subunit and two regulatory subunits, β and γ . Two distinct α isoforms ($\alpha 1$ and $\alpha 2$) exist in skeletal muscle, and the Thr¹⁷² residue in both $\alpha 1$ and $\alpha 2$ catalytic subunits is the primary phosphorylation site responsible for AMPK activation. We demonstrated recently

that caffeine stimulation (≥ 3 mM for ≥ 15 min) increases AMPK α Thr¹⁷² phosphorylation with corresponding increases in both AMPK $\alpha 1$ and $\alpha 2$ activities in incubated rat epitrochlearis and soleus muscles (Egawa et al. 2009). This activation is accompanied by an increased rate of insulin-independent glucose transport, enhanced phosphorylation of acetyl-coenzyme A carboxylase (ACC) Ser⁷⁹, and decreased phosphocreatine (PCr) concentration (Egawa et al. 2009). These results suggest that caffeine and exercise have similar actions in stimulating skeletal muscle AMPK by reducing the intracellular energy status.

In contrast, Jensen et al. (Jensen et al. 2007a) demonstrated that caffeine stimulation (3 mM for 15 min) acutely stimulates AMPK $\alpha 1$ activity but not $\alpha 2$ activity in incubated mouse and rat soleus muscles. Interestingly, Jensen et al. did not detect any changes in energy status in mouse soleus muscle. Similarly, we previously demonstrated that AMPK $\alpha 1$, but not AMPK $\alpha 2$, is activated in rat epitrochlearis muscle incubated and treated with H₂O₂ and hypoxanthine/xanthine oxidase in the absence of an increase in AMP or a decrease in PCr concentration (Toyoda et al. 2006). We also showed that AMPK $\alpha 1$ is activated in low-intensity contracting muscle, in which AMP concentration is not elevated, whereas AMPK $\alpha 1$ and AMPK $\alpha 2$ are activated in high-intensity contracting muscle, in which the AMP concentration is significantly higher than the resting value (Toyoda et al. 2006).

AMPK is activated in response to energy-depleting stresses such as muscle contraction, hypoxia, and inhibition of oxidative phosphorylation (Hayashi et al. 2000). AMPK $\alpha 2$ is more dependent on AMP than is AMPK $\alpha 1$ with respect to allosteric activation by AMP and covalent activation by upstream kinases (Salt et al. 1998, Stein et al. 2000), indicating that AMPK $\alpha 2$ is more sensitive to energy

depletion than is AMPK α 1. We hypothesized that caffeine can act on skeletal muscle and preferentially stimulate AMPK α 1 in the absence of energy deprivation. To test this hypothesis, we investigated the effects of caffeine on AMPK α 1 and α 2 activities in rat skeletal muscles stimulated by caffeine in vitro and in vivo.

Materials and Methods

Caffeine treatment in vitro

Male Sprague-Dawley rats weighing 100 g were obtained from Shimizu Breeding Laboratories (Kyoto, Japan). Animals were housed in an animal room maintained at 22-24°C with a 12:12-h light-dark cycle and fed a standard laboratory diet (Certified Diet MF; Oriental Koubo, Tokyo, Japan) and water ad libitum. Rats were fasted overnight before the experiments and were randomly assigned to the experimental groups. All protocols for animal use and euthanasia followed the Guiding Principles for the Care and Use of Animals in the Field of Physiological Sciences (Physiological Society of Japan) in accordance with international guidelines, and were reviewed and approved by the Kyoto University Graduate School of Human and Environmental Studies and Kyoto University Radioisotope Research Center.

Muscles were treated as we described previously (Hayashi et al. 1998, Toyoda et al. 2004). Rats were sacrificed by cervical dislocation without anaesthesia, and the epitrochlearis muscles of each side were rapidly removed. Both ends of each muscle were tied with sutures (silk 3-0; Natsume Seisakusho, Tokyo, Japan) and the muscles were mounted on an incubation apparatus with a tension set to 0.5 g. The buffers were continuously gassed with 95% O₂-5% CO₂ and maintained at 37°C. Muscles were preincubated in 7 mL of Krebs-Ringer bicarbonate buffer (KRB) (117 mM NaCl, 4.7 mM KCl, 2.5 mM CaCl₂, 1.2 mM KH₂PO₄, 1.2 mM MgSO₄, 24.6 mM NaHCO₃) containing 2 mM pyruvate (KRBP) for 40 min. The muscles were then incubated for 15 min in 7 ml of fresh buffer containing various concentrations (0-3

mM) of caffeine. The muscles were then used for the measurement of glucose transport, or immediately frozen in liquid nitrogen and subsequently analyzed for ATP, PCr, and glycogen concentrations, and isoform-specific AMPK activity, or used for Western blot analysis.

Caffeine treatment in vivo

We chose male Sprague-Dawley rats weighing 200 g for easy administration of caffeine into the tail vein without anaesthesia. For the dose-response changes in AMPK and ACC phosphorylation, caffeine was dissolved in saline and injected intravenously at various concentrations (0-25 mg kg⁻¹ body weight). The injection volume was 1 ml kg⁻¹ body weight. Sixty minutes after the injection rats were anaesthetized with intraperitoneal administration of pentobarbital sodium (50 mg kg⁻¹ body weight), and then epitrochlearis muscle was rapidly dissected and immediately frozen in liquid nitrogen. Some controls and samples with 5 mg kg⁻¹ caffeine stimulation were subjected to the measurement of isoform-specific AMPK activity and glucose transport. For the time-course changes in AMPK and ACC phosphorylation, caffeine (5 mg kg⁻¹ body weight) was injected intravenously into the tail vein without anaesthesia. Muscles were dissected under anaesthesia at 30, 60 and 120 min after injection, and then immediately frozen in liquid nitrogen. Control samples were collected immediately after saline injection under anaesthesia. Some controls and samples dissected at 60 min after caffeine injection were subjected to the measurement of ATP, PCr, and glycogen concentrations.

Blood sample analysis

Blood samples were collected from the tail vein using heparinized glass tube at 0 and 60 min after intravenous saline or caffeine injection. Plasma levels for glucose (Glutest-Ace; Sanwa Kagaku Kenkyusyo, Nagoya, Japan), insulin (rat insulin ELISA kit; Morinaga, Yokohama, Japan), caffeine (caffeine/pentoxifylline ELISA kit; Neogen, Lexington, KY, USA) were measured.

Western blot analysis

Sample preparation and Western blot analysis was performed as we described previously (Toyoda et al. 2004). Muscles were homogenized in ice-cold lysis buffer (1:40 wt vol⁻¹) containing 20 mM Tris·HCl (pH 7.4), 1% Triton X, 50 mM NaCl, 250 mM sucrose, 50 mM NaF, 5 mM sodium pyrophosphate, 2 mM dithiothreitol, 4 mg L⁻¹ leupeptin, 50 mg L⁻¹ trypsin inhibitor, 0.1 mM benzamidine, and 0.5 mM phenylmethylsulfonyl fluoride and centrifuged at 16,000 g for 40 min at 4°C. Lysates were solubilized in Laemmli's sample buffer containing mercaptoethanol and boiled.

The samples (10 µg of protein) were separated on either 10% polyacrylamide gel for AMPK, Ca²⁺/calmodulin-dependent protein kinase I (CaMKI), Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) or 7.5% gel for ACC. Proteins were then transferred to polyvinylidene difluoride membranes (PolyScreen; PerkinElmer, Wellesley, MA, USA) at 100 V for 1 h. Membranes were blocked for 1h at room temperature in TBS-T (TBS with 0.1% Tween 20) containing 5% nonfat dry milk and were then incubated over night at 4°C with phosphospecific AMPKα Thr¹⁷² (#2531; Cell Signaling Technology, Danvers, MA, USA), AMPKα (#2532; Cell Signaling Technology), CaMKII Thr²⁸⁶ (287) (#3361; Cell Signaling Technology,

Danvers, MA, USA), phosphospecific ACC Ser⁷⁹ (#07-303; Upstate Biotechnology, Lake Placid, NY, USA), ACC (#3662; Cell Signaling Technology) diluted 1:1000 or CaMKI Thr¹⁷⁷ (sc-28438-R; Santa Cruz Biotechnology, Santa Cruz, CA, USA) diluted 1:250. The membranes were then washed, incubated for 1h at room temperature with anti-mouse IgG antibody (GE Healthcare, Buckinghamshire, UK) diluted 1:2500, and developed with enhanced chemiluminescence reagents according to the manufacturer's instructions (Amersham Biosciences, Buckinghamshire, UK). The intensity of the signals was quantified using Multi-Analyst software (Bio-Rad, Hercules, CA, USA). The mean intensity of control samples in each membrane was used as reference for controlling gel-to-gel variation. Equal protein loading and transfer was confirmed by Coomassie brilliant blue staining of the membranes.

Isoform-specific AMPK activity assay

We have raised AMPK polyclonal antibodies in rabbit against isoform-specific peptides derived from the amino acid sequences of rat $\alpha 1$ (residues 339–358) or $\alpha 2$ (residues 490–514) (Toyoda et al. 2004). AMPK activity assay was performed as we described previously (Hayashi et al. 2000, Toyoda et al. 2004). Muscles were homogenized as described in *Western blot analysis*, and resultant supernatants (100 μ g of protein) were immunoprecipitated with the $\alpha 1$ or $\alpha 2$ AMPK antibody and protein A-Sepharose beads (Amersham Biosciences, Uppsala, Sweden). Immunoprecipitates were washed twice both in lysis buffer and in wash buffer (240 mM HEPES and 480 mM NaCl). Kinase reactions were performed in 40 mM HEPES (pH 7.0), 0.1 mM SAMS peptide (Hayashi et al. 2000, Toyoda et al. 2004), 0.2 mM AMP, 80 mM NaCl, 0.8 mM dithiothreitol, 5 mM MgCl₂, 0.2 mM ATP (2

μCi of [$\gamma\text{-}^{32}\text{P}$] ATP/sample) (PerkinElmer, Wellesley, MA, USA), in a final volume of 40 μl for 20 min at 30°C. At the end of the reaction, a 15- μl aliquot was removed and spotted onto Whatman P81 paper (Whatman International, Maidstone, UK). The papers were washed six times in 1% phosphoric acid and once in acetone. ^{32}P incorporation was quantitated with a scintillation counter, and kinase activity was expressed as fold increases relative to the control samples.

ATP and PCr assay

ATP and PCr concentrations were measured fluorometrically in perchloric acid extracts of epitrochlearis muscle according to the method of Lowry and Passonneau (Lowry and Passonneau, 1972). In brief, each frozen muscle was homogenized in 0.2 M HClO_4 (3:25 w/v) in an ethanol-dry ice bath (-20 ~ -30°C) with careful handling of the homogenate being kept unfrozen. After a centrifuge at 16,000 g for 2 min at -9°C, the supernatant of the homogenate was neutralized with a solution of 2 M KOH, 0.4 M KCl and 0.4 M imidazole and then centrifuged at 16,000 g for 2 min at -9°C, and then subjected to enzymatic analysis (Lowry and Passonneau, 1972). ATP and PCr concentrations were expressed as nanomoles per milligram wet weight of muscle.

Glycogen assay

Glycogen concentration was assayed as we described previously (Nakano et al. 2006, Miyamoto et al. 2007). Each frozen muscle was digested in 1 M NaOH at 85°C for 10 min, and the digestates were neutralized with HCl. The glycogen in the digestates was hydrolyzed by incubated in 2 M HCl for 2 h at 85°C. The digestates

were neutralized with NaOH, and the concentration of hydrolyzed glucose residues was measured enzymatically using Glucose CII Test (Wako, Osaka, Japan). Glycogen concentration was expressed as nanomoles of glucose per milligram wet weight of muscle.

3-O-methyl-D-glucose (3MG) transport

3MG transport assay was performed as we described previously (Hayashi et al. 1998, Toyoda et al. 2004, Nakano et al. 2006). To measure 3MG transport after in vitro caffeine incubation, muscles were transferred to 2 mL of KRB containing 1 mM [³H]3-MG (1.5 μ Ci mL⁻¹) (American Radiolabeled Chemicals, St. Louis, MO) and 7 mM D-[1-¹⁴C] mannitol (0.3 μ Ci mL⁻¹) (American Radiolabeled Chemicals, St. Louis, MO) at 30°C and further incubated for 10 min. To measure 3MG transport after an injection of caffeine, muscles were incubated for 10 min in 7 mL of KRBP. Muscles were then transferred to 2 mL of transport buffer and incubated for 10 min. The muscles were then blotted onto filter paper, trimmed, frozen in liquid nitrogen, and stored at -80°C. Each frozen muscle was weighed and processed by incubating them in 300 μ l of 1 M NaOH at 80°C for 10 min. Digestates were neutralized with 300 μ l of 1 M HCl, and particulates were precipitated by centrifugation at 20,000 g for 2 min. Radioactivity in aliquots of the digested protein was determined by liquid scintillation counting for dual labels, and the extracellular and intracellular spaces were calculated (Young et al. 1986).

Statistical analysis

Results are presented as means \pm SE. One-way ANOVA (Figs 1-7) or two-way

ANOVA (Table 1 and 2) was used to estimate the variance of the dose-response and time-course studies, and statistical significance of difference between control and caffeine-treated groups was evaluated by Scheffe's post hoc test. Student's t test was used to examine the significant differences between control and caffeine-treated groups (Figs 8 and 9). Differences between groups were considered statistically significant at $P < 0.05$.

Results

Caffeine preferentially increased AMPK α 1 activity in skeletal muscle incubated in vitro

Our previous study demonstrated that caffeine stimulation in vitro (3 mM for 15 min) activates both AMPK α 1 and AMPK α 2 in rat epitrochlearis muscle (Egawa et al. 2009). To investigate the dose dependency of isoform-specific AMPK activity stimulated by caffeine at concentrations lower than 3 mM, isolated epitrochlearis muscles were incubated and stimulated with 0.1, 0.5, 1, and 3 mM of caffeine for 15 min. A caffeine concentration of 1 mM clearly increased AMPK α 1 by twofold ($P < 0.05$), but AMPK α 2 activation required a caffeine concentration of 3 mM (Fig. 1).

Activation of AMPK α 1 by caffeine was not associated with energy deprivation

AMPK is activated in response to energy deprivation by allosteric modification and through a mechanism involving phosphorylation of the α subunit by upstream kinases (Hardie and Carling, 1997, Kemp et al. 1999). Our previous study using rat epitrochlearis and soleus muscles demonstrated that in vitro stimulation with 3 mM of caffeine for 15 min decreases PCr concentration by 23% (Egawa et al. 2009). To determine whether caffeine-induced AMPK α 1 activation is associated with energy deprivation, we measured concentrations of ATP, PCr, and glycogen in epitrochlearis muscle after stimulation with caffeine for 15 min. The concentrations of ATP, PCr and glycogen were not affected by stimulation with 1 mM of caffeine, but concentrations of PCr and glycogen were significantly decreased by stimulation with 3 mM of caffeine (Fig. 2).

Activation of AMPK α 1 by caffeine was associated with increased AMPK and

ACC phosphorylation, and insulin-independent glucose transport

To confirm whether caffeine activates AMPK, we measured the degree of phosphorylation of AMPK α Thr¹⁷² by Western blot analysis using a phosphospecific antibody in muscle homogenates after stimulation with caffeine. Incubation with caffeine at 1 and 3 mM clearly increased phosphorylation of AMPK α Thr¹⁷² without changing the total amount of AMPK α (Fig. 3A). ACC is a downstream target of AMPK in skeletal muscle, and the phosphorylation of the Ser⁷⁹ site of ACC reflects the total AMPK activity (Davies et al. 1990, Park et al. 2002). The marked phosphorylation of ACC paralleled the increase in AMPK phosphorylation (Fig. 3B). We also investigated whether the activation of AMPK in skeletal muscle by caffeine is associated with enhanced glucose transport. In the absence of insulin, incubation with 1 mM caffeine increased the rate of 3MG transport by 1.4-fold and incubation with 3 mM increased 3MG transport by 2.3-fold (Fig. 4).

Caffeine at a concentration of 3mM, but not 1mM, enhanced phosphorylation of CaMKII Thr²⁸⁷ in skeletal muscle incubated in vitro

Ca²⁺ has been implicated in the activation of glucose transport through signalling pathways involving AMPK (Wright et al. 2004, Hawley et al. 2005, Wright et al. 2005). CaMKII is directly regulated by Ca²⁺/calmodulin (Hudmon and Schulman, 2002, Witczak et al. 2010), and has been used as an indicator of elevated cytosolic Ca²⁺ level in skeletal muscle (Blair et al. 2009). We found that 3 mM of caffeine, but not 1mM of caffeine, increased phosphorylation of CaMKII Thr²⁸⁷ in incubated muscles (Fig. 5B). Ca²⁺/calmodulin kinase kinase (CaMKK) has been identified as an AMPK kinase in skeletal muscle (Jensen et al. 2007a, Witczak et al. 2007); CaMKI is an endogenous substrate of CaMKK (Tokumitsu et al. 1995). Incubation

with caffeine did not affect the phosphorylation status of CaMKI Thr¹⁷⁷ (Fig. 5A).

Intravenous caffeine injection increased phosphorylation of AMPK α Thr¹⁷² and ACC Ser⁷⁹ in a dose- and time-dependent manner

To determine whether caffeine preferentially activates AMPK in vivo, we measured the degree of phosphorylation of AMPK α Thr¹⁷² in muscle homogenates after stimulation with intravenous caffeine at various concentrations (0, 2.5, 5, 10, and 25 mg kg⁻¹) and for various times (0, 30, 60, and 120 min). The dose–response study revealed that phosphorylation of AMPK α Thr¹⁷² increased at caffeine concentrations of 5 mg kg⁻¹ or higher in epitrochlearis muscle (Fig. 6A). Phosphorylation of ACC Ser⁷⁹ displayed a pattern similar to that for AMPK phosphorylation (Fig. 6B). The time-course study showed that phosphorylation of AMPK α Thr¹⁷² and ACC Ser⁷⁹ increased within 60 min of caffeine stimulation in epitrochlearis muscle (Figs 7A and B). The total AMPK and ACC contents of the muscles did not change during the study (Figs 6 and 7).

Intravenous caffeine injection increased AMPK α 1 activity and insulin-independent glucose transport without affecting energy status

To determine which catalytic subunit is activated by intravenous caffeine injection, rats were injected with caffeine (5 mg kg⁻¹), and isoform-specific AMPK activity was determined 60 min later in anti- α 1 and anti- α 2 immunoprecipitates of the muscle. Caffeine increased AMPK α 1 activity by 1.7-fold, but had no effect on AMPK α 2 activity (Fig. 8). We also investigated whether caffeine-mediated in vivo activation of AMPK α 1 in skeletal muscle is associated with enhanced glucose transport. Sixty minutes after the intravenous administration of caffeine, the rate of 3MG transport was increased by 2.0-fold compared with the saline injection group

(Fig. 9). Muscle ATP, PCr, and glycogen concentrations did not differ between the saline and caffeine injection groups (Table 1).

Plasma glucose, insulin, and caffeine levels after intravenous caffeine injection

Intravenous injection of caffeine (5 mg kg^{-1}) did not alter plasma glucose and insulin levels. Plasma caffeine levels 60 min after caffeine (5 mg kg^{-1}) injection were clearly increased compared to the saline injection group (Table 2).

Discussion

Our data show three novel findings relating to the activating effect of caffeine on skeletal muscle AMPK. First, *in vitro* caffeine treatment at 1 mM activated AMPK α 1 but not AMPK α 2, and caffeine treatment at 3 mM stimulated both isoforms (Fig. 1), suggesting that AMPK α 1 is more sensitive to caffeine than is AMPK α 2. The predominant activation of AMPK α 1 was also confirmed by *in vivo* caffeine treatment (Figs 6, 7, and 8). Second, activation of AMPK α 1 by caffeine occurred in the absence of any apparent reduction in muscle fuel status (Fig. 2 and Table 1), indicating that AMPK α 1 is activated by a mechanism, at least in part, distinct from that of AMPK α 2. Third, AMPK α 1 activation was associated with increased ACC phosphorylation (Figs 3, 6, and 7) and glucose transport activity (Fig. 4 and 9), suggesting that AMPK α 2 activity is not essential for metabolic activation in skeletal muscle.

The physiological importance of AMPK α 1 has been elucidated in contraction-stimulated glucose transport using genetic animal models. Jørgensen et al. (Jørgensen et al. 2004b) examined the effect of high-intensity (tetanic) contraction, which markedly stimulated AMPK α 2 activity in muscles of a whole-body AMPK α 1-knockout mouse. Glucose transport activity was 25% lower in soleus but was normal in extensor digitorum longus muscle compared with muscles from the wild type. More recently, Jensen et al. (Jensen et al. 2008) examined the effect of low-intensity (twitch) contraction and found that glucose transport was markedly lower in muscles from AMPK α 1-knockout and kinase-dead AMPK construct (AMPK-KD) mice but not in AMPK α 2-knockout muscles. These

results suggest that AMPK α 1 plays a major role in activation of low-intensity contraction-stimulated glucose transport, and is required for full activation of high-intensity contraction-stimulated glucose transport in skeletal muscle.

It is notable that most human studies using muscle biopsy samples show that moderate-intensity exercise preferentially activates AMPK α 2 (Wojtaszewski et al. 2000, Fujii et al. 2000, Chen et al. 2000, Musi et al. 2001a, Stephens et al. 2002, Wojtaszewski et al. 2002, Wojtaszewski et al. 2003, Birk and Wojtaszewski, 2006, Sriwijitkamol et al. 2007). Similarly, treadmill running (Musi et al. 2001b, Durante et al. 2002, Klein et al. 2007) and electrical stimulation of the sciatic nerve (Vavvas et al. 1997) activate only AMPK α 2 in rodent skeletal muscles. However, we demonstrated previously that AMPK α 1, but not AMPK α 2, is activated immediately as a post-mortem artefact during the dissection procedure (Toyoda et al. 2006). Hardie and Carling (Hardie and Carling, 1997) proposed that any method of cell harvesting that leads to hypoxia or any form of stress is likely to lead a rapid, artefactual activation of AMPK and that rapid cooling is required to preserve the in vivo-activation state of AMPK. Thus, it may be difficult to measure the actual AMPK α 1 activity because it is disturbed by additional activation during dissection; only when the activation exceeds that of the dissecting stimuli, would AMPK α 1 activity be detectable. For caffeine treatment in vivo, we exposed isolated muscles to caffeine after a preincubation period sufficient to stabilize AMPK α 1 activity in oxygenated medium. After caffeine treatment in vivo, we gently and quickly removed muscles from anesthetized rats and immediately froze them in liquid nitrogen. We believe that the muscle preparations used in the present study enabled us to detect modest increases in AMPK α 1 activity in muscles treated with caffeine in

vitro and in vivo.

A study by Jensen et al. (Jensen et al. 2007a) demonstrated the causal relationship between AMPK and caffeine-stimulated glucose transport in skeletal muscle. They found that caffeine-stimulated glucose transport was reduced markedly in soleus muscle isolated from a transgenic mouse with muscle-specific expression of an AMPK-KD construct. However, because both AMPK α 1 and AMPK α 2 activities were suppressed in muscles of the AMPK-KD mouse (Jensen et al. 2007b), their findings do not allow one to draw conclusions about the relative importance of these isoforms to glucose transport regulation. Moreover, in contrast to the wild-type mouse, the AMPK-KD mouse showed significantly higher AMP concentration in soleus muscle after caffeine stimulation (Jensen et al. 2007a), and this result does not confirm the association between energy deprivation and glucose transport activity. Our study demonstrates clearly that AMPK α 1 activation occurs in the absence of AMPK α 2 activation and in the absence of any apparent reduction in muscle energy status, whereas AMPK α 1 and AMPK α 2 activation is accompanied by a significant reduction in muscle energy status.

The finding that caffeine increased AMPK α Thr¹⁷² phosphorylation in skeletal muscle (Figs 3, 6, and 7) provides evidence that caffeine induces covalent modification via an AMPK kinase. Jensen et al. (Jensen et al. 2007b) showed that the CaMKK inhibitor, STO-609, inhibits activation of AMPK α 1 and AMPK α 2 as well as AMPK α Thr¹⁷² phosphorylation in incubated mouse skeletal muscles after a low-intensity tetanic contraction in vitro. Jensen et al. (Jensen et al. 2007a) also showed that caffeine-induced AMPK α 1 activation and 2-deoxyglucose transport in mouse skeletal muscle is blocked by STO-609. However, we did not detect any

increases in the phosphorylation status of CaMKI, a downstream target of CaMKK (Fig. 5A), indicating that CaMKK is not involved in caffeine-induced AMPK activation in skeletal muscle. Moreover, 1 mM of caffeine did not significantly affect the phosphorylation status of CaMKII (Fig. 5B), indicating that Ca^{2+} signalling is not involved in the increase in glucose transport. The LKB1 complex (Sakamoto et al. 2004, Sakamoto et al. 2005) is the main kinase that regulates AMPK α 2 activity in mouse skeletal muscle during tetanic contraction in situ and in vitro (Sakamoto et al. 2005). The LKB1 complex is constitutively active and is not activated directly by AMP; binding of AMP to AMPK facilitates the phosphorylation of AMPK by the LKB1 complex (Hawley et al. 2003, Sakamoto et al. 2004). Thus, LKB1 is believed to be a crucial AMPK kinase in the response to energy deprivation in skeletal muscle during intense contraction. In the present study, AMPK α 1 activation by 1 mM caffeine was not accompanied by energy depletion (Figs 1 and 2), suggesting that LKB1 is not the main AMPK kinase involved in activation of AMPK α 1 by 1 mM caffeine. Thus, our results raise the possibility that other enzymes involved in the regulation of AMPK, such as transforming growth factor- β -activated kinase 1 (Momcilovic et al. 2006) and protein phosphatase 2C (Davies et al. 1995), are involved in caffeine-induced AMPK α 1 activation in skeletal muscle.

In summary, our results suggest (1) that caffeine preferentially activates AMPK α 1 in the absence of energy deprivation, (2) that caffeine activates both AMPK α 1 and AMPK α 2 in the presence of energy deprivation, and (3) that activation of the α 1 isoform leads to enhanced glucose transport and ACC phosphorylation in rat skeletal muscle. We propose that both AMPK α 2 and AMPK α 1 play important roles in regulating muscle metabolism and that AMPK α 2 is not essential for metabolic

activation in skeletal muscle.

Acknowledgements

We thank Licht Miyamoto and Taro Toyoda for suggestions. We also thank Department of Medicine and Clinical Science Kyoto University Graduate School of Medicine and Radioisotope Research Center of Kyoto University for instrumental support. Tatsuya Hayashi was supported by research grants from the Japan Society for the Promotion of Science (20500576), Nestle Nutrition Council Japan, and Japan Vascular Disease Research Foundation. Tatsuya Hayashi and Taku Hamada were supported by a research grant from the All Japan Coffee Association.

Conflicts of interest

The authors state that there are no conflicts of interest.

References

- Birk, J. B. & Wojtaszewski, J. F. 2006. Predominant $\alpha 2/\beta 2/\gamma 3$ AMPK activation during exercise in human skeletal muscle. *J Physiol*, **577**, 1021-1032.
- Blair, D. R., Funai, K., Schweitzer, G. G. & Cartee, G. D. 2009. A myosin II ATPase inhibitor reduces force production, glucose transport, and phosphorylation of AMPK and TBC1D1 in electrically stimulated rat skeletal muscle. *Am J Physiol Endocrinol Metab*, **296**, E993-E1002.
- Cantó, C., Chibalin, A. V., Barnes, B. R., Glund, S., Suárez, E., Ryder, J. W., Palacín, M., Zierath, J. R., Zorzano, A. & Guma, A. 2006. Neuregulins mediate calcium-induced glucose transport during muscle contraction. *J Biol Chem*, **281**, 21690-21697.
- Chen, Z. P., McConell, G. K., Michell, B. J., Snow, R. J., Canny, B. J. & Kemp, B. E. 2000. AMPK signaling in contracting human skeletal muscle: acetyl-CoA carboxylase and NO synthase phosphorylation. *Am J Physiol Endocrinol Metab*, **279**, E1202-1206.
- Davies, S. P., Helps, N. R., Cohen, P. T. & Hardie, D. G. 1995. 5'-AMP inhibits dephosphorylation, as well as promoting phosphorylation, of the AMP-activated protein kinase. Studies using bacterially expressed human protein phosphatase-2C alpha and native bovine protein phosphatase-2AC. *FEBS Lett*, **377**, 421-425.
- Davies, S. P., Sim, A. T. & Hardie, D. G. 1990. Location and function of three sites phosphorylated on rat acetyl-CoA carboxylase by the AMP-activated protein kinase.

Eur J Biochem, **187**, 183-190.

Durante, P. E., Mustard, K. J., Park, S. H., Winder, W. W. & Hardie, D. G. 2002. Effects of endurance training on activity and expression of AMP-activated protein kinase isoforms in rat muscles. *Am J Physiol Endocrinol Metab*, **283**, E178-186.

Egawa, T., Hamada, T., Kameda, N., Karaike, K., Ma, X., Masuda, S., Iwanaka, N. & Hayashi, T. 2009. Caffeine acutely activates 5'adenosine monophosphate-activated protein kinase and increases insulin-independent glucose transport in rat skeletal muscles. *Metabolism*, **58**, 1609-1617.

Fiedler, M., Zierath, J. R., Selén, G., Wallberg-Henriksson, H., Liang, Y. & Sakariassen, K. S. 2001. 5-aminoimidazole-4-carboxy-amide-1- β -D-ribofuranoside treatment ameliorates hyperglycaemia and hyperinsulinaemia but not dyslipidaemia in KKA^y-CETP mice. *Diabetologia*, **44**, 2180-2186.

Fujii, N., Hayashi, T., Hirshman, M. F., Smith, J. T., Habinowski, S. A., Kaijser, L., Mu, J., Ljungqvist, O., Birnbaum, M. J., Witters, L. A., Thorell, A. & Goodyear, L. J. 2000. Exercise induces isoform-specific increase in 5'AMP-activated protein kinase activity in human skeletal muscle. *Biochem Biophys Res Commun*, **273**, 1150-1155.

Fujii, N., Jessen, N. & Goodyear, L. J. 2006. AMP-activated protein kinase and the regulation of glucose transport. *Am J Physiol Endocrinol Metab*, **291**, E867-877.

Garcia-Roves, P. M., Osler, M. E., Holmstrom, M. H. & Zierath, J. R. 2008. Gain-of-function R225Q mutation in AMP-activated protein kinase γ 3 subunit increases mitochondrial biogenesis in glycolytic skeletal muscle. *J Biol Chem*, **283**,

35724-35734.

Hardie, D. G. & Carling, D. 1997. The AMP-activated protein kinase--fuel gauge of the mammalian cell? *Eur J Biochem*, **246**, 259-273.

Hardie, D. G., Hawley, S. A. & Scott, J. W. 2006. AMP-activated protein kinase--development of the energy sensor concept. *J Physiol*, **574**, 7-15.

Hawley, S. A., Boudeau, J., Reid, J. L., Mustard, K. J., Udd, L., Mäkelä, T. P., Alessi, D. R. & Hardie, D. G. 2003. Complexes between the LKB1 tumor suppressor, STRAD α/β and MO25 α/β are upstream kinases in the AMP-activated protein kinase cascade. *J Biol*, **2**, 28.1-28.16.

Hawley, S. A., Pan, D. A., Mustard, K. J., Ross, L., Bain, J., Edelman, A. M., Frenguelli, B. G. & Hardie, D. G. 2005. Calmodulin-dependent protein kinase kinase-beta is an alternative upstream kinase for AMP-activated protein kinase. *Cell Metab*, **2**, 9-19.

Hayashi, T., Hirshman, M. F., Fujii, N., Habinowski, S. A., Witters, L. A. & Goodyear, L. J. 2000. Metabolic stress and altered glucose transport: activation of AMP-activated protein kinase as a unifying coupling mechanism. *Diabetes*, **49**, 527-531.

Hayashi, T., Hirshman, M. F., Kurth, E. J., Winder, W. W. & Goodyear, L. J. 1998. Evidence for 5' AMP-activated protein kinase mediation of the effect of muscle contraction on glucose transport. *Diabetes*, **47**, 1369-1373.

Hudmon, A. & Schulman, H. 2002. Structure-function of the multifunctional

Ca²⁺/calmodulin-dependent protein kinase II. *Biochem J*, **364**, 593-611.

Jager, S., Handschin, C., St-Pierre, J. & Spiegelman, B. M. 2007. AMP-activated protein kinase (AMPK) action in skeletal muscle via direct phosphorylation of PGC-1 α . *Proc Natl Acad Sci U S A*, **104**, 12017-12022.

Jensen, T. E., Rose, A. J., Hellsten, Y., Wojtaszewski, J. F. & Richter, E. A. 2007a. Caffeine-induced Ca²⁺ release increases AMPK-dependent glucose uptake in rodent soleus muscle. *Am J Physiol Endocrinol Metab*, **293**, E286-292.

Jensen, T. E., Rose, A. J., Jørgensen, S. B., Brandt, N., Schjerling, P., Wojtaszewski, J. F. & Richter, E. A. 2007b. Possible CaMKK-dependent regulation of AMPK phosphorylation and glucose uptake at the onset of mild tetanic skeletal muscle contraction. *Am J Physiol Endocrinol Metab*, **292**, E1308-1317.

Jensen, T. E., Schjerling, P., Viollet, B., Wojtaszewski, J. F. & Richter, E. A. 2008. AMPK α 1 activation is required for stimulation of glucose uptake by twitch contraction, but not by H₂O₂, in mouse skeletal muscle. *PLoS ONE*, **3**, e2102.

Jørgensen, S. B., Nielsen, J. N., Birk, J. B., Olsen, G. S., Viollet, B., Andreelli, F., Schjerling, P., Vaulont, S., Hardie, D. G., Hansen, B. F., Richter, E. A. & Wojtaszewski, J. F. 2004a. The α 2-5'AMP-activated protein kinase is a site 2 glycogen synthase kinase in skeletal muscle and is responsive to glucose loading. *Diabetes*, **53**, 3074-3081.

Jørgensen, S. B., Viollet, B., Andreelli, F., Frosig, C., Birk, J. B., Schjerling, P., Vaulont, S., Richter, E. A. & Wojtaszewski, J. F. 2004b. Knockout of the α 2 but not

α 1 5'-AMP-activated protein kinase isoform abolishes 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranosidebut not contraction-induced glucose uptake in skeletal muscle. *J Biol Chem*, **279**, 1070-1079.

Kahn, B. B., Alquier, T., Carling, D. & Hardie, D. G. 2005. AMP-activated protein kinase: ancient energy gauge provides clues to modern understanding of metabolism. *Cell Metab*, **1**, 15-25.

Kemp, B. E., Mitchelhill, K. I., Stapleton, D., Michell, B. J., Chen, Z. P. & Witters, L. A. 1999. Dealing with energy demand: the AMP-activated protein kinase. *Trends Biochem Sci*, **24**, 22-25.

Klein, D. K., Pilegaard, H., Trebak, J. T., Jensen, T. E., Viollet, B., Schjerling, P. & Wojtaszewski, J. F. 2007. Lack of AMPK α 2 enhances pyruvate dehydrogenase activity during exercise. *Am J Physiol Endocrinol Metab*, **293**, E1242-1249.

Lowry, O. H. & Passonneau, J. V. 1972. A Flexible System of Enzymatic Analysis, pp. 151-156. New York, Academic Press.

Miyamoto, L., Toyoda, T., Hayashi, T., Yonemitsu, S., Nakano, M., Tanaka, S., Ebihara, K., Masuzaki, H., Hosoda, K., Ogawa, Y., Inoue, G., Fushiki, T. & Nakao, K. 2007. Effect of acute activation of 5'-AMP-activated protein kinase on glycogen regulation in isolated rat skeletal muscle. *J Appl Physiol*, **102**, 1007-1013.

Momcilovic, M., Hong, S. P. & Carlson, M. 2006. Mammalian TAK1 activates Snf1 protein kinase in yeast and phosphorylates AMP-activated protein kinase in vitro. *J Biol Chem*, **281**, 25336-25343.

Mukwevho, E., Kohn, T. A., Lang, D., Nyatia, E., Smith, J. & Ojuka, E. O. 2008. Caffeine induces hyperacetylation of histones at the MEF2 site on the *Glut4* promoter and increases MEF2A binding to the site via a CaMK-dependent mechanism. *Am J Physiol Endocrinol Metab*, **294**, E582-588.

Musi, N., Fujii, N., Hirshman, M. F., Ekberg, I., Fröberg, S., Ljungqvist, O., Thorell, A. & Goodyear, L. J. 2001a. AMP-activated protein kinase (AMPK) is activated in muscle of subjects with type 2 diabetes during exercise. *Diabetes*, **50**, 921-927.

Musi, N., Hayashi, T., Fujii, N., Hirshman, M. F., Witters, L. A. & Goodyear, L. J. 2001b. AMP-activated protein kinase activity and glucose uptake in rat skeletal muscle. *Am J Physiol Endocrinol Metab*, **280**, E677-684.

Nakano, M., Hamada, T., Hayashi, T., Yonemitsu, S., Miyamoto, L., Toyoda, T., Tanaka, S., Masuzaki, H., Ebihara, K., Ogawa, Y., Hosoda, K., Inoue, G., Yoshimasa, Y., Otaka, A., Fushiki, T. & Nakao, K. 2006. $\alpha 2$ Isoform-specific activation of 5'-adenosine monophosphate-activated protein kinase by 5-aminoimidazole-4-carboxamide-1- β -d-ribose at a physiological level activates glucose transport and increases glucose transporter 4 in mouse skeletal muscle. *Metabolism*, **55**, 300-308.

Ojuka, E. O., Jones, T. E., Nolte, L. A., Chen, M., Wamhoff, B. R., Sturek, M. & Holloszy, J. O. 2002. Regulation of GLUT4 biogenesis in muscle: evidence for involvement of AMPK and Ca^{2+} . *Am J Physiol Endocrinol Metab*, **282**, E1008-1013.

Park, H., Kaushik, V. K., Constant, S., Prentki, M., Przybytkowski, E., Ruderman, N. B. & Saha, A. K. 2002. Coordinate regulation of malonyl-CoA decarboxylase,

sn-glycerol-3-phosphate acyltransferase, and acetyl-CoA carboxylase by AMP-activated protein kinase in rat tissues in response to exercise. *J Biol Chem*, **277**, 32571-32577.

Raney, M. A. & Turcotte, L. P. 2008. Evidence for the involvement of CaMKII and AMPK in Ca²⁺-dependent signaling pathways regulating FA uptake and oxidation in contracting rodent muscle. *J Appl Physiol*, **104**, 1366-1373.

Rose, A. J., Alsted, T. J., Kobbero, J. B. & Richter, E. A. 2007. Regulation and function of Ca²⁺-calmodulin-dependent protein kinase II of fast-twitch rat skeletal muscle. *J Physiol*, **580**, 993-1005.

Sakamoto, K., Göransson, O., Hardie, D. G. & Alessi, D. R. 2004. Activity of LKB1 and AMPK-related kinases in skeletal muscle: effects of contraction, phenformin, and AICAR. *Am J Physiol Endocrinol Metab*, **287**, E310-317.

Sakamoto, K., McCarthy, A., Smith, D., Green, K. A., Grahame Hardie, D., Ashworth, A. & Alessi, D. R. 2005. Deficiency of LKB1 in skeletal muscle prevents AMPK activation and glucose uptake during contraction. *Embo J*, **24**, 1810-1820.

Salt, I., Celler, J. W., Hawley, S. A., Prescott, A., Woods, A., Carling, D. & Hardie, D. G. 1998. AMP-activated protein kinase: greater AMP dependence, and preferential nuclear localization, of complexes containing the $\alpha 2$ isoform. *Biochem J*, **334**, 177-187.

Sriwijitkamol, A., Coletta, D. K., Wajcberg, E., Balbontin, G. B., Reyna, S. M., Barrientes, J., Eagan, P. A., Jenkinson, C. P., Cersosimo, E., DeFronzo, R. A.,

Sakamoto, K. & Musi, N. 2007. Effect of acute exercise on AMPK signaling in skeletal muscle of subjects with type 2 diabetes: a time-course and dose-response study. *Diabetes*, **56**, 836-848.

Stein, S. C., Woods, A., Jones, N. A., Davison, M. D. & Carling, D. 2000. The regulation of AMP-activated protein kinase by phosphorylation. *Biochem J*, **345 Pt 3**, 437-443.

Stephens, T. J., Chen, Z. P., Canny, B. J., Michell, B. J., Kemp, B. E. & McConell, G. K. 2002. Progressive increase in human skeletal muscle AMPK α 2 activity and ACC phosphorylation during exercise. *Am J Physiol Endocrinol Metab*, **282**, E688-694.

Tanaka, S., Hayashi, T., Toyoda, T., Hamada, T., Shimizu, Y., Hirata, M., Ebihara, K., Masuzaki, H., Hosoda, K., Fushiki, T. & Nakao, K. 2007. High-fat diet impairs the effects of a single bout of endurance exercise on glucose transport and insulin sensitivity in rat skeletal muscle. *Metabolism*, **56**, 1719-1728.

Tokumitsu, H., Enslin, H. & Soderling, T. R. 1995. Characterization of a Ca²⁺/calmodulin-dependent protein kinase cascade. Molecular cloning and expression of calcium/calmodulin-dependent protein kinase kinase. *J Biol Chem*, **270**, 19320-19324.

Toyoda, T., Hayashi, T., Miyamoto, L., Yonemitsu, S., Nakano, M., Tanaka, S., Ebihara, K., Masuzaki, H., Hosoda, K., Inoue, G., Otaka, A., Sato, K., Fushiki, T. & Nakao, K. 2004. Possible involvement of the α 1 isoform of 5'AMP-activated protein kinase in oxidative stress-stimulated glucose transport in skeletal muscle. *Am J Physiol Endocrinol Metab*, **287**, E166-173.

Toyoda, T., Tanaka, S., Ebihara, K., Masuzaki, H., Hosoda, K., Sato, K., Fushiki, T., Nakao, K. & Hayashi, T. 2006. Low-intensity contraction activates the α 1-isoform of 5'-AMP-activated protein kinase in rat skeletal muscle. *Am J Physiol Endocrinol Metab*, **290**, E583-590.

Vavvas, D., Apazidis, A., Saha, A. K., Gamble, J., Patel, A., Kemp, B. E., Witters, L. A. & Ruderman, N. B. 1997. Contraction-induced changes in acetyl-CoA carboxylase and 5'-AMP-activated kinase in skeletal muscle. *J Biol Chem*, **272**, 13255-13261.

Winder, W. W. & Hardie, D. G. 1996. Inactivation of acetyl-CoA carboxylase and activation of AMP-activated protein kinase in muscle during exercise. *Am J Physiol*, **270**, E299-304.

Witczak, C. A., Fujii, N., Hirshman, M. F. & Goodyear, L. J. 2007. Ca^{2+} /calmodulin-dependent protein kinase kinase- α regulates skeletal muscle glucose uptake independent of AMP-activated protein kinase and Akt activation. *Diabetes*, **56**, 1403-1409.

Witczak, C. A., Jessen, N., Warro, D. M., Toyoda, T., Fujii, N., Anderson, M. E., Hirshman, M. F. & Goodyear, L. J. 2010. CaMKII Regulates Contraction-Induced, But Not Insulin-Induced, Glucose Uptake in Mouse Skeletal Muscle. *Am J Physiol Endocrinol Metab*, **298**, E1150-1160.

Wojtaszewski, J. F., MacDonald, C., Nielsen, J. N., Hellsten, Y., Hardie, D. G., Kemp, B. E., Kiens, B. & Richter, E. A. 2003. Regulation of 5'AMP-activated protein kinase activity and substrate utilization in exercising human skeletal muscle. *Am J*

Physiol Endocrinol Metab, **284**, E813-822.

Wojtaszewski, J. F., Mourtzakis, M., Hillig, T., Saltin, B. & Pilegaard, H. 2002. Dissociation of AMPK activity and ACC β phosphorylation in human muscle during prolonged exercise. *Biochem Biophys Res Commun*, **298**, 309-316.

Wojtaszewski, J. F., Nielsen, P., Hansen, B. F., Richter, E. A. & Kiens, B. 2000. Isoform-specific and exercise intensity-dependent activation of 5'-AMP-activated protein kinase in human skeletal muscle. *J Physiol*, **528 Pt 1**, 221-226.

Wright, D. C., Geiger, P. C., Holloszy, J. O. & Han, D. H. 2005. Contraction- and hypoxia-stimulated glucose transport is mediated by a Ca²⁺-dependent mechanism in slow-twitch rat soleus muscle. *Am J Physiol Endocrinol Metab*, **288**, E1062-1066.

Wright, D. C., Hucker, K. A., Holloszy, J. O. & Han, D. H. 2004. Ca²⁺ and AMPK both mediate stimulation of glucose transport by muscle contractions. *Diabetes*, **53**, 330-335.

Young, D. A., Uhl, J. J., Cartee, G. D. & Holloszy, J. O. 1986. Activation of glucose transport in muscle by prolonged exposure to insulin. Effects of glucose and insulin concentrations. *J Biol Chem*, **261**, 16049-16053.

Zheng, D., MacLean, P. S., Pohnert, S. C., Knight, J. B., Olson, A. L., Winder, W. W. & Dohm, G. L. 2001. Regulation of muscle GLUT-4 transcription by AMP-activated protein kinase. *J Appl Physiol*, **91**, 1073-1083.

Tables

Table 1 ATP, PCr, and glycogen concentrations in muscle after saline (Control) or Caffeine 5mg kg⁻¹ injection

		Time after injection	
		0 min	60 min
ATP (nmol mg ⁻¹)	Control	5.29 ± 0.31	5.04 ± 0.23
	Caffeine 5 mg/kg ⁻¹	5.48 ± 0.25	5.41 ± 0.23
PCr (nmol mg ⁻¹)	Control	15.9 ± 0.72	14.9 ± 0.38
	Caffeine 5 mg/kg ⁻¹	15.1 ± 0.35	14.9 ± 0.66
Glycogen (nmol mg ⁻¹)	Control	23.4 ± 1.34	24.6 ± 2.09
	Caffeine 5 mg/kg ⁻¹	23.9 ± 1.85	24.7 ± 0.98

Values are mean ± SE; n = 6-8 per group.

Table 2 Plasma glucose, insulin, and caffeine levels after saline (Control) or Caffeine 5mg kg⁻¹ injection

		Time after injection	
		0 min	60 min
Glucose (mg dL ⁻¹)	Control	56.7 ± 1.7	59.0 ± 1.9
	Caffeine 5 mg/kg ⁻¹	60.1 ± 2.0	62.5 ± 2.3
Insulin (ng mL ⁻¹)	Control	0.51 ± 0.10	0.41 ± 0.07
	Caffeine 5 mg/kg ⁻¹	0.61 ± 0.11	0.47 ± 0.05
Caffeine (μM)	Control	ND	ND
	Caffeine 5 mg/kg ⁻¹	ND	54 ± 4

Values are mean ± SE; n = 4-8 per group. ND; not detectable.

Legends to figures

Figure 1

Caffeine activates preferentially AMPK α 1 activity in incubated rat epitrochlearis muscle. Isolated muscles were incubated in the absence (Control) or presence of caffeine for 15 min. Isoform-specific AMPK activity was determined in anti-AMPK α 1 and -AMPK α 2 immunoprecipitates. Fold increases are expressed relative to the activity of muscles in the control group. Values are mean \pm SE; n = 6-14 per group. *P<0.05, **P<0.01 vs. Control.

Figure 2

ATP, PCr and glycogen concentrations in incubated rat epitrochlearis muscle. Isolated muscles were incubated in the absence (Control) or presence of caffeine for 15 min. ATP (A), PCr (B), and Glycogen (C) concentrations were measured enzymatically. Values are mean \pm SE; n = 5-10 per group. *P<0.05, **P<0.01 vs. Control.

Figure 3

Caffeine increases phosphorylation of AMPK α Thr¹⁷² (A) and ACC Ser⁷⁹ (B) in incubated rat epitrochlearis muscle. Isolated muscles were incubated in the absence (Control) or presence of caffeine for 15 min. Muscle lysates were then analyzed for phosphorylation of AMPK α Thr¹⁷² (pAMPK) and ACC Ser⁷⁹ (pACC) by Western blot. Fold increases are expressed relative to the level of muscles in the control group. Representative immunoblots are shown. Values are mean \pm SE; n = 6-12 per

group. *P<0.05, **P<0.01 vs. Control.

Figure 4

Caffeine increases 3-*O*-methyl-D-glucose (3MG) transport in incubated rat epitrochlearis muscle. Isolated muscles were incubated in the absence (Control) or presence of caffeine for 15 min, and then 3MG transport activity was determined. Values are mean \pm SE; n = 9-13 per group. *P<0.05, **P<0.01 vs. Control.

Figure 5

Caffeine does not change CaMKI Thr¹⁷⁷ phosphorylation (A), but increases CaMKII Thr²⁸⁷ phosphorylation (B) in incubated rat epitrochlearis muscle. Isolated muscles were incubated in the absence (Control) or presence of caffeine for 15 min. Muscle lysates were then analyzed for phosphorylation of CaMKI Thr¹⁷⁷ (pCaMKI,) and CaMKII Thr²⁸⁷ (pCaMKII) by Western blot. The phosphorylation of CaMKII isoforms migrated between 50 to 75 KDa was summed (Rose et al. 2007). Fold increases are expressed relative to the level of muscles in the control group. Representative immunoblots are shown. Values are mean \pm SE; n = 4-9 per group. *P<0.05 vs. Control.

Figure 6

Intravenous caffeine injection increases phosphorylation of AMPK α Thr¹⁷² (A) and ACC Ser⁷⁹ (B) in a dose-dependent manner in rat epitrochlearis muscle. Muscles were dissected 60 min after injection, and then analyzed for phosphorylation of AMPK α Thr¹⁷² (pAMPK) and ACC Ser⁷⁹ (pACC) by Western blot. Fold increases

are expressed relative to the level of muscles in the saline injection (control) group. Representative immunoblots are shown. Values are mean \pm SE; n = 8-17 per group. *P<0.05, **P<0.01 vs. Control.

Figure 7

Intravenous caffeine injection increases phosphorylation of AMPK α Thr¹⁷² (A) and ACC Ser⁷⁹ (B) in a time-dependent manner in rat epitrochlearis muscle. Muscles were dissected at indicated time points after caffeine injection (5 mg kg⁻¹). Control samples were dissected immediately after saline injection. Muscles were then analyzed for phosphorylation of AMPK α Thr¹⁷² (pAMPK) and ACC Ser⁷⁹ (pACC) by Western blot. Fold increases are expressed relative to the activity of muscles in the control group. Representative immunoblots are shown. Values are mean \pm SE; n = 7-15 per group. *P<0.05, **P<0.01 vs. Control.

Figure 8

Intravenous caffeine injection activates AMPK α 1 activity without changes of energy status in rat epitrochlearis. Muscles were dissected 60 min after caffeine (5 mg kg⁻¹) or saline injection. Isoform-specific AMPK activity was determined in anti-AMPK α 1 and -AMPK α 2 immunoprecipitates. Fold increases are expressed relative to the activity of muscles in the saline injection (control) group. Values are mean \pm SE; n = 5-8 per group. **P<0.01 vs. Control.

Figure 9

Intravenous caffeine injection increases 3-O-methyl-D-glucose (3MG) transport in

rat epitrochlearis muscle. Muscles were dissected 60 min after caffeine (5 mg kg⁻¹) or saline injection (control), and 3MG transport activity was determined. Values are mean \pm SE; n = 6-7 per group. **P<0.01 vs. Control.




Figure 1




Figure 2




Figure 3




Figure 4





Figure 5





Figure 6





Figure 7




Figure 8




Figure 9