

Zeros of partial zeta functions off the critical line

By

Yoonbok LEE*

Abstract

We extend the joint universality theorem for Artin L -functions $L(s, \chi_j, K/\mathbb{Q})$ from the previously known strip $1 - \frac{1}{2k} < \operatorname{Re} s < 1$ for $k = \#G(K/\mathbb{Q})$ to the maximal strip $\frac{1}{2} < \operatorname{Re} s < 1$ under an assumption of a weak version of the density hypothesis. Then, we study zeros of partial zeta functions $\zeta(s, \mathcal{A})$ inside the half of the critical strip as an application of universality.

§ 1. Introduction

Let $Q(m, n) = am^2 + bmn + cn^2$ be a positive definite quadratic form with its discriminant $D = b^2 - 4ac < 0$ and $a, b, c \in \mathbb{Z}$. We define the Epstein zeta function attached to Q by

$$E(s, Q) = \sum_{(m,n) \neq (0,0)} \frac{1}{Q(m, n)^s}, \quad \operatorname{Re} s > 1.$$

Then, it has a meromorphic continuation to \mathbb{C} with one simple pole at $s = 1$, and has the functional equation

$$\Phi(s) := \left(\frac{\sqrt{-D}}{2\pi} \right)^s \Gamma(s) E(s, Q) = \Phi(1 - s).$$

Voronin([9] or [4]) studied zeros of $E(s, Q)$ inside the critical strip and proved the following theorem.

Received March 16, 2011. Accepted September 16, 2011.

2000 Mathematics Subject Classification(s): 11M41

Key Words: Joint universality, Artin L -function, density hypothesis, zero, partial zeta function.

*KIAS, School of Mathematics, Hoegiro 85, Dongdaemun-gu, Seoul 130-722, Korea.

e-mail: leeyb@kias.re.kr

Present address: Department of Mathematics, University of Rochester, Rochester, NY 14627, USA.

email:lee@math.rochester.edu

Theorem 1.1 (Voronin). *Suppose that the class number of $\mathbb{Q}(\sqrt{D})$ is $h(D) > 1$. The Epstein zeta function $E(s, Q)$ described above has at least cT zeros on the rectangular region $\sigma_1 < \operatorname{Re} s < \sigma_2$, $0 < \operatorname{Im} s < T$ for each $\frac{1}{2} < \sigma_1 < \sigma_2 < 1$ and some $c > 0$ as $T \rightarrow \infty$.*

The main purpose of this paper is extending it to partial zeta functions which are natural algebraic generalizations of Epstein zeta functions.

Let \mathfrak{f} be an integral ideal of the number field K . Then, $J^\mathfrak{f}$ denotes the group of all ideals of K which are relatively prime to \mathfrak{f} and $P^\mathfrak{f}$ stands for the group of fractional principal ideals (a) such that $a \equiv 1 \pmod{\mathfrak{f}}$ with a totally positive. Choose an element \mathcal{A} of the ray class group $G^\mathfrak{f} = J^\mathfrak{f}/P^\mathfrak{f} \pmod{\mathfrak{f}}$. The partial zeta function $\zeta(s, \mathcal{A})$ attached to \mathcal{A} is defined by

$$\zeta(s, \mathcal{A}) = \sum_{\mathfrak{n} \in \mathcal{A}} \frac{1}{N(\mathfrak{n})^s} \quad \operatorname{Re} s > 1,$$

where \mathfrak{n} runs through all ideals in O_K and $N(\mathfrak{n})$ denotes the norm of \mathfrak{n} . It has a meromorphic continuation to \mathbb{C} with only one simple pole at $s = 1$ and the functional equation

$$\xi(1-s, \mathcal{A}) = \xi(s, \mathcal{A}) := D(\mathfrak{f})^s \Gamma(s)^{r_2} \prod_{m=1}^{r_1} \Gamma\left(\frac{s+a_m}{2}\right) \zeta(s, \mathcal{A}),$$

where r_1 is the number of real places of K , $2r_2$ is the number of complex places of K . The constant $D(\mathfrak{f})$ depends only on \mathfrak{f} and a_m takes the value 0 or 1. For details see Chapter 7 of [8].

The Hecke L -function attached to a ray class character $\psi : G^\mathfrak{f} \rightarrow S^1$ is defined by

$$L(s, \psi) = \sum_{\mathfrak{n} \in J^\mathfrak{f}} \frac{\psi(\mathfrak{n})}{N(\mathfrak{n})^s} = \prod_{\mathfrak{p} \in J^\mathfrak{f}} \left(1 - \frac{\psi(\mathfrak{p})}{N(\mathfrak{p})^s}\right)^{-1} \quad \operatorname{Re} s > 1.$$

Since the functional equation of $\zeta(s, \mathcal{A})$ does not depend on the choice of \mathcal{A} , we can deduce the functional equation for the Hecke L -function $L(s, \psi)$ by

$$L(s, \psi) = \sum_{\mathcal{A} \in G^\mathfrak{f}} \psi(\mathcal{A}) \zeta(s, \mathcal{A}).$$

If we take a representative $\mathfrak{a} \in \mathcal{A}$, then we have

$$\zeta(s, \mathcal{A}) = \frac{1}{h} \sum_{\psi} \bar{\psi}(\mathfrak{a}) L(s, \psi),$$

where ψ runs through all ray class characters ψ defined on $G^\mathfrak{f}$, and $h = \#G^\mathfrak{f}$.

For a positive definite quadratic form $Q(m, n)$ with its discriminant $D < 0$, it is known that

$$E(s, Q) = \omega \zeta(s, \mathcal{A}),$$

where ω is the number of units of $\mathbb{Q}(\sqrt{D})$ and \mathcal{A} is an ideal class corresponding Q . So, we have

$$E(s, Q) = \frac{\omega}{h} \sum_{\psi} \bar{\psi}(\mathfrak{a}) L(s, \psi)$$

and the condition $h(D) > 1$ of Theorem 1.1 means that $h = h(D) = \#G > 1$ for the ideal class group G of $\mathbb{Q}(\sqrt{D})$. In other words, $E(s, Q)$ is a linear combination of at least two Hecke L -functions. Voronin's proof of Theorem 1.1 is based on the joint distribution of those Hecke L -functions in the above equation and Rouché's theorem in complex analysis.

Bauer [1] proved the following theorem concerning zeros of $\zeta(s, \mathcal{A})$ inside the critical strip.

Theorem 1.2 (Bauer). *If T is sufficiently large, then there is a number $c > 0$ such that there are at least cT zeros of $\zeta(s, \mathcal{A})$ in the region with $\frac{1}{2} < \operatorname{Re} s < 1$, $|\operatorname{Im} s| < T$.*

The main ingredient of its proof is the joint universality of Artin L -functions instead of Hecke L -functions. His argument counts the number of zeros in the strip $1 - \frac{1}{2k} < \operatorname{Re} s < 1$ for $k = \#Gal(K/\mathbb{Q})$ because the inequality

$$\int_0^T |f(\sigma + it)|^2 dt \ll T$$

is required for the proof of the joint universality of Artin L -functions and known only for the strip $1 - \frac{1}{2k} < \operatorname{Re} s < 1$ for $k = \#Gal(K/\mathbb{Q})$.

§ 2. Joint distribution of Artin L -functions

Let K/\mathbb{Q} be a normal extension with number field K and G be its Galois group $G(K/\mathbb{Q})$. Let $\rho : G \rightarrow GL_m(\mathbb{C})$ be a m -dimensional representation of G in the general linear group $GL_m(\mathbb{C})$. The character $\chi : G \rightarrow \mathbb{C}$ of the representation ρ is given by

$$\chi(g) := \operatorname{tr}(\rho(g)).$$

The Artin L -function of χ and G is defined by the Euler product

$$L(s, \chi, K/\mathbb{Q}) = \prod_{p:\text{unr.}} L_p(s, \chi), \quad \operatorname{Re} s > 1,$$

where $L_p(s, \chi) = \det(I - \rho(\sigma_p)p^{-s})^{-1}$ and σ_p denotes one of the conjugate Frobenius automorphisms over p . This definition is independent of the specific representation ρ of the character χ and the chosen conjugate of the Frobenius σ_p .

Brauer’s theorem states that every character χ of a finite group G is a \mathbb{Z} -linear combination of characters ψ_{l*} induced from characters ψ_l of degree 1 associated to subgroups H_l of G . Thus, for $j \leq J$ we have

$$\chi_j = \sum_{l \leq l_0} n_{j,l} \psi_l^*,$$

where ψ_l^* are deduced from characters ψ_l of degree 1 associated to subgroups H_l of G and $n_{j,l} \in \mathbb{Z}$. As a consequence, we have

$$(2.1) \quad L(s, \chi_j, K/\mathbb{Q}) = \prod_{l \leq l_0} L(s, \psi_l)^{n_{j,l}}$$

and $L(s, \psi_l) = L(s, \psi_l, H_l)$ are Hecke L -functions over number fields contained in K . Note that (2.1) shows that the Artin L -function $L(s, \chi_j, K/\mathbb{Q})$ has a meromorphic continuation to \mathbb{C} .

Conjecture 2.1. *Let $L(s, \chi, K/\mathbb{Q})$ be an Artin L -function and write*

$$L(s, \chi, K/\mathbb{Q}) = \prod_{l \leq l_0} L(s, \psi_l)^{n_l}$$

for some $n_l \in \mathbb{Z}$. Define $N_\psi(\sigma, T)$ by the number of zeros of Hecke L -function $L(s, \psi)$ on the region $\text{Re } s > \sigma$, $0 < \text{Im } s < T$. Then, there is a constant $c > 0$ such that

$$N_{\psi_l}(\sigma, T) \ll T^{1+c(\frac{1}{2}-\sigma)} \log T$$

uniformly for $\sigma \geq \frac{1}{2}$ and $l \leq l_0$.

Now, we are ready to state the joint universality of Artin L -functions.

Theorem 2.2. *Let K be a finite Galois extension of \mathbb{Q} and let χ_1, \dots, χ_J be \mathbb{C} -linearly independent characters of the group $G = \text{Gal}(K/\mathbb{Q})$. Assume Conjecture 2.1 for $L(s, \chi_j, K/\mathbb{Q})$, $j \leq J$. Let $D = D_r(s_0)$ be the closed disc with center s_0 and radius r such that D is contained in the vertical strip $\frac{1}{2} < \text{Re } s < 1$. Suppose that $h_1(s), \dots, h_J(s)$ are analytic and nonvanishing on $s \in \text{int}D$, and continuous on $s \in D$. Then, for every $\epsilon > 0$ we have*

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \left| \left\{ \tau \in [T, 2T] : \max_{j \leq J} \max_{s \in D} |L(s + i\tau, \chi_j, K/\mathbb{Q}) - h_j(s)| < \epsilon \right\} \right| > 0.$$

We modify several lemmas from [1] and [6] for the proof of Theorem 2.2.

Lemma 2.3. *Assume Conjecture 2.1 for $L(s, \chi_j, K/\mathbb{Q})$, $j \leq J$. Let $\frac{1}{2} < \sigma \leq 1$ and $X = T^\kappa$. Then,*

$$\int_T^{2T} \left| \log L(\sigma + it, \chi_j, K/\mathbb{Q}) - \sum_{p \leq X} \log L_p(\sigma + it, \chi_j) \right|^2 dt = O\left(T^{1+c(\frac{1}{2}-\sigma)}\right)$$

for some $c > 0$ and small enough $\kappa > 0$ and all $j \leq J$.

Lemma 2 of [6] and (2.1) imply Lemma 2.3.

Lemma 2.4. *Assume Conjecture 2.1 for $L(s, \chi_j, K/\mathbb{Q})$, $j \leq J$. Let $D = D_r(s_0) \subset \{s \in \mathbb{C} : \frac{1}{2} < \operatorname{Re} s < 1\}$.*

$$\max_{j \leq J} \max_{s \in D} \left| \log L(s + i\tau, \chi_j, K/\mathbb{Q}) - \sum_{p \leq X} \log L_p(s + i\tau, \chi_j) \right| \leq T^{-c_2}$$

for $\tau \in [T, 2T] \setminus A_T$, $|A_T| \leq T^{1-c_1}$.

Lemma 2.4 is a simple consequence of Lemma 4 of [6] and Lemma 2.3.

Define $L_M(s, \chi, \theta) = \prod_{p \in M} f_p(p^{-s} e(-\theta_p))$ and $f_p(t) = \det(I - \rho(\sigma_p)t)^{-1}$. Lemma 2.2 of [1] is on the disc $D_r(1 - \frac{1}{4k})$ with $0 < r < \frac{1}{4k}$, but its proof in fact proves more than written. So, we restate Lemma 2.2 of [1] based on its proof.

Lemma 2.5. *Let χ_1, \dots, χ_J be \mathbb{C} -linearly independent characters of the Galois group $G = \operatorname{Gal}(K/\mathbb{Q})$, where K is a finite normal algebraic extension of \mathbb{Q} . Let $D = D_r(s_0) \subset \{s \in \mathbb{C} : \frac{1}{2} < \operatorname{Re} s < 1\}$. Suppose that $h_1(s), \dots, h_J(s)$ are analytic and nonvanishing on $s \in \operatorname{int} D$, and continuous on $s \in D$. Then for every pair $\epsilon > 0$ and $y > 0$, there exists a finite set of primes M containing all primes smaller than y and a vector $\theta \in \mathbb{R}^{\mathbb{P}}$ such that*

$$\max_{j \leq J} \max_{s \in D} |L_M(s, \chi_j, \theta) - h_j(s)| < \epsilon.$$

Lemma 2.6. *Let $f(s)$ be an analytic function on a region containing $|s| \leq R$ and $\alpha > 0$. Then, we have*

$$|f(0)|^\alpha \leq \frac{1}{\pi R^2} \int \int_{|s| \leq R} |f(s)|^\alpha d\sigma dt.$$

This is a property of subharmonic function $|f(s)|^\alpha$ and its proof can be found in Lemma 3 of [6].

Lemma 2.7.

$$\int_0^T \left| \sum_{n \leq N} a_n n^{-it} \right|^2 dt = \sum_{n \leq N} |a_n|^2 (T + O(n)).$$

Lemma 2.7 is well-known and we may refer [7] for its proof. Now, we are ready to prove Theorem 2.2.

Proof of Theorem 2.2. By Lemma 2.4 and 2.5, it is enough to show that

$$\max_{j \leq J} \max_{s \in D} \left| \log L_M(s, \chi_j, \theta) - \sum_{p \leq X} \log L_p(s + i\tau, \chi_j) \right| < \epsilon$$

for a positive proportion of $\tau \in [T, 2T] \setminus A_T$, where $X = T^\kappa$ and M is a finite set of primes containing all primes smaller than $y > 0$.

Define the sets $C(\delta, M, T)$ and $C(\delta, M)$ by

$$C(\delta, M, T) = \left\{ \tau \in [T, 2T] : \left| \theta_p - \frac{\tau}{2\pi} \log p \right| < \frac{\delta}{2} \text{ for all } p \in M \right\}$$

and

$$C(\delta, M) = \left\{ (\vartheta_p) \in \Omega : \left| \vartheta_p - \theta_p \right| < \frac{\delta}{2} \text{ for all } p \in M \right\}$$

where $\theta = (\theta_p) \in \mathbb{R}^{\mathbb{P}}$ is as given in Lemma 2.5 and $\|x\| = \min\{|x - n| : n \in \mathbb{Z}\}$. We also use the short expression $C(\delta, X) = C(\delta, \{p \leq X\})$ for a real number $X > 0$. By uniform continuity, there exists $\delta > 0$ such that for $\tau \in C(\delta, M, T)$

$$(2.2) \quad \max_{j \leq J} \max_{s \in D} \left| \log L_M(s, \chi_j, \theta) - \sum_{p \in M} \log L_p(s + i\tau, \chi_j) \right| < \epsilon$$

and we have

$$|C(\delta, M, T)| \sim |C(\delta, M)|T = \delta^{|M|}T, \quad T \rightarrow \infty$$

by Kronecker's theorem.

Let $D' = D_R(s_0)$ be the disc containing $D = D_r(s_0)$ with $R > r$ and contained in the strip $\frac{1}{2} < \operatorname{Re} s < 1$. Let $\sigma_0 = \min\{\operatorname{Re} s : s \in D'\}$. Take $Y > \max\{y\delta^{|M|(1-2\sigma_0)^{-1}}, \max\{p \in M\}\}$ and let P be the largest prime $\leq Y$. We write $Y \setminus M$ for the set $\{p \leq Y : p \notin M\}$. By Kronecker's theorem, we have

$$\begin{aligned} & \lim_{T \rightarrow \infty} \frac{1}{T} \int_{C(\delta, M, T)} \left| \sum_{p \in M} \log L_p(s + i\tau, \chi_j) - \sum_{p \leq Y} \log L_p(s + i\tau, \chi_j) \right|^2 d\tau \\ &= \int_{C(\delta, M)} \left| \sum_{p \in Y \setminus M} \log L_p(s, \chi_j, \vartheta) \right|^2 d\vartheta_2 \cdots d\vartheta_P \leq \\ &\leq \delta^{|M|} \int_0^1 \cdots \int_0^1 \left| \sum_{p \in Y \setminus M} \log L_p(s, \chi_j, \vartheta) \right|^2 \prod_{p \in Y \setminus M} d\vartheta_p = \delta^{|M|} \sum_{p \in Y \setminus M} \sum_{m=1}^{\infty} \frac{|a_{p,j,m}|^2}{p^{2m\sigma}}, \end{aligned}$$

where $\log L_p(s, \chi_j) = \sum_{m=1}^{\infty} a_{p,j,m} p^{-ms}$. Since $|a_{p,j,m}| \leq [K : \mathbb{Q}]$ for all p, j, m , we have

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_{C(\delta, M, T)} \left| \sum_{p \in M} \log L_p(s + i\tau, \chi_j) - \sum_{p \leq Y} \log L_p(s + i\tau, \chi_j) \right|^2 d\tau \leq c_3 \delta^{|M|} y^{1-2\sigma}$$

for some $c_3 > 0$. Thus, we have

(2.3)

$$\int_{C(\delta, M, T)} \int \int_{D'} \left| \sum_{p \in M} \log L_p(s + i\tau, \chi_j) - \sum_{p \leq Y} \log L_p(s + i\tau, \chi_j) \right|^2 d\sigma dt d\tau \leq c_4 \delta^{|M|} y^{1-2\sigma_0} T$$

for some $c_4 > 0$ and sufficiently large T .

By lemma 2.7, we have

$$\begin{aligned} \int_T^{2T} \left| \sum_{Y < p \leq X} \log L_p(s + i\tau, \chi_j) \right|^2 d\tau &\leq 2 \int_T^{2T} \left| \sum_{Y < p \leq X} \frac{\chi_j(\sigma_p)}{p^{s+i\tau}} \right|^2 d\tau + O(TY^{2-4\sigma}) \\ &\leq 2 \sum_{Y < p \leq X} \frac{d^2}{p^{2\sigma}} (T + O(p)) + O(TY^{2-4\sigma}) \\ &\leq c_5 TY^{1-2\sigma} \end{aligned}$$

for some $c_5 > 0$ and all $s \in D'$ and as a consequence

(2.4)

$$\int_{C(\delta, M, T)} \int \int_{D'} \left| \sum_{Y < p \leq X} \log L_p(s + i\tau, \chi_j) \right|^2 d\sigma dt d\tau \leq c_6 TY^{1-2\sigma_0} < c_6 \delta^{|M|} y^{1-2\sigma_0} T.$$

Then (2.3) and (2.4) yield

$$\int_{C(\delta, M, T)} \int \int_{D'} \left| \sum_{p \in X \setminus M} \log L_p(s + i\tau, \chi_j) \right|^2 d\sigma dt d\tau \leq c_7 \delta^{|M|} y^{1-2\sigma_0} T,$$

where $X \setminus M$ denotes the set $\{p \leq X : p \notin M\}$ and $c_7 > 0$ is some constant. From the simple inequality $\max_{n \leq N} |\alpha_n| \leq \sum_{n \leq N} |\alpha_n|$, we have

$$\int_{C(\delta, M, T)} \max_{j \leq J} \int \int_{D'} \left| \sum_{p \in X \setminus M} \log L_p(s + i\tau, \chi_j) \right|^2 d\sigma dt d\tau \leq c_7 J \delta^{|M|} y^{1-2\sigma_0} T.$$

As a consequence, we have

$$\left| \left\{ \tau \in C(\delta, M, T) : \max_{j \leq J} \int \int_{D'} \left| \sum_{p \in X \setminus M} \log L_p(s + i\tau, \chi_j) \right|^2 d\sigma dt \leq y^{\frac{1}{2} - \sigma_0} \right\} \right| > \frac{1}{2} \delta^{|M|} T$$

by taking y satisfying $c_7 J y^{\frac{1}{2} - \sigma_0} < \frac{1}{2}$. By Lemma 2.6, we have

$$\left| \left\{ \tau \in C(\delta, M, T) : \max_{j \leq J} \max_{s \in D} \left| \sum_{p \in X \setminus M} \log L_p(s + i\tau, \chi_j) \right| \leq \frac{1}{\sqrt{\pi}(R-r)} y^{\frac{1}{4} - \frac{1}{2}\sigma_0} \right\} \right| > \frac{1}{2} \delta^{|M|} T.$$

By taking a real number y satisfying $\frac{1}{\sqrt{\pi}(R-r)}y^{\frac{1}{4}-\frac{1}{2}\sigma_0} < \epsilon$, we have

$$|E_T| > \frac{1}{2}\delta^{|M|}T,$$

where

$$E_T = \left\{ \tau \in C(\delta, M, T) : \max_{j \leq J} \max_{s \in D} \left| \sum_{p \in M} \log L_p(s + i\tau, \chi_j) - \sum_{p \leq X} \log L_p(s + i\tau, \chi_j) \right| \leq \epsilon \right\}.$$

Therefore, there exists a $\delta > 0$ and a finite set M such that

$$\max_{j \leq J} \max_{s \in D} |\log L(s + i\tau, \chi_j, K/\mathbb{Q}) - \log h_j(s)| < 4\epsilon$$

for $\tau \in E_T \setminus A_T$ with $|E_T \setminus A_T| \geq |E_T| - |A_T| \geq \frac{1}{2}\delta^{|M|}T + o(T)$. □

§ 3. Zeros of the partial zeta functions

We extend Theorem 1.1 to partial zeta functions subject to Conjecture 2.1 as an application of joint universality of Artin L -functions. Let K be a number field and G^\dagger be its ray class group. By class field theory, there is a unique Abelian extension L of K with $G^\dagger \simeq G(L/K)$. Thus, every Abelian Artin L -function is a Hecke L -function, and vice versa. There is a unique minimal normal extension N of \mathbb{Q} containing L .

Theorem 3.1. *Assume Conjecture 2.1 for $L(s, \chi, N/\mathbb{Q})$ for all characters χ defined on $G(N/\mathbb{Q})$, where the field N is as described above. Suppose that $\#G^\dagger > 1$ and $\mathcal{A} \in G^\dagger$. Then, the number of zeros of the partial zeta function $\zeta(s, \mathcal{A})$ on the rectangular region $\sigma_1 < \operatorname{Re} s < \sigma_2$, $0 < \operatorname{Im} s < T$ is bigger than*

$$\gg T$$

for any fixed $\frac{1}{2} < \sigma_1 < \sigma_2 < 1$.

Proof. Suppose that $\chi \neq 1$ is an irreducible character of $G(L/K) \simeq G^\dagger$. By Frobenius reciprocity, we know that

$$(\chi^*, 1)_{G(N/\mathbb{Q})} = (\chi, 1|_{G(N/K)})_{G(N/K)} = (\chi, 1)_{G(N/K)} = 0.$$

If we denote the irreducible characters of $G(N/\mathbb{Q})$ by $\phi_1 := 1, \phi_2, \dots, \phi_k$, then for every non-trivial character of $G(L/K)$ we have

$$\chi^* = \sum_{j=2}^k m_j \phi_j, \quad m_j \in \mathbb{Z}_{\geq 0}.$$

For the induced character 1^* of the trivial character 1 defined on $G(L/K)$, we get

$$(1^*, 1)_{G(N/\mathbb{Q})} = (1, 1|_{G(N/K)})_{G(N/K)} = (1, 1)_{G(N/K)} = 1.$$

Therefore, we have

$$1^* = \phi_1 + \sum_{j=2}^k n_j \phi_j, \quad n_j \in \mathbb{Z}_{\geq 0}.$$

So we get

$$(3.1) \quad L(s, 1) = L(s, 1^*, N/\mathbb{Q}) = L(s, \phi_1, N/\mathbb{Q}) \prod_{j=2}^k L(s, \phi_j, N/\mathbb{Q})^{n_j}$$

and for the non-trivial Abelian characters χ of $G(L/K)$

$$(3.2) \quad L(s, \chi) = \prod_{j=2}^k L(s, \phi_j, N/\mathbb{Q})^{m_j}.$$

Since the irreducible characters ϕ_j are linearly independent, we apply Theorem 2.2 to $L(s, \chi_j, N/\mathbb{Q})$ with $1 \leq j \leq k$. For any $\epsilon > 0$, there exists a set $A_\epsilon \subset [T, 2T]$ with

$$(3.3) \quad \liminf_{T \rightarrow \infty} \frac{1}{T} |A_\epsilon| > 0$$

such that

$$\left| L(s + i\tau, \phi_1, N/\mathbb{Q}) - (s - s_0 - \sum_{\chi \neq 1} \bar{\chi}(\mathfrak{a})) \right| < \epsilon$$

and

$$|L(s + i\tau, \phi_j, N/\mathbb{Q}) - 1| < \epsilon$$

for any $2 \leq j \leq k$, $s \in D_{s_0}(r) \subset \{z \in \mathbb{C} : \sigma_1 < \operatorname{Re} z < \sigma_2\}$ and $\tau \in A_\epsilon$ and for an integral ideal $\mathfrak{a} \in \mathcal{A}$. By (3.1) and (3.2), we have

$$\left| L(s + i\tau, 1) - (s - s_0 - \sum_{\chi \neq 1} \bar{\chi}(\mathfrak{a})) \right| < \epsilon$$

and for $\chi \neq 1$ we find

$$|L(s + i\tau, \chi) - 1| < \epsilon$$

for all $s \in D_{s_0}(r)$ and $\tau \in A_\epsilon$.

Note that

$$(3.4) \quad \zeta(s, \mathcal{A}) = \frac{1}{h} \sum_{\chi} \bar{\chi}(\mathfrak{a}) L(s, \chi)$$

for some $\mathfrak{a} \in \mathcal{A}$ and $h = \#G^f$. We have

$$\begin{aligned} & \left| \zeta(s + i\tau, \mathcal{A}) - \frac{s - s_0}{h} \right| \\ & \leq \frac{1}{h} \left(\sum_{\chi \neq 1} |\bar{\chi}(\mathfrak{a})L(s + i\tau, \chi) - \bar{\chi}(\mathfrak{a})| + \left| L(s + i\tau, 1) - (s - s_0 - \sum_{\chi \neq 1} \bar{\chi}(\mathfrak{a})) \right| \right) < \epsilon \end{aligned}$$

for all $s \in D_{s_0}(r)$ and all $\tau \in A_\epsilon$. Suppose that $\epsilon < \frac{r}{h}$, then

$$\left| \zeta(s + i\tau, \mathcal{A}) - \frac{s - s_0}{h} \right| < \left| \frac{s - s_0}{h} \right|$$

on the circle $|s - s_0| = r$. Inside the disc $|s - s_0| < r$, there is exactly one zero of $\zeta(s + i\tau, \mathcal{A})$ by Rouché Theorem for each $\tau \in A_\epsilon$. By (3.3), we complete the proof of Theorem 3.1. \square

Let $N_{\zeta(s, \mathcal{A})}(\sigma_1, \sigma_2; T)$ be the number of zeros of $\zeta(s, \mathcal{A})$ on the rectangular region $\sigma_1 < \operatorname{Re} s < \sigma_2$, $0 < \operatorname{Im} s < T$. Theorem 3.1 gives a lower bound for $N_{\zeta(s, \mathcal{A})}(\sigma_1, \sigma_2; T)$ on the assumption of Conjecture 1. What can we say about an upper bound for $N_{\zeta(s, \mathcal{A})}(\sigma_1, \sigma_2; T)$? The following theorem gives an answer.

Theorem 3.2. *Let K be a number field and G^f be its ray class group. Assume Conjecture 1 for all Hecke L -functions $L(s, \chi)$ with $\chi : G^f \rightarrow S^1$. Suppose that $\#G^f > 1$ and $\mathcal{A} \in G^f$. Then, the number of zeros of the partial zeta function $\zeta(s, \mathcal{A})$ on the rectangular region $\operatorname{Re} s > \sigma_0$, $0 < \operatorname{Im} s < T$ is less than*

$$\ll T$$

for any fixed $\sigma_0 > \frac{1}{2}$.

Proof. By Littlewood's lemma, we have

$$2\pi \int_{\sigma}^{\infty} N(u, T) du = \int_0^T \log |\zeta(\sigma + it, \mathcal{A})| dt + O(\log T),$$

where $N(u, T)$ denotes the number of zeros of $\zeta(s, \mathcal{A})$ on the region $\operatorname{Re} s > u$, $0 < \operatorname{Im} s < T$. Thus, it is enough to show that the integral on the right is less than $\ll T$.

We are going to use a simple inequality. First,

$$\left| \frac{1}{J} \sum_{j \leq J} z_j \right| \leq \frac{1}{J} \sum_{j \leq J} |z_j| \leq \max_{j \leq J} |z_j|.$$

Take logarithms on both sides, then

$$(3.5) \quad \log \left| \frac{1}{J} \sum_{j \leq J} z_j \right| \leq \max_{j \leq J} \log |z_j|.$$

By (3.4) and (3.5), we have

$$\int_0^T \log |\zeta(\sigma + it, \mathcal{A})| dt \ll \int_0^T \max_{\chi} \log |L(\sigma + it, \chi)| dt.$$

Apply Lemma 2 of [6], then we have

$$\begin{aligned} \int_0^T \max_{\chi} \log |L(\sigma + it, \chi)| dt &\leq \int_0^T \max_{\chi} \left| \log |L(\sigma + it, \chi)| dt - \operatorname{Re} \sum_{p \leq X} \frac{a(p, \chi)}{p^{\sigma + it}} \right| dt \\ &\quad + \int_0^T \max_{\chi} \left| \operatorname{Re} \sum_{p \leq X} \frac{a(p, \chi)}{p^{\sigma + it}} \right| dt \\ &\leq \sum_{\chi} \int_0^T \left| \log |L(\sigma + it, \chi)| dt - \operatorname{Re} \sum_{p \leq X} \frac{a(p, \chi)}{p^{\sigma + it}} \right| dt \\ &\quad + \sum_{\chi} \int_0^T \left| \sum_{p \leq X} \frac{a(p, \chi)}{p^{\sigma + it}} \right| dt, \end{aligned}$$

where $X = T^{\kappa}$, $0 < \kappa < \frac{1}{2}$, $a(p, \chi) = \sum_{N\mathfrak{p}=p, \mathfrak{p}|p} \chi(\mathfrak{p})$, $|a(p, \chi)| \leq [K : \mathbb{Q}]$. By Cauchy's inequality and Lemma 2 of [6], we have

$$\begin{aligned} \int_0^T \left| \log |L(\sigma + it, \chi)| dt - \operatorname{Re} \sum_{p \leq X} \frac{a(p, \chi)}{p^{\sigma + it}} \right| dt \\ \ll \sqrt{T} \left(\int_0^T \left| \log |L(\sigma + it, \chi)| dt - \operatorname{Re} \sum_{p \leq X} \frac{a(p, \chi)}{p^{\sigma + it}} \right|^2 dt \right)^{\frac{1}{2}} \ll T, \end{aligned}$$

and by Lemma 2.7

$$\begin{aligned} \int_0^T \left| \sum_{p \leq X} \frac{a(p, \chi)}{p^{\sigma + it}} \right| dt &\ll \sqrt{T} \left(\int_0^T \left| \sum_{p \leq X} \frac{a(p, \chi)}{p^{\sigma + it}} \right|^2 dt \right)^{\frac{1}{2}} \\ &= \sqrt{T} \left(\sum_{p \leq X} \frac{|a(p, \chi)|^2}{p^{2\sigma}} (T + O(p)) \right)^{\frac{1}{2}} \ll T. \end{aligned}$$

Thus, the proof is complete. □

§ 4. Concluding remarks

The author [5] improved Theorem 1.1 by obtaining asymptotic formula $cT + o(T)$ for the number of zeros of Epstein zeta function $E(s, Q)$ on the rectangular region $\frac{1}{2} < \sigma_1 < \operatorname{Re} s < \sigma_2$, $0 < \operatorname{Im} s < T$ with the constant c has an integral formula $c = \int_{\sigma_1}^{\sigma_2} \mu(\sigma) d\sigma$ for some density function $\mu(\sigma)$. The main ingredient of the proof is the method given by Borchsenius and Jessen [2]. Based on Theorems 3.1 and 3.2, we expect the following statement.

Conjecture 4.1. *Let K be a number field and let G^f be its ray class group. Let $\mathcal{A} \in G^f$. Then, the number of zeros of partial zeta function $\zeta(s, \mathcal{A})$ on the region $\frac{1}{2} < \sigma_1 < \operatorname{Re} s < \sigma_2$, $0 < \operatorname{Im} s < T$ is*

$$= cT + o(T),$$

where $c = \int_{\sigma_1}^{\sigma_2} \mu(\sigma) d\sigma$ for some density function $\mu(\sigma)$ depending on \mathcal{A} .

The simplest case $K = \mathbb{Q}$ is considered and completed by the author with Haseo Ki [3].

References

- [1] Bauer, H., The value distribution of Artin L -series and zeros of zeta-functions, *J. Number Theory*, **98** (2003), 254–279.
- [2] Borchsenius, V. and Jessen, B., Mean Motions and values of the Riemann zeta function, *Acta Math.*, **80** (1948), 97–166.
- [3] Ki, H. and Lee, Y., On the zeros of degree one L -functions from the extended Selberg class, *Acta Arith.*, **149** (2011), 23–36.
- [4] Karatsuba, A. A. and Voronin, S. M., The Riemann Zeta-Function, *Berlin, New York: Walter de Gruyter*, 1992.
- [5] Lee, Y., On the zeros of Epstein zeta functions, *Forum Math.*, (to appear), available at <http://arxiv.org/abs/1204.6297>.
- [6] Lee, Y., The Universality Theorem for Hecke L -functions, *Math Z.* online first, DOI: 10.1007/s00209-011-0895-6.
- [7] Montgomery, H. L. and Vaughan, R. C. Hilbert's Inequality, *J. London Math. Soc. (2)*, **8** (1974), 73–82.
- [8] Neukirch, J., Algebraic number theory, *Grundlehren der Mathematischen Wissenschaften* 322, Springer-Verlag, Berlin, 1999.
- [9] Voronin, S. M., The zeros of zeta-functions of quadratic forms. (Russian), *Trudy Mat. Inst. Steklov.* **142** (1976), 135–147.