TITLE:

Flash generation of α -(trifluoromethyl)vinyl lithium and application to continuous flow three-component synthesis of α -trifluoromethyl amides.

AUTHOR(S):

Nagaki, Aiichiro; Tokuoka, Shinya; Yoshida, Jun-ichi

CITATION:

Nagaki, Aiichiro ...[et al]. Flash generation of α -(trifluoromethyl)vinyl lithium and application to continuous flow three-component synthesis of α -trifluoromethyl amides.. Chemical communications 2014, 50(95): 15079-15081

ISSUE DATE:

2014-10-21

URL:

<http://hdl.handle.net/2433/198580>

RIGHT:

© The Royal Society of Chemistry 2014; 許諾条件により本文ファイルは 2015-10-21 に公開.; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

COMMUNICATION

Flash Generation of α -(Trifluoromethyl)vinyl lithium and an Application to Continuous Flow Three-Component Synthesis of α -Trifluoromethylamides†

Aiichiro Nagaki,^a Shinya Tokuoka,^a and Jun-ichi Yoshida^{*a}

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX
DOI: 10.1039/b000000x

α -(Trifluoromethyl)vinyl lithium was generated and used for the reaction with electrophiles at $-78\text{ }^{\circ}\text{C}$ in a flow microreactor although the batch reaction should be carried out at ca. $-100\text{ }^{\circ}\text{C}$, and the method was applied to continuous flow three-component synthesis of α -trifluoromethylamides.

Trifluoromethyl-substituted vinylmetals¹ such as α -(trifluoromethyl)vinyl lithium serve as powerful building blocks² for constructing CF_3 -containing molecules.³ However, because α -(trifluoromethyl)vinyl lithium is very unstable,⁴ it should be generated and reacted at extremely low temperatures if we use batch macro reactors. In fact, Tarrant^{4a} and Ichikawa^{2c} reported independently that the generation of α -(trifluoromethyl)vinyl lithium followed by the reaction with an electrophile should be carried out at ca. $-100\text{ }^{\circ}\text{C}$ because of rapid elimination of LiF to give 1,1-difluoroallene. Notably, batch reactions are often carried out using an excess amount of α -(trifluoromethyl)vinyl lithium, and the yields are usually reported based on the amounts of electrophiles. Here, we show that flash chemistry^{5,6} using flow microreactor systems^{7,8,9} enables the generation and reactions of α -(trifluoromethyl)vinyl lithium at more easily accessible temperatures to give the desired products in good yields without using an excess amount of the lithium reagent. We also report here that the method enables continuous flow three-component synthesis^{10,11,12} of α -trifluoromethylamides,¹³ which serve as fascinating motifs in peptidomimetics.

A flow microreactor system consisting of three T-shaped micromixers (M1, M2 and M3) and three microtube reactors (R1, R2 and R3) shown in Figure 1 was used for Br/Li exchange of 2-bromo-3,3,3-trifluoropropene with $s\text{-BuLi}$ to generate α -(trifluoromethyl)vinyl lithium followed by reactions with electrophiles.

First, we examined the reaction using benzaldehyde as an electrophile, and the desired product was obtained in a good yield (79%) based on 2-bromo-3,3,3-trifluoropropene at $-78\text{ }^{\circ}\text{C}$ (Table 1). The reactions with other aldehydes also gave the corresponding products in good yields as shown in Table 1. Thus, the flash method enables efficient generation and use of α -(trifluoromethyl)vinyl lithium at easily accessible conditions.

Fig. 1 A flow microreactor system for generation of α -(trifluoromethyl)vinyl lithium followed by reactions with electrophiles and the subsequent quenching with methanol or acetic acid. T-shaped micromixers: M1, M2 and M3, microtube reactors: R1, R2 and R3.

Table 1 Reactions of α -(trifluoromethyl)vinyl lithium with various electrophiles.

electrophile	quenching agent	product	yield (%)
PhCHO	MeOH		79
(<i>p</i> - CF_3)PhCHO	MeOH		72
	MeOH		90
$\text{O}=\text{C}=\text{N}-\text{Bn}$	MeOH		80
	AcOH		62
$\text{O}=\text{C}=\text{N}-\text{Ph}$	MeOH		65
	AcOH		63
$\text{O}=\text{C}=\text{N}-\text{Bu}$	MeOH		83

The reactions with isocyanates are interesting. Quenching with methanol gave the compounds having a methoxy group, although the use of acetic acid as a quenching agent gave the desired alkenes having CF₃ and amide groups in good yields (Table 1).

The formation of the compounds having a methoxy group can be explained by deprotonation of methanol by the initially formed lithium amide followed by the addition of the methoxide ion to the carbon-carbon double bond activated by the electron withdrawing CF₃ and amide groups. This explanation leads to the idea of using carbon nucleophiles having an acidic proton instead of methanol (Figure 2). The lithium amide deprotonates a subsequently added carbon nucleophile (Nu-H), and the resulting carbanion (Nu⁻ Li⁺) adds to the electron-deficient carbon-carbon double bond. Protonation gives the corresponding three-component coupling products.

Fig. 2 The reaction of α -(trifluoromethyl)vinyllithium with an isocyanate followed by the reaction with a subsequently added nucleophile.

We chose to use malonate esters as carbon nucleophiles. Thus, the Br/Li exchange of 2-bromo-3,3,3-trifluoropropene, the reaction of the resulting α -(trifluoromethyl)vinyllithium with an isocyanate, and the subsequent reaction with a malonate ester were integrated¹⁴ using a flow microreactor system consisting of four T-shaped micromixers (M1, M2, M3 and M4) and four microtube reactors (R1, R2, R3 and R4) shown in Figure 3.

Fig. 3 An integrated flow microreactor system for generation of α -(trifluoromethyl)vinyllithium followed by the reactions with isocyanates and subsequently added malonate esters. T-shaped micromixers: M1, M2, M3 and M4, microtube reactors: R1, R2, R3 and R4.

As shown in Table 2, the integration of the reactions was successfully achieved with diethyl malonate, dibenzyl malonate, and di-*tert*-butyl malonate to obtain the corresponding addition products in good yields, although the use of 1,3-diketones such as 2,4-pentanedione did not give the desired products.

Table 2 Three-component coupling of α -(trifluoromethyl)vinyllithium, isocyanate, and malonate esters.

malonate ester	isocyanate	product	yield (%)
	O=C=N-Bn		80
	O=C=N-Ph		65
	O=C=N-Bu		73
	O=C=N-Bn		51
	O=C=N-Bn		66

In conclusion, flash chemistry enables efficient generation of α -(trifluoromethyl)vinyllithium and its reactions with electrophiles at -78 °C. Space integration of reactions enabled three-component synthesis of α -trifluoromethylamides by successive reactions of α -(trifluoromethyl)vinyllithium with isocyanates and malonate esters. The key to the success of the three-component coupling is deprotonation of malonate esters by the lithium amide intermediates followed by the nucleophilic attack of the resulting carbanions on the carbon-carbon double bond activated by the CF₃ and the amide groups. This mode of reaction integration adds a new dimension to continuous flow multicomponent coupling.¹⁵ Further applications of the present method to the continuous-flow synthesis of fluorine-containing organic compounds¹⁶ are currently in progress in our laboratory.

Acknowledgements

This work was partially supported by the Grant-in-Aid for Scientific Research (S) (no. 26220804) and Scientific Research (B) (no. 26288049). We also thank Taiyo Nippon Sanso for providing a low temperature cooling device and partial financial support.

Notes and references

- Reviews on reactions of α -(trifluoromethyl)vinyl compounds: (a) K. Uneyama, *Organofluorine Chemistry*, Blackwell, Oxford, 2006; (b) M. Shimizu and T. Hiyama, *Angew. Chem., Int. Ed.*, 2005, **44**, 214; (c) J. M. Percy, *Top. Curr. Chem.*, 1997, **193**, 131; (d) P. Lin and J. Jiang, *Tetrahedron*, 2000, **56**, 3635; (e) H. Ito, *Adv. Polym. Sci.*, 2005, **172**, 37; (f) H. Ito, H. D. Truong, M. Okazaki and R. A. DiPietro, *J.*

- Photopolym. Sci. Technol.*, 2003, **16**, 523.
- ² (a) K. Iseki, Y. Kuroki, T. Nagai and Y. Kobayashi, *J. Fluorine Chem.*, 1994, **69**, 5; (b) Q. Chen and F. L. Qing, *Tetrahedron*, 2007, **63**, 11965; (c) R. Nadano, K. Fuchibe, M. Ikeda, H. Takahashi and J. Ichikawa, *Chem. Asian J.*, 2010, **5**, 1875; (d) H. Lebel, M. Davi and G. T. Stoklosa, *J. Org. Chem.*, 2008, **73**, 6828; (e) V. De Matteis, F. L. van Delft, J. Tiebes and F. P. J. T. Rutjes, *Synlett*, 2008, 351; (f) T. Yamazaki and N. Ishikawa, *Chem. Lett.*, 1984, **13**, 521; (g) B. Jiang and Y. Xu, *J. Org. Chem.*, 1991, **56**, 7336; (h) Y. Xu, F. Jin and W. Huang, *J. Org. Chem.*, 1994, **59**, 2638; (i) B. Jiang, Q. Wang, C. Yang and M. Xu, *Tetrahedron Lett.*, 2001, **42**, 4083; (j) R. Pan, X. Liu and M. Deng, *J. Fluorine Chem.*, 1999, **95**, 167; (k) C. M. Hu, F. Hong and Y. Y. Xu, *J. Fluorine Chem.*, 1993, **64**, 1; (l) S. Peng and F. L. Qing, *J. Chem. Soc. Perkin Trans. 1*, 1999, 3345; (m) F. Hong, X. Tang and C. Hu, *J. Chem. Soc. Chem. Commun.*, 1994, 289; (n) Q. Chen and F. L. Qing, *Tetrahedron*, 2007, **63**, 11965; (o) I. Nowak and M. J. Robins, *J. Org. Chem.*, 2007, **72**, 2678; (p) T. Konno, J. Chae, T. Tanaka, T. Ishihara and H. Yamanaka, *J. Fluorine Chem.*, 2006, **127**, 36; (q) T. Hanamoto, N. Morita and K. Shindo, *Eur. J. Org. Chem.*, 2003, 4279; (r) T. Kobayashi, T. Eda, O. Tamura and H. Ishibashi, *J. Org. Chem.*, 2002, **67**, 3156; (s) X. Liu, M. Shimizu and T. Hiyama, *Angew. Chem., Int. Ed.*, 2004, **43**, 879; (t) T. Yamazaki, T. Kawasaki-Takasuka, A. Furuta and S. Sakamoto, *Tetrahedron*, 2009, **65**, 5945.
 - ³ (a) Modern Fluoroorganic Chemistry: Synthesis Reactivity, Applications (Ed.: P. Kirsch), Wiley-VCH: Weinheim, 2004; (b) Fluorine and Health: Molecular Imaging, Biomedical Materials and Pharmaceuticals (Eds.: A. Tressaud and G. Haufe), Elsevier: Amsterdam, 2008.
 - ⁴ (a) F. G. Drakesmith, O. J. Stewart and P. Tarrant, *J. Org. Chem.*, 1968, **33**, 280; (b) W. R. Dolbier, Jr. C. R. Burkholder and C. A. Piedrahita, *J. Fluorine Chem.*, 1982, **20**, 637.
 - ⁵ Flash chemistry is defined as a field of chemical synthesis where extremely fast reactions are conducted in a highly controlled manner to produce the desired compounds with high selectivity: (a) J. Yoshida, *Flash Chemistry. Fast Organic Synthesis in Microsystems*; Wiley-Blackwell, 2008; (b) J. Yoshida, A. Nagaki and T. Yamada, *Chem. Eur. J.*, 2008, **14**, 7450; (c) J. Yoshida, *Chem. Rec.*, 2010, **10**, 332; (d) J. Yoshida, Y. Takahashi and A. Nagaki, *Chem. Commun.*, 2013, **49**, 9896.
 - ⁶ (a) H. Usutani, Y. Tomida, A. Nagaki, H. Okamoto, T. Nokami and J. Yoshida, *J. Am. Chem. Soc.*, 2007, **129**, 3046; (b) A. Nagaki, H. Kim and J. Yoshida, *Angew. Chem., Int. Ed.*, 2008, **47**, 7833; (c) A. Nagaki, H. Kim and J. Yoshida, *Angew. Chem., Int. Ed.*, 2009, **48**, 8063; (d) A. Nagaki, E. Takizawa and J. Yoshida, *J. Am. Chem. Soc.*, 2009, **131**, 1654; (e) Y. Tomida, A. Nagaki and J. Yoshida, *J. Am. Chem. Soc.*, 2011, **133**, 3744; (f) H. Kim, A. Nagaki and J. Yoshida, *Nat. Commun.*, 2011, **2**, 264.
 - ⁷ Books on flow microreactor synthesis: (a) W. Ehrfeld, V. Hessel and H. Löwe, *Microreactors*; Wiley-VCH: Weinheim, 2000; (b) V. Hessel, S. Hardt and H. Löwe, *Chemical Micro Process Engineering*; Wiley-VCH Verlag: Weinheim, 2004; (c) V. Hessel, A. Renken, J. C. Schouten and J. Yoshida, *Micro Process Engineering*; Wiley-Blackwell, 2009.
 - ⁸ Reviews on flow microreactor synthesis: (a) K. Jähnisch, V. Hessel, H. Löwe and M. Baerns, *Angew. Chem., Int. Ed.*, 2004, **43**, 406; (b) G. N. Doku, W. Verboom, D. N. Reinhoudt and A. van den Berg, *Tetrahedron*, 2005, **61**, 2733; (c) J. Yoshida, A. Nagaki, T. Iwasaki and S. Suga, *Chem. Eng. Tech.*, 2005, **3**, 259; (d) P. Watts and S. J. Haswell, *Chem. Soc. Rev.*, 2005, **34**, 235; (e) K. Geyer, J. D. C. Codée and P. H. Seeberger, *Chem. Eur. J.*, 2006, **12**, 8434; (f) A. J. deMello, *Nature*, 2006, **442**, 394; (g) H. Song, D. L. Chen and R. F. Ismagilov, *Angew. Chem., Int. Ed.*, 2006, **45**, 7336; (h) J. Kobayashi, Y. Mori and S. Kobayashi, *Chem. Asian J.*, 2006, **1**, 22; (i) M. Brivio, W. Verboom and D. N. Reinhoudt, *Lab Chip*, 2006, **6**, 329; (j) B. P. Mason, K. E. Price, J. L. Steinbacher, A. R. Bogdan and D. T. McQuade, *Chem. Rev.*, 2007, **107**, 2300; (k) B. Ahmed-Omer, J. C. Brandt and T. Wirth, *Org. Biomol. Chem.*, 2007, **5**, 733; (l) P. Watts and C. Wiles, *Chem. Commun.*, 2007, 443; (m) T. Fukuyama, M. T. Rahman, M. Sato and I. Ryu, *Synlett*, 2008, 151; (n) R. L. Hartman and K. F. Jensen, *Lab Chip*, 2009, **9**, 2495; (o) J. P. McMullen and K. F. Jensen, *Annu. Rev. Anal. Chem.*, 2010, **3**, 19; (p) J. Yoshida, H. Kim and A. Nagaki, *ChemSusChem*, 2011, **4**, 331; (q) C. Wiles and P. Watts, *Green Chem.*, 2012, **14**, 38; (r) A. Kirschning, L. Kupracz and J. Hartwig, *Chem. Lett.*, 2012, **41**, 562; (s) D. T. McQuade and P. H. Seeberger, *J. Org. Chem.*, 2013, **78**, 6384; (t) K. S. Elvira, X. C. Solvas, R. C. R. Wootton and A. J. deMello, *Nat. Chem.*, 2013, **5**, 905; (u) J. C. Pastre, D. L. Browne and S. V. Ley, *Chem. Soc. Rev.*, 2013, **42**, 8849; (v) I. R. Baxendale, *J. Chem. Technol. Biotechnol.*, 2013, **88**, 519; (w) L. Malet-Sanz and F. Susanne, *J. Med. Chem.*, 2012, **55**, 4062.
 - ⁹ Some selected recent examples: (a) D. Cantillo, M. Baghbanzadeh and C. O. Kappe, *Angew. Chem., Int. Ed.*, 2012, **51**, 10190; (b) W. Shu and S. L. Buchwald, *Angew. Chem., Int. Ed.*, 2012, **51**, 5355; (c) F. Lévesque and P. H. Seeberger, *Angew. Chem., Int. Ed.*, 2012, **51**, 1706; (d) K. C. Basavaraju, S. Sharma, R. A. Maurya and D. P. Kim, *Angew. Chem., Int. Ed.*, 2013, **52**, 6735; (e) C. Brancour, T. Fukuyama, Y. Mukai, T. Skrydstrup and I. Ryu, *Org. Lett.*, 2013, **15**, 2794; (f) J. D. Nguyen, B. Reiß, C. Dai and C. R. J. Stephenson, *Chem. Commun.*, 2013, **49**, 4352; (g) C. Battilocchio, J. M. Hawkins and S. V. Ley, *Org. Lett.*, 2013, **15**, 2278; (h) A. S. Kleinke and T. F. Jamison, *Org. Lett.*, 2013, **15**, 710; (i) K. Asano, Y. Uesugi and J. Yoshida, *Org. Lett.*, 2013, **15**, 2398; (j) A. Nagaki, D. Ichinari and J. Yoshida, *Chem. Commun.*, 2013, **49**, 3242; (k) L. Guetzoyan, N. Nikbin, I. R. Baxendale and S. V. Ley, *Chem. Sci.*, 2013, **4**, 764; (l) S. Fuse, Y. Mifune and T. Takahashi, *Angew. Chem., Int. Ed.*, 2014, **53**, 851; (m) A. Nagaki, Y. Takahashi and J. Yoshida, *Chem. Eur. J.*, 2014, **20**, 7931.
 - ¹⁰ Reviews on domino, tandem and cascade reactions: (a) L. F. Tietze, *Chem. Rev.*, 1996, **96**, 115; (b) I. Ryu and N. Sonoda, *Chem. Rev.*, 1996, **96**, 177; (c) P. J. Parsons, C. S. Penkett and A. J. Shell, *Chem. Rev.*, 1996, **96**, 195.
 - ¹¹ Reviews on multicomponent coupling reactions: (a) A. Dömling and I. Ugi, *Angew. Chem., Int. Ed.*, 2000, **39**, 3168; (b) H. Bienaymé, C. Hulme, O. Oddon and P. Schmitt, *Chem. Eur. J.*, 2000, **6**, 3321; (c) I. Ugi, *Pure Appl. Chem.*, 2001, **73**, 187; (d) B. Ganem, *Acc. Chem. Res.*, 2009, **42**, 463.
 - ¹² Selected examples of one-pot sequential reactions: (a) A. Orita, J. Yaruva and J. Otera, *Angew. Chem., Int. Ed.*, 1999, **38**, 2267; (b) A. Orita, N. Yoshioka, P. Struwe, A. Braier, A. Beckmann and J. Otera, *Chem. Eur. J.*, 1999, **5**, 1355; (c) P. A. Clarke, S. Santos and W. H. C. Martin, *Green Chem.*, 2007, **9**, 438; (d) T. Nokami, H. Tsuyama, A. Shibuya, T. Nakatsutsumi and J. Yoshida, *Chem. Lett.*, 2008, **37**, 942; (e) Y. Numata, J. Kawashima, T. Hara and Y. Tajima, *Chem. Lett.*, 2008, **37**, 1018; (f) K. Yamaguchi, M. Kotani, K. Kamata and N. Mizuno, *Chem. Lett.*, 2008, **37**, 1258.
 - ¹³ CF₃-substituted amides: (a) K. Uneyama, O. Morimoto and H. Nanbu, *Tetrahedron Lett.*, 1989, **30**, 109; (b) J. J. Morris, L. R. Hughes, A. T. Glen and P. J. Taylor, *J. Med. Chem.*, 1991, **34**, 447; (c) J. Walkowiak, M. Tomas-Szwaczkyk, G. Haufe and H. Koroniak, *J. Fluorine Chem.*, 2012, **143**, 189.
 - ¹⁴ Space integration of reactions: (a) S. Suga, D. Yamada and J. Yoshida, *Chem. Lett.*, 2010, **39**, 404; (b) A. Nagaki, A. Kenmoku, Y. Moriwaki, A. Hayashi and J. Yoshida, *Angew. Chem., Int. Ed.*, 2010, **49**, 7543; (c) J. Yoshida, K. Saito, T. Nokami and A. Nagaki, *Synlett*, 2011, **9**, 1189; (d) A. Nagaki, C. Matsuo, S. Kim, K. Saito, A. Miyazaki and J. Yoshida, *Angew. Chem., Int. Ed.*, 2012, **51**, 3245; (e) A. Nagaki, Y. Moriwaki and J. Yoshida, *Chem. Commun.*, 2012, **48**, 11211.
 - ¹⁵ (a) W. Stacy and M. G. Organ, *J. Comb. Chem.*, 2007, **9**, 14; (b) P. R. D. Murray, D. L. Browne, J. C. Pastre, C. Butters, D. Guthrie and S. V. Ley, *Org. Process Res. Dev.*, 2013, **17**, 1192; (c) D. T. McQuade and P. H. Seeberger, *J. Org. Chem.*, 2013, **78**, 6384; (d) K. S. Nalivela, M. Tilley, M. A. McGuire, and M. G. Organ, *Chem. Eur. J.*, 2014, **20**, 6603; (e) J. Wu, J. A. Kozak, F. Simeon, T. A. Hatton, and T. F. Jamison, *Chem. Sci.*, 2014, **5**, 1227.
 - ¹⁶ (a) H. Amii, A. Nagaki, and J. Yoshida, *Beilstein J. Org. Chem.*, 2013, **9**, 2793. (b) C. B. McPake and G. Sandford, *Org. Process. Res. Dev.*, 2012, **16**, 844.