TITLE:

Ecogeographical and phylogenetic effects on craniofacial variation in macaques.

AUTHOR(S):

Ito, Tsuyoshi; Nishimura, Takeshi; Takai, Masanaru

CITATION:

Ito, Tsuyoshi ...[et al]. Ecogeographical and phylogenetic effects on craniofacial variation in macaques.. American journal of physical anthropology 2014, 154(1): 27-41

ISSUE DATE:

2014-01-22

URL:

<http://hdl.handle.net/2433/198636>

RIGHT:

This is the peer reviewed version of the following article: Ito, T., Nishimura, T. and Takai, M. (2014), Ecogeographical and phylogenetic effects on craniofacial variation in macaques. Am. J. Phys. Anthropol., 154: 27-41, which has been published in final form at <http://dx.doi.org/10.1002/ajpa.22469>; この論文は著者最終稿です。内容が印刷版と異なることがありますので、引用の際には出版社版をご確認ご利用ください。 This is the Accepted Author Manuscript. Please cite only the published version.

Ecogeographical and phylogenetic effects on craniofacial variation in macaques

Tsuyoshi Ito, Takeshi Nishimura, Masanaru Takai

Primate Research Institute, Kyoto University, Inuyama, Aichi 484-8506, Japan

44 text pages, 7 figures, 6 tables.

Abbreviated title: Craniofacial variation in macaques

Key words: biogeography; adaptive convergence; allometry; Bergmann's rule; geometric morphometrics

Corresponding author: Tsuyoshi Ito

Primate Research Institute, Kyoto University, Inuyama, Aichi 484-8506, Japan

Tel & Fax: +81-568-63-0536

E-mail: ito.tsuyoshi.3r@kyoto-u.ac.jp

This study was financially supported by AS-HOPE and ITP-HOPE Programs from JSPS (to T. I.), Grant-in-Aid for JSPS Fellows (to T. I.), Grant-in-Aid for Challenging Exploratory Research (to T. N.), and Global COE Program from MEXT (Grant A06 to Kyoto University).

ABSTRACT

The widespread and complex ecogeographical diversity of macaques may have caused adaptive morphological convergence among four phylogenetic subgroups, making their phylogenetic relationships unclear. We used geometric morphometrics and multivariate analyses to test the null hypothesis that craniofacial morphology does not vary with ecogeographical and phylogenetic factors. As predicted by Bergmann's rule, size was larger for the *fascicularis* and *sinica* groups in colder environments. No clear size cline was observed in the *silenus* and *sylvanus* groups. An allometric pattern was observed across macaques, indicating that as size increases, rounded faces become more elongated. However, the elevation was differentiated within each of the former two groups and between the *silenus* and *sylvanus* groups, and the slope decreased in each of the two northern species of the *fascicularis* group. All allometric changes resulted in the similar situation of the face being more rounded in animals inhabiting colder zones and/or in animals having a larger body size than that predicted from the overarching allometric pattern. For non-allometric components, variations in prognathism were significantly correlated with dietary differences; variations in localized shape components in zygomatics and muzzles were significantly correlated with phylogenetic differences among the subgroups. The common allometric pattern was probably influenced directly or indirectly by climate-related factors, which are pressures favoring a more rounded face in colder environments and/or a more elongated face in warmer

environments. Allometric dissociation could have occurred several times in *Macaca* even within a subgroup because of their wide latitudinal distributions, critically impairing the taxonomic utility of craniofacial elongation.

INTRODUCTION

Craniofacial morphology widely varies among extant macaque monkeys (*Macaca* spp.) (Albrecht, 1978; Pan et al., 1998; Pan and Oxnard, 2000, 2001, 2002). Craniofacial features have been used to taxonomically identify fossil macaques (Schlosser, 1924; Iwamoto, 1975; Delson, 1977; Fooden, 1990; Rook and O'Higgins, 2005); however, classification is occasionally controversial (Delson, 1980; Jablonski and Pan, 1988; Jablonski, 2002), possibly because of some homoplastic features related to their salient and complicated ecogeographical diversity.

Macaques are distributed across a wide range of climates from tropical to temperate zones. Such a wide distribution is not found in any other primate genus except humans (Fleagle, 1999). Approximately 20 species of extant macaques are phylogenetically classified into four species groups (i.e., *sylvanus*, *silenus*, *sinica*, and *fascicularis* groups) based on genitalia, cranial morphology, biogeography, and fossil records (Fooden, 1976; Delson, 1980) (Fig. 1A; Table 1). This traditional classification is supported by recent molecular studies (Tosi et al., 2000, 2003; Li et al., 2009) (Fig. 1B–D). The *sylvanus* group (*M. sylvanus*) is distributed in a small temperate region of northern Africa (Fooden, 2007). The *silenus* group is found in tropical areas of southern India and from Sundaland to Indochina (Fooden, 1969, 1975). The *sinica* group occupies tropical India and Sri Lanka and also inhabits subtropical and temperate areas between Nepal and eastern China (Fooden, 1985, 1990). The *fascicularis*

group has the widest continuous distribution, ranging from tropical Indonesia to subtropical and temperate areas between eastern Afghanistan and Japan (Fooden, 2006). Notably, the latitudinal distribution ranges of the four species groups considerably overlap, and the species groups often share similar climatic conditions. This observation suggests that geographic dispersal among various climatic conditions has independently occurred for each species group rather than accompanying species group divergence.

Such a complex ecogeographical diversity, which is independent of phylogenetic (among the species groups) divergence, inevitably obscures phylogenetic value of some morphological features. For example, each species group would have independently modified its craniofacial morphology in response to the varied environmental conditions, which may have caused convergent evolution among the species groups.

One of the major convergent factors is a latitudinal gradient in body mass (Albrecht, 1980; Fooden, 1988; Fooden and Albrecht, 1993; Hamada et al., 1996; Fooden, 2006). This gradient is usually explained by Bergmann's rule, which predicts that animals living in northern, colder environments have a larger body size than those living in southern, warmer regions, as a physiological adaptation to effectively maintain body temperature. This latitudinal gradient in body mass is strongly correlated with craniofacial shape variation across macaques, except for Sulawesi macaques, which have an extremely unusual craniofacial shape (Albrecht, 1978). Positive facial allometry, indicating that the face

elongates relatively more with size, has been reported in several macaque species (Pan et al., 2003). Such allometric scaling probably caused significant convergence among the species groups. Despite belonging to different species groups, small-bodied macaques (i.e., *M. sinica*, *M. radiata*, and *M. fascicularis*) have numerous craniofacial shape similarities and are distinguished from large-bodied species (Pan and Oxnard, 2000, 2001). Thus, allometric scaling amplified by the effect of Bergmann's rule is probably the most effective factor in the formation of homoplastic craniofacial features among the species groups.

Other aspects of the effect of latitudinal environmental differences may also contribute to a convergence among the species groups with respect to craniofacial morphology. For example, because Allen's rule predicts that a more elongated body part is more liable to lose body heat than a more rounded one, animals in northern and colder environments may be under a selection pressure that favors a more rounded face with a shorter muzzle (as shown in the short muzzle of the arctic fox; Prestrud, 1991). A dietary shift is another possible cause of craniofacial modifications with latitude (Antón, 1996). Thus, even if the allometric effect is adjusted for, craniofacial shape possibly reflects latitudinal environmental differences that obscure phylogenetic information.

Overall, craniofacial shape variations among extant macaque species groups potentially contain a considerable number of homoplastic features caused by convergent evolution along latitudinal environmental gradients. In the present study, we investigated how environmental

factors influence craniofacial morphology in extant macaques in order to obtain a better understanding of the factors that underlie phenotypic variation and to evaluate the potential use of craniofacial shape in the taxonomic (species group) identification of fossil specimens. We distinguished among size, allometric variations, and non-allometric variations because environmental pressures can affect each of these three components independently and differently. Our null hypothesis is that none of the three components varies with ecogeographical or phylogenetic factors. We tested this null hypothesis for a better understanding of the factors that underlie phenotypic variations in extant macaques.

MATERIALS AND METHODS

Cranial samples and landmark coordinates

We examined 12 species of macaques, excluding Sulawesi macaques. We used 549 (243 female and 306 male) adult dry crania with erupted third molars (Table 2). The animals were shot or sampled in the wild. *M. fascicularis*, *M. mulatta*, *M. fuscata*, *M. assamensis*, and *M. nemestrina* individuals were divided into two or three populations based on geography and/or genetic distances, and the populations were regarded as sampling units (Table 2; Fig. 2) because they occupy wide ranges and/or have considerable genetic differentiation among populations (for references refer to Supporting information). Specimen samples for the other

species were not subdivided, and the species themselves were regarded as sampling units. We randomly selected specimens by a sampling unit (i.e., population or species) from as many localities as possible and reduced the number to approximately 20 per sex to avoid statistical artifacts related to sample size.

Three-dimensional cranial surface landmark coordinates were measured using a three-dimensional digitizer (Microscribe MX; Immersion Corp., San Jose, CA, USA). Of the 33 landmarks defined in Table 3, 23 were from the face and eight were from the neurocranium (the glabella was included in both sets of the face and the neurocranium) (Fig. 3). Only measurements on the left side were used to avoid redundancy, but horizontal reversals of the right-side measurements were used for specimens with broken left sides. Missing landmarks were estimated using mean sampling unit coordinate values by sex following registration procedures (see Slice, 1999; Cardini and Elton, 2008).

Ecogeographical and phylogenetic data

The geographic coordinate for each specimen was defined as the approximate center of the province, state, region, prefecture, district, or island that encompassed the sampling location. If the approximate center of the political jurisdiction or island was outside the actual distribution range of the species/population, the geographic coordinate was redefined as the approximate center of the overlapping area between the actual distribution range and the

political jurisdiction or island. Annual mean temperatures were taken from the BIOCLIM database on the WorldClim website (Hijmans et al., 2005; <http://www.worldclim.org/>) using the DIVA-GIS software (<http://www.diva-gis.org/>). The absolute value of the latitudinal coordinate was used in the following analysis, substituted for annual mean temperature because annual mean temperature can potentially reflect microscale regional outliers and localization artifacts that do not have evolutionary significance. Nevertheless, the results using annual mean temperature were similar and therefore not shown.

For phylogenetic data, the phylogenetic relationship revealed by *Alu* elements {*sylvanus* group, [*silenus* group, (*sinica* group, *fascicularis* group)]} (Fig. 1D; Table 1), was used because it is robust; this robustness is because the *Alu* elements show very little homoplasy and their ancestral state is known (Li et al., 2009). The phylogenetic relationship within a species group was not considered because of its uncertainty and was treated as a polytomy.

Categorical data were taken from the literature and summarized in Table 1 (refer to Supporting information).

Size variation

Centroid size (CS) for each specimen was calculated as the square root of the sum of the squared Euclidean distances from each landmark to the centroid. CS was calculated for each landmark set from the face, neurocranium, and entire cranium.

We regressed the natural logarithm of species average (split sex) body mass (g) against the natural logarithm of species average (split sex) CS for each of the three cranial regions to evaluate which cranial region was the most appropriate body mass predictor. Among the three regressions, the one that best explained the body mass data was selected (Fig. 4A). The coefficient (slope) for the selected regression was used to predict each individual's body mass, where an intercept was defined respectively for each species (split sex) so that a regression line passed through the coordinate of each species average (split sex). A body size surrogate was calculated as a predicted value by substituting each individual's lnCS for the independent variable in the regression equation that was defined for the species (split sex) to which the individual belonged (Fig. 4B). Body mass averages were primarily obtained from Delson et al. (2000); the data for *M. cyclopis* were from Fooden and Wu (2001).

A multivariate regression analysis was performed for the entire sample to evaluate factors influencing body size; the dependent variable was the body size surrogate, and the independent variables were sex, species group, habitat use type, and annual mean temperature. Sex and species group were nominal and thus respectively coded as dummy variables (e.g., if an individual belonged to the *sinica* group, it was coded as 1; if not, it was coded as 0). Habitat use type was an ordinal scale variable, where "arboreal" was coded as 0, "intermediate or uncertain" as 1, and "terrestrial" as 2. Diet was excluded as an independent variable because it was highly correlated with annual mean temperature ($r = -0.71$, $p < 0.001$).

and was not likely to be a direct factor affecting body size. We performed the same analyses for each species group.

Allometric variation

Standard geometric morphometric methodological approaches (Mitteroecker and Gunz, 2009; Zelditch et al., 2012) were employed. A generalized Procrustes analysis (Rohlf and Slice, 1990) was conducted to register landmark configurations of the face (23 landmarks). The registered data, i.e., Procrustes shape coordinates, were used as shape variables in the following analyses.

Multivariate allometric patterns of craniofacial shape variation have been studied using several methods. One of the most commonly used is based on the principal component analysis (PCA) of the covariance matrix of Procrustes shape coordinates. In this case, the first principal component (PC1), which accounts for the maximum variance, is usually plotted and/or regressed against a size variable to explore an allometric pattern (O'Higgins and Jones, 1998; Singleton, 2002). This approach is adequate when an allometric (size-correlated) component is relatively dominant in total variation, as in an ontogenetic sample or in a sample that has great size variation and is relatively homogeneous. However, depending on the nature of the sample, there is a risk that an allometric component is unexpectedly shared with PCs other than PC1, whereas PC1 contains some of the non-allometric variation. Because an

allometric component may not be dominant in total variation in our data, we instead applied the multiple regression of Procrustes shape coordinates (Monteiro, 1999). The multiple regression detects shape variation that the independent variable (size, in this case) best explains, which suits our aim and samples.

The multiple regression of the Procrustes shape coordinates for the face on the body size surrogate with a 10,000 times permutation test was performed for the entire sample to explore a common allometric pattern across macaques (MorphoJ; Klingenberg, 2011). Herein, shape scores, which are analogous to the PC1 scores in the study of allometry, were computed for all individuals. The shape score was defined as $s = yb^T(b^Tb)^{-0.5}$, where b^T is the transposed matrix of b (MorphoJ; Drake and Klingenberg, 2008; Klingenberg, 2011) when considering the regression equation $y = bx + e$, where y is the vector of shape variables, b is the regression vector, x is the size variable, and e is the error term. This is a projection of the shape variable vector onto regression vector b and can be interpreted as the shape variable that is most strongly associated with the size variable. Nevertheless, the shape score includes the residual variation in that direction. We examined the allometric pattern and its residual variation by plotting the shape scores against the body size surrogates. To evaluate the factors influencing differences in the allometric pattern (i.e., factors contributing to residual variations), the homogeneity of slopes and elevations for the least-squares regression of the shape scores on the body size surrogates was tested between the two climatic zones, among the four species

groups, among species within a species group, and between each species and a general pattern (a climatic zone).

The same analyses were conducted by regression of the PC1 scores on the body size surrogates following PCA of the covariance matrix for the Procrustes shape coordinates for the face to compare with previous studies.

Non-allometric (and sex-adjusted) variation

We performed multiple multivariate regression of the Procrustes shape coordinates for the face on the body size surrogate and sex, where sex was coded as a dummy variable (i.e., 0 or 1) to correct for the effect of sex and size on face shape. Residuals from the regression were calculated and used as size- and sex-adjusted shape variables (MorphoJ; Klingenberg, 2011).

The Mantel test (Mantel, 1967) was performed among all 19 sampling units to evaluate the association between shape and ecogeographical/phylogenetic factors. This test examines the correlation between the distance matrices of two of these variables. Shape distances were calculated as Mahalanobis squared distances of the size- and sex-adjusted shape variables. Climatic distances were calculated as Euclidean distances of the unit average for annual mean temperature, which was standardized. Dietary distances were calculated as Gower's distances (Gower, 1971) because these data are binary (i.e., the distances between “hard food eaters”

and “the others” were 1, and between the same values were 0). Phylogenetic distances were calculated as ultrametric distances (i.e., distances within a species group, between the *sinica* and *fascicularis* groups, between the *silenus* and *sinica/fascicularis* groups, and between the *sylvanus* group and the other groups were 1, 2, 3, and 4, respectively). The Mantel test with 100,000 random permutations was performed between the shape distances and the phylogenetic/climatic/dietary distances. Furthermore, a three-way Mantel test with 100,000 random permutations was performed (Smouse et al., 1986) to calculate the partial correlation between the shape distances and the climatic/dietary distances, while holding the effect of phylogeny constant. This was done to detect the true effect of climate/diet on shape by factoring out the confounding phylogenetic effect.

Canonical variate analysis (CVA) with 10,000 random permutations was performed for the size- and sex-adjusted shape variables to identify the shape features that best distinguished the four species groups and dietary differences.

Shape variations assessed by the multiple regression and CVA were represented by deformations of the Cartesian transformation grids calculated from triplets of thin plate splines (TPS; Bookstein, 1989; O'Higgins and Jones, 1998). The grids derived from TPS show how the space surrounding a reference shape is deformed into the space surrounding a target shape using minimum bending energy. The space deformations were further visualized

by reconstructing hypothetical forms of a wireframe and by smooth surface rendering of polygons.

These analyses were performed using Morphueus et al. (Slice, 1999), MorphoJ (Klingenberg, 2011), Morphologika ver. 2.5 (O'Higgins and Jones, 1998, 2006), NTSYSpc ver. 2.21a (Rohlf, 2005), and R software with the “cluster” and package.

RESULTS

Size variation

The natural logarithm of species average (split sex) body mass was better explained by neurocranial $\ln CS$ ($R^2 = 0.84$, $p < 0.001$, $y = 5.0191x - 13.763$; Fig. 4A) than by facial $\ln CS$ ($R^2 = 0.75$, $p < 0.001$) or entire cranial $\ln CS$ ($R^2 = 0.78$, $p < 0.001$). Thus, the body size surrogate was calculated for each individual on the basis of the neurocranial $\ln CS$ (Fig. 4B).

The factors influencing body size are listed in Table 4. The size variation in the entire sample was significantly explained by annual mean temperature as well as differences in sex, species group, and habitat use type. The (standard) partial regression coefficient indicated that size was larger under colder climatic conditions, in males than in females, in terrestrial animals than in arboreal animals, and in the *silenus* group > the *sinica* group > the *fascicularis* group (the *sylvanus* group was less significant). According to the standard partial regression

coefficient values for the *fascicularis* and *sinica* groups, climatic factors were considerably more effective predictors than sex and habitat use type. In contrast, climatic factors were less effective than sex and habitat use type for the *silenus* group. The contribution of sex was tested (40% was explained) for the *sylvanus* group, but the other factors were not tested because of the group's limited distribution and single habitat use type.

Allometric variation

Multiple regression analysis revealed that 8.8% of the total variance of the Procrustes shape coordinates was explained by the body size surrogate for the entire sample ($p < 0.001$), indicating that this regression vector represented a common allometric pattern across macaques. Increasing shape scores represented shape modifications; the upper face became smaller in width and length relative to the lower face, the muzzle was relatively longer, and the anterior portion of the face tilted more inferiorly, a feature known as klinorhynch (Fig. 5A). Taken together, these findings indicate that as size increases, a rounded face becomes more elongated.

The homogeneity of slopes and elevations was tested for the least-squares regression of the shape scores on the body size surrogates. The slopes were not different between the two climatic zones ($df = 1$, $F = 0.415$, $p = 0.52$; Fig. 5B), but elevations were significantly higher for animals in tropical zones than for animals in subtropical/temperate zones ($t = 14.69$, $p <$

0.001; Fig. 5B). In contrast, the slopes were significantly different among the four species groups ($df = 3$, $F = 8.401$, $p < 0.001$; Fig. 5C). The slope of the *fascicularis* group was significantly lower than those of the *sinica* group ($df = 1$, $F = 7.969$, $p = 0.005$) and the *silenus* group ($df = 1$, $F = 19.61$, $p < 0.001$). The slope of the *sinica* group was significantly lower than that of the *silenus* group ($df = 1$, $F = 6.603$, $p = 0.011$), although this difference was not significant either at adjusted Bonferroni $p < 0.05$ or after removing *M. sinica*, whose regression was not significant ($df = 1$, $F = 2.549$, $p = 0.112$). The slope of the *sylvanus* group appeared to be identical to that of the *silenus* group and higher than those of the other two groups (Fig. 5C), although these differences did not reach statistical significance, probably in part because of the limited sample size. The elevation of the *silenus* group was significantly higher than that of the *sylvanus* group ($t = -6.852$, $p < 0.001$).

Regressions for each species are summarized in Table 5. The slopes were significantly different among species within the *fascicularis* group ($df = 3$, $F = 3.996$, $p = 0.008$; Fig. 5D), with *M. fascicularis* > *M. mulatta* ($df = 1$, $F = 6.951$, $p = 0.009$) and *M. fascicularis* > *M. fuscata* ($df = 1$, $F = 7.133$, $p = 0.008$). The slopes were not different between *M. fascicularis* and *M. cyclopis* ($df = 1$, $F = 0.012$, $p = 0.915$), but the elevation was significantly higher for the former than for the latter ($t = -8.188$, $p < 0.001$). The slopes were not different among species within the *sinica* group after removing *M. sinica* ($df = 3$, $F = 0.715$, $p = 0.545$). Elevations seemed to be higher for *M. radiata* and *M. assamensis* than for *M. arctoides* and *M.*

thibetana (Fig. 5E); they were significantly different between *M. assamensis* and *M. thibetana* ($t = -3.768$, $p < 0.001$), between *M. assamensis* and *M. arctoides* ($t = 6.173$, $p < 0.001$), and between *M. radiata* and *M. arctoides* ($t = -2.683$, $p = 0.01$), although the last one was not significant at adjusted Bonferroni $p < 0.05$. Most *M. sinica* individuals tended to be scattered along or above the *M. radiata* and *M. assamensis* regression lines, although without statistical significance (Fig. 5E). The two species within the *silenus* group had similar slopes ($df = 1$, $F = 0.855$, $p = 0.358$) and elevations ($t = 0.396$, $p = 0.693$) (Fig. 5F).

The slopes of most species were not significantly different from the general one, i.e., the slope of the climatic zone to which a species belongs (Table 5). However, slopes were significantly lower in *M. mulatta* and *M. fuscata* than in the general one. Slopes appeared to be higher in *M. assamensis*, *M. thibetana*, and *M. nemestrina* than in the general one, although these differences did not reach statistical significance, probably in part because of the limited sample size.

Similar results, including the pattern of shape change, the parallel shift in the allometric trajectory between the two climatic zones, the pattern of slope dissociation among the four species groups, and their levels of statistical significance were also obtained from the regression analysis of PC1 scores on body size surrogates (detailed results not shown). However, PC1, which accounted for 29.1% of the total variance, demonstrated a larger amount of residual variation and thus was indicative of the inclusion of some non-allometric

components.

Non-allometric (and sex-adjusted) variation

The size- and sex-adjusted shape variables (i.e., residuals from the multiple multivariate regression of the Procrustes shape coordinates against the body size surrogate and sex; 15.6% was explained, $p < 0.001$) were tested for a correlation with ecogeographical and phylogenetic factors. The Mantel test revealed that the correlations between shape distances and phylogenetic distances were higher ($r = 0.52$, $p = 0.001$) than those with dietary distances ($r = 0.30$, $p = 0.039$) and climatic distances ($r = 0.19$, $p = 0.077$). The three-way Mantel test revealed that when controlling for phylogenetic effects, the correlation between shape distances and climatic distances was not significant ($r = 0.13$, $p = 0.147$), whereas that with dietary distances was weak but significant ($r = 0.27$, $p = 0.037$).

CVA demonstrated that the first three canonical variates accounted for 54.3%, 26.6%, and 19.1% of the total among-group variance, respectively. The first canonical variate (CV1) distinguished the *fascicularis* and *sylvanus* groups from the *sinica* and *silenus* groups (Fig. 6A). The former two groups represented relatively higher and wider upper muzzles than the latter two groups (Fig. 6B). CV2 distinguished the *sinica* group from the *silenus* group, whereas the *fascicularis* and *sylvanus* groups were intermediate between them (Fig. 6A). The *silenus* group exhibited a relatively narrow zygomatic and anteriorly convex nasal profile line,

and the *sinica* group showed the opposite features (Fig. 6C). CV3 distinguished the *sylvanus* group from the others (Fig. 6D). The zygomatic was positioned more posterolaterally in the *sylvanus* group as compared to the others (Fig. 6E). Overall, the *sylvanus* group consistently showed the largest Mahalanobis and Procrustes distances from the other groups; distances between the *fascicularis* and *sinica* groups were smaller than the distances between the *silenus* group and either the *fascicularis* or *sinica* groups (Table 6).

CVA also revealed that the hard-food eaters exhibited an anteroposteriorly shorter, narrower, and vertically higher (i.e., less prognathic) face than the others (Mahalanobis distance: $r = 2.2179$, $p < 0.001$; Procrustes distance: $r = 0.0339$, $p < 0.001$; Fig. 7).

DISCUSSION

We used geometric morphometrics and multivariate analyses to comprehensively assess ecogeographical and phylogenetic factors influencing size, allometric variations, and non-allometric variations in the facial cranium of extant macaques. Our null hypothesis was rejected, indicating that each of the three components is significantly influenced by ecogeographical and phylogenetic factors. Prior studies also applied geometric morphometrics to distinguish between allometric and non-allometric patterns in craniofacial shape but on a limited number of species (Pan et al., 2003; Rook and O'Higgins, 2005; Ito et al., 2011;

1
2
3
4 Singleton, 2012). With its large sample size encompassing 12 species, our analysis provides
5
6
7 greater insight into craniofacial variation and its potential use in taxonomic (species group)
8
9
10 identification of fossil macaques.

11
12 Body size was estimated by cranial measurements and is the most obvious limitation of
13
14 our analysis. Nevertheless, cranial measurements, including CS and length, have been used as
15
16
17 a surrogate for body size in numerous primate studies (e.g., Singleton, 2002; Schillaci et al.,
18
19
20 2009; Cardini et al., 2013). We estimated body size using neurocranial lnCS because species
21
22
23 average body mass was more highly correlated with neurocranial size than with facial or
24
25
26 entire cranial sizes (see also Delson et al., 2000). Although the estimated values could
27
28
29 potentially overestimate or underestimate individual body size, the values were reasonable
30
31
32 surrogates for body size considering the limited availability of individual body mass data.
33
34
35

36 Geographic variation in body size has been studied in several primate taxa. Evidence for
37
38 Bergmann's rule, namely latitude- or temperature-related size variation, has been detected in
39
40
41 some high taxonomic (Subfamily to Order) levels in primates (Harcourt and Schreier, 2009).
42
43
44 However, this was driven by Old World monkeys, including macaques, and was not
45
46
47 significant when phylogenetic effects were controlled for in Colobinae (Harcourt and Schreier,
48
49
50 2009) and Malagasy primates (Kamilar et al., 2012). Bergmann's rule has been reported for
51
52
53 the genus *Nycticebus* (Ravosa, 1998), *Aotus azarai* (Fernandez-Duque, 2011), and
54
55
56 *Microcebus murinus* (Lahann et al., 2006). These taxa are small in body size and therefore are
57
58
59
60

probably sensitive to ambient temperature in terms of regulating body temperature. However, Bergmann's rule does not seem to play a significant role in producing intrageneric or intraspecific size variation in middle- to large-sized African monkeys; rather, the size variation reflects their phylogenetic relationships or rainfall gradients (as a proxy for resource productivity) in a longitudinal direction (Dunbar, 1990; Turner et al., 1997; Cardini et al., 2007; Jolly, 2012; Cardini et al., 2013).

In contrast, size variation in macaques largely reflects latitudinal climatic gradients, which is peculiar among middle- to large-sized Cercopithecine monkeys; this distinction is attributable to the wider latitudinal distribution of the macaques. Nevertheless, the relative contribution of latitudinal climatic gradients as well as other factors, including differences in sex, species group, and habitat use type, is different among the four species groups. Size variation largely reflects latitudinal climatic gradients in the *fascicularis* and *sinica* groups, indicating that animals are larger in colder environments. This finding is in accordance with previous studies (Fooden, 1988, 2006) and is reasonably explained by Bergmann's rule. In contrast, latitudinal climatic gradients have less of an effect on body size in the *silenus* group (Albrecht, 1980), probably because of its restricted distribution in tropical zones; rather, body size variation reflects differences in habitat use type, as demonstrated by the fact that the terrestrial *M. nemestrina* is larger than the arboreal *M. leonina* (Fooden, 1975, 1982). In addition, size variation in the *sylvanus* group is probably influenced by other factors,

including sex, rather than latitudinal climatic gradients, as these animals inhabit restricted temperate regions. These findings suggest that the *silenus* and *sylvanus* groups are primarily under the control of climate-unrelated pressures, and that the *fascicularis* and *sinica* groups are under strong pressures from climatic conditions along a latitudinal gradient, which significantly amplifies body size variations.

Size variation was strongly correlated with facial elongation across macaques; as size increases, a rounded face becomes more elongated. This scaling pattern is probably conserved in the tribe Papionini because a similar pattern was observed in other clades of this tribe, including the genus *Papio* (Leigh, 2006), large-bodied papionins (Frost et al., 2003), and the entire Papionini tribe (Collard and O'Higgins, 2001; Singleton, 2002; Leigh et al., 2003; Gilbert and Grine, 2010). This common allometric pattern is largely explained by truncation or extension of a common ontogenetic trajectory (O'Higgins and Collard, 2002; Leigh et al., 2003; Leigh, 2006), suggesting that variation in facial elongation of macaques is largely a passive consequence of body size modifications along a common allometry. Thus, facial elongation indirectly reflects various factors influencing body size.

Although differences in the factors influencing body size variation change the amount of shape variation, namely facial elongation, along an allometric trajectory, they cannot be the direct cause of changing the nature (i.e., slope and elevation) of the common allometric pattern. The nature of the common allometric pattern is phylogenetically conserved because

the slope of the scaling relationship is often homogeneous in papionins, even at the intrageneric level, despite significant body size differences (Singleton, 2002). Nevertheless, some allometric (including ontogenetic) studies have detected the dissociation of common allometric trajectories, including differences in slope and/or elevation that sometimes produce curvilinearity, among papionin genera (Collard and O'Higgins, 2001; Singleton, 2002, 2004, 2005) and even within the genus *Papio* (Leigh, 2006). These findings suggest that facial elongation in some taxa is not completely explained by the simple extension or truncation of the ancestral ontogenetic trajectory. Such allometric dissociations may be a consequence of functional and adaptive modifications, possibly reflecting dietary differences or sexual selection for increasing canine height and larger gape (Singleton, 2005; Leigh, 2006).

Our analyses revealed a dissociation of the common allometric pattern within the genus *Macaca*. The slopes of the allometric lines of the *fascicularis* and *sinica* groups appeared to be lower than those of the *silenus* and *sylvanus* groups. The analysis comparing slopes and elevations among species suggested that the lower slope of the *sinica* group can be attributed to almost parallel shifts in the allometric lines between tropical and subtropical/temperate species, although *M. assamensis* does not follow this pattern. The deviation pattern of *M. sinica* was similar to another tropical species, *M. radiata*, and definitely contributed to the lower slope of the *sinica* group, but it was not tested because of the non-significant regression of *M. sinica*. The lower slope of the *fascicularis* group was probably caused not only by the

parallel shifts but also by the decreased slopes in *M. fuscata* and *M. mulatta*, which occupy the most widespread range consisting of the coldest climatic conditions. Parallel shifts in the allometric lines, i.e., higher elevations for the tropical zone than the subtropical/temperate zone, were also observed in the entire sample and between the *silenus* and *sylvanus* groups. A similar pattern, namely an allometric dissociation among several macaque species, has been reported in previous studies (Fooden, 1988; Mouri, 1996; Rook and O'Higgins, 2005; Fooden, 2006; Singleton, 2012). Our findings suggest that allometric changes have occurred several times in *Macaca*, including the elevation differences between the species groups, the elevation differences among species within each of the *fascicularis* and *sinica* species groups, the decreased slopes in *M. mulatta* and *M. fuscata*, and the possible increased slope in *M. assamensis*, *M. thibetana*, and *M. nemestrina*. Notably, many allometric changes except for the latter result in the similar situation that the face is more rounded in colder climatic zones and/or larger-bodied animals (thus reflecting colder environments under the effect of Bergmann's rule) than predicted from the overarching allometric pattern. This strongly suggests the influence of climate-related factors either directly or indirectly, which is a pressure favoring a more rounded face in colder environments or a more elongated face in warmer environments.

Specification of climate-related factors affecting the allometric changes in macaques is pending, but three possible candidates can be proposed. One is the effect of Allen's rule,

which predicts that shorter appendages reduce body heat loss in colder conditions. Treating a face as an appendage, there may be selection pressure that favors a more rounded face with a shorter muzzle in colder conditions. Although the effect of Allen's rule on facial morphology has been rejected in human experimental studies (Stegmann, 1970, 1972), the possibility remains that animals naturally possessing a relatively long muzzle, such as macaques, follow this rule (as in the arctic fox; Prestrud, 1991). Further experimental and theoretical studies are required to verify this supposition. The other two possible factors are indirectly related to climatic conditions. One is the functional adaptation that is related to dietary differences along a latitudinal gradient. A non-tropical seasonal condition reduces annual fruit productivity (Hanya et al., 2013); therefore, some species in this zone eat larger amounts of leaves instead of mature fruit as compared with those in a tropical zone (e.g., Goldstein and Richard, 1989; Ahsan, 1994; Zhou et al., 2011). If the leaves they eat require larger biting force than fruits, it is reasonable to consider that subtropical/temperate species are under selection pressure to have a more rounded face with a shorter muzzle (Singleton, 2005). In addition, intraspecific dietary variations, i.e., relatively larger amounts of leaves and/or barks in colder environments, have also been reported in *M. mulatta* (Goldstein and Richard, 1989) and *M. fuscata* (Tsuji, 2010), which can explain their decreased slopes considering that a shorter face probably permits larger bite force (Singleton, 2005). The other possible factor is sexual selection related to differences in reproductive seasonality. Reproductive seasonality is generally weak

1
2
3
4 in a tropical zone compared with that in a subtropical/temperate zone, although with some
5
6
7 exceptions (Thierry et al., 2000). Weak seasonality means that females in estrus are observed
8
9
10 throughout the year, and therefore the number of estrus females at a given time could be
11
12
13 limited when compared with species with estrus females that are seasonally constrained
14
15
16 within a certain period. Limitations on estrus females may enhance male-male competition,
17
18
19 thus causing selection pressure for a more elongated face that allows the males to possess
20
21
22 longer canines (Mouri, 1996). Tropical macaque species tend to have relatively longer male
23
24
25 canines than do subtropical/temperate species (Plavcan, 1990; Ito, personal observation). Thus,
26
27
28 each or a combination of these selection pressures may have caused the allometric
29
30
31 dissociation in macaques, as suggested in previous papionin studies (Singleton, 2005; Leigh,
32
33
34 2006). In contrast to the findings for other papionin taxa, these possible factors may be
35
36
37 significantly related to latitudinal environmental differences in macaques.

38
39 All allometric changes found in this study cannot be explained by the climate-related
40
41
42 factors. *M. assamensis*, despite living in non-tropical regions, shares allometric elevation with
43
44
45 tropical species, which means possessing a relatively elongated face. The possible increased
46
47
48 slope in *M. assamensis*, *M. thibetana*, and *M. nemestrina* also needs explanations. It may be
49
50
51 that the male-male competition was particularly and exceptionally enhanced in these species
52
53
54 when compared with their relatives in the same climatic zone, favoring a longer face (see
55
56
57 above). Whether it changes slope and/or elevation possibly depends on the strength of other
58
59
60

selection pressures and on the percentage of sex difference in the total intraspecific body size variation. Furthermore, the fact that regression was insignificant in *M. sinica* would be noteworthy. Although it may be caused just by the limited sample size, it possibly reflects the wide intraspecific shape diversity considering the small adult size range. However, these interpretations lack statistical support or evidence and are pure suppositions.

Much of the non-allometric (and sex-adjusted) shape variation reflected phylogenetic (among species group) differences, whereas some reflected dietary differences. An anteroposteriorly shorter and vertically higher face observed in the hard-food eaters such as *M. fuscata*, *M. thibetana*, and *M. sylvanus* is probably advantageous for producing a greater occlusal bite force at the anterior dentition (Singleton, 2004; Koyabu and Endo, 2009) and is desirable for dissipating larger occlusal loads (Antón, 1996). However, it is not conclusive whether this feature solely reflects a dietary adaptation or is a consequence of other climate-related factors (as mentioned above) because dietary differences are highly correlated with climatic conditions (Tsuji et al., 2013). In any case, the three species classified into different species groups may have independently acquired this feature as an adaptation to severe conditions in high-latitude regions, resulting in convergent evolution. In contrast, we showed that some localized shape components often observed in the zygomatic and muzzle regions provide well-conserved phylogenetic information that distinguishes the four species groups. This finding suggests that localized shape components are less affected by latitudinal

1
2
3
4 environmental differences, and therefore are useful for taxonomic (species group)
5
6
7 identification of fossil specimens.
8

9
10 The species group identification of fossil macaques is intriguing in terms of
11
12 understanding their paleobiogeography and adaptation to varied climates (Delson, 1977;
13
14 Jablonski and Pan, 1988), but our analyses suggest that the phylogenetic information on
15
16 overall facial configuration, degree of elongation, and prognathism is obscured by convergent
17
18 evolution and the effect of the common allometric pattern. This suggests that conflicts
19
20 regarding the taxonomic identification of fossil macaques are possibly due to the overall
21
22 impression of homoplasy in craniofacial shape, as suggested for African papionins (Collard
23
24 and O'Higgins, 2001; Leigh, 2007). In particular, the phylogenetic utility of the degree of
25
26 facial elongation is seriously impaired by allometric dissociation even within a species group,
27
28 reflecting latitudinal environmental differences. This means that even if the common
29
30 allometric effect is controlled for, the degree of facial elongation may not reflect phylogenetic
31
32 relationships among the species groups. Because the wider latitudinal distribution of
33
34 macaques compared with other non-human primates may enhance the direct and/or indirect
35
36 effects of climatic environments on craniofacial variations, caution should be exercised when
37
38 evaluating macaque fossil specimens. Further phylogenetic appraisal is required using the
39
40 localized shape components observed in zygomatics and muzzles, which are probably less
41
42 affected by environmental differences.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ACKNOWLEDGEMENTS

We thank Terutake Hayashi, Tetsu Hayashi, Linda K. Gordon, Eric Delson, Judith Chupasko, Lawrence R. Heaney, William Stanley, Tracy Damitz, Frieder Mayer, Saskia Jancke, Nora Brinkmann, Katrin Krohmann, Richard Kraft, Georges Lenglet, Paula Jenkins, Louise Tomsett, Jiang Xue-Long, Song Li, Vu The Long, Vu Ngoc Than, and Lim Kok Peng Kelvin for their help with data collection. We would like to extend our gratitude to Christopher Ruff, an anonymous associate editor, and two anonymous referees for their constructive comments that greatly improved this paper; to Toshio Mouri, Yuzuru Hamada, and Naoko Egi for their comments on an earlier version of this manuscript; and to Chris Klingenberg and Wataru Yano for their supportive comments on geometric morphometric analysis. This study was financially supported in part by the ITP-HOPE and AS-HOPE Programs from JSPS (to T. I.), Grant-in-Aid for JSPS Fellows (to T. I.), Grant-in-Aid for Challenging Exploratory Research (to T. N.), and the Global COE Program from MEXT (Grant A06 to Kyoto University).

LITERATURE CITED

- 1
- 2
- 3
- 4
- 5
- 6
- 7 Ahsan MF. 1994. Feeding ecology of the primates of Bangladesh. In: Thierry B, Anderson JR,
- 8
- 9 Roeder JJ, and Herrenschmidt N, editors. Current primatology Volume 1 Ecology and
- 10
- 11 evolution. Strasbourg: Universite Louis Pasteur. p 79-86.
- 12
- 13
- 14
- 15 Albrecht GH. 1978. The craniofacial morphology of the Sulawesi macaques: Multivariate
- 16
- 17 approaches to biological problems. In: Szalay FS, editor. Contrib Primatol. Basel: S.
- 18
- 19 Karger. p 1-151.
- 20
- 21
- 22
- 23
- 24 Albrecht GH. 1980. Latitudinal, taxonomic, sexual, and insular determinants of size variation
- 25
- 26 in pigtail macaques, *Macaca nemestrina*. Int J Primatol 1(2):141-152.
- 27
- 28
- 29
- 30 Antón SC. 1996. Cranial adaptation to a high attrition diet in Japanese macaques. Int J
- 31
- 32 Primatol 17(3):401-427.
- 33
- 34
- 35
- 36 Bookstein FL. 1989. Principal warps: Thin-plate splines and the decomposition of
- 37
- 38 deformations. ITPAM 11(6):567-585.
- 39
- 40
- 41
- 42 Cardini A, Dunn J, O'Higgins P, and Elton S. 2013. Clines in Africa: does size vary in the
- 43
- 44 same way among widespread sub-Saharan monkeys? J Biogeogr 40(2):370-381.
- 45
- 46
- 47 Cardini A, and Elton S. 2008. Does the skull carry a phylogenetic signal? Evolution and
- 48
- 49 modularity in the guenons. Biol J Linn Soc Lond 93:813-834.
- 50
- 51
- 52
- 53 Cardini A, Jansson A-U, and Elton S. 2007. A geometric morphometric approach to the study
- 54
- 55 of ecogeographical and clinical variation in vervet monkeys. J Biogeogr
- 56
- 57
- 58
- 59
- 60

- 1
- 2
- 3
- 4 34:1663-1678.
- 5
- 6
- 7 Collard M, and O'Higgins P. 2001. Ontogeny and homoplasy in the papionin monkey face.
- 8
- 9
- 10 Evol Dev 3:322-331.
- 11
- 12
- 13 Delson E. 1977. Vertebrate paleontology, especially of nonhuman primates, in China. In:
- 14
- 15 Howells WW, and Tsuchitani PJ, editors. Paleoanthropology in the People's Republic
- 16
- 17 of China. Washington, D.C.: National Academy of Science. p 40-65.
- 18
- 19
- 20
- 21 Delson E. 1980. Fossil macaques, phyletic relationships and a scenario of deployment. In:
- 22
- 23 Lindburg DG, editor. The macaques: Studies in ecology, behavior and evolution. New
- 24
- 25 York: Nostrand-Reinhold. p 10-30.
- 26
- 27
- 28
- 29
- 30 Delson E, Terranova CJ, Jungers WL, Sargis EJ, Jablonski NG, and Dechow PC. 2000. Body
- 31
- 32 mass in Cercopithecidae (Primates, Mammalia): estimation and scaling in extinct and
- 33
- 34 extant taxa. Anthropol Pap Am Mus Nat Hist(83):1-159.
- 35
- 36
- 37
- 38 Drake AG, and Klingenberg CP. 2008. The pace of morphological change: historical
- 39
- 40 transformation of skull shape in St Bernard dogs. Proc R Soc B-Biol Sci
- 41
- 42 275(1630):71-76.
- 43
- 44
- 45
- 46
- 47 Dunbar RIM. 1990. Environmental determinants of intraspecific variation in body-weight in
- 48
- 49 baboons (*Papio* spp). J Zool 220:157-169.
- 50
- 51
- 52
- 53 Fernandez-Duque E. 2011. Rensch's rule, Bergmann's effect and adult sexual dimorphism in
- 54
- 55 wild monogamous Owl monkeys (*Aotus azarai*) of Argentina. Am J Phys Anthropol
- 56
- 57
- 58
- 59
- 60

1
2
3
4 146(1):38-48.
5

6
7 Fleagle JG. 1999. Primate adaptation and evolution, 2nd edition. New York: Academic Press.
8

9
10 Fooden J. 1969. Taxonomy and evolution of the monkeys of Celebes. *Bibliotheca Primatol*
11
12 10:1-148.
13

14
15 Fooden J. 1975. Taxonomy and evolution of liontail and pigtail macaques. *Fieldiana Zool*
16
17 67:1-169.
18

19
20
21 Fooden J. 1976. Provisional classification and key to living species of macaques (Primates:
22
23 *Macaca*). *Folia Primatol* (Basel) 25:225-236.
24

25
26
27 Fooden J. 1982. Ecogeographic segregation of macaque species. *Primates* 23(4):574-579.
28

29
30 Fooden J. 1985. Taxonomy and evolution of the *sinica* group of macaques: 5. Overview of
31
32 natural history. *Fieldiana Zool* 29:1-22.
33

34
35
36 Fooden J. 1988. Taxonomy and Evolution of the *sinica* group of macaques: 6. Interspecific
37
38 comparisons and synthesis. *Fieldiana Zool* 45:1-44.
39

40
41
42 Fooden J. 1990. The bear macaques, *Macaca arctoides*: a systematic review. *J Hum Evol*
43
44 19:607-686.
45

46
47 Fooden J. 2006. Comparative review of *fascicularis*-group species of macaques (Primates:
48
49 *Macaca*). *Fieldiana Zool* 107(16):1-44.
50

51
52
53 Fooden J. 2007. Systematic review of the Barbary macaque, *Macaca sylvanus* (Linnaeus,
54
55 1758). *Fieldiana Zool* 113(63):1-58.
56

- 1
- 2
- 3
- 4 Fooden J, and Albrecht GH. 1993. Latitudinal and insular variation of skull size in crab-eating
- 5
- 6 macaques (primates, Cercopithecidae: *Macaca fascicularis*). Am J Phys Anthropol
- 7
- 8 92:521-538.
- 9
- 10
- 11
- 12 Fooden J, and Wu HY. 2001. Systematic review of the Taiwanese macaque, *Macaca cyclopis*
- 13
- 14 Swinhoe, 1863. Fieldiana Zool 98(30):1-70.
- 15
- 16
- 17
- 18 Frost SR, Marcus LF, Bookstein FL, Reddy DP, and Delson E. 2003. Cranial allometry,
- 19
- 20 phylogeography, and systematics of large-bodied papionins (primates:
- 21
- 22 Cercopithecinae) inferred from geometric morphometric analysis of landmark data.
- 23
- 24 Anat Rec A Discov Mol Cell Evol Biol 275A:1048-1072.
- 25
- 26
- 27
- 28
- 29
- 30 Gilbert CC, and Grine FE. 2010. Morphometric variation in the papionin muzzle and the
- 31
- 32 biochronology of the South African Plio-Pleistocene karst cave deposits. Am J Phys
- 33
- 34 Anthropol 141:418-429.
- 35
- 36
- 37
- 38
- 39 Goldstein SJ, and Richard AF. 1989. Ecology of rhesus macaques (*Macaca mulatta*) in
- 40
- 41 northwest Pakistan. Int J Primatol 10(6):531-567.
- 42
- 43
- 44
- 45 Gower JC. 1971. A general coefficient of similarity and some of its properties.
- 46
- 47 Biometrics:857-871.
- 48
- 49
- 50 Hamada Y, Watanabe T, and Iwamoto M. 1996. Morphological variations among local
- 51
- 52 populations of Japanese macaque (*Macaca fuscata*). In: Shotake T, and Wada K,
- 53
- 54 editors. Variations in the Asian macaques. Tokyo: Tokyo University Press. p 97-115.
- 55
- 56
- 57
- 58
- 59
- 60

- 1
- 2
- 3
- 4 Hanya G, Tsuji Y, and Grueter CC. 2013. Fruiting and flushing phenology in Asian tropical
- 5
- 6 and temperate forests: implications for primate ecology. *Primates*:1-10.
- 7
- 8
- 9
- 10 Harcourt AH, and Schreier BM. 2009. Diversity, body mass, and latitudinal gradients in
- 11
- 12 primates. *Int J Primatol* 30(2):283-300.
- 13
- 14
- 15 Hijmans RJ, Cameron SE, Parra JL, Jones PG, and Jarvis A. 2005. Very high resolution
- 16
- 17 interpolated climate surfaces for global land areas. *IJCLI* 25(15):1965-1978.
- 18
- 19
- 20
- 21 Ito T, Nishimura T, and Takai M. 2011. Allometry and interspecific differences in the facial
- 22
- 23 cranium of two closely related macaque species. *Anat Res Int* 849751.
- 24
- 25
- 26
- 27 Iwamoto M. 1975. On a skull of a fossil macaque from the Shikimizu limestone quarry in the
- 28
- 29 Shikoku district, Japan. *Primates* 16(1):83-94.
- 30
- 31
- 32
- 33 Jablonski N. 2002. Fossil old world monkeys: the late Neogene radiation. In: Hartwig WC,
- 34
- 35 editor. *The primate fossil record*. Cambridge: Cambridge University Press. p 255-299.
- 36
- 37
- 38
- 39 Jablonski NG, and Pan Y. 1988. The evolution and palaeobiogeography of monkeys in China.
- 40
- 41 In: Whyte P, Aigner J, Jablonski NG, Taylor G, Walker D, and Wang PX, editors. *The*
- 42
- 43 *palaeoenvironment of East Asia from the Mid-Tertiary*. Hong Kong: Centre of Asian
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51 Jolly CJ. 2012. Rainfall is not a genus-wide predictor of mean body mass in baboon
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- p>the resource seasonality hypothesis in Malagasy primates using GIS-based climate
-
- data. Am J Phys Anthropol 147(3):401-408.
-
- Klingenberg CR. 2011. MorphoJ: an integrated software package for geometric
-
- morphometrics. Mol Ecol Res 11(2):353-357.
-
- Koyabu DB, and Endo H. 2009. Craniofacial variation and dietary adaptations of African
-
- colobines. J Hum Evol 56:525-536.
-
- Lahann P, Schmid J, and Ganzhorn JU. 2006. Geographic variation in populations of
-
- Microcebus murinus*
- in Madagascar: Resource seasonality or Bergmann's rule? Int J
-
- Primatol 27(4):983-999.
-
- Leigh SR. 2006. Cranial ontogeny of
- Papio*
- baboons (
- Papio hamadryas*
-). Am J Phys
-
- Anthropol 130:71-84.
-
- Leigh SR. 2007. Homoplasy and the evolution of ontogeny in papionin primates. J Hum Evol
-
- 52(5):536-558.
-
- Leigh SR, Shah NF, and Buchanan LS. 2003. Ontogeny and phylogeny in papionin primates.
-
- J Hum Evol 45:285-316.
-
- Li J, Han K, Xing J, Kim HS, Rogers J, Ryder OA, Disotell T, Yue B, and Batzer MA. 2009.
-
- Phylogeny of the macaques (Cercopithecidae:
- Macaca*
-) based on Alu elements. Gene
-
- 448:242-249.
-
- Mantel N. 1967. The detection of disease clustering and a generalized regression approach.

- Cancer Res 27(2 Part 1):209-220.
- Mitteroecker P, and Gunz P. 2009. Advances in geometric morphometrics. *Evol Biol* 36(2):235-247.
- Monteiro LR. 1999. Multivariate regression models and geometric morphometrics: the search for causal factors in the analysis of shape. *Syst Biol* 48(1):192-199.
- Morales JC, and Melnick DJ. 1998. Phylogenetic relationships of the macaques (*Cercopithecidae: Macaca*), as revealed by high resolution restriction site mapping of mitochondrial ribosomal genes. *J Hum Evol* 34:1-23.
- Mouri T. 1996. Multivariate cranial ontogenetic allometries in Crab-eating Rhesus and Japanese macaques. *Anthropol Sci* 104(4):281-303.
- O'Higgins P, and Collard M. 2002. Sexual dimorphism and facial growth in papionin monkeys. *J Zool* 257(2):255-272.
- O'Higgins P, and Jones N. 1998. Facial growth in *Cercocebus torquatus*: an application of three-dimensional geometric morphometric techniques to the study of morphological variation. *J Anat* 193:251-272.
- O'Higgins P, and Jones N. 2006. Tools for statistical shape analysis. Hull York Medical School; <http://sites.google.com/site/hymsfme/resources>.
- Pan R, Jablonski NG, Oxnard C, and Freedman L. 1998. Morphometric analysis of *Macaca arctoides* and *M. thibetana* in relation to other macaque species. *Primates*

- 1
- 2
- 3
- 4 39(4):519-537.
- 5
- 6
- 7 Pan R, and Oxnard CE. 2000. Craniodental variation of macaques (*Macaca*): size, function
- 8
- 9 and phylogeny. Zool Res 21(4):308-322.
- 10
- 11
- 12 Pan R, and Oxnard CE. 2001. Radiation and evolution of three macaque species, *Macaca*
- 13
- 14 *fascicularis*, *M. radiata* and *M. sinica*, as related to geographic changes in the
- 15
- 16 Pleistocene of Southeast Asia. In: Metcalfe I, Smith JMB, Morwood M, and Hewson
- 17
- 18 K, editors. Faunal and floral migrations and evolution in SE Asia-Australasia.
- 19
- 20 Netherlands: Swets and Zeitlinger. p 337-355.
- 21
- 22
- 23
- 24
- 25
- 26
- 27 Pan R, and Oxnard CE. 2002. Craniodental variation among Macaques (*Macaca*), nonhuman
- 28
- 29 primates. BMC Evol Biol 2:1-12.
- 30
- 31
- 32
- 33 Pan R, Wei F, and Li M. 2003. Craniofacial variation of the Chinese macaques explored with
- 34
- 35 morphologica. J Morphol 256:342-348.
- 36
- 37
- 38
- 39 Plavcan JM. 1990. Sexual dimorphism in the dentition of extant anthropoid primates. PhD
- 40
- 41 Thesis. North Carolina: Duke University.
- 42
- 43
- 44
- 45 Prestrud P. 1991. Adaptations by the arctic fox (*Alopex lagopus*) to the polar winter.
- 46
- 47 Arctic:132-138.
- 48
- 49
- 50
- 51 Ravosa MJ. 1998. Cranial allometry and geographic variation in slow lorises (*Nycticebus*).
- 52
- 53 Am J Primatol 45:225-243.
- 54
- 55
- 56
- 57 Rohlf FJ. 2005. NTSYSpc, Numerical taxonomy system, version 2.21a. Exeter Software.
- 58
- 59
- 60

- New York.
- Rohlf FJ, and Slice D. 1990. Extensions of the procrustes method for the optimal superimposition of landmarks. Syst Zool 39(1):40-59.
- Rook L, and O'Higgins P. 2005. A comparative study of adult facial morphology and its ontogeny in the fossil macaque *Macaca majori* from Capo Figari, Sardinia, Italy. Folia Primatol (Basel) 76:151-171.
- Schillaci M, Meijaard E, and Clark T. 2009. The effect of island area on body size in a primate species from the Sunda shelf island. J Biogeogr 36:362-371.
- Schlosser M. 1924. Fossil primates from China. Palaeontologia Sinica, Series C 1(2):1-16.
- Singleton M. 2002. Patterns of cranial shape variation in the Papionini (Primates: Cercopithecinae). J Hum Evol 42:547-578.
- Singleton M. 2004. Geometric morphometric analysis of functional divergence in mangabey facial form. J Anthropol Sci 82:29-46.
- Singleton M. 2005. Functional shape variation in the Cercopithecine masticatory complex. In: Slice DE, editor. Modern morphometrics in Physical Anthropology. New York: Kluwer Academic/Plenum Publishers. p 319-348.
- Singleton M. 2012. Postnatal cranial development in papionin primates: An alternative model for hominin evolutionary development. Evol Biol 39(4):499-520.
- Slice DE. 1999. Morpheus et al. (beta version). Department of ecology and evolution, State

- 1
- 2
- 3
- 4 University. New York, Stony Brook.
- 5
- 6
- 7 Smouse PE, Long JC, and Sokal RR. 1986. Multiple regression and correlation extensions of
- 8
- 9
- 10 the Mantel test of matrix correspondence. *Syst Zool* 35(4):627-632.
- 11
- 12 Steegmann AT. 1970. Cold adaptation and the human face. *Am J Phys Anthropol* 32:243-250.
- 13
- 14 Steegmann AT. 1972. Cold response, body form, and craniofacial shape in two racial groups
- 15
- 16
- 17
- 18 of Hawaii. *Am J Phys Anthropol* 37(2):193-221.
- 19
- 20
- 21 Thierry B, Iwaniuk AN, and Pellis SM. 2000. The influence of phylogeny on the social
- 22
- 23
- 24 behavior of macaques (Primates: Cercopithecidae, genus *Macaca*). *Ethology*
- 25
- 26
- 27 106:713-728.
- 28
- 29
- 30 Tosi AJ, Morales JC, and Melnick DJ. 2000. Comparison of Y chromosome and mtDNA
- 31
- 32
- 33 phylogenies leads to unique inferences of macaque evolutionary history. *Mol*
- 34
- 35
- 36 *Phylogen Evol* 17:133-144.
- 37
- 38
- 39 Tosi AJ, Morales JC, and Melnick DJ. 2003. Paternal, maternal, and biparental molecular
- 40
- 41
- 42 markers provide unique windows onto the evolutionary history of macaque monkeys.
- 43
- 44
- 45 *Evolution* 57:1419-1435.
- 46
- 47
- 48 Tsuji Y. 2010. Regional, temporal, and interindividual variation in the feeding ecology of
- 49
- 50
- 51 Japanese macaques. *The Japanese macaques*: Springer. p 99-127.
- 52
- 53
- 54 Tsuji Y, Hanya G, and Grueter CC. 2013. Feeding strategies of primates in temperate and
- 55
- 56
- 57 alpine forests: comparison of Asian macaques and colobines. *Primates* 54(3):201-215.
- 58
- 59
- 60

Turner TR, Anapol F, and Jolly CJ. 1997. Growth, development, and sexual dimorphism in
vervet monkeys (*Cercopithecus aethiops*) at four sites in Kenya. Am J Phys Anthropol
103(1):19-35.

Zelditch ML, Swiderski DL, and Sheets HD. 2012. Geometric morphometrics for biologists: a
primer: Academic Press.

Zhou QH, Wei H, Huang ZH, and Huang CM. 2011. Diet of the Assamese macaque *Macaca*
assamensis in limestone habitats of Nonggang, China. Curr Zool 57(1):18-25.

FIGURE LEGENDS

Fig. 1 Cladogram of macaques: phylogenetic relationship among the four species groups (Table 1). A: Modified after Delson (1980), which is based on genitalia, cranial morphology, biogeography, and fossil records (see also Fooden, 1976). B: Modified after Tosi et al. (2003), which was based on mitochondrial DNA (see also Morales and Melnick, 1998). This resolved the phylogenetic relationship among the four species groups but the phylogenetic position of *M. arctoides* was different from the others. C: Modified after Tosi et al. (2003), which was based on Y-chromosome DNA. This supports the monophyly of each of the four species groups but cannot resolve the intergroup relationships. D: Modified after Li et al. (2009), which was based on *Alu* elements. This provides strong statistical support for the monophyly of each of the four species groups and resolved the intergroup phylogenetic relationships.

Fig. 2 Geographic distribution of local populations of macaques used in this study (Table 2). Dashed lines indicate border between populations belonging to same species. A: the *fascicularis* group. Circle, *M. fascicularis*; triangle, *M. mulatta*; cross, *M. cyclopis*; square, *M. fuscata*. B: the *sinica* group. Circle, *M. arctoides*; triangle, *M. assamensis*; square, *M. thibetana*; cross, *M. radiata*; star, *M. sinica*. C: the *silenus* group. Circle, *M. nemestrina*; triangle, *M. leonina*. D: the *sylvanus* group. Circle, *M. sylvanus*.

Fig. 3 Landmarks on facial regions used in this study. A: Lateral view. B: Frontal view.
C: Inferior view.

Fig. 4 A: Relationship between the natural logarithms of species average (split sex) neurocranium centroid size (CS) and species average (split sex) body mass. B: The estimation of individual's body size. Both sexes are indicated by the same symbol; for all species, males are larger than females.

Fig. 5 Allometric pattern of craniofacial shape across macaques. A: Shape variations along the axis of the shape score. Shape changes are shown as the smooth surface rendering of polygons, Cartesian transformation grids, and wireframes. The deformation of Cartesian transformation grids indicates transformation from the grand mean to positive and negative extremes. The wireframe of the solid line and right images represent the positive extreme (+1.0), and that of the dashed line and left images represent the negative extreme (-1.0). Arrows indicate the relative size of the upper face against the lower face and the relative elongation of the muzzle and the facial profile line. B: Relationship between body size surrogates and shape scores. Symbols represent climatic zones. Solid line represents a regression line for subtropical/temperate animals ($y = 0.074x - 0.680$, $R^2 = 0.57$, $p < 0.001$); dashed line represents a regression line for tropical animals ($y = 0.078x - 0.671$, $R^2 = 0.54$, p

< 0.001). C: Same as B, but symbols represent species groups. Solid line represents a regression line for the *fascicularis* group ($y = 0.043x - 0.340$, $R^2 = 0.32$, $p < 0.001$); dashed line represents a regression line for the *sinica* group ($y = 0.061x - 0.545$, $R^2 = 0.54$, $p < 0.001$); dotted line represents a regression line for the *silenus* group ($y = 0.087x - 0.752$, $R^2 = 0.51$, $p < 0.001$); dot-dashed line represents a regression line for the *sylvanus* group ($y = 0.077x - 0.725$, $R^2 = 0.34$, $p = 0.058$). D-F: Same as B, but for the *fascicularis*, *sinica*, and *silenus* groups, respectively (Table 5). Symbols represent species.

Fig. 6 Shape changes associated with species group differences. A: Scatterplot of CV1 against CV2. B: Shape changes along the axis of CV1. Arrows indicate the relative height and width of the upper muzzle. C: Shape changes along the axis of CV2. Arrows indicate the convexity (and concavity) of the nasal profile line and the relative width of the zygomatic. D: Scatterplot of CV3 against CV2. E: Shape changes along the axis of CV3. Arrows indicate the relative position of the zygomatic. The shapes of positive (+4.0, but +8.0 for CV3) and negative (-4.0) extremes are represented for each axis.

Fig. 7 Shape changes associated with dietary difference. A: Histogram of CV1 scores. B: Shape changes along the axis of CV1. Arrows indicate relative height, width, and length of the face.

Table 1. Ecogeographical and phylogenetic categories.

Species	Climatic zones	Dietary habits	Habitat use types	Species groups
<i>M. fascicularis</i>	tropical zone	non-hard food eaters	arboreal	the <i>fascicularis</i> group
<i>M. mulatta</i>	subtropical/temperate zone	non-hard food eaters	intermediate/uncertain	the <i>fascicularis</i> group
<i>M. cyclopis</i>	subtropical/temperate zone	non-hard food eaters	intermediate/uncertain	the <i>fascicularis</i> group
<i>M. fuscata</i>	subtropical/temperate zone	hard food eaters	intermediate/uncertain	the <i>fascicularis</i> group
<i>M. sinica</i>	tropical zone	non-hard food eaters	intermediate/uncertain	the <i>sinica</i> group
<i>M. radiata</i>	tropical zone	non-hard food eaters	intermediate/uncertain	the <i>sinica</i> group
<i>M. arctoides</i>	subtropical/temperate zone	non-hard food eaters	terrestrial	the <i>sinica</i> group
<i>M. assamensis</i>	subtropical/temperate zone	non-hard food eaters	arboreal	the <i>sinica</i> group
<i>M. thibetana</i>	subtropical/temperate zone	hard food eaters	intermediate/uncertain	the <i>sinica</i> group
<i>M. nemestrina</i>	tropical zone	non-hard food eaters	terrestrial	the <i>silenus</i> group
<i>M. leonina</i>	tropical zone	non-hard food eaters	arboreal	the <i>silenus</i> group
<i>M. sylvanus</i>	subtropical/temperate zone	hard food eaters	terrestrial	the <i>sylvanus</i> group

Details are described in the text.

Table 2. Samples used in this study.

Species	Population (Sampling unit)	Region	Sample size*					The number of geographic points
			F		M		Total	
<i>M. fascicularis</i>	Sunda Islands	Sunda islands ¹	20				20	20
	<i>fascicularis</i>	Indochinese	20				40	25
	<i>fascicularis</i>	Indochina						
	Philippine	Philippine	20				40	14
	<i>fascicularis</i>	islands ²						
<i>M. mulatta</i>	Indochinese	Northern	20		19	(1)	39	15
	<i>mulatta</i>	Indochina and northeastern India (Manipur)						
	South Asian	Northern India and Pakistan	12	(1)	19	(1)	31	14
	<i>mulatta</i>							
	Chinese	Southern and	16	(1)	15	(1)	31	5
	<i>mulatta</i>	eastern China						
<i>M. cyclopis</i>	<i>cyclopis</i>	Taiwan	11		12		23	1
<i>M. fuscata</i>	<i>fuscata</i>	Yaku island	20		20		40	1
	<i>yakui</i>							
	<i>fuscata</i>	Japan	20		20		40	13
	<i>fuscata</i>	excluding Yaku island						
<i>M. sinica</i>	<i>sinica</i>	Ceylon island	6		12	(1)	18	6
<i>M. radiata</i>	<i>radiata</i>	Southern India	7	(1)	10		17	4
<i>M. arctoides</i>	<i>arctoides</i>	Northeastern	12	(3)	20	(1)	32	13
		India, southern China, and Indochina						
<i>M. assamensis</i>	<i>assamensis</i>	Nepal and	5		10	(1)	15	3
	<i>pelops</i>	northeastern India						

		(Arunachal Pradesh)						
	<i>assamensis</i>	Southern	11		20	(1)	31	14
	<i>assamensis</i>	China, northern Indochina, and northeastern India (Nagaland)						
<i>M. thibetana</i>	<i>thibetana</i>	Southern and eastern China	4		12		16	3
<i>M. nemestrina</i>	Borneo	Borneo	7	(1)	12	(1)	19	5
	<i>nemestrina</i>	Sumatra and southern Malay peninsula	12	(2)	20		32	13
<i>M. leonina</i>	<i>leonina</i>	Northeastern India, Southern China, and northern Indochina	14	(3)	20	(1)	34	12
<i>M. sylvanus</i>	<i>sylvanus</i>	Morocco and Gibraltar	6	(2)	5		11	2

The specimens are housed at the Hakusan Nature Conservation Center, Hakusan, Japan; the Tochigi Prefectural Museum, Utsunomiya, Japan; the Primate Research Institute of Kyoto University, Inuyama, Japan; the Kunming Institute of Zoology, Kunming, China; the Zoological Museum of National University of Hanoi, Hanoi, Vietnam; the Raffles Museum of Biodiversity Research, Singapore; the Museum für Naturkunde of Humboldt University, Berlin, Germany; the Naturmuseum Senckenberg, Frankfurt, Germany; the Zoologische Staatssammlung München, Munich, Germany; the Natural History Museum, London, UK; the National Museum of Natural History, Washington DC, USA; the American Museum of Natural History, New York, NY, USA; the Museum of Comparative Zoology of Harvard University, Cambridge, MA, USA; and the Field Museum of Natural History, Chicago, IL, USA. *Number in parentheses indicates the number of specimens lacking geographic data. ¹Bali, Borneo, Bunguran, Flores, Java, Lombok, Nias, Simeulue, Sumatra, Sumbawa, and Timor islands. ² Distribution region of *M. f. philippinensis* and contact region between this subspecies and *M. f. fascicularis* (Fooden 1990b).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Landmarks used in this study.

Region	Abbreviation	Definition
Face	PRS	Prosthion: antero-inferior point on projection of premaxilla between central incisors. ¹
	IFR	Posterior-most point of incisive foramen. ¹
	MXP	Meeting point of maxilla and palatine along midline. ¹
	PNS	Tip of posterior nasal spine. ¹
	GPF	Most posterior point on the margin of greater palatine foramen. ¹
	NSP	Nasospinale: inferior-most midline point of piriform aperture. ¹
	WPA	Point corresponding to largest width of piriform aperture. ¹
	NPM	Meeting point of nasal and pre-maxilla on margin of piriform aperture. ¹
	RHI	Rhinion: most anterior midline point on nasals. ¹
	NAS	Nasion: midline point on fronto-nasal suture. ¹
	DCR	Dacryon: most superior point of the lacrimomaxillary suture (intersection with frontal bone). ²
	IST	Point on inferior margin of supraorbital torus (superior margin of orbit) at middle of orbit. ³
	FRO	Frontomale orbitale: where frontozygomatic suture crosses inner orbital rim. ¹
	ZMS	Zygo-max superior: antero-superior point of zygomaticomaxillary suture taken at orbit rim. ¹
	FRT	Frontomale temporale: where frontozygomatic suture crosses lateral edge of zygoma. ¹
	ZMI	Zygo-max inferior: antero-inferior point of zygomaticomaxillary suture. ¹
	CZA	Maximum curvature of anterior upper margin of zygomatic arch. ¹
	PMS	The point where premaxillary suture crosses alveolar margin. ³
	MP3	Mesial P3: most mesial point on P ³ alveolus, projected labially onto alveolar margin. ¹
	MM1	Mesial M1: contact points between P ⁴ and M ¹ , projected labially onto alveolar margin. ¹
	MM3	Mesial M3: contact points between M ² and M ³ , projected labially onto alveolar margin. ¹
	DM3	Distal M3: posterior midpoint onto alveolar margin of M ³ . ¹
Face/Neurocrani	GLA	Glabella: most forward projecting midline point of frontals at the level of the supraorbital ridges. ¹

um

Neurocranium	BRG	Bregma: junction of coronal and sagittal sutures, on sagittal crest if necessary. ³
	INI	Inion: most posterior point of cranium, when viewed in the Frankfurt horizontal, be it on sagittal/nuchal crest or not. ³
	OPS	Opisthion: posterior most point of foramen magnum. ³
	BAS	Basion: anterior most point of foramen magnum. ³
	HOR	Hormion: midpoint of the posterosuperior border of the vomer. ²
	POR	Porion: top of auditory meatus. ³
	ZAP	Meeting point of zygomatic arch and alisphenoid on superior margin of pterygomaxillary fissure. ¹
Other	ZTS	Zygo-temp superior: superior point of zygomatico-temporal suture on lateral face of zygomatic arch. ³
	ZTI	Zygo-temp inferior: inferior point of zygomatico-temporal suture on lateral face of zygomatic arch. ³
	ZPT	Posterior-most point on curvature of anterior margin of zygomatic process of temporal bone. ¹

¹Cardini et al. (2007), ²Cobb and O'Higgins (2007), ³Frost et al. (2003).

Table 4. Multivariate regression of body size surrogate against climate, sex, habitat use type, and species group.

Sample	R ²	p	independent variables	partial regression coefficient	Std. Error	t value	p	standard partial regression coefficient*
total sample	0.78	<0.001	intercept	9.99	0.056	179.6	<0.001	0.01
			temperature ¹	-0.01	0.000	-30.4	<0.001	-0.70
			male ²	0.37	0.021	17.9	<0.001	0.38
			habitat use type	0.18	0.016	11.1	<0.001	0.26
			<i>sinica</i> group ³	0.22	0.025	8.7	<0.001	0.19
			<i>silenus</i> group ³	0.31	0.033	9.2	<0.001	0.23
			<i>sylvanus</i> group ³	-0.21	0.082	-2.5	0.012	-0.06
<i>fascicularis</i> group	0.81	<0.001	intercept	10.06	0.091	110.1	<0.001	0.00
			temperature ¹	-0.01	0.000	-21.0	<0.001	-0.73
			male ²	0.34	0.024	14.2	<0.001	0.35
			habitat use type	0.14	0.035	3.9	<0.001	0.13
<i>sinica</i> group	0.69	<0.001	intercept	10.47	0.141	74.5	<0.001	0.02
			temperature ¹	-0.01	0.001	-13.2	<0.001	-0.73
			male ²	0.44	0.054	8.2	<0.001	0.43
			habitat use type	0.14	0.035	3.9	<0.001	0.22

<i>silenus</i> group	0.81	<0. 001	intercept	9.76	0.209	46.8	<0. 001	-0.02
			temperature ¹	-0.01	0.001	-6.5	<0. 001	-0.39
			male ²	0.40	0.041	9.6	<0. 001	0.54
			habitat use type	0.25	0.021	11.9	<0. 001	0.68
<i>sylvanu</i> s group	0.40	<0. 001	intercept	9.20	0.096	95.8	<0. 001	0.00
			male ²	0.35	0.142	2.5	0.0 35	0.64

*The largest contribution is indicated by bold. ¹The annual mean temperature. ²Sex.
³Species group. ^{2,3}These are nominal and thus coded as dummy variables. One of the
dummy variables ("female" for sex and "*fascicularis* group" for species group) was
eliminated from the regression equation because the information of one variable can be
readily obtained from that of the other variables (Suits, 1957). The eliminated variable can
be regarded as a reference.

Table 5. Scaling pattern for each species.

Equations for the least-squares regression of the shape scores on the body size surrogates.					Slope divergences from a general trend.			
Species	Slope	Elevation	R ²	p	Angles between the slopes for each species and a general trend	df	F	p
<i>M. fascicularis</i>	0.081	-0.688	0.41	<0.001	0.003	1	0.058	0.810
<i>M. mulatta</i>	0.052	-0.491	0.42	<0.001	-0.022	1	8.008	0.005
<i>M. cyclopis</i>	0.083	-0.778	0.54	<0.001	0.009	1	0.162	0.688
<i>M. fuscata</i>	0.043	-0.377	0.18	<0.001	-0.032	1	7.262	0.007
<i>M. sinica</i>	0.009	-0.102	0.01	0.629	-	-	-	-
<i>M. radiata</i>	0.080	-0.703	0.36	0.011	0.002	1	0.004	0.952
<i>M. arctoides</i>	0.072	-0.667	0.54	<0.001	-0.002	1	0.014	0.905
<i>M. assamensis</i>	0.098	-0.870	0.63	<0.001	0.023	1	3.416	0.065
<i>M. thibetana</i>	0.103	-0.955	0.48	0.003	0.029	1	0.562	0.454
<i>M. nemestrina</i>	0.099	-0.860	0.53	<0.001	0.021	1	1.931	0.166
<i>M. leonina</i>	0.078	-0.673	0.35	<0.001	0.000	1	0	0.990
<i>M. sylvanus</i>	0.077	-0.725	0.34	0.058	0.002	1	0.006	0.940

Table 6. Mahalanobis squared distances (below the diagonal) and Procrustes distances (above the diagonal) of the size- and sex-adjusted Procrustes shape coordinates between the species groups.

	<i>fascicularis</i> group	<i>sinica</i> group	<i>silenus</i> group	<i>sylvanus</i> group
<i>fascicularis</i> group		0.036	0.062	0.078
<i>sinica</i> group	3.226		0.057	0.091
<i>silenus</i> group	3.862	3.512		0.122
<i>sylvanus</i> group	6.684	7.620	8.148	

All the differences were statistically significant ($p < 0.001$).

Figure 1
196x158mm (300 x 300 DPI)

Figure 2
286x196mm (300 x 300 DPI)

Figure 3
202x200mm (300 x 300 DPI)

Figure 4
209x106mm (300 x 300 DPI)

Figure 5A-C
293x717mm (300 x 300 DPI)

Figure 5D-F
293x792mm (300 x 300 DPI)

Figure 6
188x120mm (300 x 300 DPI)

Figure 7
129x208mm (300 x 300 DPI)

Supporting information for Table 1: Ecogeographical and phylogenetic categories

Fooden (1982a) suggested that Asian macaques are ecogeographically segregated into Southern tropical and Northern subtropical species, with a boundary occurring at approximately 18°N in both India and the Indochinese Peninsula. Based on this criterion, we defined five tropical species and seven subtropical/temperate species. Although the geographic range of *M. arctoides* extends southward beyond this boundary, we classified this species as subtropical/temperate (see Eudey, 1980; Fa, 1989; Fooden, 1990a).

The species distributed in the northern extremes of the macaques' distribution or in high mountain regions, namely *M. fuscata*, *M. thibetana*, and *M. sylvanus*, feed on considerable amounts of tough and/or fibrous foods (e.g., mature leaves, bark, and nuts) in certain seasons when the availability of more easily obtainable and nutritious food items (e.g., fruit) is limited (Izawa and Nishida, 1963; Suzuki, 1965; Zhao et al., 1991; Hanya, 2004; Hanya et al., 2011). These species were categorized as hard food eaters and were distinguished from the others.

For habitat use types, *M. arctoides*, *M. nemestrina*, and *M. sylvanus* were categorized as “terrestrial”; *M. fascicularis*, *M. assamensis*, and *M. leonina* were categorized as “arboreal”; and the others were categorized as intermediate or uncertain (Sugiyama, 1971; Fooden, 1975, 1979; Crockett and Wilson, 1980; Fooden, 1982a, 1985, 2000; Fooden and Wu, 2001; Chatani, 2003; Fooden, 2007).

The species group classification used in this study was based on that of Delson (1980) (Fig. 1A). Although there has been some debate concerning whether *M. arctoides* is more closely related to the *sinica* or *fascicularis* group (Morales and Melnick, 1998; Tosi et al., 2000, 2003) (Figs. 1B, C), a recent molecular study by Li et al. (2009) demonstrated strong statistical support for the placement of *M. arctoides* in the *sinica* group, confirming the classification by Delson (1980) (Fig. 1D). This result is robust because *Alu* elements used in this study showed very little homoplasy and their ancestral state is known (Li et al., 2009).

Supporting information for Table 2: Samples used in this study

For the determination of sampling units on the basis of the distribution pattern and genetic divergence, we referred to the following literature: Fooden (1975, 1982b, 1990b, 1995, 2000), Tosi et al. (2002), Evans et al. (2003), Marmi et al. (2004), Fooden and Aimi (2005), Tosi and Coke (2007), Ziegler et al. (2007), Shiina et al. (2010).

Supporting information references

- Chatani K. 2003. Positional behavior of free-ranging Japanese macaques (*Macaca fuscata*).
Primates 44:13-23.
- Crockett CM, and Wilson WL. 1980. The ecological separation of *Macaca nemestrina* and *M. fascicularis* in Sumatra. In: Lindburg DG, editor. The macaques: Studies in ecology, behavior and evolution. New York: Nostrand-Reinhold. p 148-181.
- Delson E. 1980. Fossil macaques, phyletic relationships and a scenario of deployment. In: Lindburg DG, editor. The macaques: Studies in ecology, behavior and evolution. New York: Nostrand-Reinhold. p 10-30.
- Eudey AA. 1980. Pleistocene glacial phenomena and the evolution of Asian macaques. In: Lindburg DG, editor. The macaques: Studies in ecology, behavior and evolution. New York: Nostrand Reinhold Ltd. p 52-83.
- Evans BJ, Supriatna J, Andayani N, and Melnick DJ. 2003. Diversification of Sulawesi macaque monkeys: decoupled evolution of mitochondrial and autosomal DNA. *Evolution* 57:1931-1946.
- Fa J. 1989. The genus *Macaca*: a review of taxonomy and evolution. *Mamm Rev* 19(2):45-81.
- Fooden J. 1975. Taxonomy and evolution of liontail and pigtail macaques. *Fieldiana Zool* 67:1-169.
- Fooden J. 1979. Taxonomy and evolution of the *sinica* group of macaques: 1. species and subspecies accounts of *Macaca sinica*. *Primates* 20(1):109-140.

Fooden J. 1982a. Ecogeographic segregation of macaque species. *Primates* 23(4):574-579.

Fooden J. 1982b. Taxonomy and evolution of the *sinica* group of macaques: 3. species and subspecies accounts of *Macaca assamensis*. *Fieldiana Zool* 10:1-52.

Fooden J. 1985. Taxonomy and evolution of the *sinica* group of macaques: 5. Overview of natural history. *Fieldiana Zool* 29:1-22.

Fooden J. 1990a. The bear macaques, *Macaca arctoides*: a systematic review. *J Hum Evol* 19:607-686.

Fooden J. 1990b. Systematic review of Philippine macaques (Primates, Cercopithecidae: *Macaca fascicularis* subsp.). *Fieldiana Zool* 64:1-44.

Fooden J. 1995. Systematic review of southeast Asian longtail macaques, *Macaca fascicularis* (Raffles, 1821). *Fieldiana Zool* 81(30):1-206.

Fooden J. 2000. Systematic review of the rhesus macaques, *Macaca mulatta* (Zimmermann, 1780). *Fieldiana Zool* 96(30):1-180.

Fooden J. 2007. Systematic review of the Barbary macaque, *Macaca sylvanus* (Linnaeus, 1758). *Fieldiana Zool* 113(63):1-58.

Fooden J, and Aimi M. 2005. Systematic review of Japanese macaques, *Macaca fuscata* (Gray, 1870). *Fieldiana Zool* 104(14):1-200.

Fooden J, and Wu HY. 2001. Systematic review of the Taiwanese macaque, *Macaca cyclopis* Swinhoe, 1863. *Fieldiana Zool* 98(30):1-70.

- Hanya G. 2004. Diet of a Japanese macaque troop in the coniferous forest of Yakushima. *Int J Primatol* 25(1):55-69.
- Hanya G, Menard N, Qarro M, Ibn Tattou M, Fuse M, Vallet D, Yamada A, Go M, Takafumi H, Tsujino R et al. . 2011. Dietary adaptations of temperate primates: comparisons of Japanese and Barbary macaques. *Primates* 52(2):187-198.
- Izawa K, and Nishida T. 1963. Monkeys living in the northern limits of their distribution. *Primates* 4(2):67-88.
- Li J, Han K, Xing J, Kim HS, Rogers J, Ryder OA, Disotell T, Yue B, and Batzer MA. 2009. Phylogeny of the macaques (Cercopithecidae: *Macaca*) based on Alu elements. *Gene* 448:242-249.
- Marmi J, Bertranpetit J, Terradas J, Takenaka O, and Domingo-Roura X. 2004. Radiation and phylogeography in the Japanese macaque, *Macaca fuscata*. *Mol Phylogen Evol* 30:676-685.
- Morales JC, and Melnick DJ. 1998. Phylogenetic relationships of the macaques (Cercopithecidae: *Macaca*), as revealed by high resolution restriction site mapping of mitochondrial ribosomal genes. *J Hum Evol* 34:1-23.
- Shiina T, Tanaka K, Katsuyama Y, Otabe K, Sakamoto K, Kurata M, Nomura M, Yamanaka H, Nakagawa H, Inoko H et al. 2010. Mitochondrial DNA diversity among three subpopulations of cynomolgus macaques (*Macaca fascicularis*) originating from the

- Indochinese region. *Exp Anim* 5:567-578.
- Sugiyama Y. 1971. Characteristics of the social life of bonnet macaques (*Macaca radiata*).
Primates 12(3-4):247-266.
- Suzuki A. 1965. An ecological study of wild Japanese monkeys in snowy areas: focused on
their food habits. *Primates* 6(1):31-72.
- Tosi AJ, and Coke CS. 2007. Comparative phylogenetics offer new insights into the
biogeographic history of *Macaca fascicularis* and the origin of the Mauritian
macaques. *Mol Phylogen Evol* 42:498-504.
- Tosi AJ, Morales JC, and Melnick DJ. 2000. Comparison of Y chromosome and mtDNA
phylogenies leads to unique inferences of macaque evolutionary history. *Mol
Phylogen Evol* 17:133-144.
- Tosi AJ, Morales JC, and Melnick DJ. 2002. Y-chromosome and mitochondrial markers in
Macaca fascicularis indicate introgression with Indochinese *M. mulatta* and a
biogeographic barrier in the isthmus of Kra. *Int J Primatol* 23(1):161-178.
- Tosi AJ, Morales JC, and Melnick DJ. 2003. Paternal, maternal, and biparental molecular
markers provide unique windows onto the evolutionary history of macaque monkeys.
Evolution 57:1419-1435.
- Zhao QK, Deng ZY, and Xu JM. 1991. Natural foods and their ecological implications for
Macaca thibetana at Mount Emei, China. *Folia Primatol (Basel)* 57(1):1-15.

Ziegler T, Abegg C, Meijaard E, Perwitasari-Farajallah D, Walter L, Hodges JK, and Roos C.

2007. Molecular phylogeny and evolutionary history of Southeast Asian macaques

forming the *M. silenus* group. Mol Phylogen Evol 42:807-816.