TITLE:

Regulatory expression of components in the BMP pathway in white adipose tissues of cattle

AUTHOR(S):

Qiao, Yuhang; Yamada, Tomoya; Kanamori, Yohei; Kida, Ryosuke; Shigematsu, Mei; Fujimoto, Yusuke; Tomonaga, Shozo; Matsui, Tohru; Funaba, Masayuki

CITATION:

Qiao, Yuhang ...[et al]. Regulatory expression of components in the BMP pathway in white adipose tissues of cattle. *Livestock Science* 2015, 174: 144-149

ISSUE DATE:

2015-04

URL:

<http://hdl.handle.net/2433/198736>

RIGHT:

© 2015 Elsevier B.V. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>. NOTICE: this is the author's version of a work that was accepted for publication in [*Livestock Science*]. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in [*Livestock Science*, Volume 174, Pages 144-149], doi:10.1016/j.livsci.2015.02.008.; 許諾条件により本文ファイルは2015-02-20に公開.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

Regulatory expression of components in the BMP pathway in white
adipose tissues of cattle

Yuhang Qiao^{1*}, Tomoya Yamada^{2*}, Yohei Kanamori¹, Ryosuke Kida¹,
Mei Shigematsu¹, Yusuke Fujimoto¹, Shozo Tomonaga¹,
Tohru Matsui¹ and Masayuki Funaba^{1†}

¹Division of Applied Biosciences, Graduate School of Agriculture, Kyoto University,
Kyoto 606-8502, Japan.

²National Institute of Livestock and Grassland Science, Nasushiobara 329-2793, Japan.

*These authors equally contributed to this study.

†To whom correspondence should be addressed:

Masayuki Funaba, Ph.D.

Division of Applied Biosciences

Graduate School of Agriculture

Kyoto University

Kitashirakawa Oiwakecho, Kyoto 606-8502, Japan

Tel: +81-75-753-6055

Fax: +81-75-753-6344

E-mail: mfunaba@kais.kyoto-u.ac.jp

Abstract

The BMP pathway is known to positively regulate murine brown adipogenesis. The present study examined the mRNA levels of BMPs and activin β B as well as receptors for the BMP pathway in the adipose tissues of cattle fed diets with differential ratio of concentrate to roughage or vitamin A-deficient diet. The expression of activin β B was significantly increased in the subcutaneous fat depot of animals fed the concentrate diet, while the vitamin A-deficient diet significantly increased the expression of BMP4 in the mesenteric fat depot. The expression of receptors for the BMP pathway, ALK2, ALK3, ActRIIA, and BMPR2, showed a similar pattern to that of BMP4 and activin β B in response to the dietary treatments. The results of the present study suggest that diet modulates expression of BMP pathway and may be responsible for the regulatory expression of brown/beige adipocyte-related genes in the adipose tissues of cattle.

Keywords: BMP; brown adipocyte; beige adipocyte; cattle

1. Introduction

Brown/beige adipocytes dissipate chemical energy in the form of heat (Cannon and Nedergaard, 2004; Shabalina et al., 2013); the expression of uncoupling protein 1 (Ucp1), a proton channel located on the inner mitochondrial membrane, enables brown/beige adipocytes to expedite energy (Cannon and Nedergaard, 2004). In mice, brown adipose tissue located in the interscapular region is composed of brown adipocytes (Smorlesi et al., 2012), whereas beige adipocytes, also known as brite adipocytes, are sparsely distributed in various white adipose tissues including subcutaneous, mesenteric and epididymal fat depots (Ishibashi and Seale, 2010). Molecular signatures revealed a distinct pattern between brown adipocytes and beige adipocytes, suggesting that the origin of these two adipocytes differ (Wu et al., 2013).

We previously reported Ucp1-positive adipocytes in the white adipose tissues of cattle (Asano et al., 2013). Furthermore, the diet affected the expression levels of genes predominantly expressed in brown/beige adipocytes more than those expressed in white adipocytes (Seale et al., 2007; Sharp et al., 2012) in the fat depots of cattle (Asano et al., 2013; Kanamori et al., 2014); the genes included Ucp1, Cidea, Dio2, Cox1 and Cox8b (Asano et al., 2013; Kanamori et al., 2014). Based on the close relationship between the expression levels of brown/beige adipocyte-related genes and brown/beige adipocyte activity (Seale et al., 2008; Tseng et al., 2008; Barbatelli et al., 2010; Yadav et al., 2011; Whittle et al., 2012), these findings suggest that diet may modulate energy expenditure in the brown/beige adipocytes of cattle. Beef cattle are raised as industrial animals, and fattening efficiency is one of the determining factors of the economy of beef production. Therefore, the activation of brown/beige adipocytes is desirably avoided because their activation may decrease fattening efficiency, the ratio of body weight gain to feed consumed.

The differentiation of brown/beige preadipocytes is regulated by the bone morphogenetic protein (BMP) pathway (Tseng et al., 2008; Schulz et al., 2011; Elsen et al., 2014; Xue et al., 2014). BMP7 was previously shown to stimulate brown adipogenesis (Tseng et al., 2008; Schulz et al., 2011), and BMP4 was recently reported to induce differentiation into brown/beige adipocytes (Elsen et al., 2014; Xue et al., 2014). We hypothesized that the diet-induced modulation of the expression of brown/beige adipocyte-related genes is involved in the activity of the BMP pathway. The induction of ligands and their signaling components often occurs as a positive regulation of BMP signaling (Miyazono et al., 2000). Therefore, the present study examined the expression levels of components that elicit BMP signaling in the adipose tissues of cattle fed a roughage diet or concentrate diet, and a control or vitamin A-deficient diet. In previous studies, the expression levels of brown/beige adipocyte-related genes were modulated by the diets (Asano et al., 2013; Kanamori et al., 2014).

2. Materials and methods

2.1. *Animals and feeds*

Sixteen Japanese Black steers were used in the present study. Dietary treatments were the same as those reported by Yamada and Nakanishi (2012) and Asano et al. (2013) for experiment (Exp) 1 and Yamada et al. (2013) and Kanamori et al. (2014) for Exp 2.

Briefly, in Exp 1, 10-month-old steers were allotted by body weight to one of two groups: a roughage diet group (n = 4) or concentrate diet group (n = 4). The roughage diet consisted of 35% roughage (orchard grass hay) and 65% concentrate mixture consisting of corn, barley, wheat bran, rice bran, and soybean meal (Nasuno for Wagyu Fattening; JA Higashi-nihon Kumiai Shiryō, Ota, Gunma, Japan), whereas the

concentrate diet contained 10% roughage and 90% concentrate mixture on a total digestible nutrients (TDN) basis.

In Exp 2, 10-month-old steers were allotted by body weight to one of two groups: a control group (n = 4) or vitamin A-deficient group (n = 4). Two kinds of fermented total mixed ration (TMR) containing whole crop rice silage, fermented byproducts (beer cake and tofu cake), and concentrate, i.e., TMR-A and TMR-B, were prepared in a TMR center, and used as a control diet. The content of β -carotene was 7 mg/kg and 16 mg/kg for TMR-A and TMR-B, respectively. TMR-A and TMR-B were given to steers aged 10-20 months and 21-30 months, respectively. The steers in the vitamin A-deficient group were fed orchard grass hay (0.5 mg/kg of β -carotene) and a concentrate mixture consisting of corn, barley, wheat bran, rice bran, and soybean meal (0.1 mg/kg of β -carotene, Nasuno for Wagyu Fattening; JA Higashi-nihon Kumiai Shiryou, Ota, Gunma, Japan). The supplied dry matter amount of orchard grass hay in the vitamin A-deficient group was matched to that of whole crop rice silage in the control diet to equalize the ratio of roughage to concentrate between the two groups.

Feeds were provided to steers to equalize TDN intake for 20 months between groups in both experiments. Subcutaneous and mesenteric adipose tissues were collected at 30 months of age (Yamada and Nakanishi, 2012; Yamada et al., 2013). All animals received humane care as outlined in the Guide for the Care and Use of Experimental Animals (National Institute of Livestock and Grassland Science, #09032638 for Exp 1 and #10032404 for Exp 2).

2.2. RNA isolation and RT-quantitative PCR

Total RNA isolation, cDNA synthesis, and real-time RT-quantitative PCR (RT-qPCR) were conducted as described by Asano et al. (2013). The cDNA synthesized from 5 or

10 ng of total RNA was subjected to RT-qPCR analyses as the template. Ct values were determined, and the abundance of gene transcripts was analyzed by the $\Delta\Delta\text{Ct}$ method using Hprt1 as the corrected gene (Duran et al., 2005). Gene expression in the roughage diet group (Exp 1) or control group (Exp 2) in each fat depot was set to 100. The nucleotide sequence of qPCR primers, size of qPCR product, and mean Ct value in the subcutaneous fat depot are shown in Table 1.

2.3. Statistical analyses

Data are expressed as the mean $\pm$ SE. Data on gene expression were log-transformed to provide an approximation of a normal distribution before analyses. Differences between dietary groups were examined by the unpaired *t*-test using Microsoft Excel®. Differences of $P < 0.05$ were considered significant. A slight difference was considered to be present when $0.05 \leq P < 0.10$.

3. Results

Effects of dietary ratio of roughage to concentrate on the gene expression of BMPs and activin βB in the adipose tissues of cattle were examined in Exp. 1 (Fig. 1A). The expression of activin βB in the subcutaneous fat depot was significantly higher ($P = 0.007$) in the concentrate diet group than in the roughage diet group. The expression of BMP4 tended to be higher in the concentrate diet group ($P = 0.06$). In contrast, no significant differences were observed in the expression of these ligands in the mesenteric fat depot between the dietary groups. Exp 2 examined the effects of a dietary vitamin A deficiency (Fig. 1B). The expression of BMP4 in the mesenteric fat depot was significantly increased by feeding the vitamin A-deficient diet ($P = 0.02$). In addition, the expression of activin βB tended to be higher in the vitamin A-deficient diet group ($P = 0.07$). The expression of BMP7 was 20-fold higher in the vitamin

A-deficient diet group than in the control diet group, although not significantly so because of large variations in the vitamin A-deficient diet group ($P = 0.18$). The vitamin A-deficient diet did not affect the expression levels of BMPs or activin β B in the subcutaneous fat depot. The diet-related changes in gene expression were not a non-specific event, because the expression of fatty acid synthase (FAS) was not significantly changed by the diets (Fig. 2). Expression of TGF- β 1 in the fat depots was also unchanged, and similar changes were observed in the expression levels by the diets when β 2-macroglobulin was used as a corrected gene (data not shown). The Ct value of BMPs and activin β B was relatively larger than that of FAS (Table 1), suggesting lower expression of BMPs and activin β B than that of FAS.

BMP signals through complex formation with the type I receptor and type II receptor (Miyazono et al., 2010); ALK2 and ALK3 are BMP type I receptors, while ActRIIA, ActRIIB and BMPR2 are BMP type II receptors (Miyazono et al., 2010). The expression of ALK2 and ALK3 in the subcutaneous fat depot tended to be higher in the concentrate diet group than in the roughage diet group ($P = 0.07$ and $P = 0.09$, respectively), whereas no significant differences were observed in expression levels in the mesenteric fat depot between the groups (Fig. 3A). The vitamin A deficiency did not affect the expression of BMP receptors in the subcutaneous fat depot (Fig. 3B). However, the vitamin A-deficient diet significantly or tended to increase the expression of ALK2 ($P = 0.03$), ALK3 ($P = 0.03$), ActRIIA ($P = 0.04$), and BMPR2 ($P = 0.05$) in the mesenteric fat depot. Significant changes in the expression of ActRIIB were not detected in the adipose tissues (data not shown).

4. Discussion

Brown adipogenesis is stimulated by the activation of the BMP pathway (Tseng et al., 2008; Schulz et al., 2011; Elsen et al., 2014; Xue et al., 2014). Signal activity of the BMP pathway is often regulated at the mRNA level of components involved in the signal (Miyazono, 2000). Previously, we reported that the expression of Ucp1 in the subcutaneous fat depot, but not the mesenteric fat depot was higher in cattle fed a concentrate diet than in those fed a roughage diet (Asano et al., 2013). Furthermore, the expression levels of brown/beige adipocyte-related genes (Seale et al., 2007) in the mesenteric fat depot were generally higher in cattle fed the vitamin A-deficient diet than in those fed the control diet, whereas these expression levels in the subcutaneous fat depot was similar between the groups (Kanamori et al., 2014). Taken the evidence, we hypothesized that expression levels of the components of the BMP pathway are changed in the fat depots of cattle fed diets with different ratio of roughage to concentrate or vitamin A-deficient diet. We evaluated the mRNA levels of molecules that elicit BMP signaling. In addition to BMP4 and BMP7, we examined the expression level of activin β B. The signal transduction of activin B, a homodimer of the activin β B subunit, has not yet been elucidated in detail; activin B signals in a similar manner to activin A, a structurally related molecule (Tsuchida et al., 2004), but also transmits BMP-mediated signals (Besson-Fournier et al., 2012). The present study revealed that the expression levels of BMP4, activin β B and BMP receptors were up-regulated in the subcutaneous fat depot of cattle fed the concentrate diet and in the mesenteric fat depot of cattle fed the vitamin A-deficient diet. The present results suggest that the modulation of expression levels of brown/beige adipocyte-related genes in the fat depots of cattle shown in previous studies (Asano et al., 2013; Kanamori et al., 2014) was achieved by altered activity of the BMP pathway.

In humans, expression levels of the components of the BMP pathway in adipose tissue appear to be related to adiposity. The expression level of activin β B in adipose tissue

was previously shown to be higher in obese humans than in lean humans, and was decreased by reductions in body weight (Sjoholm et al., 2006). A negative relationship has been reported between BMP4 expression levels in adipose tissue and body mass index (Qian et al., 2013). Furthermore, the expression levels of ALK3 and BMPR2 in adipose tissue were found to be higher in obese humans than in lean humans (Böttcher et al., 2009; Schleintz et al., 2011). The carcass composition of cattle used in this study was similar between the dietary groups because they were pair-fed to equalize TDN intake (Yamada and Nakanishi, 2012; Yamada et al., 2013). Nevertheless, these diets significantly affected the expression levels of molecules involved in the BMP pathway in the adipose tissues of cattle. This altered gene expression may reflect an intrinsic effect of the diet, but not adiposity.

Size of white adipocytes was changed in fattening cattle used in this study (Yamada and Nakanishi, 2012; Yamada et al., 2013); white adipocytes were smaller in cattle fed the concentrate diet and the vitamin A-deficient diet. Considering that the number of beige adipocytes related to the decrease in cell size of white adipocytes in mice (Qian et al., 2013), the stimulation of the BMP pathway through increased expression of the signaling components may induce emergence or activation of brown/beige adipocytes, leading to decrease in adipocyte size of cattle fed the concentrate diet or vitamin A-deficient diet. Generally, adiposity is closely related to the increased size of white adipocytes (Tchermof and Despres, 2013). However, as described above, the carcass composition of cattle was not significantly altered by the diets (Yamada and Nakanishi, 2012; Yamada et al., 2013). It is possible that fat deposition evaluated by carcass analyses is relatively insensitive to clarify effects of diet under the condition of pair-fed on TDN intake. In fact, expression level of FAS was not affected by the diets.

The present study clarified that expression of components of the BMP pathway was

changed by the diets in an anatomical location-dependent manner. The reason why the diet effect was different between fat depots is unclear. Development of adipose tissue in ruminants depends on its anatomical location (Alexander, 1978; Bonnet et al., 2010). Growth of adipose tissue is also dependent on breed of cattle (Landis et al., 2002). Basically, brown adipose tissue is developed during fetal period in cattle, whereas white adipose tissue grows mainly after birth (Bonnet et al., 2010). The differential development of adipose tissue depending on its anatomical location may be responsible for diet-related modulation of expression levels of genes involved in the BMP pathway. As compared with modulation of the ratio of roughage to concentrate, feeding the vitamin A-deficient diet induces severer nutritional restriction. It is possible that the extent in nutritional modulation is responsible for the differences of the affected fat depot.

Feeding vitamin A-deficient diet increased plasma concentration of 8-isoprostane, a marker of oxidative stress, in cattle (Yamada et al., 2013). Mitochondrial uncoupling mediated by Ucp1 expression reduces production of reactive oxygen species, suggesting that Ucp1 expression in brown adipocytes has a role in prevention of oxidative stress (Oelkrog et al., 2014). It is possible that increased expression of genes involved in the BMP pathway reflects defensive reaction to oxidative stress induced by vitamin A-deficient diet.

Prdm16 is a master regulator of the development of brown adipocytes as well as the emergence of beige adipocytes (Seale et al., 2008, 2011). The expression level of Prdm16 in the subcutaneous, but not mesenteric fat depot was shown to be higher in the concentrate diet group than in the roughage diet group (Asano et al., 2013). The vitamin A-deficient diet also increased Prdm16 expression in the mesenteric fat depot, but not in the subcutaneous fat depot (Kanamori et al., 2014). The up-regulation of Prdm16

expression by BMP4 and BMP7 in brown preadipocytes as well as mesenchymal stem cells (Tseng et al., 2008; Xue et al., 2014) suggests that activation of the BMP pathway may induce browning, which is defined as an increase in the mRNA levels of brown/beige adipocyte-related genes, including Ucp1 (Fisher et al., 2012; Nedergaard and Cannon, 2014), in the fat depots of cattle through the up-regulation of Prdm16 expression. Further studies are needed to clarify dietary factor(s) affecting the expression levels of molecules for the BMP pathway and cells affected by dietary factor(s).

Acknowledgement

This work was supported by a Grant-in-Aid for Scientific Research (23580368 and 26292137) from The Japan Society for the Promotion of Science.

Conflict of interest statement

Authors of this manuscript certify that there is no conflict of interest.

References

- Alexander, G., 1978. Quantitative development of adipose tissue in foetal sheep. *Aust. J. Biol. Sci.* 31, 489-504.
- Asano, H., Yamada, T., Hashimoto, O., Umemoto, T., Sato, R., Ohwatari, S., Kanamori, Y., Terachi, T., Funaba, M., Matsui, T., 2013. Diet-induced changes in Ucp1 expression in bovine adipose tissues. *Gen. Comp. Endocrinol.* 184, 87-92.
- Barbatelli, G., Murano, I., Madsen, L., Hao, Q., Jimenez, M., Kristiansen, K., Giacobino, J.P., De Matteis, R., Cinti, S., 2010. The emergence of cold-induced brown adipocytes in mouse white fat depots is determined predominantly by white to brown adipocyte transdifferentiation. *Am. J. Physiol.*

- 281 Endocrinol. Metab. 298, E1244-E1253.
- 282 Besson-Fournier, C., Latour, C., Kautz, L., Bertrand, J., Ganz, T., Roth, M.P., Coppin,
283 H., 2012. Induction of activin B by inflammatory stimuli up-regulates expression
284 of the iron-regulatory peptide hepcidin through Smad1/5/8 signaling. Blood 120,
285 431-439.
- 286 Bonnet, M., Cassar-Malek, I., Chilliard, Y., Picard, B., 2010. Ontogenesis of muscle
287 and adipose tissues and their interactions in ruminants and other species. Animal 4,
288 1093-1109.
- 289 Böttcher, Y., Unbehauen, H., Klötting, N., Ruschke, K., Körner, A., Schleinitz,
290 D., Tönjes, A., Enigk, B., Wolf, S., Dietrich, K., Koriath, M., Scholz, G.H., Tseng,
291 Y.H., Dietrich, A., Schön, M.R., Kiess, W., Stumvoll, M., Blüher, M., Kovacs, P.,
292 2009. Adipose tissue expression and genetic variants of the bone morphogenetic
293 protein receptor 1A gene (BMPRI1A) are associated with human obesity.
294 Diabetes 58, 2119-2128.
- 295 Cannon, B., Nedergaard, J., 2004. Brown adipose tissue: function and physiological
296 significance. Physiol. Rev. 84, 277-359.
- 297 Duran, E.M., Shapshak, P., Worley, J., Minagar, A., Ziegler, F., Haliko, S.,
298 Moleon-Borodowsky, I., Haslett, P.A., 2005. Presenilin-1 detection in brain
299 neurons and FOXP3 in peripheral blood mononuclear cells: normalizer gene
300 selection for real time reverse transcriptase pcr using the deltadeltaCt method.
301 Front Biosci. 10, 2955-2965.
- 302 Elsen, M., Raschke, S., Tennagels, N., Schwahn, U., Jelenik, T., Roden, M., Romacho,
303 T., Eckel, J., 2014. BMP4 and BMP7 induce the white-to-brown transition of
304 primary human adipose stem cells. Am. J. Physiol. Cell Physiol. 306, C431-C440.
- 305 Fisher, F.M., Kleiner, S., Douris, N., Fox, E.C., Mepani, R.J., Verdeguer, F., Wu, J.,
306 Kharitonov, A., Flier, J.S., Maratos-Flier, E., Spiegelman, B.M., 2012.
307 FGF21 regulates PGC-1 α and browning of white adipose tissues in adaptive

- 308 thermogenesis. *Genes Dev.* 26, 271-281.
- 309 Ishibashi, J., Seale, P., 2010. Beige can be slimming. *Science* 328, 1113-1114.
- 310 Kanamori, Y., Yamada, T., Asano, H., Kida, R., Qiao, Y., Abd Eldaim, M.A., Tomonaga,
311 S., Matsui, T., Funaba, M., 2014. Effects of vitamin A status on expression of
312 Ucp1 and brown/beige adipocyte-related genes in white adipose tissues of beef
313 cattle. *J. Vet. Med. Sci.* 76, 1261-1265.
- 314 Landis, M.D., Carstens, G.E., McPhail, E.G., Randel, R.D., Green, K.K., Slay,
315 L., Smith, S.B., 2002. Ontogenic development of brown adipose tissue in Angus
316 and Brahman fetal calves. *J. Anim. Sci.* 80, 591-601.
- 317 Miyazono, K., 2000. Positive and negative regulation of TGF- β signaling. *J. Cell*
318 *Sci.* 113, 1101-1109.
- 319 Miyazono, K., Kamiya, Y., Morikawa, M., 2010. Bone morphogenetic protein receptors
320 and signal transduction. *J. Biochem.* 147, 35-51.
- 321 Nedergaard, J., Cannon, B., 2014. The browning of white adipose tissue: some burning
322 issues. *Cell Metab.* 20, 396-407.
- 323 Oelkrug, R., Götze, N., Meyer, C.W., Jastroch, M., 2014. Antioxidant properties of
324 UCP1 are evolutionarily conserved in mammals and buffer mitochondrial reactive
325 oxygen species. *Free Radic. Biol. Med.* 77, 210-216.
- 326 Qian, S.W., Tang, Y., Li, X., Liu, Y., Zhang, Y.Y., Huang, H.Y., Xue, R.D., Yu,
327 H.Y., Guo, L., Gao, H.D., Liu, Y., Sun, X., Li, Y.M., Jia, W.P., Tang, Q.Q., 2013.
328 BMP4-mediated brown fat-like changes in white adipose tissue alter glucose and
329 energy homeostasis. *Proc. Natl. Acad. Sci. USA* 110, E798-E807.
- 330 Schleinitz, D., Klötting, N., Böttcher, Y., Wolf, S., Dietrich, K., Tönjes, A., Breitfeld,
331 J., Enigk, B., Halbritter, J., Körner, A., Schön, M.R., Jenkner, J., Tseng,
332 Y.H., Lohmann, T., Dressler, M., Stumvoll, M., Blüher, M., Kovacs, P., 2011.
333 Genetic and evolutionary analyses of the human bone morphogenetic protein
334 receptor 2 (BMPR2) in the pathophysiology of obesity. *PLoS One* 6, e16155.

- 335 Schulz, T.J., Huang, T.L., Tran, T.T., Zhang, H., Townsend, K.L., Shadrach, J.L.,
336 Cerletti, M., McDougall, L.E., Giorgadze, N., Tchkonina, T., Schrier, D., Falb, D.,
337 Kirkland, J.L., Wagers, A.J., Tseng, Y.H., 2011. Identification of inducible brown
338 adipocyte progenitors residing in skeletal muscle and white fat. *Proc. Natl. Acad.*
339 *Sci. USA* 108, 143-148.
- 340 Seale, P., Bjork, B., Yang, W., Kajimura, S., Chin, S., Kuang, S., Scimè, A.,
341 Devarakonda, S., Conroe, H.M., Erdjument-Bromage, H., Tempst, P., Rudnicki,
342 M.A., Beier, D.R., Spiegelman, B.M., 2008. PRDM16 controls a brown
343 fat/skeletal muscle switch. *Nature* 454, 961-967.
- 344 Seale, P., Conroe, H.M., Estall, J., Kajimura, S., Frontini, A., Ishibashi, J., Cohen,
345 P., Cinti, S., Spiegelman, B.M., 2011. Prdm16 determines the thermogenic
346 program of subcutaneous white adipose tissue in mice. *J. Clin. Invest.* 121, 96-105.
- 347 Seale, P., Kajimura, S., Yang, W., Chin, S., Rohas, L.M., Uldry, M., Tavernier, G.,
348 Langin, D., Spiegelman, B.M., 2007. Transcriptional control of brown fat
349 determination by PRDM16. *Cell Metab.* 6, 38-54.
- 350 Shabalina, I.G., Petrovic, N., de Jong, J.M., Kalinovich, A.V., Cannon, B., Nedergaard,
351 J., 2013. UCP1 in brite/beige adipose tissue mitochondria is functionally
352 thermogenic. *Cell Rep.* 5, 1196-1203.
- 353 Sharp, L.Z., Shinoda, K., Ohno, H., Scheel, D.W., Tomoda, E., Ruiz, L., Hu, H., Wang,
354 L., Pavlova, Z., Gilsanz, V., Kajimura, S., 2012. Human BAT possesses molecular
355 signatures that resemble beige/brite cells. *PLoS One* 7, e49452.
- 356 Sjöholm, K., Palming, J., Lystig, T.C., Jennische, E., Woodruff, T.K., Carlsson, B.,
357 Carlsson, L.M., 2006. The expression of inhibin β B is high in human adipocytes,
358 reduced by weight loss, and correlates to factors implicated in metabolic disease.
359 *Biochem. Biophys. Res. Commun.* 344, 1308-1314.

- 360 Smorlesi, A., Frontini, A., Giordano, A., Cinti, S., 2012. The adipose organ:
361 white-brown adipocyte plasticity and metabolic inflammation. *Obes. Rev.* 13
362 Suppl 2, 83-96.
- 363 Tchernof A, Després JP. Pathophysiology of human visceral obesity: an update. *Physiol*
364 *Rev.* 2013 93:359-404.
- 365 Tseng, Y.H., Kokkotou, E., Schulz, T.J., Huang, T.L., Winnay, J.N., Taniguchi, C.M.,
366 Tran, T.T., Suzuki, R., Espinoza, D.O., Yamamoto, Y., Ahrens, M.J., Dudley, A.T.,
367 Norris, A.W., Kulkarni, R.N., Kahn, C.R., 2008. New role of bone morphogenetic
368 protein 7 in brown adipogenesis and energy expenditure. *Nature* 454, 1000-1004.
- 369 Tsuchida, K., Nakatani, M., Yamakawa, N., Hashimoto, O., Hasegawa, Y., Sugino, H.,
370 2004. Activin isoforms signal through type I receptor serine/threonine kinase
371 ALK7. *Mol. Cell. Endocrinol.* 220, 59-65.
- 372 Whittle, A.J., Carobbio, S., Martins, L., Slawik, M., Hondares, E., Vázquez, M.J.,
373 Morgan, D., Csikasz, R.I., Gallego, R., Rodriguez-Cuenca, S., Dale, M., Virtue, S.,
374 Villarroya, F., Cannon, B., Rahmouni, K., López, M., Vidal-Puig, A., 2012.
375 BMP8B increases brown adipose tissue thermogenesis through both central and
376 peripheral actions. *Cell* 149, 871-885.
- 377 Wu, J., Cohen, P., Spiegelman, B.M. 2013. Adaptive thermogenesis in adipocytes: is
378 beige the new brown? *Genes Dev.* 27, 234-250.
- 379 Xue, R., Wan, Y., Zhang, S., Zhang, Q., Ye, H., Li, Y., 2014. Role of bone
380 morphogenetic protein 4 in the differentiation of brown fat-like adipocytes. *Am. J.*
381 *Physiol. Endocrinol. Metab.* 306, E363-E372.
- 382 Yadav, H., Quijano, C., Kamaraju, A.K., Gavrilova, O., Malek, R., Chen, W., Zervas, P.,
383 Zhigang, D., Wright, E.C., Stuelten, C., Sun, P., Lonning, S., Skarulis, M., Sumner,
384 A.E., Finkel, T., Rane, S.G., 2011. Protection from obesity and diabetes by
385 blockade of TGF- β /Smad3 signaling. *Cell Metab.* 14, 67-79.
- 386 Yamada, T., Higuchi, M., Nakanishi, N., 2013. Plasma 8-isoprostane concentrations and

387 adipogenic and adipokine gene expression patterns in subcutaneous and
388 mesenteric adipose tissues of fattening Wagyu cattle. J. Vet. Med. Sci. 75,
389 1021-1027.

390 Yamada, T., Nakanishi, N., 2012. Effects of the roughage/concentrate ratio on the
391 expression of angiogenic growth factors in adipose tissue of fattening Wagyu
392 steers. Meat Sci. 90, 807-813.

393

394

Figure legends

Fig. 1. Expression of BMP4, BMP7, and activin β B in white fat depots of cattle

(A) In Exp 1, cattle were fed either a roughage diet or concentrate diet for 20 months.
(B) In Exp 2, cattle were fed either a control diet or vitamin A-deficient diet for 20 months. At 30 months of age, subcutaneous and mesenteric white fat depots were collected, and the mRNA levels of BMP4, BMP7, and activin β B were examined by RT-qPCR. Expression was normalized to that of Hprt1, and gene expression in each fat depot in the roughage diet group (A) or control diet group (B) was set to 100. Data are shown as the mean $\pm$ SE ($n = 4$). $\dagger$, * and **: $P < 0.10$, 0.05 and $P < 0.01$, respectively.

Fig. 2. Expression of FAS in white fat depots of cattle

(A) In Exp 1, cattle were fed either a roughage diet or concentrate diet for 20 months.
(B) In Exp 2, cattle were fed either a control diet or vitamin A-deficient diet for 20 months. At 30 months of age, subcutaneous and mesenteric white fat depots were collected, and the mRNA level of FAS was examined by RT-qPCR. Expression was normalized to that of Hprt1, and gene expression in each fat depot in the roughage diet group (A) or control diet group (B) was set to 100. Data are shown as the mean $\pm$ SE ($n = 4$).

Fig. 3. Expression of receptors transmitting BMP signals in white fat depots of cattle

(A) In Exp 1, cattle were fed either a roughage diet or concentrate diet for 20 months.
(B) In Exp 2, cattle were fed either a control diet or vitamin A-deficient diet for 20 months. At 30 months of age, subcutaneous and mesenteric white fat depots were collected, and the mRNA levels of receptors for the BMP pathway were examined by RT-qPCR. Expression was normalized to that of Hprt1, and gene expression in each fat depot in the roughage diet group (A) or control diet group (B) was set to 100. Data are

422 shown as the mean $\pm$ SE ($n = 4$). † and *: $P < 0.10$ and 0.05, respectively.

423

424 Table 1. Nucleotide sequence of primers used in RT-qPCR analyses, size of PCR product and mean Ct value

425	Gene	Primer	Sequence	GenBank	Size (bp)	Ct value ¹
426				accession number		
427	Activin β B	5'	5'-cgtctccgagatcatcagc-3'	NM_176852	89	28.67 $\pm$ 0.45
428		3'	5'-ggttgccttcgttgagat-3'			
429	ActRIIA	5'	5'-gctcgtccaactcaagacc-3'	NM_174227	75	27.41 $\pm$ 0.97
430		3'	5'-tctagtaactgtaatggcttcaaacct-3'			
431	ActRIIB	5'	5'-tttgagccagggaacctcc-3'	NM_174495	125	BDL ²
432		3'	5'-tacatgtcgcgcagaaa-3'			
433	ALK2	5'	5'-agtgagtgaacggagccttg-3'	NM_176663	96	25.88 $\pm$ 0.22
434		3'	5'-ccatccaccattgtaaaacttaga-3'			
435	ALK3	5'	5'-acactgcccagatgatgcta-3'	NM_001076800	103	24.31 $\pm$ 0.30
436		3'	5'-ccctgaagctaattgtggttct-3'			
437	BMP4	5'	5'-gggcatcggtctgggagtat-3'	NM_001045877	90	26.70 $\pm$ 0.33
438		3'	5'-gggatgttctccagatgttctt-3'			
439	BMP7	5'	5'-ccaggtgttgcaagagcac-3'	NM_001206015	107	36.69 $\pm$ 0.68
440		3'	5'-tggtgtgatatcaaaaacgag-3'			

441	BMPR2	5'	5'-tcagagccctctcttgacct-3'	XM_002685492	106	29.10 ± 0.61
442		3'	5'-cagcaactggacgttcattctaa-3'			
443	FAS ³	5'	5'-cgggtgtggacatggtgac-3'	NM_001012669	64	21.33 ± 0.70
444		3'	5'-ccgaggcaggccatatagt-3'			
445	Hprt1	5'	5'-gtgattagcgatgatgaaccag-3'	NM_001034035	95	22.99 ± 0.20
446		3'	5'-ccatgaggaataaacaccttctc-3'			
447	β2-microglobulin	5'	5'-catgtccatgtttgaccttc-3'	NM_173893	70	19.13 ± 0.34
448		3'	5'-tcttccccacctctaagatgc-3'			
449	TGF-β1	5'	5'-cctgtgaggctcaagttaa-3'	NM_001166068	78	26.44 ± 0.59
450		3'	5'-aggtagcgccaggaattgt-3'			

451 ¹Mean ± SE of expression in the subcutaneous fat depot. ²BDL: below detection limit. ³For RT-qPCR analyses, cDNA
452 synthesized from 5 ng of total RNA was used as the template; for the other genes, cDNA corresponding to 10 ng of total
453 RNA was used.