TITLE:

Analysis of methane production pathways in a riparian wetland of a temperate forest catchment, using $\delta_{13}\text{C}$ of pore water CH_4 and CO_2

AUTHOR(S):

Itoh, Masayuki; Ohte, Nobuhito; Koba, Keisuke; Sugimoto, Atsuko; Tani, Makoto

CITATION:

Itoh, Masayuki ...[et al]. Analysis of methane production pathways in a riparian wetland of a temperate forest catchment, using $\delta_{13}\text{C}$ of pore water CH_4 and CO_2 . *Journal of Geophysical Research* 2008, 113(G3): G03005.

ISSUE DATE:

2008-07-12

URL:

<http://hdl.handle.net/2433/198846>

RIGHT:

Copyright 2008 by the American Geophysical Union.

Analysis of methane production pathways in a riparian wetland of a temperate forest catchment, using $\delta^{13}\text{C}$ of pore water CH_4 and CO_2

Masayuki Itoh,¹ Nobuhito Ohte,² Keisuke Koba,³ Atsuko Sugimoto,⁴ and Makoto Tani¹

Received 3 November 2007; revised 23 February 2008; accepted 7 May 2008; published 12 July 2008.

[1] To clarify how hydrological processes affect biogenic methane (CH_4) production and emission from soil surfaces, we analyzed the $\delta^{13}\text{C}$ of CH_4 and CO_2 and chemical constituents dissolved in groundwater at a wetland in the headwater catchment of a temperate forest in Japan. We estimated the contribution of acetate fermentation using the $\delta^{13}\text{C}$ isotope mass balance of dissolved CH_4 and CO_2 . CH_4 production pathways (e.g., acetate fermentation and carbonate reduction) changed temporally and spatially with hydrologically controlled redox conditions. The proportion of methanogenesis attributable to acetate fermentation usually decreased with temperature, suggesting that carbonate reduction dominated under conditions of high CO_2 concentration. In particular, the groundwater table and summer temperatures were key controlling factors in the interannual and intra-annual changes in CH_4 production pathways, controlling oxygen supply and consumption and, therefore, redox conditions in the soil. Under high temperature and high water table conditions during summer, the soil was strongly reduced and the proportion of carbonate reduction increased. Acetate fermentation also increased episodically, resulting in sporadic increases in $\delta^{13}\text{C}\text{-CH}_4$. The calculated acetate contribution obviously decreased in periods of low water table and high temperature when the soil surface was relatively oxic, implying deactivation of acetoclastic methanogenesis under oxic conditions. Thus, hydrological processes control the supply of these electron donors and acceptors and therefore play an important role in determining the relative proportions of CH_4 -producing pathways. Our results also indicate that an increase in acetate contribution under highly reducing conditions stimulates CH_4 production and emission from the soil surface.

Citation: Itoh, M., N. Ohte, K. Koba, A. Sugimoto, and M. Tani (2008), Analysis of methane production pathways in a riparian wetland of a temperate forest catchment, using $\delta^{13}\text{C}$ of pore water CH_4 and CO_2 , *J. Geophys. Res.*, 113, G03005, doi:10.1029/2007JG000647.

1. Introduction

[2] Methane (CH_4) is a key greenhouse gas, and its infrared radiative heating effect is 26 times greater than that of carbon dioxide on a mole-per-mole basis [Lelieveld and Crutzen, 1992]. Soil functions as both a main source and a main sink of CH_4 . In anoxic environments such as wetland soils, CH_4 is produced by methanogenic bacteria that are active only under anoxic and strongly reducing conditions [Takai, 1970; Schütz *et al.*, 1989]. In contrast, CH_4 is usually oxidized by methanotrophic bacteria in oxic soils. Over the last 150 years, the mixing ratio of CH_4 in the

atmosphere has more than doubled [Etheridge *et al.*, 1998], and improved estimates of the strength of each source and sink are of high importance. However, much uncertainty regarding the CH_4 production and consumption mechanisms in soils still leads to uncertainty in estimating the levels of atmospheric CH_4 .

[3] Because biospheric sources of CH_4 are highly variable, stable isotope ratios of CH_4 have been used to constrain the global CH_4 budget, as microbe-produced CH_4 has a significantly different isotopic signal than CH_4 from other sources [Whiticar, 1999]. In particular, the ^{13}C compositions of CH_4 in background tropospheric air and of the major CH_4 sources have added further constraint to the individual CH_4 source strengths by isotope mass balance using the $\delta^{13}\text{C}$ value of each source [Stevens and Rust, 1982; Cicerone and Oremland, 1988; Stevens and Engelkemeir, 1988; Whalen *et al.*, 1989; Quay *et al.*, 1991; Lowe *et al.*, 1994; Gupta *et al.*, 1996; Bräunlich *et al.*, 2001; Fletcher *et al.*, 2004] and have revealed that 70–80% of atmospheric CH_4 is of biogenic origin, with natural wetlands as the largest source [Bartlett and Harriss, 1993; Khalil and Shearer, 1993]. However, estimating the representative $\delta^{13}\text{C}\text{-CH}_4$ of each source

¹Laboratory of Forest Hydrology, Division of Environmental Science and Technology, Graduate School of Agriculture, Kyoto University, Kyoto, Japan.

²Department of Forest Science, Graduate School of Agricultural and Life Sciences, University of Tokyo, Tokyo, Japan.

³Institute of Symbiotic Science and Technology, Tokyo University of Agriculture and Technology, Tokyo, Japan.

⁴Division of Earth System Science, Faculty of Environmental Earth Science, Hokkaido University, Hokkaido, Japan.

Figure 1. (a) Location of KEW. (b) Topographic map of KEW and the locations of wetlands. Shading indicates riparian wetland areas. (c) Locations of the observation plots. Each number indicates an observation plot (1: KW2-edge, 2: KW2-center, 3: KW2-downstream).

remains challenging because $\delta^{13}\text{C}$ -CH₄ values are highly variable, especially in rice paddies and wetlands [Quay *et al.*, 1991], reflecting the multiple processes involved in CH₄ production and consumption in these ecosystems. For example, methanogenesis from carbonate results in a larger fractionation against ¹³C and, thus, lower $\delta^{13}\text{C}$ -CH₄ values than methanogenesis from acetate [Games *et al.*, 1978; Krzycki *et al.*, 1987; Gelwicks *et al.*, 1994]. However, previous studies have shown that the fractionation factors vary with site and conditions (e.g., reviewed by Conrad [2005]). More data under various environmental conditions are required for the explicit determination of fractionation factors.

[4] Among the various CH₄ sinks and sources, forests are assumed to be a major sink of atmospheric CH₄ by microbial oxidation in aerobic soils [Reeburgh *et al.*,

1993; IPCC, 2001]. However, Itoh *et al.* [2005, 2007] suggested that wet riparian areas in forests can function as ‘hot spots’ of CH₄ emission and that these ‘hot spots’ can also significantly affect the total budget of trace gas emissions on larger scales, such as whole forest ecosystems. Soil hydrological conditions in forest catchments are spatially variable. Riparian wetlands in small headwater catchments are characterized by high CH₄ production, and production and emission rates are strongly affected by changes in hydrological processes and temperature [Itoh *et al.*, 2007]. This hydrological variability can also affect CH₄ production pathways that mainly consist of acetate fermentation [Zeikus *et al.*, 1975] and carbonate reduction [Takai, 1970; Crill and Martens, 1986; Martens *et al.*, 1986; Burke *et al.*, 1988; Schütz *et al.*, 1989]. Redox conditions in such wetland soils

Table 1. Mean Annual Air Temperature and Annual and Summer Precipitation in the Kiryu Experimental Watershed^a

	2003	2004	2005	1996–2005
Mean annual air temperature (°C)	12.9	13.9	13.0	13.2 ^b
Annual precipitation (mm year ⁻¹)	1946.8	1796.8	1150.5	1574.1
Summer precipitation (mm) ^c	804.0	500.3	474.0	550.6

^aData from Itoh *et al.* [2007].

^bAverage of values obtained from 2000 to 2005.

^cFrom June to August.

change on a shorter temporal scale with hydrological conditions (precipitation patterns and water movement in soil [Mitsch and Gosselink, 2000; Itoh *et al.*, 2007]) than in ombrotrophic wetlands, where most previous work has been conducted. CH₄ production pathways can change drastically on small temporal and spatial scales in such riparian wetlands. Thus, for a more reliable estimate of the CH₄ budget on a larger scale, such as an entire forest catchment, an understanding of CH₄ production mechanisms, including production pathways, is strongly required.

[5] In this study, we used the $\delta^{13}\text{C}$ isotope mass balance of pore water CH₄ and CO₂ [Sugimoto and Wada, 1993] to determine temporal and spatial changes in CH₄ production pathways and to understand what processes control pathway changes in riparian wetlands. In addition, we considered the effects of climate differences, such as characteristic precipitation patterns in Asian monsoon climates, on CH₄ production pathways.

2. Materials and Methods

2.1. Site Description and Hydrological and Biogeochemical Features

[6] We studied forested wetlands in the Kiryu Experimental Watershed (KEW; 35°N, 136°E; 190–255 m above sea level; 5.99 ha), located in southeastern Shiga Prefecture, central Japan (Figure 1). The KEW comprises about 99.3% forest floor, with 0.67% (400.6 m²) distinct wetland riparian zones. The wetland studied here, Kiryu Wetland 2 (KW2), is located upstream of a check dam constructed across the mainstream of the watershed about 100 years ago to prevent soil erosion. There are other natural wetlands in KEW, but all are located in riparian zones along streams. The wetland soils are either always submerged or periodically submerged. The entire watershed is on a base of weathered granitic rock with an abundance of albite. In the 1960s, Japanese cypress (*Chamaecyparis obtusa*) was planted on the hillslope over the watershed. Although several tree species (*Eurya japonica*, *Alnus japonica*, *Clethra barvinervis*, *Evodiapanax innovans*, and *Rhus trichocarpa*) and sphagnum grow in and around the wetlands, vegetation in the wetlands is sparse, probably because of occasional sediment transport with surface flow.

[7] Precipitation was measured at a meteorological station within the watershed. Mean annual air temperature and precipitation were recorded during the observation period (Table 1). Surface soil temperatures at depths of 0.02 and 0.10 m were also continuously monitored (Figure 2a) at each groundwater-sampling plot. The water levels in KW2 were measured in wells installed at each groundwater-sampling plot. We also installed a capacitance water level sensor and data recorder (CR-10x, Campbell, USA) 0.50 m from the KW2-downstream plot.

[8] The study site is affected by the Asian monsoon system, which usually results in a summer rainy season. Variation was observed in summer (June, July, and August) precipitation; a large amount of precipitation was observed in 2003, and less in 2004 and 2005 (Table 1 and Figure 2b). Summer precipitation accounted for 146.0, 90.9, and 86.1% of the mean annual precipitation (1574.1 mm for 1996–2005) in 2003, 2004, and 2005, respectively.

[9] Strongly reducing conditions (e.g., activation in denitrification and manganese-, iron-, and sulfate-reductions) were formed in high temperature periods in all three sampling plots [Itoh *et al.*, 2007]. However, relatively dissolved oxygen (DO)-rich stream and subsurface flow from the hillslope due to heavy precipitation induced more oxic conditions than usual in the surface and bottom layers of wetlands in 2003. In contrast, a decrease in the water table formed oxic conditions in the surface soil layer in summer 2005. In these summers, the CH₄ concentration in the surface soil was much lower than in the summer of 2004, when the water table and soil temperature were high. These changes in redox conditions, depending on hydrologic conditions, strongly affect CH₄ production in the soil and CH₄ emission from the soil surface [Itoh *et al.*, 2007].

2.2. Groundwater Collection and Analysis

[10] Pore water samples were collected vertically to measure the concentration (of CH₄ and CO₂) and isotopic composition of $\delta^{13}\text{C}$ -CH₄ and $\delta^{13}\text{C}$ -CO₂ and water chemistry in each of three observation plots, KW2-edge (0.55 m soil depth), KW2-center (1.77 m), and KW2-downstream (1.38 m; Figure 1c). KW2-edge was near the hillslope and KW2-downstream was along a perennial stream (Figure 1c). The surface soils of all three sampling plots were silty with much undecomposed litter. We used double-walled pore water samplers [Itoh *et al.*, 2007] which collected pore water without degassing and high decompression. The pore water samplers were placed at KW2-edge (at soil depths of 0.10 and 0.25 m), KW2-center, and KW2-downstream (at 0.10, 0.20, 0.30, 0.50, and 0.70 m). Pore water samples were injected into 20-, 30-, or 50-mL pre-evacuated vials for the measurement of dissolved CH₄ and CO₂ concentrations and their carbon isotope ratios without exposure to the atmosphere, and into plastic bottles for other chemical analyses; the vials and plastic bottles were stored in a cooler (around 4°C) in the field. Surface water (KW2-edge and KW2-center) and stream water (KW2-downstream) were also sampled.

[11] In situ measurements, including pH and electrical conductivity (EC), and laboratory measurements of dissolved components including DO, CH₄, and CO₂ were conducted. Detailed information on the methods and water chemical constituent results were given by Itoh *et al.* [2007].

Figure 2. Seasonal variations in (a) soil temperature at KW2-downstream, (b) monthly precipitation measured at the nearby meteorological station, (c) groundwater CH₄ concentration and water table level, and (d) $\delta^{13}\text{C}$ of CH₄ in each sampling plot. The shaded areas indicate low temperature (<10°C) periods.

[12] Dissolved CH₄ and CO₂ concentrations were determined within 8 h of sampling by multiple equilibrations with a headspace of ultra high purity (UHP) helium [McAulliffe, 1971]. The headspace was prepared in a vial by replacing sample water with He (>99.999% purity). The vials were vigorously shaken for 2 min to drive gases from the pore water into the headspace. The headspace gas was withdrawn using a gas-tight syringe, and CH₄ concentration was determined using a gas chromatograph (GC; GC-14BPF, Shimadzu, Japan) equipped with a flame ionization detector (FID [Itoh et al., 2007]). For CO₂ concentration measurements, samples of the same gas were collected from the headspace and injected into a GC (GC-8APT, Shimadzu, Japan) equipped with a thermal conductivity detector (TCD [Itoh et al., 2007]). Dissolved CO₂ concentrations in water samples were also measured in situ from June 2003 to August 2005 using a portable pCO₂ meter (CGP-1, DKK-TOA, Japan [Ohte et al., 1995]). GC measurements of dissolved CO₂ concentrations were carried out from February 2005 and confirmed the data obtained by in situ dissolved pCO₂ measurements.

[13] Carbon isotopic compositions of dissolved CH₄ and CO₂ were analyzed using a gas chromatograph/combustion/

isotope ratio mass spectrometer (GCCMS) MAT 252 equipped with an HP G1530A system [Sugimoto, 1996] at the Center for Ecological Research at Kyoto University. Because of the detection limit, isotopic measurement could not be conducted on samples with low CH₄ or CO₂ concentrations, such as stream and surface water.

2.3. Soil Sampling and Analysis

[14] Mineral and organic soil samples were collected in triplicate in each plot in January 2006. Topsoil and 10–20-cm interval samples underlying the thin litter layer (0–1 cm thick) were collected from the surface to depths of 0.25 m (KW2-edge) or 1.00 m (KW2-center and KW2-downstream). The depth of organic soil was approximately 0.15, 0.40, and 0.50 m in KW2-edge, KW2-center, and KW2-downstream, respectively. Soils were sieved through a 2-mm mesh sieve to remove coarse fragments and then homogenized. The total C and total N concentrations of soil samples were measured using the combustion method [Bremner, 1996] in an NC-analyzer (Sumigraph NC-900, Sumigraph Co., Japan).

[15] $\delta^{13}\text{C}$ analyses were also carried out on soil samples. The samples were dried in an oven at 40°C for 48 h prior to

Table 2. C and N Content, C/N Ratio, and $\delta^{13}\text{C}$ of Litter and Soil at Each Sampling Depth

Plot	Depth (m)	C(%)	N(%)	C/N	$\delta^{13}\text{C}(\text{‰})$
Litter		46.4	1.05	45.5	-29.0
KW2-edge	0	5.89	0.226	25.4	-28.0
	0.10	5.72	0.221	26.3	-28.1
	0.25	3.14	0.131	25.0	-28.0
KW2-center	0	10.4	0.547	20.1	-29.0
	0.10	11.3	0.617	18.6	-29.3
	0.20	7.25	0.384	19.2	-29.0
	0.30	2.30	0.114	19.0	-28.5
	0.50	0.65	0.0316	20.4	-27.9
	0.70	1.29	0.0630	19.2	-28.2
KW2-downstream	0	1.73	0.0753	22.2	-28.6
	0.10	2.37	0.0967	24.6	-28.5
	0.20	5.53	0.229	24.3	-28.5
	0.30	6.30	0.280	22.5	-28.5
	0.50	4.27	0.220	19.6	-28.1
	0.70	2.06	0.100	20.7	-28.2

$\delta^{13}\text{C}$ analysis. $\delta^{13}\text{C}$ was determined using a mass spectrometer (Delta plus XP, Thermo Electron) coupled with an elemental analyzer (Flash EA, Thermo Electron).

3. Results

3.1. C and N Concentrations and $\delta^{13}\text{C}$ of Wetland Soils

[16] C and N contents, the C/N ratio, and $\delta^{13}\text{C}$ of litter and soil at each sampling depth are shown in Table 2. At KW2-edge and KW2-center, litter accumulates in the surface soils because of slow degradation under wet and anoxic conditions, resulting in higher soil C concentrations at the surface than in the bottom layer. At KW2-center, in particular, the upper soil layer contained much more organic matter than the bottom layer or the other plots. At KW2-downstream, soil C concentration was highest at 0.30 m. At all sampling plots, soil $\delta^{13}\text{C}$ values ranged between -29.8 and -28.3‰, and there was no obvious trend in $\delta^{13}\text{C}$ with depth or plot (Table 2).

3.2. CH₄ and CO₂ Concentrations and Carbon Isotopic Composition

[17] CH₄ and CO₂ concentrations in groundwater increased with temperature at all three plots (Figures 2c, 3c,

Figure 3. Seasonal variations in (a) soil temperature at KW2-downstream, (b) monthly precipitation measured at the nearby meteorological station, (c) groundwater CO₂ concentration and water table level, and (d) $\delta^{13}\text{C}$ of CO₂ in each sampling plot. The shaded areas indicate low temperature (<10°C) periods.

Table 3. Mean, Minimum, and Maximum $\delta^{13}\text{CH}_4$ and $\delta^{13}\text{CO}_2$ ^a

Plot	Depth (m)	Mean CH ₄			$\delta^{13}\text{C-CH}_4$ (‰)			Mean CO ₂			$\delta^{13}\text{C-CO}_2$ (‰)			n
		Concentration (mmol/L)	n		Mean	Min	Max	Concentration (mmol/L)	n		Mean	Min	Max	
KW2-edge	0	0.01	32		-55.4	-65.2	-49.3	0.17	29		-19.7	-22.7	-15.4	17
	0.10	0.69	36		-60.3	-67.1	-51.6	0.91	35		-11.7	-20.8	-4.1	33
	0.25	0.42	45		-60.8	-65.2	-53.9	1.24	44		-17.1	-23.8	-8.4	42
KW2-center	0	0.03	15		-61.7	-66.9	-54.6	0.21	17		-19.7	-22.5	-16.5	13
	0.10	0.42	19		-63.7	-69.0	-54.8	0.94	17		-17.7	-24.4	-11.7	18
	0.20	0.69	27		-62.8	-83.0	-41.9	1.72	27		-13.8	-21.6	-5.6	26
	0.30	0.86	27		-64.1	-85.0	-56.0	1.99	27		-12.2	-18.8	-5.3	26
	0.50	0.58	27		-62.9	-72.4	-58.7	1.48	27		-15.5	-19.8	-12.1	26
	0.70	0.34	27		-64.0	-89.1	-55.7	1.16	26		-20.3	-22.8	-17.3	24
KW2-downstream	0	5.3E-04	46		-55.4	-58.8	-51.8	0.05	44		-18.6	-21.7	-16.3	10
	0.10	0.33	46		-60.4	-70.3	-50.8	0.64	44		-17.3	-26.9	-5.9	36
	0.20	0.37	46		-61.0	-72.3	-49.5	0.66	43		-17.3	-27.2	-5.0	39
	0.30	0.42	46		-60.6	-66.7	-52.5	0.73	44		-15.9	-24.5	-4.0	41
	0.50	0.36	43		-60.7	-66.8	-50.8	0.62	41		-16.6	-24.6	-3.6	39
	0.70	0.29	43		-60.8	-66.3	-48.4	0.60	39		-16.8	-24.1	-4.5	40

^aData from June 2003 to June 2006 at KW2-edge and KW2-downstream, and from August 2004 to June 2006 at KW2-center.

and 5). The $\delta^{13}\text{C}$ of pore water CH₄ usually increased from summer to autumn at KW2-edge and KW2-downstream. At KW2-center, variation was relatively small in summer (Figure 2d). CH₄ production generally increased with temperature; however, Figures 2a and 2c suggest a time lag between temperature change and methanogenic activity (approximately 1–2 months).

[18] At KW2-edge, CH₄ was higher at 0.10 m than at 0.25 m, except in summer 2005 when the water table dropped (Figure 2c), suggesting that the most reducing conditions were usually at 0.10 m [Itoh *et al.*, 2007]. CO₂ in groundwater increased in high temperature periods and was highest in summer 2005, when the surface soil was drier than usual (Figure 3c). $\delta^{13}\text{C-CH}_4$ values at 0.10 m were slightly higher than those at 0.25 m in the hot and wet summer of 2004 and lower in low temperature periods (Figure 2d). In this plot, the maximum $\delta^{13}\text{C-CO}_2$ value was -4.1‰ (Table 3). Rapid increases in CH₄ and CO₂ concentrations, $\delta^{13}\text{C-CH}_4$, and $\delta^{13}\text{C-CO}_2$ in groundwater were observed at the beginning of summer (Figures 2 and 3). This indicates that microbial activity increased rapidly with temperature in this plot.

[19] At KW2-center, CH₄ in groundwater was highest at 0.30 m through the sampling period (Figure 2c and Table 3) and maintained a high level even in low temperature periods (<10°C), when it decreased dramatically in other plots (Figure 2c). In this plot, mean and maximum CO₂ in groundwater were higher than those in the other plots. CO₂ was also highest at 0.30 m, where the soil C concentration was highest (Table 2 and Figure 3c). The range of $\delta^{13}\text{C-CH}_4$ values was much larger than in any other plot (Figure 2d and Table 3). That is, we observed lower $\delta^{13}\text{C-CH}_4$ values in low temperature periods (e.g., -83.0‰ at 0.30 m depth, 9 March 2005; -85.0‰ at 0.50 m and -89.1‰ at 0.70 m, 9 February 2005; Figure 2d and Table 3) and higher values in high temperature periods (e.g., -41.9‰ at 0.20 m, 10 September 2004), relative to other plots. $\delta^{13}\text{C-CH}_4$ was highest at 0.20 m in summer 2004, and no clear vertical trend was seen in other summers (Figure 2d). A temporal increase in $\delta^{13}\text{C-CO}_2$ was seen in the high tem-

perature period, but this trend was unclear in the other two plots. Comparisons of the vertical distributions of CH₄, CO₂, and $\delta^{13}\text{C-CO}_2$ between the summers of 2004 and 2005 (Figures 2c, 3c, and 3d) all showed convex distributions, peaking at 0.30 m in both summers.

[20] At KW2-downstream, no clear vertical distribution pattern in CH₄ concentration was observed below 0.20 m. Seasonal (low in late winter and high in late summer) and yearly variations in CH₄ were clearer than in the other two plots. Maximum groundwater CH₄ and CO₂ in this plot were much lower in 2003 than in 2004 and 2005, especially at 0.10 m, indicating less microbial activity at low temperatures and high levels of DO in the rainy summer of 2003 (Figures 2c and 3c [Itoh *et al.*, 2007]). Maximum $\delta^{13}\text{C-CH}_4$ (Figure 2d) and $\delta^{13}\text{C-CO}_2$ (Figure 3d) were also much lower in 2003, especially at 0.10 m. For example, maximum $\delta^{13}\text{C-CO}_2$ was much lower in 2003 (-10.9‰ at 0.50 m, 18 September 2003) than in 2004 (-4.0‰ at 0.30 m, 21 August 2004) and 2005 (-3.6‰ at 0.50 m, 15 September 2005). $\delta^{13}\text{C-CO}_2$ in groundwater showed clear seasonal changes, ranging from -27 to -4‰, and increased dramatically in high temperature periods (Figure 3d). $\delta^{13}\text{C-CH}_4$ showed a similar seasonal change, though not as obvious as $\delta^{13}\text{C-CO}_2$ (Figure 2d). In this plot, no obvious vertical difference in $\delta^{13}\text{C-CH}_4$ was observed (Figure 2d). $\delta^{13}\text{C-CH}_4$ increased dramatically in the summers of 2004 (at all depths) and 2005 (especially at 0.30 m).

4. Discussion

4.1. Effects of Temperature and Hydrology on Changes in CH₄ and CO₂ Dynamics

[21] The measured soil $\delta^{13}\text{C}$ at KW2 ranged from -27.9 to -29.3‰ (Table 2). The measured $\delta^{13}\text{C}$ of leaves ranges from -24.6 to -28.9‰ in cypress [Matsuo, 2003], which is the major vegetation in KEW, and from -33.3 to -32.5‰ in forest floor vegetation (*Eurya japonica*, N. Matsuo, personal communication). CO₂ is produced as a by-product of respiration under both oxic and anoxic conditions in soils. CO₂ produced from the metabolism of lactate is

Figure 4. Relationships between CH₄ concentration and $\delta^{13}\text{C}\text{-CH}_4$ and $\delta^{13}\text{C}\text{-CO}_2$ at (a) KW2-edge, (b) KW2-center, and (c) KW2-downstream.

depleted in ^{13}C by as much as 26‰ [Smejkal *et al.*, 1971]. Also, CO₂ respired during heterotrophic microbial metabolism is 3.4‰ depleted in ^{13}C relative to the glucose used as the carbon source [Blair *et al.*, 1985]. Negative isotopic enrichment occurs during the decomposition process, whatever the substrate [Mary *et al.*, 1992]. The $\delta^{13}\text{C}$ of CO₂ respired by decomposing litter and soil organic matter in KEW is expected to be lower than the $\delta^{13}\text{C}$ of soil organic matter ($= -29\text{‰}$ in surface soil at KW2; Table 2), which is similar to the ^{13}C of cypress.

[22] However, we observed much heavier $\delta^{13}\text{C}\text{-CO}_2$ values in groundwater than in soil, litter, and vegetation. At KW2-edge and KW2-downstream, $\delta^{13}\text{C}\text{-CO}_2$ dramatically increased in the summer when dissolved CH₄ concentrations were high (Figures 2c and 3d). At KW2-center, $\delta^{13}\text{C}\text{-CO}_2$ was also much higher than soil $\delta^{13}\text{C}$ (Table 3). Processes that might induce such an increase in $\delta^{13}\text{C}\text{-CO}_2$ are (1) preferential use of $^{12}\text{CO}_2$ as a substrate for CH₄ by carbonate reduction or (2) ^{13}C -enriched CO₂ as a by-product of methanogenesis via acetate fermentation under anoxic conditions [Sugimoto and Wada, 1993]. CO₂ produced in this manner, i.e., by acetate fermentation, is strongly enriched in ^{13}C and may have a $\delta^{13}\text{C}$ value as high as -5‰ [Charman *et al.*, 1994] or $+9\text{‰}$ [Waldron *et al.*, 1999].

[23] The significant positive regression of CH₄ concentration on $\delta^{13}\text{C}\text{-CO}_2$ (Figures 4 and 5 and Table 4) in all

plots suggests that much of the CH₄ produced originates from carbonate reduction. Heavy $\delta^{13}\text{C}\text{-CO}_2$ from acetate fermentation may also contribute to CO₂ enrichment. The fact that the CO₂ concentration does not decrease in this zone indicates that CO₂ is continuously added by organic remineralization to the pool in the methanogenic zone, although ^{13}C enrichment by methanogenesis does occur. Here, the increase in the proportion of carbonate reduction with temperature is confirmed by an increase in the apparent α in high temperature periods (Figures 6, 7, and 8; discussed below). These results support the idea that carbonate reduction becomes dominant in high temperature periods. As for carbonate (CO₂/H₂) reduction, although we have no data on hydrogen concentrations in groundwater, hydrogen concentration can increase with soil temperature by stimulating litter decomposition [Sugimoto and Fujita, 2006]. A less obvious increase in $\delta^{13}\text{C}\text{-CO}_2$ in the rainy summer of 2003 at KW2-downstream (Figure 3d) was attributed to lower CH₄ production than in a typical summer because of the DO supplied by increased water flow [Itoh *et al.*, 2007]. These results show that preferential use of $^{12}\text{CO}_2$ for methanogenesis can drastically change $\delta^{13}\text{C}\text{-CO}_2$. Although the coefficients were not statistically significant at the deep zone of KW2-center, where CH₄ concentration was high throughout the year (Table 4), our results suggest that CO₂ contributed largely to CH₄ production in our site and residual CO₂ was enriched with ^{13}C during this process.

Figure 5. Relationships between soil temperature and CH₄ concentration and $\delta^{13}\text{C}\text{-CH}_4$ at (a) KW2-edge, (b) KW2-center, and (c) KW2-downstream.

In addition, changes in both temperature and hydrologic conditions (rainfall and the water table) played important roles in the variations in CH₄ and CO₂ concentrations and $\delta^{13}\text{C}\text{-CO}_2$. However, acetate fermentation must affect CH₄ production, and we consider the effects of acetate contributions to methanogenesis in the next section.

4.2. Acetate Contribution Determined by Isotope Mass Balance

[24] Similar seasonal variations in $\delta^{13}\text{C}$ of CO₂ and CH₄ indicated that CO₂ contributes much as a substrate for CH₄ production. However, $\delta^{13}\text{C}\text{-CH}_4$ varied seasonally and fluctuated more wildly than $\delta^{13}\text{C}\text{-CO}_2$ at KW2-downstream

(Figures 2d and 3d). Remarkably high $\delta^{13}\text{C}\text{-CH}_4$ values were occasionally observed in summer (e.g., KW2-downstream: -48.4‰ on 28 June 2004 at 0.70 m; -50.8‰ on 10 September 2004 at 0.10 m; -49.5‰ on 8 November 2004 at 0.20 m; and -52.5‰ on 15 September 2005 at 0.30 m; Figure 2d) when high CH₄ production occurred. Also at KW2-edge, heavy $\delta^{13}\text{C}\text{-CH}_4$ values were observed in summer (i.e., -51.6‰ at 0.10 m on 7 October 2004 and -57.3‰ at 0.25 m on 12 October 2005). The crossplots of $\delta^{13}\text{C}\text{-CH}_4$ and $\delta^{13}\text{C}\text{-CO}_2$ are shown in Figures 6a–6c, with the apparent fractionation between CO₂ and CH₄ (α) calculated by the

Table 4. Results of Regression Analyses Between CH₄ Concentration and $\delta^{13}\text{CH}_4$ or $\delta^{13}\text{CO}_2$ at Each Sampling Depth^a

Plot	Depth (m)	$\delta^{13}\text{CH}_4$				$\delta^{13}\text{CO}_2$			
		<i>n</i>	<i>F</i>	<i>p</i>	<i>r</i> ²	<i>n</i>	<i>F</i>	<i>p</i>	<i>r</i> ²
KW2-edge	0.10	32	21.7	<0.001	0.41	33	22.7	<0.001	0.41
	0.25	42	5.67	<0.05	0.12	41	75.3	<0.001	0.65
KW2-center	0.10	18	4.81	<0.05	0.22	18	46.1	<0.001	0.73
	0.20	26	2.35	0.137	0.09	26	35.6	<0.001	0.59
	0.30	26	2.13	0.157	0.08	26	18.8	<0.01	0.43
	0.50	26	2.91	0.100	0.10	26	0.164	0.689	0.01
	0.70	24	0.479	0.495	0.02	24	2.06	0.164	0.08
KW2-downstream	0.10	33	21.7	<0.001	0.40	36	352	<0.001	0.91
	0.20	39	25.3	<0.001	0.40	38	205	<0.001	0.85
	0.30	42	60.9	<0.001	0.60	41	320	<0.001	0.89
	0.50	39	19.5	<0.001	0.34	39	229	<0.001	0.86
	0.70	39	27.5	<0.001	0.42	40	191	<0.001	0.83

^aValues of *F* and *p* are observed variance ratios and probabilities in the analyses, respectively.

Figure 6. Crossplots of δ¹³C data from CH₄ and CO₂ in groundwater at (a) KW2-edge, (b) KW2-center, and (c) KW2-downstream. The dashed lines give the apparent fractionation between CO₂ and CH₄ (α) calculated by the ratio of (δ¹³C-CO₂ + 10³)/(δ¹³C-CH₄ + 10³).

ratio (δ¹³C-CO₂ + 1000)/(δ¹³C-CH₄ + 1000). From this, the apparent α was widely distributed from 1.032 to 1.083 and was usually between 1.04 and 1.06. This range is lower than reported values for ¹³C fractionation during CO₂ reduction to CH₄ (α_c = 1.06–1.07) according to a review of experimentally determined α_c of natural wetlands [Conrad, 2005]. Here,

$$\alpha_c = \frac{(\delta^{13}\text{C-CO}_2 + 1000)}{(\delta^{13}\text{C-CH}_4 + 1000)} \quad (1)$$

where δ¹³C-CO₂ is the δ¹³C value of the CO₂ pool at the sampling time and δ¹³C-CH_{4(CO2)} is the δ¹³C value of CH₄ from CO₂.

[25] Our results indicate that acetate fermentation, which has a smaller isotopic fractionation than carbonate reduction, affected the change in apparent α in our wetland. Here, we calculate the acetate contribution (F_{Ac}) to CH₄ production by means of isotope mass balance [Sugimoto and Wada, 1993]. In this calculation, only two substrates, CO₂ and acetate, are considered for methanogenesis, because CH₄ from other substrates such as methanol and trimethylamine has so far not been found to play a major role in freshwater environments [Lovley and Klug, 1983b; Conrad and Claus, 2005]. The fractional Ac contribution is expressed by F_{Ac} as:

$$F_{Ac} = \frac{\text{CH}_4 \text{ from acetate}}{\text{CH}_4 \text{ from CO}_2 + \text{CH}_4 \text{ from acetate}} \quad (2)$$

[26] The isotope mass balance for produced CH₄ is expressed by the following equation:

$$\delta^{13}\text{C-CH}_4(\text{Ac})F_{Ac} + \delta^{13}\text{C-CH}_4(\text{CO}_2)(1 - F_{Ac}) = \delta^{13}\text{C-CH}_4 \quad (3)$$

where δ¹³C-CH_{4(Ac)} and δ¹³C-CH_{4(CO2)} are the δ¹³C values of CH₄ produced from acetate and CO₂, respectively, and δ¹³C-CH₄ is that of CH₄ produced during the indicated period. When the δ¹³C values of CH₄, δ¹³C-CH_{4(Ac)}, and δ¹³C-CH_{4(CO2)} are obtained, F_{Ac} can be calculated from this equation. δ¹³C-CH_{4(CO2)} was calculated using equation (1).

[27] Because we have no data on δ¹³C of acetate and CH₄ from acetate, we used possible δ¹³C values of CH₄ from acetate ranging from −44 to −27‰, considering the values obtained from previous studies [e.g., Sugimoto and Wada, 1993; Avery et al., 1999; Nakagawa et al., 2002], and δ¹³C data from soil collected from KW2 (Table 2). During acetate fermentation, CH₄ is produced primarily from the methyl carbon of acetate [Pine and Barker, 1956; Krzycki et al., 1982]. Sugimoto and Wada [1993] incubated Japanese rice paddy field soil (δ¹³C = −26.5‰) with BES (a methanogenesis inhibitor) and measured the δ¹³C of both methyl carbon and carboxyl carbon of acetate. The δ¹³C of methyl carbon ranged from −36 to −30‰, which was lower than that of carboxyl carbon (−21 to −15‰). From this and the δ¹³C of our wetland soils (approximately −29‰), we assume δ¹³C-CH_{4(Ac)} = −35‰. We must also assume values for the ¹³C fractionation coefficient during CO₂ reduction to CH₄ (α_c). When assuming the α_c value, the temperature dependence of α_c [e.g., Whiticar et al., 1986; Whiticar, 1999; Conrad, 2005] is considered. Blair et al. [1993] found that α_c in marine sediment decreased with increasing temperature according to:

$$\ln \alpha_c = (23.0/T) - 0.022 \quad (4)$$

[28] A similar relationship was found for methanogenesis cultures [Botz et al., 1996], $\ln \alpha_c = (29/T) - 0.030$. We used the slope of equation (4) with the soil temperature range of KEW (0°C to 23°C at a depth of 0.10 m at KW2-downstream). As a first approximation, we assumed $\ln \alpha_c = (23/T) - 0.022$ according to the value of 1.06–1.07 based on a review of

Figure 7. Seasonal variation in (a) soil temperature and (b) F_{Ac} (acetate contribution to methanogenesis) in each sampling plot. The shaded areas indicate low temperature (<10°C) periods.

Figure 8. Relationships between soil temperature and F_{Ac} at (a) KW2-edge, (b) KW2-center, and (c) KW2-downstream.

experimentally determined α_c of natural wetlands [Conrad, 2005], resulting in a range of 1.068 at 23°C to 1.075 at 0°C (α_{c1}) for α_c . Next, if the F_{Ac} result was negative, we assumed a larger α_c value in order to obtain positive F_{Ac} results (α_{c2}).

[29] Figure 7b shows the seasonal variation of F_{Ac} calculated from the above equations, assuming α_{c1} and $\delta^{13}\text{CH}_4(\text{Ac}) = -35\text{‰}$ for KW2-edge and KW2-downstream. Table 5 shows the possible ranges of F_{Ac} with varying parameters. For KW2-center, however, calculated F_{Ac} values using α_{c1} in winter 2004 were negative for depths of 0.20 and 0.30 m regardless of the $\delta^{13}\text{CH}_4(\text{Ac})$ value (Table 5). Therefore, we assumed α_{c2} [$\ln \alpha_{c2} = (23/T) - 0.0015$], resulting in a range of 1.079 at 23°C to 1.086 at 0°C, and used these values to calculate Ac for the three plots (Figure 7b and Table 5).

[30] Larger F_{Ac} values result from larger α_c (at most 14.5% given the same $\delta^{13}\text{CH}_4(\text{Ac})$) and $\delta^{13}\text{CH}_4(\text{Ac})$ (at most 18.5% given the same α_c). Also, smaller α_c and larger $\delta^{13}\text{CH}_4(\text{Ac})$ values result in wider ranges of F_{Ac} (Figure 7b and Table 5). This indicates that the setting of these values changes the result significantly. However, the seasonal trend of F_{Ac} was characteristic in each plot and did not change with changes to the parameters. As mentioned above, for purposes of discussion, here we used the results based on an appropriate $\delta^{13}\text{CH}_4(\text{Ac})$ value for KEW of -35‰ [Sugimoto and Wada, 1993]. In addition, some attention should be paid to the expected systematic change in $\delta^{13}\text{CH}_4(\text{Ac})$ with temperature, although the degree of change is unknown because this has not been studied in detail. The range of possible $\delta^{13}\text{CH}_4(\text{Ac})$ is most

Table 5. Mean, Minimum, and Maximum Ac Contributions Calculated Using Equation (3)^a

			Ac Contribution (%) $\delta^{13}\text{CH}_4$ (Ac) = -27	Ac Contribution (%) $\delta^{13}\text{CH}_4$ (Ac) = -35	Ac Contribution (%) $\delta^{13}\text{CH}_4$ (Ac) = -44
Plot	Depth (m)	α_c	Mean (Min. – Max.)	Mean (Min. – Max.)	Mean (Min. – Max.)
Edge	0.10	α_{c1}	33.1 (22.3 – 43.1)	39.5 (26.5 – 52.9)	50.6 (33.6 – 71.1)
	0.25	α_{c1}	38.2 (22.5 – 51.9)	44.7 (27.2 – 59.8)	55.4 (35.5 – 72.1)
	0.10	α_{c2}	44.1 (34.8 – 53.6)	51.0 (40.1 – 63.1)	62.0 (48.5 – 78.9)
	0.25	α_{c2}	47.5 (35.8 – 58.4)	54.2 (41.8 – 65.9)	64.6 (51.3 – 77.1)
Center	0.10	α_{c1}	34.6 (24.7 – 53.4)	40.3 (28.9 – 61.7)	49.6 (35.6 – 74.7)
	0.20	α_{c1}	30.9 (6.8 – 66.9)	36.6 (8.1 – 81.3)	46.2 (10.3 – 107)
	0.30	α_{c1}	26.4 (20.5 – 37.7)	31.3 (24.7 – 45.4)	39.8 (32.3 – 59.1)
	0.50	α_{c1}	32.9 (16.2 – 42.3)	38.7 (19.0 – 48.9)	48.2 (23.6 – 59.2)
	0.70	α_{c1}	38.3 (27.2 – 50.6)	44.4 (31.3 – 58.7)	54.1 (37.7 – 71.5)
	0.10	α_{c2}	44.2 (35.8 – 59.9)	50.3 (40.8 – 67.7)	59.3 (49.0 – 75.1)
	0.20	α_{c2}	41.9 (9.8 – 72.9)	48.2 (11.2 – 85.3)	58.0 (13.5 – 105)
	0.30	α_{c2}	38.3 (0.3 – 48.3)	44.2 (0.4 – 56.3)	53.4 (0.5 – 69.1)
	0.50	α_{c2}	43.1 (28.8 – 50.3)	49.4 (32.9 – 56.9)	59.0 (39.3 – 67.2)
	0.70	α_{c2}	45.3 (6.2 – 57.6)	51.4 (7.1 – 65.3)	60.5 (8.4 – 76.9)
Downstream	0.10	α_{c1}	31.9 (14.9 – 43.6)	38.0 (18.6 – 53.8)	48.6 (25.9 – 72.9)
	0.20	α_{c1}	37.7 (22.2 – 52.0)	44.1 (27.5 – 62.6)	54.8 (33.7 – 81.5)
	0.30	α_{c1}	37.3 (21.5 – 51.7)	43.8 (26.8 – 59.8)	54.8 (37.0 – 72.5)
	0.50	α_{c1}	36.7 (16.2 – 57.3)	43.0 (20.0 – 65.9)	53.4 (27.3 – 79.1)
	0.70	α_{c1}	37.8 (22.1 – 57.0)	44.3 (26.4 – 67.9)	55.1 (33.9 – 86.6)
	0.10	α_{c2}	46.4 (35.2 – 56.5)	53.2 (41.5 – 66.1)	60.9 (42.4 – 76.4)
	0.20	α_{c2}	47.1 (34.5 – 60.2)	53.8 (39.1 – 70.1)	64.2 (45.8 – 86.0)
	0.30	α_{c2}	47.0 (36.7 – 58.4)	53.8 (43.6 – 66.1)	64.5 (55.4 – 77.6)
	0.50	α_{c2}	46.4 (32.0 – 63.0)	53.1 (38.0 – 71.0)	63.4 (47.9 – 82.8)
	0.70	α_{c2}	46.2 (35.6 – 64.2)	53.0 (41.2 – 74.1)	63.5 (49.9 – 89.7)

^aData obtained from June 2003 to June 2006 at KW2-edge and KW2-downstream, and from August 2004 to June 2006 at KW2-center.

Figure 9. Crossplot of $\delta^{13}\text{C}$ data from CH_4 and CO_2 in groundwater at KW2-downstream. The dashed lines give the apparent fractionation between CO_2 and CH_4 . Arrows indicate temporal change in $\delta^{13}\text{C}-\text{CH}_4$ and $\delta^{13}\text{C}-\text{CO}_2$ during an incubation experiment using rice paddy soil at different temperatures [Fey *et al.*, 2004]. Arrow 1 is from the results of experiments conducted under the condition that CO_2 reduction solely contributed to methanogenesis. Arrow 2 is provided under the condition that acetate fermentation was the dominant form of methanogenesis. Figure 9b shows the data at KW2-downstream in summer 2004 labeled by sampling month.

likely within the ranges reported in Table 5. Therefore, in our discussion, this effect would be covered by the ranges used for calculating F_{Ac} with various parameters.

[31] Mean acetate contributions during sampling periods varied between 30 and 44% with α_{c1} and 44 and 54% with α_{c2} (Figure 7b and Table 5). As mentioned above, the calculated F_{Ac} at KW2-center fell below 0% in winter 2004/2005 with α_{c1} , indicating that α_{c2} was more appropriate for this plot. At KW2-edge, F_{Ac} increased in late summer 2004 and decreased from winter to summer 2005 (Figure 7b). At KW2-center, F_{Ac} was usually stable except in winter 2004/2005 ($\approx 0\%$, Figure 7b). At KW2-downstream, the seasonal trend was a lower F_{Ac} in high temperature periods (40.3% with α_{c1} and 52.7% with α_{c2} , averaged from 0.10 m to 0.70 m) and higher F_{Ac} in low temperature periods (45.3% with α_{c1} and 54.4% with α_{c2}); however, F_{Ac} showed episodic increases especially in summer 2004 when the

most reducing conditions were established (Figure 9b) [Itoh *et al.*, 2007]. These episodic and short-lived increases suggest that acetate is easily exhausted, probably due to its availability. This episodic increase in F_{Ac} was supported by various results. First, incubation experiments showed that active consumption of added acetate induces a rapid increase in $\delta^{13}\text{C}-\text{CH}_4$ [Sugimoto and Wada, 1993]; second, field observations showed rapid decreases in acetate with increases in temperature [Shannon and White, 1996; Avery *et al.*, 1999]. Third, pore water acetate concentrations increased during transition phases between acetate consumption by SO_4^{2-} reducers and methanogens [Alperin *et al.*, 1992; Sugimoto and Wada, 1993], suggesting a high availability of acetate. On the other hand, F_{Ac} clearly decreased at KW2-edge and downstream in the dry summer of 2005.

[32] Figure 9a shows a crossplot of $\delta^{13}\text{C}-\text{CH}_4$ and $\delta^{13}\text{C}-\text{CO}_2$, with arrows indicating temporal changes in $\delta^{13}\text{C}-\text{CH}_4$ and $\delta^{13}\text{C}-\text{CO}_2$, from an incubation experiment of rice paddy soil at different temperatures [Fey *et al.*, 2004]. Using ^{14}C -labeled bicarbonate, the incubation experiment found CH_4 production to be solely the result of CO_2 reduction at high temperature (50°C) [Fey and Conrad, 2000; Fey *et al.*, 2001]. Arrow 1 is provided from the results of experiments conducted under the same conditions, suggesting that $\alpha_{\text{c}} = 1.073$ [Fey *et al.*, 2004]. The results of Fey *et al.* [2004] also suggest that acetate with heavy $\delta^{13}\text{C}$ initially increased and then decreased with increases in CH_4 concentration and $\delta^{13}\text{C}-\text{CH}_4$ at temperatures of 10–37°C, hence arrow 2. This suggests that when acetate fermentation is dominant, only $\delta^{13}\text{C}-\text{CH}_4$ becomes heavier. Chan *et al.* [2005] also showed larger α_{c} , i.e., 1.0864–1.0885 and 1.0811–1.0892, for lake sediment cores sampled in May and August, respectively, by using an inhibitor of acetotrophic methanogenesis (CH_3F). The α_{c} values shown by Chan *et al.* [2005] are much larger than the apparent α in the present study. Figure 9b shows the summer 2004 data labeled by sampling month. In summer 2004, when CH_4 production dramatically increased, both $\delta^{13}\text{C}-\text{CO}_2$ and $\delta^{13}\text{C}-\text{CH}_4$ increased simultaneously in early summer, followed by a sharp increase only in $\delta^{13}\text{C}-\text{CH}_4$. According to the results of Fey *et al.* [2004], the simultaneous increase in $\delta^{13}\text{C}-\text{CO}_2$ and $\delta^{13}\text{C}-\text{CH}_4$ indicates that CO_2 reduction usually is dominant in periods of high temperature, and the contribution from acetate fermentation is variable. This is shown in the linear relationship between $\delta^{13}\text{C}-\text{CH}_4$ and $\delta^{13}\text{C}-\text{CO}_2$, composed of arrows 1 and 2 in Figure 9a. This increase in the proportion of carbonate reduction is also suggested by the increase in apparent α in this season. The outliers (high $\delta^{13}\text{C}-\text{CH}_4$) from this linear relationship suggest the sporadic and short-lived increase in acetate fermentation (shown as arrow 2 in Figure 9a). It is unlikely that increases in $\delta^{13}\text{C}-\text{CH}_4$ due to re-oxidation of CH_4 [Roslev and King, 1994, 1996] occurred in this period under low DO conditions [Itoh *et al.*, 2007]. Furthermore, only two of nine values with $\delta^{13}\text{C}-\text{CH}_4$ above -55‰ are from 0.10 m (CH_4 oxidation may be expected to occur because oxidation is usually observed from the soil surface downward), while during the same period, high CH_4 concentrations were measured up to that depth. Thus, our results suggest that the unusually high $\delta^{13}\text{C}-\text{CH}_4$ values observed in high temperature periods were due to increases in acetate fermentation under high soil temperature (Figures 8b and 8c). This implies that acetate

fermentation requires more time to be activated than does CO₂ reduction, in agreement with *Vogels et al.* [1988], who showed that doubling times for acetoclastic methanogens were longer than those for CO₂-reducing methanogens.

[33] In addition to the effect of increased temperature, episodic increases in F_{Ac} were also observed in low temperature periods (e.g., KW2-downstream, 0.50 m: -53.3% , 9 March 2006), implying that there is another factor influencing F_{Ac} , such as supply of acetate or competition for acetate with another acetate utilizer, e.g., sulfate-reducing bacteria (SRB) that are active under almost the same reducing conditions as methanogenesis. SRB may compete for acetate with acetoclastic methanogens under mildly reducing conditions (with some SO_4^{2-} [e.g., *Winfrey and Zeikus*, 1977; *Schönheit et al.*, 1982]). This phenomenon is usually observed in marine sediments, but can also occur in freshwater sediments; SRB can only outcompete methanogenesis when sulfate concentrations are increased to 60 μM [*Lovley and Klug*, 1983a]. In our wetland, the SO_4^{2-} concentration was usually low in summer at all plots (≈ 0 [Itoh et al., 2007]). In particular, SO_4^{2-} was lowest at all plots in the hot and wet season of 2004. This agrees with the increase in CH₄ production. Also, SO_4^{2-} was higher in rainy (2003; $\approx 50 \mu\text{M}$ at 0.10 m) and dry (2005 at 0.50 m) seasons; under such conditions, the activity of SRB may be higher, thereby reducing acetate fermentation. Under highly reducing conditions, SRB activity becomes limited by a decrease in available sulfate; thus, acetate may become available for methanogenesis under highly reducing summer conditions.

[34] F_{Ac} decreased more in the dry summer of 2005 than in 2004 at KW2-downstream (Figure 7b; summer to fall 2005), indicating that acetate fermentation may be inhibited by a decrease in acetate supply with low water transport in a low precipitation summer. In fact, the water table in the entire KW2 area was much lower in summer 2005 than in 2004 (Figure 2c).

[35] The relationship between $\delta^{13}\text{C}\text{-CH}_4$ and $\delta^{13}\text{C}\text{-CO}_2$ was unclear at KW2-center (Figure 6b), where strong reducing conditions were maintained during the year. A low level of water exchange with streams indicated by a small hydraulic head [Itoh et al., 2007] and high soil C concentration (Table 2) in this plot may affect the stable contribution of acetate fermentation.

[36] *Conrad et al.* [1989] and *Chin and Conrad* [1995] showed that the contribution of carbonate reduction decreased when temperature decreased (from 30°C to 10°C) and that CH₄ is mainly produced from acetate. Our results indicating a higher acetate contribution in periods of low temperature agree with their report. However, much lower levels of $\delta^{13}\text{C}\text{-CH}_4$ (and therefore, calculated F_{Ac}) in deeper soil were observed only in the mild winter of 2004/2005 (Figure 2d). The reported range of $\delta^{13}\text{C}\text{-CH}_4$ from biogenic sources is -41 to -86% [*Quay et al.*, 1988] and that from peatlands is -46% [*Martens et al.*, 1992] to -83% [*Lansdown et al.*, 1992]. The low $\delta^{13}\text{C}$ values observed here usually fell in these ranges; however, we do not know why such low values were observed only in the winter of 2004/2005. This winter was warmer than normal, especially in December (monthly average air temperature at meteorological station was 6.3°C; average value from 1996 to 2005 was 5.1°C). In addition, such a large decrease in $\delta^{13}\text{C}\text{-CH}_4$

was not observed in the much colder winter of 2005/2006 (Figure 2a), suggesting that some factor other than temperature affects this phenomenon. Although statistical analysis showed that hydrological conditions such as rainfall [including the antecedent precipitation index (API)], runoff, and water table are not directly related to this phenomenon (data not shown), we assume that acetate becomes exhausted under conditions of high methanogenic activity, even in early winter 2004, because CH₄ production remained much higher from late summer to autumn 2004 than in other years (Figure 2b). This may have induced the large depletion of $\delta^{13}\text{C}\text{-CH}_4$ and F_{Ac} , especially at KW2-center where water exchange is slow.

4.3. Relationship Between Changes in Contribution of Acetate Fermentation and CH₄ Flux

[37] At KW2-downstream, the patterns of $\delta^{13}\text{C}\text{-CH}_4$ and $\delta^{13}\text{C}\text{-CO}_2$ differed each summer (Figures 2d and 3d) depending on reducing conditions controlled by precipitation, runoff, water table, and thus DO (Table 1 [Itoh et al., 2007]). As described above, based on isotope signatures, the calculated contribution of acetate fermentation usually decreased in high temperature periods; however, it episodically increased under the highly reducing conditions in summer 2004 (Figures 7b and 9b). Itoh et al. [2007] reported a dramatic increase in CH₄ emission from the soil surface with increased CH₄ concentration in the surface groundwater in summer 2004. These observations suggest that CH₄ emission from the soil surface results from increased CH₄ production and accumulation in soil. The activation of acetate fermentation under highly reducing conditions probably contributed to this large CH₄ emission. A similar pattern was observed in KW2-edge in summer 2004. These results are consistent with previous observations of high acetate concentrations during early spring, followed by a large CH₄ flux, and then a sharp decrease in acetate concentration in freshwater peatlands in southern Michigan, USA [Shannon and White, 1996].

[38] Our results indicate that detecting increases in acetate fermentation using isotopic signatures can provide information on changes in CH₄ production and emission in summers with different redox conditions. Large variations in CH₄ production pathways, as described above, are probably more applicable in regions with large variations in temperature and precipitation, such as the Asian monsoon region. In such regions, the methanogenic pathway can vary widely in spatial and temporal scale with variations in redox conditions. Our observations indicate that the amount of acetate available for methanogenesis is controlled by supply, and therefore both water movement and microbial consumption are important factors.

[39] Schematics of the effects of changes in environmental conditions (e.g., precipitation, water table, and runoff) on CH₄ dynamics (formation pathways, groundwater concentrations, and emissions from soil surfaces) in summer are presented in Figure 10. Under “low temperature” and “high water table” conditions (e.g., the rainy summer of 2003 at our study site), less-reduced conditions than normal for summer were maintained by (1) increased oxygen supply from more frequent incoming water due to high runoff and precipitation and (2) low oxygen consumption due to low microbial activity. In such years, the acetate contribution to CH₄ production is stable and methanogenic activity from

Figure 10. Schematics of the effects of environmental changes on CH₄ dynamics in summer. The thickness of an arrow indicates its degree of contribution.

acetate is low. The amount of CH₄ produced and retained in groundwater is also low; consequently, CH₄ emissions from the soil surface are small (Figure 10).

[40] In a highly reducing summer, under conditions of “high temperature” and “high water table” (e.g., 2004 at our study site), CH₄ production from both carbonate reduction and acetate fermentation becomes active, and the proportion of carbonate reduction is generally increased under high CO₂ concentrations. In addition, acetate fermentation episodically increases under high microbial activity, resulting in heavier ¹³C-CH₄. A large quantity of CH₄ is retained in surface soils and contributes to high CH₄ emissions from the soil surface.

[41] Under “high temperature” and “low water table” conditions that usually occur in low-precipitation summers (e.g., 2005 at our study site), water table depletion induces the formation of an aerobic (oxic) layer in surface soils. On the other hand, highly reducing conditions form in the bottom layer because of the decreased oxygen levels caused by the reduced water movement [Itoh *et al.*, 2007]. The proportion of acetate fermentation decreases, probably because of low water movement under such conditions. CH₄ emissions from the soil surface are suppressed because of low anoxic microbial activity in the aerobic surface layer. Re-oxidation of CH₄ may occur in this layer. Our results suggest that changes in the CH₄ production pathway are influenced by changes in hydrologically controlled redox conditions and, consequentially, affect emissions from the soil surface.

5. Conclusions

[42] Our results show that changes in environmental conditions, especially hydrological conditions (water table and temperature), affect the CH₄ production pathways in riparian wetlands in temperate forests because hydrological conditions control the spatial and temporal variation in

redox conditions. Under “high water table” and “low temperature” conditions (rainy summers with high levels of runoff), high levels of DO create more oxic conditions, and the proportion of carbonate reduction and acetate fermentation is stable and activity levels are low. In contrast, in a highly reducing summer with “high water table” and “high temperature” conditions, the proportion of carbonate reduction increases with soil temperature, with episodic increases in acetate fermentation. In such a period, the decreased SRB activity associated with limited SO₄²⁻ levels can also induce acetate fermentation. This induces higher CH₄ production and CH₄ accumulation in soil and, consequently, high CH₄ emissions from the soil surface. In drier summers with “low water table” and “high temperature” conditions, the contribution of acetate fermentation is lower probably due to oxic conditions and decreased acetate supply. Acetate fermentation also decreases sharply in winter following a highly reducing summer. These patterns indicate that the amount of acetate available for methanogenesis is controlled by supply, and both water movement and microbial consumption are important factors regulating this supply.

[43] The results of this study demonstrate the effects of hydrological changes in wetlands on changes in CH₄ production pathways. Our findings indicate the importance of considering hydrologic effects when assessing CH₄ production in wetlands where redox conditions vary on shorter temporal and smaller spatial scales. In addition, our results that acetate fermentation was highly variable compared to carbonate reduction suggest that a more detailed understanding of what processes affect acetate supply and consumption is required.

[44] **Acknowledgments.** We thank anonymous reviewers for their valuable comments on an earlier draft of this manuscript; N. Matsuo (Mie University) for providing data; Y. Kosugi (Kyoto University) and

M. Katsuyama (Research Institute for Humanity and Nature) for helpful discussions; and the members of the Laboratory of Forest Hydrology for helping with the observations and analyses. This work was supported by CREST, JST, the Ministry of Education, Culture, Sports, Science, and Technology for Science Research (18380093, 18201036, 20780117), and by the Global Environment Research Program of the Ministry of the Environment, Japan.

References

- Alperin, M. J., N. E. Blair, D. B. Albert, T. M. Hoehler, and C. S. Martens (1992), Factors that control the stable carbon isotopic composition of methane produced in anoxic marine sediments, *Global Biogeochem. Cycles*, **6**, 271–291.
- Avery, G. B., Jr., R. D. Shannon, J. R. White, C. S. Martens, and M. J. Alperin (1999), Effect of seasonal changes in the pathways of methanogenesis on the $\delta^{13}\text{C}$ values of pore water methane in a Michigan peatland, *Global Biogeochem. Cycles*, **13**, 475–484.
- Bartlett, K. B., and R. C. Harriss (1993), Review and assessment of methane emissions from wetlands, *Chemosphere*, **26**, 261–320.
- Blair, N., A. Leu, E. Muñoz, J. Olsen, E. Kwong, and D. D. Marais (1985), Carbon isotopic fractionation in heterotrophic microbial metabolism, *Appl. Environ. Microbiol.*, **50**, 996–1001.
- Blair, N. E., S. E. Boehme, and W. D. Carter (1993), The carbon isotope biogeochemistry of methane production in anoxic sediments: 1. Field observations, in *The Biogeochemistry of Global Change: Radiative Trace Gases*, edited by R. S. Oremrand, pp. 574–593, Chapman and Hall, New York.
- Botz, R., H. D. Pokojski, M. Schmitt, and M. Thomm (1996), Carbon isotope fractionation during bacterial methanogenesis by CO₂ reduction, *Org. Geochem.*, **25**, 255–262.
- Bräunlich, M., O. Aballain, T. Marik, P. Jöckel, C. A. M. Brenninkmeijer, J. Chappellaz, J. M. Barnola, R. Mulvaney, and W. T. Sturges (2001), Changes in the global atmospheric methane budget over the last decades inferred from ^{13}C and D isotopic analysis of Antarctic firn air, *J. Geophys. Res.*, **106**, 20,465–20,481.
- Bremner, J. M. (1996), N-total, in *Methods of soil analysis. Part 3, Soil Sci. Soc. Am. Book Ser.*, vol. 5, edited by D. L. Sparks, pp. 1085–1121, Soil Sci. Soc. Am., Madison, Wis.
- Burke, R. A., C. S. Martens, and W. M. Sackett (1988), Seasonal variations of D/H and $^{13}\text{C}/^{12}\text{C}$ ratios of microbial methane in surface sediments, *Nature*, **332**, 829–831.
- Chan, O. C., P. Claus, P. Casper, A. Ulrich, T. Lueders, and R. Conrad (2005), Vertical distribution of structure and function of the methanogenic archaeal community in Lake Dagow sediment, *Environ. Microbiol.*, **7**(8), 1139–1149.
- Charman, D. J., R. Aravena, and B. G. Warner (1994), Carbon dynamics in a forested peatland in north-eastern Ontario, Canada, *J. Ecol.*, **82**, 55–62.
- Chin, K. J., and R. Conrad (1995), Intermediary metabolism in methanogenic paddy soil and the influence of temperature, *FEMS Microbiol. Ecol.*, **18**, 85–102.
- Cicerone, R. J., and R. S. Oremland (1988), Biogeochemical aspects of atmospheric methane, *Global Biogeochem. Cycles*, **2**, 299–327.
- Conrad, R. (2005), Quantification of methanogenic pathways using stable carbon isotopic signatures: A review and a proposal, *Org. Geochem.*, **36**, 739–752.
- Conrad, R., and P. Claus (2005), Contribution of methanol to the production of methane and its ^{13}C -isotopic signature in anoxic rice field soil, *Biogeochemistry*, **73**, 381–393.
- Conrad, R., H. P. Mayer, and M. Wüst (1989), Temporal change of gas metabolism by hydrogen-syntrophic methanogenic bacterial associations in anoxic paddy soil, *FEMS Microbiol. Lett.*, **62**, 265–273.
- Crill, P. M., and C. S. Martens (1986), Methane production from bicarbonate and acetate in an anoxic marine sediment, *Geochim. Cosmochim. Acta*, **50**, 2089–2097.
- Etheridge, D. M., L. P. Steele, R. J. Francey, and R. L. Langenfelds (1998), Atmospheric methane between 1000 A.D. and present: Evidence of anthropogenic emissions and climatic variability, *J. Geophys. Res.*, **103**, 15,979–15,993.
- Fey, A., and R. Conrad (2000), Effect of temperature on carbon and electron flow and on the archaeal community in methanogenic rice field soil, *Appl. Environ. Microbiol.*, **66**, 4790–4797.
- Fey, A., K. J. Chin, and R. Conrad (2001), Thermophilic methanogens in rice field soil, *Environ. Microbiol.*, **3**(5), 295–303.
- Fey, A., P. Claus, and R. Conrad (2004), Temporal change of ^{13}C -isotope signatures and methanogenic pathways in rice field soil incubated anoxically at different temperatures, *Geochim. Cosmochim. Acta*, **68**, 293–306.
- Fletcher, S. E. M., P. P. Tans, L. M. Bruhwiler, J. B. Miller, and M. Heimann (2004), CH₄ sources estimated from atmospheric observations of CH₄ and its $^{13}\text{C}/^{12}\text{C}$ isotopic ratios: 1. Inverse modeling of source processes, *Global Biogeochem. Cycles*, **18**, GB4004, doi:10.1029/2004GB002223.
- Games, L. M., J. M. Hayesand, and R. P. Gunsalus (1978), Methane-producing bacteria: Natural fractionations of the stable carbon isotopes, *Geochim. Cosmochim. Acta*, **42**, 1295–1297.
- Gelwicks, J. T., J. B. Risatti, and J. M. Hayes (1994), Carbon isotope effects associated with aceticlastic methanogenesis, *Appl. Environ. Microbiol.*, **60**, 467–472.
- Gupta, M., S. Tyler, and R. Cicerone (1996), Modeling atmospheric $\delta^{13}\text{C}$ and the causes of recent changes in atmospheric CH₄ amounts, *J. Geophys. Res.*, **101**, 22,923–22,932.
- Intergovernmental Panel on Climate Change (2001), *The Scientific Basis*, edited by J. T. Houghton et al., pp. 10–64, Cambridge Univ. Press, New York.
- Itoh, M., N. Ohte, M. Katsuyama, K. Koba, M. Kawasaki, and M. Tani (2005), Temporal and spatial variability of methane flux in a temperate forest watershed (in Japanese with English abstract), *J. Jpn. Soc. Hydrol. Water Resour.*, **18**, 244–256.
- Itoh, M., N. Ohte, K. Koba, M. Katsuyama, K. Hayamizu, and M. Tani (2007), Hydrologic effects on methane dynamics in riparian wetlands in a temperate forest catchment, *J. Geophys. Res.*, **112**, G01019, doi:10.1029/2006JG000240.
- Khalil, M. A. K., and M. J. Shearer (1993), Sources of methane, in *An Overview. Atmospheric Methane, Sources, Sinks and Role in Global Change*, edited by M. A. K. Khalil, Springer, Berlin.
- Krzycki, J. A., R. H. Wolkin, and J. G. Zeikus (1982), Comparison of unitrophic and mixotrophic substrate metabolism by an acetate-adapted strain of *Methanosarcina barkeri*, *J. Bacteriol.*, **149**, 247–254.
- Krzycki, J. A., W. R. Kenealy, M. J. Deniro, and J. G. Zeikus (1987), Stable carbon isotope fractionation by *Methanosarcina barkeri* during Methanogenesis from acetate, methanol, or carbon dioxide-hydrogen, *Appl. Environ. Microbiol.*, **53**, 2597–2599.
- Lansdown, J. M., P. D. Quay, and S. L. King (1992), CH₄ production via CO₂ reduction in a temperate bog: A source of ^{13}C -depleted CH₄, *Geochim. Cosmochim. Acta*, **56**, 3493–3503.
- Lelieveld, J., and P. J. Crutzen (1992), Indirect chemical effects of methane on climate warming, *Nature*, **355**, 339–342.
- Lovley, D. R., and M. J. Klug (1983a), Sulfate reducers can outcompete methanogens at freshwater sulfate concentrations, *Appl. Environ. Microbiol.*, **45**, 187–192.
- Lovley, D. R., and M. J. Klug (1983b), Methanogenesis from methanol and methylamines and acetogenesis from hydrogen and carbon dioxide in the sediments of a eutrophic lake, *Appl. Environ. Microbiol.*, **45**, 1310–1315.
- Lowe, D. C., C. A. M. Brenninkmeijer, G. W. Brailsford, K. R. Lassey, A. J. Gomez, and E. G. Nisbet (1994), Concentration and ^{13}C records of atmospheric methane in New Zealand and Antarctica: Evidence for changes in methane sources, *J. Geophys. Res.*, **99**, 16,913–16,926.
- Martens, C. S., N. E. Blair, C. D. Green, and D. J. Des Marais (1986), Seasonal variations in the stable carbon isotope signature of biogenic methane in a coastal sediment, *Science*, **233**, 1300–1303.
- Martens, C. S., C. A. Kelley, J. P. Chanton, and W. J. Showers (1992), Carbon and hydrogen isotopic characterization of methane from wetlands and lakes of the Yukon-Kuskokwim Delta, Western Alaska, *J. Geophys. Res.*, **97**, 16,689–16,701.
- Mary, B., A. Mariotti, and J. L. Morel (1992), Use of ^{13}C variations at natural abundance for studying the biodegradation of root mucilage, roots and glucose in soil, *Soil Biol. Biochem.*, **24**, 1065–1072.
- Matsuo, N. (2003), Evaluation of leaf gas exchange using carbon isotopic analysis, Doctoral dissertation, Kyoto Univ., Kyoto, Japan.
- McAuliffe, C. (1971), GC determination of solutes by multiple phase equilibration, *Chemtech*, **1**, 46–51.
- Mitsch, W. J., and J. G. Gosselink (2000), *Wetland biogeochemistry*, in *Wetlands*, 3rd ed., edited by W. J. Mitsch and J. G. Gosselink, pp. 155–204, John Wiley, New York.
- Nakagawa, F., N. Yoshida, Y. Nojiri, and V. N. Makarov (2002), Production of methane from alasses in eastern Siberia: Implications from its ^{14}C and stable isotopic compositions, *Global Biogeochem. Cycles*, **16**(3), 1041, doi:10.1029/2000GB001384.
- Ohte, N., N. Tokuchi, and M. Suzuki (1995), Biogeochemical influences on the determination of water chemistry in a temperate forest basin: Factors determining the pH value, *Water Resour. Res.*, **31**, 2823–2834.
- Pine, M. J., and H. A. Barker (1956), Studies on the methane fermentation XII.: The pathway of hydrogen in the acetate fermentation, *J. Bacteriol.*, **71**, 644–648.
- Quay, P. D., S. L. King, J. M. Lansdown, and D. O. Wilbur (1988), Isotopic composition of methane released from wetlands: Implications for the increase in atmospheric methane, *Global Biogeochem. Cycles*, **2**, 385–397.
- Quay, P. D., et al. (1991), Carbon isotopic composition of atmospheric CH₄: Fossil and biomass burning source strengths, *Global Biogeochem. Cycles*, **5**, 25–47.
- Reeburgh, W. S., S. C. Whalen, and M. J. Alpern (1993), The role of methylophony in the global methane budget, in *Microbial Growth on*

G03005

ITO ET AL.: ANALYSIS OF CH₄ PRODUCTION USING δ¹³C

G03005

- Cl Compounds*, edited by J. C. Murrell and D. P. Kelly, pp. 1–14, Intercept, Andover, UK.
- Roslev, P., and G. M. King (1994), Survival and recovery of methanotrophic bacteria starved under oxic and anoxic conditions, *Appl. Environ. Microbiol.*, **60**, 2602–2608.
- Roslev, P., and G. M. King (1996), Regulation of methane oxidation in a freshwater wetland by water table changes and anoxia, *FEMS Microbiol. Ecol.*, **19**, 105–115.
- Schönheit, P., J. K. Kristjansson, and R. K. Thauer (1982), Kinetic mechanism for the ability of sulfate reducers to out-compete methanogens for acetate, *Arch. Microbiol.*, **132**, 285–288.
- Schütz, H., W. Seiler, and R. Conrad (1989), Process evolved in formation and emission of methane in rice paddies, *Biogeochemistry*, **7**, 33–53.
- Shannon, R., and J. White (1996), The effects of spatial and temporal variations in acetate and sulfate on methane cycling in two Michigan peatlands, *Limnol. Oceanogr.*, **41**, 435–443.
- Smejkal, V., F. D. Cook, and H. R. Krouse (1971), Studies of sulfur and carbon isotope fractionation with microorganisms isolated from springs of western Canada, *Geochim. Cosmochim. Acta*, **35**, 787–800.
- Stevens, C. M., and A. Engelkemeir (1988), Stable carbon isotopic composition of methane from some natural and anthropogenic sources, *J. Geophys. Res.*, **93**, 725–733.
- Stevens, C. M., and F. E. Rust (1982), The carbon isotopic composition of atmospheric methane, *J. Geophys. Res.*, **87**, 4879–4882.
- Sugimoto, A. (1996), GC/GC/C/IRMS system for carbon isotope measurement of low level methane concentration, *Geochem. J.*, **30**, 195–200.
- Sugimoto, A., and N. Fujita (2006), Hydrogen concentration and stable isotopic composition of methane in bubble gas observed in a natural wetland, *Biogeochemistry*, **81**, 33–44.
- Sugimoto, A., and E. Wada (1993), Carbon isotopic composition of bacterial methane in a soil incubation experiment: Contributions of acetate and CO₂/H₂, *Geochim. Cosmochim. Acta*, **57**, 4015–4027.
- Takai, Y. (1970), The mechanism of methane fermentation in flooded paddy soil, *Soil Sci. Plant Nutr.*, **6**, 238–244.
- Vogels, G. D., K. T. Keltjens, and C. Van der Drift (1988), Biochemistry of methane production, in *Biology of Anaerobic Microorganisms*, edited by A. J. B. Zehnder, pp. 707–770, John Wiley, New York.
- Waldron, S., A. J. Hall, and A. E. Fallick (1999), Enigmatic stable isotope dynamics of deep peat methane, *Global Biogeochem. Cycles*, **13**, 93–100.
- Whalen, M., N. Tanaka, R. Henry, B. Deck, J. Zeglen, J. S. Vogel, J. Southon, A. Shemesh, R. Fairbanks, and W. Broecker (1989), Carbon-14 in methane sources and in atmospheric methane: The contribution from fossil carbon, *Science*, **245**, 286–290.
- Whiticar, M. J. (1999), Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane, *Chem. Geol.*, **161**, 291–314.
- Whiticar, M. J., E. Faber, and M. Schoell (1986), Biogenic methane formation in marine and freshwater environments: CO₂ reduction vs. acetate fermentation-isotope evidence, *Geochim. Cosmochim. Acta*, **50**, 693–709.
- Winfrey, M. R., and J. G. Zeikus (1977), Effect of sulfate on carbon and electron flow during microbial methanogenesis in freshwater sediments, *Appl. Environ. Microbiol.*, **33**, 275–281.
- Zeikus, J. G., P. J. Weimer, D. R. Nelson, and L. Daniels (1975), Bacterial methanogenesis: Acetate as a methane precursor in pure culture, *Arch. Microbiol.*, **104**, 129–134.

M. Itoh and M. Tani, Laboratory of Forest Hydrology, Division of Environmental Science and Technology, Graduate School of Agriculture, Kyoto University, Kitashirakawa Oiwake-Cho, Sakyo-Ku, Kyoto 606-8502, Japan. (itoma@kais.kyoto-u.ac.jp)

K. Koba, Institute of Symbiotic Science and Technology, Tokyo University of Agriculture and Technology, Saiwaicho, 3-5-8, Fuchu-city, Tokyo 183-8509, Japan.

N. Ohte, Department of Forest Science, Graduate School of Agricultural and Life Sciences, University of Tokyo, 1-1-1, Yayoi, Bunkyo-Ku, Tokyo 113-8657, Japan.

A. Sugimoto, Division of Earth System Science, Faculty of Environmental Earth Science, Hokkaido University, N10W5 Sapporo, Hokkaido 060-0810, Japan.