

Title	Synthetic Study of Matrine-Type Alkaloids: Stereoselective Construction of the AB Rings of the Quinolizidine Skeleton
Author(s)	Tsukano, Chihiro; Oimura, Atsuko; Enkhtaivan, Iderbat; Takemoto, Yoshiji
Citation	Synlett (2014), 25(5): 653-656
Issue Date	2014-03-05
URL	http://hdl.handle.net/2433/199887
Right	© Georg Thieme Verlag Stuttgart · New York; This is not the published version. Please cite only the published version. この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。
Type	Journal Article
Textversion	author

Synthetic Study of Matrine-type Alkaloids: Stereoselective Construction of the AB rings of the Quinolizidine Skeleton

Chihiro Tsukano, Atsuko Oimura, Iderbat Enkhtaivan, Yoshiji Takemoto*

Graduate School of Pharmaceutical Sciences, Kyoto University, Yoshida, Sakyo-ku, Kyoto, 606-8501, Japan

Fax: +81(75)7534532; E-mail: Takemoto@pharm.kyoto-u.ac.jp

Received: The date will be inserted once the manuscript is accepted.

Abstract: A new method has been developed for the stereoselective construction of the AB rings of the quinolizidine skeleton of matrine-type alkaloids with a *cis-cis* stereochemistry. The key features of this method involve: (i) the construction of the quinolizidine by the reduction of an acylpyridinium cation; and (ii) the late stage introduction of methoxypyridine by sequential Stille coupling and diastereoselective hydrogenation reactions.

Keywords: Quinolizidines, stereoselective synthesis, alkaloids, matrine, pyridines

Matrine is representative of the alkaloids derived from *Sophora flavescens* Ait., and has been used for many years as a traditional Chinese medicine (KuShen).¹ The structure of matrine is characterized by a tetracyclic core consisting of two quinolizidines and four contiguous stereogenic centers (Figure 1).² To date, over 30 matrine-type alkaloids have been reported in the literature, and these compounds usually only differ from each other in terms of their oxidation level and relative stereochemistry.³ Alkaloids belonging to this structural class also exhibit a variety of biological properties, including antitumor^{4a,e}, anti-viral^{4b,c}, and anti-inflammatory^{4d} activities. Because of their interesting and diverse biological activities, these compounds have attracted considerable attention from medicinal chemists.





Figure 1 Matrine and related quinolizidine alkaloids.

Although there are many congeners in this structural class, only four racemic total syntheses and three semi-syntheses have been reported to date, including those of matrine^{6-8,9a}, leontine^{6a}, sophoramine^{9b} and isosophoramine.^{9c,10} These syntheses revealed that the stereoselective construction of the tetracyclic core of

these compounds represents a significant synthetic challenge. Brown *et al.*¹¹ recently reported the synthesis of (+)-allomatrine, where the tetracyclic core was elegantly constructed in a diastereoselective manner using an imino-aldol reaction and *N*-acyliminium cyclization. Methods for the stereoselective synthesis of tetracyclic cores with a *cis-cis* configuration, however, such as those found in sophoramine and matrine, remain scarce. We recently reported a procedure for the synthesis of quinolizidines by the reduction of acylpyridinium cations under mild conditions.⁵ It was envisaged that this method could be used as a powerful tool for the concise synthesis of quinolizidine alkaloids, because the resulting products could be readily derivatized. With this in mind, we became interested in investigating the application of our method to the stereoselective synthesis of the AB rings of matrine type alkaloids with *cis-cis* stereochemistry.


Retrosynthetically, alcohol **1** was set as a suitable synthetic target, because it was envisaged that this compound would provide comprehensive access to matrine and several related alkaloids via the cyclization of the C ring and the adjustment of the oxidation levels (Scheme 1). The *cis-cis* stereochemistry of **1** could be successfully installed by the hydrogenation of compound **2** or **5**. Compound **2**



Scheme 1 Retrosynthetic analysis of matrine type alkaloids.

could be accessed via the introduction of a C1 unit to the quinolizidine core of compound **3**, which could be constructed via the reduction of the acylpyridinium cation derived from carboxylic acid **4** (route a). Alternatively, the addition of a pyridyl moiety to compound **6**, which could be derived from **7**, could also be used to provide access alcohol **1** (route b). Herein, we report the development of a stereoselective synthesis of the key *cis-cis* intermediate **1** based on our reductive cyclization strategy by examining both possibilities.


To begin, we examined route a, which required the challenging selective reduction of one of two pyridines. The synthesis of the cyclization precursor **4** started with the Stille coupling of 2-iodopicolinate methyl ester **8**¹² and pyridylstannane **9**¹³ using catalytic amounts of Pd(PPh₃)₄ and CuI (Scheme 2). CuI was essential for the production of bipyridine **10**.¹⁴ Compound **10** was converted to carboxylic acid **4** by the Horner-Wadsworth-Emmons reaction of **11** with ethyl glyoxylate, followed by sequential hydrogenation and hydrolysis reactions, which gave the cyclization precursor **4** in 50% yield from the coupling product **10**.¹⁵ The quinolizidine skeleton was constructed by the reduction of the acylpyridinium cation intermediate, which was produced by the activation of one of the two pyridine rings of **4**. Treatment of **4** with Ghosez's reagent followed by Hantzsch ester gave the desired product **3** in 36% yield. This low yield was attributed to the unfavorable nucleophilic attack of the nitrogen of the other pyridine on the *in situ* generated acid chloride. Hydrogenation of the cyclized product **3** proceeded stereoselectively to give the desired compound **12** as a



Scheme 2 Synthesis of bipyridine **4** and its cyclization. (a) Pd(PPh₃)₄, CuI, DMF, 110 °C, 74%; (b) *n*BuLi, MeP(O)(OMe)₂, THF, -78 °C; (c) *t*BuOK, EtO₂CCHO, DME, -20 °C, 57% (2 steps); (d) H₂, Pd/C, AcOEt; (e) LiOH, THF-H₂O, 88% (2 steps); (f) Ghosez's reagent, MS 4Å then Hantzsch ester, ClCH₂CH₂Cl, 36%; (g) H₂, Pd/C, EtOH, 70%.

single isomer. The newly generated stereochemistry was confirmed by NOE experiments. Whilst ketone **12** was accessed in stereoselective manner, the quinolizidine ring formation resulted in a low yield and we decided to then focus our efforts on the other route for accessing the *cis-cis* intermediate **1**.

The second route started from 2-bromo-3-hydroxymethylpyridine (**13**)¹⁶, which was converted to the corresponding benzyl ether **14** using standard techniques (Scheme 3). Subsequent treatment of **14** with *n*-butyl lithium followed by the addition of succinic anhydride **15** gave carboxylic acid **7**, albeit in low yield (45%). To improve the yield and reproducibility of this step, compound **14** was first converted to the 2-TMS-pyridine species, which was treated with succinic anhydride **15** to give carboxylic acid **7** in 65% yield over two steps.^{15,17} Compound **7** was then successfully converted to compound **6** by treatment with Ghosez's reagent followed by Hantzsch ester.¹⁸ Compound **6** was found to be unstable, and was immediately hydrogenated over Pd/C to give ketone **16**, whose stereochemistry was confirmed by NOE experiments and the coupling constants in its ¹H NMR, as shown in Figure 2. The second route performed more effectively than the first route in terms of the number of steps required and the yield of the key cyclization step.



Scheme 3 Synthesis of quinolizidine **16**. (a) BnBr, NaH, DMF, 68%; (b) *n*BuLi then succinic anhydride **15**, Et₂O, -78 °C to RT, 45%; (c) *n*BuLi, TMSCl, Et₂O, -78 °C to RT; (d) **15**, ClCH₂CH₂Cl, 100 °C, 65% (two steps); (e) Ghosez's reagent, MS 4Å then Hantzsch ester, ClCH₂CH₂Cl, 0 °C to RT, 61%; (f) H₂, Pd/C, EtOH, 53%.





Figure 2 ¹H NMR coupling constant and NOESY experiments of compound **16** (left) and a stable conformation calculated by Spartan at the B3LYP/6-31+G(d) level of theory (DFT) (right).

We then proceeded to investigate the stereoselective introduction of the pyridyl moiety into compound **16**. Preliminary studies revealed that the structurally

related ketone **12** was readily enolized rather than attacked by nucleophiles such as TMSCH_2Li and $\text{Ph}_3\text{P}=\text{CH}_2$.¹⁹ The treatment of compound **16** with NaHMDS and Comins' reagent gave enol triflate **17** together with a significant amount of its regioisomer **18** in a combined yield of 61% (**17**:**18** = 1.3:1) (Scheme 4). A variety of different bases (i.e., LDA , LiTMP , TrLi , KHMDS , NaHMDS , LiHMDS , Et_3N) and triflating reagents (i.e., TF_2O , TF_2NPh , and Comins' reagent²⁰) were evaluated for this transformation, but none of these combinations led to an improvement in the low regioselectivity. Pleasingly, it was possible to separate these two compounds by silica gel column chromatography, and this allowed for sufficient quantities of the enol triflate **17** to be obtained for the subsequent Stille coupling. Following an extensive investigation of the conditions required for the coupling of **17** with **9** using several copper salts such as CuTC ,²¹ $\text{CuBr}\cdot\text{SMe}_2$, and CuDPP ,²² it was found that the reaction proceeded smoothly in the presence of catalytic $\text{Pd}(\text{PPh}_3)_4$ with CuDPP (copper(I) diphenylphosphinate) and LiCl in THF at 50°C to give the coupling product **5** in good yield. Hydrogenation of **5** gave compound **19** as a single isomer, because the α face was shielded by the hydroxymethyl group (Figure 3, left). The newly generated stereochemistry was confirmed by NOE experiments (Figure 3, right). Finally, reduction with LiAlH_4 gave the common intermediate **1** with the required *cis-cis* stereochemistry.²³



Scheme 4 Synthesis of common intermediate **1**. (a) NaHMDS , Comins' reagent, THF , -78 to 0°C , 61%, (**17**:**18** = 1.3:1); (b) **9**, $\text{Pd}(\text{PPh}_3)_4$, CuDPP , LiCl , THF , 50°C , 67%; (c) Pd/C , H_2 (5 atm), EtOH , 70°C , 65%; (d) LiAlH_4 , THF , 50°C , 80%.




Figure 3 Regioselectivity of hydrogenation of compound **5** (left), a stable conformation calculated by Spartan at the B3LYP/6-31+G(d) level of theory (DFT) (middle), NOESY experiments of compound **19** (right).

In summary, we have developed a method for the concise synthesis of quinolizidine **1** in a stereoselective manner. The key features of this method include: (i) the construction of the quinolizidine ring by the reduction of an acylpyridinium cation; and (ii) the late stage introduction of a pyridyl group by sequential Stille coupling and hydrogenation reactions. This synthetic strategy represents a reasonable technique for the stereoselective construction of quinolizidine rings with *cis-cis* stereochemistry. The synthesis of matrine-type alkaloids such as sophoramine, sophocarpine and matrine from quinolizidine **1** via the construction of the C ring and subsequent adjustment of the oxidation levels is currently underway in our laboratory.

Supporting Information for this article is available online at <http://www.thieme-connect.com/ejournals/toc/synlett>.

Acknowledgment

This work was supported by a Grant-in-Aid for Young Scientists (B) (C. T.) from the Japan Society for the Promotion of Science.

References

- (1) Nagai, N.; Kondo, H. *Yakugaku Zasshi* **1903**, 260, 993.
- (2) (a) Tsuda, K.; Saeki, S.; Imura, S.-I.; Okuda, S.; Sato, Y.; Mishima, H. *J. Org. Chem.* **1956**, 21, 1481. (b) Ochiai, E.; Okuda, S.; Minato, H. *Yakugaku Zasshi* **1952**, 72, 781.
- (3) (a) Micheal, J. P. *Nat. Prod. Rep.* **2008**, 25, 139. (b) K. A. Aslanov, Y. K. Kushmuradov, S. Sadykov, In *The Alkaloids*, Vol. 31; Brossi, A., Ed.; Academic Press: **1987**, 117, and references therein.
- (4) For recent examples of biological studies, see (a) Wang, L.; You, Y.; Wang, S.; Liu, X.; Liu, B.; Wang, J.; Lin, X.; Chen, M.; Liang, Y.; Yang, H. G. *Bioorg. Med. Chem. Lett.* **2012**, 22, 4100. (b) Yang, Y.; Xiu, J.; Zhang, X.; Zhang, L.; Yan, K.; Qin, C.; Liu, J. *Molecules*, **2012**, 17, 10370. (c) Gao, L.-M.; Han, Y.-X.; Wang, Y.-P.; Li, Y.-H.; Shan, Y.-Q.; Li, X.; Peng, Z.-G.; Bi, C.-W.; Zhang, T.; Du, N.-N.; Jiang, J.-D.; Song, D.-Q. *J. Med. Chem.* **2011**, 54, 869. (d) Hu, H.; Wang, S.; Zhang, C.; Wang, L.; Ding, L.; Zhang, J.; Wu, Q. *Bioorg. Med. Chem. Lett.* **2010**, 20, 7537. (e) Ma, L.; Wen, S.; Zhan, Y.; He, Y.; Liu, X.; Jiang, J. *Planta Med.* **2008**, 74, 245.
- (5) Tsukano, C.; Oimura, A.; Enkhtaivan, I.; Takemoto, Y. *Org. Lett.* **2012**, 14, 1902.
- (6) (a) Mandell, L.; Singh, K. P.; Gresham, J. T.; Freeman, W. J. *J. Am. Chem. Soc.* **1965**, 87, 5234. (b) Mandell, L.; Piper, J. U.; Singh, K. P. *J. Org. Chem.* **1963**, 28, 3440. (c) Mandell, L.; Singh, K. P. *J. Am. Chem. Soc.* **1961**, 83, 1766.
- (7) Chen, J.; Browne, L. J.; Gonnella, N. C. *J. Chem. Soc., Chem. Commun.*, **1986**, 905.
- (8) Boiteau, L.; Boivin, J.; Liard, A.; Quiclet-Sire, B.; Zard, S. Z. *Angew. Chem. Int. Ed.* **1998**, 37, 1128.
- (9) (a) Okuda, S.; Kamata, H.; Tsuda, K.; Murakoshi, I. *Chem. & Ind. (London)*, **1962**, 1326. (b) Okuda, S.; Yoshimoto, M.; Tsuda, K. *Chem. Pharm. Bull.* **1966**, 14, 275. (c) Okuda, S.; Kamata, H.; Tsuda, K. *Chem. Pharm. Bull.* **1963**, 11, 1349.

- (10) Wenkert, E.; Chauncy, B.; Dave, K. G.; Jeffcoat, A. R.; Schell, F. M.; Schenk, H. P. *J. Am. Chem. Soc.* **1973**, *95*, 8427.
- (11) Watkin, S. V.; Camp, N. P.; Brown, R. C. D. *Org. Lett.* **2013**, *15*, 4596.
- (12) Carson, M. W.; Giese, M. W.; Coghlan, M. J. *Org. Lett.* **2008**, *10*, 2701.
- (13) Honda, T.; Takahashi, R.; Namiki, H. *J. Org. Chem.* **2005**, *70*, 499.
- (14) Liebeskind, L. S.; Fengl, R. W. *J. Org. Chem.* **1990**, *55*, 5359.
- (15) Tsukano, C.; Zhao, L.; Takemoto, Y.; Hiramama, M. *Eur. J. Org. Chem.* **2010**, 4198.
- (16) (a) Ashimori, A.; Ono, T.; Uchida, T.; Ohtaki, Y.; Fukaya, C.; Watanabe, M.; Yokoyama, K. *Chem. Pharm. Bull.*, **1990**, *38*, 2446. (b) Srinivasan, J. M.; Burks, H. E.; Smith, C. R.; Visvanathan, R.; Johnston, J. N. *Synthesis*, **2005**, 330.
- (17) Pinkerton, F. H.; Thames, S. F. *J. Organomet. Chem.* **1970**, *24*, 623.
- (18) **Synthesis of 1,4-Dihydropyridine 6:** To a solution of carboxylic acid **7** (2.20 g, 7.34 mmol) and MS4A (ca. 5 g) in CH₂Cl₂ (50 mL) at 0 °C was added Ghosez's reagent (1.00 ml, 7.41 mmol). The mixture was stirred at 0 °C for 30 min, then Hantzsch ester (5.58 g, 22.0 mmol) was added to the reaction mixture. After stirring at room temperature for 2 h, the mixture was filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (2-3% ether/toluene) to afford dihydropyridine **6** (1.28 g, 4.52 mmol, 61%) as a solid; IR (ATR) 2979, 2721, 1689, 1235, 1089, 893 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.35-7.25 (m, 5H), 7.08 (dt, *J* = 8.0, 1.7 Hz, 1H), 5.13 (dt, *J* = 8.3, 3.5 Hz, 1H), 4.60-4.57 (m, 2H), 4.49 (s, 2H), 3.19-3.17 (m, 2H), 2.74 (ddd, *J* = 8.3, 6.0, 1.5 Hz, 2H), 2.67 (ddd, *J* = 8.3, 6.0, 1.5 Hz, 2H); ¹³C NMR (126 MHz, CDCl₃) δ 192.3, 165.3, 138.1, 132.8, 128.9, 128.5, 127.8, 127.7, 122.7, 108.1, 73.0, 69.7, 35.9, 29.7, 25.9; MS (FAB) *m/z* 284 [(M+H)⁺]; HRMS (FAB) calcd for C₁₇H₁₈NO₃ [(M+H)⁺] 284.1287, found 284.1296.
- (19) We also found that the related ketone, which did not have a substituent on the quinolizidine ring, enolized under the conditions of the Grignard and Wittig reactions. See ref. 5.
- (20) Comins, D. L.; Dehghani, A. *Tetrahedron Lett.* **1992**, *33*, 6299.
- (21) Allred, G. D.; Liebeskind, L. S. *J. Am. Chem. Soc.* **1996**, *118*, 2748.
- (22) Wittenberg, R.; Srogl, J.; Egi, M.; Liebeskind, L. S. *Org. Lett.* **2003**, *5*, 3033.
- (23) **Synthesis of quinolizidine 1:** To a solution of alcohol **19** (11.3 mg, 0.0390 mmol) in THF (1 mL) was added dropwise a solution of LiAlH₄ (2.2 mg, 0.058 mmol, 1.5 eq) in anhydrous THF (0.6 mL) at 0 °C under argon. The resulting mixture was stirred at 50 °C for 30 min. After careful hydrolysis with 3 M aq. NaOH (1 mL), EtOAc (1 mL) was added and the organic layer was separated. The aqueous layer was extracted with EtOAc (1 x 2 mL). The combined organic layers were dried over Na₂SO₄, filtered, and concentrated. The crude residue was purified by column chromatography (CHCl₃-MeOH, 20:1) to give alcohol **1** (10.8 mg, 0.035 mmol, 80%) as an oil. IR (ATR) 3356, 2928, 2857, 2754, 2683, 1578, 1466 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.47 (dd, *J* = 8.0, 7.4 Hz, 1H), 7.00 (d, *J* = 7.4 Hz, 1H), 6.57 (d, *J* = 8.0 Hz, 1H), 3.91 (s, 3H), 3.65-3.54 (m, 2H), 3.46 (dd, *J* = 11.2, 3.0 Hz, 1H), 3.20 (dd, *J* = 8.5, 6.3, 4.0 Hz, 1H), 2.98 (br d, *J* = 11.7 Hz, 1H), 2.87-2.83 (m, 2H), 2.30-2.14 (m, 3H), 2.10-2.02 (m, 2H), 1.86-1.81 (m, 2H), 1.59-1.43 (m, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 163.1, 160.8, 138.3, 116.2, 107.7, 67.2, 64.9, 57.7, 53.3, 45.2, 37.9, 31.9, 30.7, 29.6, 21.7, 21.6; MS (FAB) *m/z* 277.2 [(M+H)⁺]; HRMS (FAB) calcd for C₁₆H₂₅N₂O₂ [(M+H)⁺]; 277.1911, found 277.1913.

Synthetic Study of Matrine-type Alkaloids: Stereoselective Construction of the AB rings of the Quinolizidine Skeleton

