TITLE:

Effects of litter type, origin of isolate, and temperature on decomposition of leaf litter by macrofungi

AUTHOR(S):

Osono, Takashi

CITATION:

Osono, Takashi. Effects of litter type, origin of isolate, and temperature on decomposition of leaf litter by macrofungi. Journal of Forest Research 2014, 20(1): 77-84

ISSUE DATE:

2014-09-05

URL:

<http://hdl.handle.net/2433/200282>

RIGHT:

The final publication is available at Springer via <http://dx.doi.org/10.1007/s10310-014-0462-1>; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

1 Effects of litter type, origin of isolate, and temperature on decomposition of leaf

2 litter by macrofungi

3

4 Takashi Osono

5

6 T. Osono

7 Center for Ecological Research, Kyoto University, Otsu, Shiga 520-2113 Japan

8 e-mail: tosono@ecology.kyoto-u.ac.jp

9

10 **Abstract**The dependence of hyphal growth and litter decomposition on litter type
11 and incubation temperature used as substratum was compared for
12 litter-decomposing macrofungi (LDM) originating from subtropical (ST), cool
13 temperate (CT), and subalpine forests (SA) in Japan. In the first series of pure
14 culture decomposition tests using a total of 39 litter types as substrata inoculated
15 with six fungal isolates from the three climatic regions, the fungal decomposition
16 of litter was negatively affected by the content of AUR or extractives and
17 positively by N content in the litter. Secondly, cross-inoculation tests were

performed to examine the mass loss of leaf litter of broad-leaved trees from ST, CT, and SA, each inoculated with three *Mycena* species from the three climates and incubated at seven temperatures between 5°C and 35°C. Fungal isolate, litter type, incubation temperature, and their interactions significantly affected the mass loss of litter during the incubation. The greatest values of mass loss were found at 20°C or 25°C, and were generally consistent with the optimum temperatures of colony diameter growth rate of these isolates. Isolates from cooler regions were more sensitive to higher temperature than isolates from warmer regions. The decomposition of recalcitrant compounds (as acid-unhydrolyzable residues, AUR) by *Mycena* sp. from ST was also affected by litter type and incubation temperature, but the degree of selective decomposition of AUR relative to other components, such as cellulose, was insensitive to the range of temperature tested.

Keywords Acid-unhydrolyzable residue · Temperature · Leaves · Litter-decomposing macrofungi · Selective delignification

35 Introduction

36

37 Litter-decomposing macrofungi (LDM) include active ligninolytic species in

38 Basidiomycota and Ascomycota and play central roles in leaf-litter decomposition

39 processes (Osono 2007; Lindahl and Boberg 2008; van der Wal et al. 2013).

40 Ligninolytic abilities of LDM have been examined with the pure culture test and

41 compared for multiple species that were commonly found at a single site

42 (Lindeberg 1946; Miyamoto et al. 2000; Steffen et al. 2007; Valášková et al. 2007;

43 Boberg et al. 2011; Žifčáková et al. 2011). Few studies have been available that

44 compared the diversity and functioning of LDM across multiple sites of different

45 climatic regions. Recently, I reported that the species richness of LDM varied with

46 climatic region, declining from subtropical to temperate forest, and then from

47 temperate to subalpine forest in Japan (Osono in press b). Then, I compared the

48 potential abilities of diverse LDM from those forests to decompose leaf litter and

49 recalcitrant compounds, such as lignin, under constant laboratory conditions and

50 found that the decomposing abilities were variable among LDM species within a

51 climatic region but that the variability in decomposing ability was relatively

similar among climatic regions (Osono submitted). These results led to a further hypothesis that the decomposing ability of LDM originating from different climates varies in its sensitivity to temperature and litter quality. Several studies have examined the dependence of fungal decomposition on incubation temperature and/or the quality of leaf litter (Lindeberg 1946; Mikola 1956; Miyamoto et al. 2000; Osono and Takeda 2006; Osono et al. 2011c). To the knowledge of the author, however, few studies have explored chemical components that can limit fungal decomposition and compared the dependence of decomposition on temperature and litter types between major LDM originating from regions with different climatic conditions (Osono 2011).

The purpose of the present study was to assess under pure culture conditions the dependence of fungal growth and decomposition on litter types used as substrata, incubation temperature, and the origin of the isolate. I used six isolates of LDM species, five of which belonged to *Mycena*, obtained in subtropical, cool temperate, and subalpine forests in Japan because (i) they occurred as fruiting bodies most frequently on the forest floor of the respective forests (Osono in press b) and (ii) they exhibited the greatest activity of the LDM isolates to

decompose leaf litter (Osono submitted). I performed two pure culture tests that manipulated litter types and incubation temperature. First, a total of 39 litter types and six fungal isolates from the three climatic regions were used to examine possible limiting factors of fungal decomposition (denoted as the litter types test). Secondly, the pure culture decomposition tests were designed to evaluate the effects of litter type, incubation temperature, *Mycena* isolates of different origins, and the interactions of these factors on the decomposition of leaf litter and recalcitrant compounds in the litter (denoted as the cross inoculation test).

Materials and methods

Study site

Isolates of macrofungi and litter materials used in pure culture decomposition tests were collected at three sites in Japan: a subtropical (ST), a cool temperate (CT), and a subalpine (SA) forest. ST was located in an evergreen broadleaved forest in the northern part of Okinawa Island. CT was located in a deciduous

broadleaved forest in Kyoto. SA was located in an evergreen coniferous forest in Gifu. The mean annual temperature was 22°C, 10°C, and 2°C in ST, CT, and SA, respectively. The study sites received similar amounts of precipitation annually (approximately 2500 mm). Further details about the location, climatic condition, vegetation, and properties of the forest floor are given in Osono (in press a, in press b).

Fungal isolates

Six isolates were used, two isolates from each of the three study sites: two unidentified *Mycena* species (denoted as *Mycena* sp.1 MAFF241586 and *Mycena* sp.2 MAFF241604) from ST, *Mycena polygramma* IFO33011 and an unidentified *Clitocybe* species from CT, and *Mycena aurantiidisca* O2_07101503b and *M. epipterygia* O9_07101508 from SA. The *Mycena* species were the macrofungal taxa most frequently encountered as fruiting bodies at each study site (Osono in press b) and were active decomposers of leaf litter (Osono submitted). *Clitocybe* sp. fruited at low frequency in CT (Osono in press b) and was a selective decomposer

103 of recalcitrant compounds on the forest floor (Osono et al. 2011a). These
104 macrofungal isolates were obtained from tissues of fruiting bodies collected at the
105 study sites (Osono submitted), and maintained on slants of 1% malt extract agar
106 medium [MEA, malt extract 1% and agar 2% (w/v)] at 20°C in darkness until the
107 tests were performed. *Mycena* sp.1 and sp.2 from ST were analyzed for base
108 sequences of the rRNA gene 28S D1/D2 (accession numbers: AB512383 for
109 *Mycena* sp.1 and AB512392 for *Mycena* sp.2), but identification to species level
110 was not successful.

111

112 Litter materials

113

114 Newly shed leaves of 12, 15, and 12 plant species without obvious fungal or faunal
115 attack were collected from the surface of the forest floor in January to April 2008,
116 November 2000, and October 2009 in ST, CT, and SA, respectively (see Table S1 in
117 Electronic Supplementary Material) and used for the litter type test. For the cross
118 inoculation test, newly shed leaves of *Castanopsis sieboldii*, *Fagus crenata*, and
119 *Betula ermanii* without obvious fungal or faunal attack were collected from the

120 surface of the forest floor of ST, CT, and SA at peak periods of litter fall in April
121 2009, October 2009, and October 2009, respectively. The litters were oven-dried at
122 40°C for one week and preserved in vinyl bags for one to two months until the
123 experiments were started.

124

125 Litter type test

126

127 The effect of litter type on decomposition was examined under pure culture
128 conditions in the litter type test. Two isolates each from ST, CT, or SA were
129 inoculated onto 12, 15, or 12 litter types that differed in chemical composition
130 collected at ST, CT, or SA, respectively.

131 An individual pure culture decomposition test consisted of one fungal
132 isolate inoculated to one litter type. Litter (0.3 g) was sterilized by exposure to
133 ethylene oxide gas at 60°C for 6 hours and used in the tests, according to the
134 methods described in Osono et al. (2011c). The sterilized litters were placed on the
135 surface of Petri dishes (9 cm diameter) containing 20 ml of 2% agar. Inocula for
136 each assessment were cut out of the margin of previously inoculated Petri dishes

137 on 1% MEA with a sterile cork borer (6 mm diameter) and placed on the agar
138 adjacent to the litter, one plug per plate. The plates were incubated for 12 weeks
139 in the dark at 20°C. The plates were sealed firmly with laboratory film during
140 incubation so that moisture did not limit decomposition on the agar. After
141 incubation the litters were retrieved, oven-dried at 40°C for 1 week, and weighed.
142 The initial, undecomposed litters were also sterilized, oven-dried at 40°C for 1
143 week, and weighed to determine the original mass. Three to four plates were
144 prepared for each isolate, and four uninoculated plates served as a control. Mass
145 loss of litter was determined as a percentage of the original mass, taking the mass
146 loss of litter in the uninoculated and incubated control treatment into account,
147 and the mean values were calculated for each plate. Prior to the tests, the
148 sterilized litters were placed on 1% MEA, and after 8 weeks of incubation at 20°C
149 in darkness, no microbial colonies had developed on the plates. Thus, the
150 effectiveness of the sterilization method used in the present study was verified.
151 The initial litters of multiple tree species were analyzed for the contents of
152 acid-unhydrolyzable residues (AUR), total carbohydrates, extractives, and
153 nitrogen (N) as described below.

154

155 Cross-inoculation test

156

157 Cross-inoculation tests were carried out to examine the effects of litter type,
158 incubation temperature, and origin of *Mycena* isolates on fungal decomposition.
159 Three fungal isolates (*Mycena* sp.2 from ST, *M. polygramma* from CT, and *M.*
160 *aurantiidisca* from SA) were inoculated onto leaves of three litter types
161 (*Castanopsis sieboldii* from ST, *Fagus crenata* from CT, and *Betula ermanii* from
162 SA), and each combination of the three isolates × three litter types was incubated
163 at seven temperatures (5, 10, 15, 20, 25, 30, and 35°C).

164 Prior to this decomposition test, colony diameter growth rates were
165 measured for the isolates of *Mycena* sp.2, *M. polygramma*, and *M. aurantiidisca*.
166 Mycelial disks of the three isolates, 6 mm in diameter, were taken from the edge of
167 cultures on Petri dishes containing 1% MEA and incubated at 20°C for 2 weeks.
168 They were transferred to the center of another Petri dish (9 cm in diameter)
169 containing 20 ml 1% MEA. The plates were incubated at one of the seven
170 temperatures in the dark. Colony diameter was measured in two directions at

right angles from each other three to five times at given intervals during a 3-month incubation, and colony diameter growth rate was calculated by regressing the colony diameter against the days after inoculation. Four plates were prepared for each isolate.

The decomposition tests were performed as described above, except that the plates were incubated at one of the seven temperatures. The initial litter, the control litter, and the litter with at least 5.0% mass loss were used for the analysis of AUR as described below.

Chemical analyses

Litter materials were combined to make one sample for each test and ground in a laboratory mill (0.5 mm screen). The amount of AUR in the samples was estimated by means of gravimetry as acid-insoluble residue, using hot sulfuric acid digestion (King and Heath 1967). Samples were extracted with alcohol-benzene at room temperature (15-20°C), and the residue was treated with 72% sulfuric acid (v/v) for 2 h at room temperature with occasional stirring. The

mixture was diluted with distilled water so that the concentration of sulfuric acid reached 2.5% and autoclaved at 120°C for 60 min. After cooling, the residue was filtered and washed with water through a porous crucible (G4), dried at 105°C and weighed as acid-insoluble residue. The filtrate (autoclaved sulfuric acid solution) was subjected to total carbohydrate analysis. The amount of carbohydrate in the filtrate was estimated by means of the phenol-sulfuric acid method (Dubois et al. 1956). One milliliter of 5% phenol (v/v) and 5 ml of 98% sulfuric acid (v/v) were added to the filtrate. The optical density of the solution was measured using a spectrophotometer at 490 nm, using known concentrations of D-glucose as standards. Total N content was measured by automatic gas chromatography (NC analyzer SUMIGRAPH NC-900, Sumitomo Chemical Co., Osaka, Japan).

Mass loss of AUR was determined as a percentage of the original mass, taking the mass loss of AUR in the uninoculated and incubated control treatment into account. AUR/litter mass (AUR/L) loss ratio is a useful index of the selective delignification caused by each fungal species (Osono and Hirose 2009). AUR/L loss ratio of each fungal species was calculated using the equation:

$$\text{AUR/L loss ratio} = \text{mass loss of AUR (\% of original AUR mass)} / \text{mass loss}$$

205 of litter (% of original litter mass)

206

207 Statistical analysis

208

209 In the litter type test, Pearson's correlation coefficients were calculated for linear
210 relationships between chemical properties of litter and mass loss of the litter
211 caused by macrofungal isolates. In the cross inoculation test, factors affecting the
212 mass loss of litter were analyzed with generalized linear models (GLMs) with
213 fungal species, litter type, incubation temperature, and their interactions as
214 categorical predictors. Factors affecting the mass loss of AUR and AUR/L loss
215 ratio were also analyzed with GLMs with litter type and incubation temperature
216 as categorical predictors, but only the data of *Mycena* sp.2 were used because the
217 numbers of samples of *M. polygramma* and *M. aurantiidisca* used for AUR
218 analysis were too low. Tukey's HSD test was used for multiple comparisons. JMP
219 6.0 for Macintosh was used to perform these analyses.

220

221 **Results**

222

223 Litter mass loss in the litter type test

224

225 To examine the effect of litter types and possible effects of chemical components
226 on fungal decomposition, six fungal isolates were inoculated onto multiple litter
227 types that differed in chemical properties and incubated at 20°C, and the mass
228 loss of litter during the incubation was related to the chemical composition of the
229 litters. The initial content of AUR in leaf litter of a total of 39 litter types used in
230 the tests ranged from 8.8% to 45.6%, that of total carbohydrates from 20.1% to
231 44.9%, that of extractives from 5.3% to 20.6%, and that of N from 0.41% to 1.75%
232 (see Table S1).

233 Overall, the mean mass loss of litter caused by six fungal isolates ranged
234 from -2.8% to 50.7% (see Table S1), indicating that there were large variations of
235 decomposition between litter types caused by fungal isolates. Three significant
236 correlation coefficients were detected out of a total of 24 combinations (six fungal
237 isolates × four chemical properties) (Table 1). Mass loss of litter caused by *Mycena*
238 sp.2 from ST and *M. polygramma* from CT was significantly and negatively

correlated with AUR content in litter; and that caused by *M. epipterygia* from SA was significantly and negatively correlated with the content of extractives and significantly and positively correlated with N content (Fig. 1).

Colony diameter growth rate as related to temperature

The colony diameter growth rate increased linearly with temperature from 5 to 25°C for *Mycena* sp.2 from ST and from 5 to 20°C for *M. polygramma* from CT and *M. aurantiidisca* from SA (Fig. 2). The optimal growth rate occurred at 25°C for *Mycena* sp.2 and at 20°C for *M. polygramma* and *M. aurantiidisca*. The growth rate was lower at temperatures above the optimal temperature. At 25°C, *M. aurantiidisca* displayed more severely reduced growth than *M. polygramma*. *Mycena* sp.2 was the only isolate that grew at 30°C. No growth occurred at 35°C for the three isolates.

Litter mass loss in the cross inoculation test

Overall, the mean value of mass loss of litter ranged from -1.4% to 54.4% (Fig. 3, upper). Fungal isolate, litter type, incubation temperature, and their interactions significantly affected the mass loss of litter during the incubation (Table 2A). The mass loss of litter was in the order: *Mycena* sp.2 > *M. polygramma* > *M. aurantiidisca*, and *Betula* > *Castanopsis*, *Fagus*. The mass loss of litter increased with temperature: from 5°C to 25°C for *Mycena* sp.2 and *M. polygramma* and from 10°C to 20°C for *M. aurantiidisca*. The greatest values of mass loss were found at 25°C for *Mycena* sp.2 and at 20°C and 25°C for *M. polygramma*, and at 20°C for *M. aurantiidisca*, which were generally consistent with the optimum temperatures of colony diameter growth rate of these isolates (Fig. 2). Mass loss of litter was negligible at 5°C for all three isolates and at 35°C for *Mycena* sp.2, at 30°C to 35°C for *M. polygramma*, and at 25°C to 35°C for *M. aurantiidisca* (Fig. 3).

Incubation temperature affected the mass loss of litter caused by three *Mycena* isolates, and the interactions between temperature and fungal isolate and between temperature and litter type were also significant (Fig. 3; Table 2), indicating that the temperature effect depended on the fungal isolates and litter types. For example, the changes in the mass loss of litter in relation to

273 temperature were larger for *Mycena* sp.2, which had greater ability to cause mass
274 loss than *M. polygramma* and *M. aurantiidisca*, while these changes were less
275 obvious when the fungal isolates were inoculated onto *Fagus* litter, which was
276 more recalcitrant than *Castanopsis* and *Betula* litters.

277

278 AUR decomposition in the cross inoculation test

279

280 The litters with more than 5.0% mass loss were analyzed for the mass loss of AUR.

281 The mean value of mass loss of AUR ranged from 2.6% to 68.9% (Fig. 3, middle).

282 Litter type and incubation temperature significantly affected the mass loss of

283 AUR caused by *Mycena* sp.2 during the incubation (Table 2B). Mass loss of AUR

284 by *Mycena* sp.2 was lower at 10°C than at 15 to 30°C and was higher on *Betula*

285 than on *Fagus* and *Castanopsis*.

286 The mean values of AUR/L loss ratio ranged from 0.3 to 2.6 (Fig. 3, lower).

287 Litter type significantly affected AUR/L loss ratio for *Mycena* sp.2, whereas

288 incubation temperature had no significant effect on AUR/L loss ratio (Table 2B).

289

Discussion

The decomposition of litter by *Mycena* isolates varied with litter type (Tables S1, 2), and the extent of decomposition in various litter types was correlated negatively with the contents of AUR or extractives and positively with N content (Fig. 1; Table 1). Recalcitrant compounds in leaves designated as AUR, such as lignin and tannin, have often been shown to limit the rate of decomposition in forest soils (Geng et al. 1993; Stump and Binkley 1993; Murphy et al. 1998; Osono and Takeda 2005) and of litter decomposition by ligninolytic fungi under pure culture conditions (Osono et al. 2011b, 2011c; Hagiwara et al. 2012). Extractives include soluble polyphenols, hydrocarbons, and pigments that are often released rapidly from decomposing litter (Osono et al. 2014), and can include inhibitory substances for fungal growth (Hughes et al. 2007). Nitrogen is a major essential element limiting fungal growth and enzyme production and was found to be positively related to mass loss caused by ligninolytic fungi (Osono et al. 2011b).

The findings that growth and decomposition were maximal at 20°C and 25°C (Figs. 2 and 3), respectively, indicate that the *Mycena* isolates used are

mesophilic. The isolate from cooler regions was more sensitive to higher temperature than the isolate from warmer regions (Fig. 2), suggesting physiological adaptations of these fungi to the climatic conditions of the respective study sites (subtropical versus cool temperate versus subalpine). The optimal temperature of hyphal growth was generally consistent with that of decomposition, which agrees with the finding of Osono et al. (2011c) for three isolates of *Xylaria* sp. from CT.

Incubation temperature affected the mass loss of litter by the three *Mycena* isolates and the mass loss of AUR by *Mycena* sp.2 (Fig. 3; Table 2). However, AUR/L loss ratio of *Mycena* sp.2 was not affected by incubation temperatures (Table 2), suggesting that the degree of selective AUR decomposition by *Mycena* sp.2 was insensitive to the range of temperature adopted in the present study. Previously, Adaskaveg et al. (1995) and Osono et al. (2011c) reported that at temperatures above the optimum growth temperatures, the ligninolytic activity of fungi increased at the expense of cellulolytic activity, resulting in suppressed overall decomposition of leaf litter. My finding here for *Mycena* sp.2 from ST suggested that the sensitivity of selective delignification to

temperature could vary with fungal species. Osono and Takeda (2006) also showed that the degree of selective delignification caused by four ligninolytic fungi did not differ between incubation at 10°C or 20°C, whereas *Gymnopus dryophilus* decomposed AUR more selectively at 10°C than at 20°C. More studies are needed to evaluate the sensitivity to temperature of ligninolytic activity of litter-decomposing macrofungi, including those from cooler regions, such as *M. polygramma* and *M. aurantiidisca*.

The three *Mycena* isolates from different climates differed significantly in their ability to decompose litter and in AUR and AUR/L loss ratio (Fig. 3, Table 2). This difference in decomposing ability was within the range of variability found for a suite of *Mycena* species encountered at these sites. That is, Osono (submitted) compared the ability of 32 *Mycena* isolates from ST, CT, and SA to decompose leaf litter and demonstrated a similar variation in decomposing ability among the climatic regions. The results of the present study were consistent with those of Osono (submitted) showing that some *Mycena* species generally are active decomposers of AUR. *Mycena polygramma* IFO33011 from CT is an outlier as this isolate has been reported to decompose cellulose selectively over AUR

341 (Osono and Takeda 2002; Osono et al. 2003).

342 The effects of litter type, origin of isolate, and temperature on the
343 decomposition of leaf litter by ligninolytic *Mycena* and other LDM may have
344 implications regarding the changes in fungal decomposition of leaf litter in
345 relation to climatic conditions. First, the variation of decomposition of litter
346 between the fungal isolates was larger than the variations with temperature and
347 litter type (Table 2), suggesting that a shift in fungal species composition can
348 affect the decomposition at the level of LDM assemblages more than changes in
349 temperature and/or litter. Studying the species composition of fungal assemblages,
350 decomposing abilities of individual fungal species, and their geographical
351 distributions is thus crucial for predicting the response of fungal decomposition to
352 possible climate changes. However, studies on the geographical distribution of
353 litter-decomposing fungi are still scarce (e.g. Iwamoto and Tokumasu 2001;
354 Tokumasu 2001; Hosoya et al. 2010). Further studies are needed to examine the
355 geographical distribution of fungi in conjunction with their decomposing abilities.

356 Secondly, the ability of individual *Mycena* species to decompose litter and
357 AUR also varied with temperature and litter type (Table 2; Fig. 1). This suggests

that the functioning of LDM could change along with a possible future increase in temperature worldwide, which in temperate regions is expected to be an increase of 2-3°C (Manabe et al. 1991; Boer et al. 1992; Russell et al. 1995) and/or with concomitant changes in vegetation (Tsukada 1983; Matsui et al. 2004). For example, an increase of litter temperature to optimum growth temperatures could lead to enhanced decomposition by LDM, resulting in positive feedback to the atmospheric CO₂ level. An increase in the relative abundance of tree species with low AUR content in litter might have a similar positive feedback. These predictions are obviously oversimplified, but data such as the results of the present study will provide useful insights into the possible effects of future climate changes on fungal decomposition in forest soils.

Acknowledgments I thank Mr. Y. Hagiwara for help with chemical analyses; and Dr. Elizabeth Nakajima for critical reading of the manuscript. This study received partial financial support from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (MEXT) (No. 19780114), The Sumitomo Foundation, Nissan Global Foundation, Nippon Life Inst. Foundation, and the

375 Grants for Excellent Graduate Schools, MEXT, Japan (12-01) to Kyoto University.

376

377 **References**

378

379 Adaskaveg JE, Gilbertson RL, Dunlap MR (1995) Effects of incubation time and

380 temperature on in vitro selective delignification of silver leaf oak by

381 *Ganoderma colossum*. Appl Environ Microbiol 61:138-144

382 Boberg JB, Ihrmark K, Lindahl BD (2011) Decomposing capacity of fungi

383 commonly detected in *Pinus sylvestris* needle litter. Fungal Ecol

384 4:110-114

385 Boer GJ, McFarlane NA, Lazare M (1992) Greenhouse gas-induced climate

386 change simulated with the CCC second-generation general circulation

387 model. J Climate 5:1045-1077

388 Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F (1956) Colorimetric

389 method for determination of sugars and related substances. Anal Chem

390 28:350-356

391 Geng X, Pastor J, Dewey B (1993) Decay and nitrogen dynamics of litter from

- 392 disjunct, congeneric tree species in old-growth stands in northeastern
393 China and Wisconsin. *Can J Bot* 71:693–699
- 394 Hagiwara Y, Osono T, Ohta S, Agus W, Hardjono A (2012) Colonization and
395 decomposition of leaf litter by ligninolytic fungi in *Acacia mangium*
396 plantations and adjacent secondary forests. *J For Res* 17:51-57
- 397 Hosoya T, Hirose D, Fujisaki M, Osono T, Kubono T, Tokumasu S, Kakishima M
398 (2010) Internal transcribed spacer haplotype diversity and their
399 geographical distribution in *Dasyscyphella longistipitata*
400 (Hyaloscyphaceae, Helotiales) occurring on *Fagus crenata* cupules in
401 Japan. *Mycoscience* 51:116-122
- 402 Hughes KA, Bridge P, Clark MS (2007) Tolerance of Antarctic soil fungi to
403 hydrocarbons. *Sci Total Environ* 372:539-548
- 404 Iwamoto S, Tokumasu S (2001) Dematiaceous hyphomycetes inhabiting decaying
405 blackish needles of *Abies firma* and their distribution in the Kanto
406 district, Japan. *Mycoscience* 42:273-279
- 407 King HGC, Heath GW (1967) The chemical analysis of small samples of leaf
408 material and the relationship between the disappearance and

- 409 composition of leaves. *Pedobiologia* 7:192-197
- 410 Lindahl B, Boberg J (2008) Distribution and function of litter basidiomycetes in
411 coniferous forests. In: Boddy L, Frankland JC, van West P (eds) *Ecology*
412 of Saprotrophic Basidiomycetes. Academic Press, London, pp. 183-196
- 413 Lindeberg G (1946) On the decomposition of lignin and cellulose in litter caused
414 by soil-inhabiting hymenomycetes. *Ark Bot* 33a:1-16
- 415 Manabe S, Nissinen A, Erhard M, Bryan K (1991) Transient responses of a
416 coupled ocean-atmosphere model to gradual changes of atmospheric CO₂.
417 Part I: Annual mean response. *J Climate* 4:785-818
- 418 Matsui T, Yagihashi T, Nakaya T, Taoda H, Yishinaga S, Daimaru H, Tanaka N
419 (2004) Probability distributions, vulnerability and sensitivity in *Fagus*
420 *crenata* forests following predicted climate changes in Japan. *J Veg Sci*
421 15:605-614
- 422 Mikola P (1956) Studies on the decomposition of forest litter by basidiomycetes.
423 *Commun Inst For Fenn* 48:4-48
- 424 Miyamoto T, Igarashi T, Takahashi K (2000) Lignin-degrading ability of
425 litter-decomposing basidiomycetes from *Picea* forests of Hokkaido.

- 426 Mycoscience 41:105-110
- 427 Murphy KL, Klopatek JM, Klopatek CC (1998) The effects of litter quality and
- 428 climate on decomposition along an elevational gradient. Ecol Appl
- 429 8:1061–1071
- 430 Osono T (2007) Ecology of ligninolytic fungi associated with leaf litter
- 431 decomposition. Ecol Res 22:955-974
- 432 Osono T (2011) Diversity and functioning of fungi associated with leaf litter
- 433 decomposition in Asian forests of different climatic regions. Fun Ecol
- 434 4:375-385
- 435 Osono T. Mycelial biomass in the forest floor and soil of subtropical, temperate,
- 436 and subalpine forests. J For Res in press a
- 437 Osono T. Diversity, resource utilization, and phenology of fruiting bodies of
- 438 litter-decomposing macrofungi in subtropical, temperate, and subalpine
- 439 forests. J For Res in press b
- 440 Osono T. Decomposing ability of diverse litter-decomposer macrofungi in
- 441 subtropical, temperate, and subalpine forests. J For Res submitted
- 442 Osono T, Takeda H (2002) Comparison of litter decomposing ability among diverse

- 443 fungi in a cool temperate deciduous forest in Japan. *Mycologia*
- 444 94:421-427
- 445 Osono T, Takeda H (2005) Decomposition of organic chemical components in
- 446 relation to nitrogen dynamics in leaf litter of 14 tree species in a cool
- 447 temperate forest. *Ecol Res* 20:41-49
- 448 Osono T, Takeda H (2006) Fungal decomposition of *Abies* needle and *Betula* leaf
- 449 litter. *Mycologia* 98:172-179
- 450 Osono T, Hirose D (2009) Effects of prior decomposition of *Camellia japonica* leaf
- 451 litter by an endophytic fungus on the subsequent decomposition by
- 452 fungal colonizers. *Mycoscience* 50:52-55
- 453 Osono T, Fukasawa Y, Takeda H (2003) Roles of diverse fungi in larch needle litter
- 454 decomposition. *Mycologia* 95:820-826
- 455 Osono T, Hobara S, Hishinuma T, Azuma JI (2011a) Selective lignin
- 456 decomposition and nitrogen mineralization in forest litter colonized by
- 457 *Clitocybe* sp. *Eur J Soil Biol* 47:114-121
- 458 Osono T, To-Anun C, Hagiwara Y, Hirose D (2011b) Decomposition of wood, petiole
- 459 and leaf litter by *Xylaria* species from northern Thailand. *Fun Ecol*

- 460 4:210-218
- 461 Osono T, Hagiwara Y, Masuya H (2011c) Effects of temperature and litter type on
- 462 fungal growth and decomposition of leaf litter. *Mycoscience* 52:327-332
- 463 Osono T, Azuma JI, Hirose D (2014) Plant species effect on the decomposition and
- 464 chemical changes of leaf litter in grassland and pine and oak forest soils.
- 465 *Plant Soil* 376:411-421
- 466 Russell GL, Miller JR, Rind D (1995) A coupled atmosphere-ocean model for
- 467 transient climate change studies. *Atmos-Ocean* 33:683-730
- 468 Steffen KT, Cajthaml T, Šnajdr J, Baldrian P (2007) Differential degradation of
- 469 oak (*Quercus petraea*) leaf litter by litter-decomposing basidiomycetes.
- 470 *Res Microbiol* 158:447-455
- 471 Stump LM, Binkley D (1993) Relationships between litter quality and nitrogen
- 472 availability in Rocky Mountain forests. *Can J For Res* 23:492-502
- 473 Tokumasu S (2001) Geographical distribution of *Sporidesmium goidanichii* in pine
- 474 forests of Japan. *Mycoscience* 42:575-589
- 475 Tsukada M (1983) Vegetation and climate during the Last Glacial Maximum in
- 476 Japan. *Quatern Res* 19:212-235

- 477 Valášková V, Šnajdr J, Bittner B, Cajthaml T, Merhautová V, Hofrichter M,
478 Baldrian P (2007) Production of lignocellulose-degrading enzymes and
479 degradation of leaf litter by saprotrophic basidiomycetes isolated from a
480 *Quercus petraea* forest. Soil Biol Biochem 39:2651-2660
- 481 Van der Wal A, Geydan TD, Kuyper TW, de Boer W (2013) A thready affair:
482 linking fungal diversity and community dynamics to terrestrial
483 decomposition processes. FEMS Microbiol Rev 37:477-494
- 484 Žifčáková L, Dobiášová P, Kolářová Z, Koukol O, Baldrian P (2011) Enzyme
485 activities of fungi associated with *Picea abies* needles. Fungal Ecol
486 4:427-436

Osono Table 1

Table 1. Pearson's correlation coefficients for linear relationship between contents of organic chemical components and nitrogen in leaf litter and mass loss of leaf litter caused by isolates of macrofungi. *** P<0.001, * P<0.05, ns not significant.

Fungus		Number of tree species used	AUR		Total carbohydrates		Extractives		Nitrogen	
ST	<i>Mycena</i> sp.1	12	-0.51	ns	0.06	ns	0.05	ns	0.21	ns
	<i>Mycena</i> sp.2	12	-0.78	**	0.02	ns	0.13	ns	0.35	ns
CT	<i>Mycena polyramma</i>	15	-0.80	***	0.26	ns	0.17	ns	-0.25	ns
	<i>Clitocybe</i> sp.1	15	-0.36	ns	0.22	ns	-0.27	ns	-0.14	ns
SA	<i>Mycena aurantiidisca</i>	12	0.21	ns	0.14	ns	-0.52	ns	0.52	ns
	<i>Mycena epipterygia</i>	12	0.29	ns	0.15	ns	-0.60	*	0.57	*

Osono Table 2

Table 2. Results of generalized linear models (GLMs) examining the effect of fungal species, litter species, and temperature on litter mass loss (A), and examining the effect of litter species and temperature on AUR mass loss and AUR/litter mass (AUR/L) loss ratio for *Mycena* sp.2. *** P<0.001, ** P<0.01, * P<0.05, ns not significant.

(A) Litter mass loss	d.f.	χ^2	
Model	62	504.7	***
Fungal isolate	2	313.8	***
Litter type	2	100.5	***
Temperature	6	280.1	***
Fungal isolate × Litter type	4	25.5	***
Fungal isolate × Temperature	12	251.9	***
Litter type × Temperature	12	63.5	***
Fungal isolate × Litter type × Temperature	24	37.6	*

(B)	AUR mass loss			AUR/L loss ratio		
	d.f.	χ^2		d.f.	χ^2	
Model	6	41.2	***	6	13.1	*
Litter type	2	22.7	***	2	7.2	*
Temperature	4	37.3	***	4	8.6	ns

1 Figure legends.

2

3 Fig. 1. Mass loss of leaf litter caused by isolates of macrofungi as related to
4 contents of organic chemical components and nitrogen. ST, mass loss caused by
5 *Mycena* sp.2 was negatively correlated with AUR content in leaf litter of 12 tree
6 species. CT, mass loss caused by *Mycena polygramma* was negatively correlated
7 with AUR content in leaf litter of 15 tree species. SA, mass loss caused by *Mycena*
8 *epipterygia* was negatively correlated with extractive content and positively
9 correlated with nitrogen content in 12 litter types. Correlation coefficients for the
10 relationships are shown in Table 1.

11

12 Fig. 2. Colony diameter growth rate of isolates of *Mycena* sp.2 (□, from ST), *M.*
13 *polygramma* (●, from CT), and *M. aurantiidisca* (▲, from SA) as related to
14 temperature. Bars indicate standard errors (n=4).

15

16 Fig. 3. Mass loss of leaf litter (upper) and AUR (middle) and AUR/litter (AUR/L)
17 mass loss rate (lower) in leaf litter of *Castanopsis sieboldii* (□, from ST), *Fagus*
18 *crenata* (●, from CT), and *Betula ermanii* (▲, from SA) caused *in vitro* by isolates
19 of *Mycena* sp.2 (left, from ST), *M. polygramma* (center, from CT), and *M.*
20 *aurantiidisca* (right, from SA) at seven temperatures for 12 weeks in the dark.
21 Bars indicate standard errors (n=3 or 4).

22

1 Osono Fig. 1

1 Osono Fig. 2

3

1 Osono Fig. 3

Electronic Supplementary Material

Effects of litter type, origin of isolate, and temperature on decomposition of leaf litter by macrofungi

Takashi Osono

Table S1. Contents of organic chemical components and nitrogen (% w/w) and mass loss of leaf litter (% original mass) caused by isolates of macrofungi in leaf litter of multiple litter types. AUR, acid unhydrolyzable residue; TCH, total carbohydrates; Extr, extractives; N, nitrogen.

Litter type	AUR	TCH	Extr	N	Mass loss	Mass loss
Subtropical forest					<i>Mycena sp.1</i>	<i>Mycena sp.2</i>
<i>Schefflera octophylla</i>	21.7	32.0	20.6	0.69	35.6	39.3
<i>Sapium japonicum</i>	13.0	43.5	9.3	0.58	35.3	50.7
<i>Dendropanax trifidus</i>	22.0	35.6	11.3	0.55	33.1	49.6
<i>Malotus japonicus</i>	17.0	33.8	10.2	1.03	32.1	43.4
<i>Heterosmilax japonica</i>	40.0	36.7	5.9	0.72	28.5	33.0
<i>Pinus luchuensis</i>	41.2	44.9	6.0	0.41	26.4	22.3
<i>Schima wallichii</i>	35.8	29.7	10.7	0.52	26.3	34.9
<i>Castanopsis sieboldii</i>	26.2	34.5	11.2	0.62	23.1	30.8
<i>Syzygium buxifolium</i>	41.1	27.0	11.8	0.47	22.5	23.7
<i>Podocarpus macrophyllus</i>	33.1	43.3	6.5	0.47	19.9	21.4
<i>Quercus miyagii</i>	37.8	30.3	14.9	0.71	14.3	21.0
<i>Daphniphyllum teijsmannii</i>	26.1	39.2	12.5	0.61	12.0	23.7
Cool temperate forest					<i>Mycena polygramma</i>	<i>Clitocybe sp.</i>
<i>Malotus japonicus</i>	15.4	27.1	17.3	1.20	34.0	3.9
<i>Benthamidia kousa</i>	8.8	32.8	11.3	1.04	33.9	8.9
<i>Carpinus laxiflora</i>	17.8	32.6	11.5	1.25	28.3	10.2
<i>Acer micranthum</i>	15.3	32.3	12.5	0.78	27.7	9.6

<i>Quercus serrata</i>	37.4	26.3	8.0	0.94	21.1	5.4
<i>Betula grossa</i>	36.2	26.7	7.0	1.12	19.9	11.8
<i>Weigela hortensis</i>	35.5	28.6	9.9	0.95	17.8	0.4
<i>Acer rufinerve</i>	33.7	27.0	15.0	0.78	15.1	8.0
<i>Castanea crenata</i>	32.2	29.6	8.6	1.14	12.3	9.9
<i>Quercus crispula</i>	35.6	29.6	9.7	0.99	11.2	6.6
<i>Malus tschonoskii</i>	25.1	37.6	12.4	0.79	9.6	2.5
<i>Magnolia obovata</i>	31.0	31.4	10.4	1.04	7.1	9.5
<i>Fagus crenata</i>	41.5	28.9	8.8	1.68	4.2	1.1
<i>Pterocarya rhoifolia</i>	35.9	25.0	11.5	1.75	3.4	4.8
<i>Aesculus turbinata</i>	38.0	23.5	15.1	0.89	1.9	1.1
Subalpine forest					<i>Mycena aurantiidisca</i>	<i>Mycena epipterygia</i>
<i>Betula ermanii</i>	45.6	29.1	8.2	1.10	36.7	30.4
<i>Sorbus commixta</i>	30.6	33.0	11.2	1.00	35.9	29.8
<i>Sorbus japonicum</i>	31.9	37.0	8.5	1.38	22.2	27.2
<i>Reynoutria japonica</i>	44.8	31.1	5.3	0.81	20.3	20.5
<i>Aralia cordata</i>	23.9	25.9	9.6	1.03	16.6	13.4
<i>Viburnum furcatum</i>	29.8	36.5	9.1	1.19	12.9	24.7
<i>Sasa kurilensis</i>	23.1	41.1	7.3	0.48	8.3	6.6
<i>Salix sachaliensis</i>	42.6	20.1	14.4	1.01	7.7	15.3
<i>Pinus pentaphylla</i>	42.1	36.1	9.3	0.43	6.3	17.6
<i>Picea jezoensis</i> var. <i>hondoensis</i>	34.4	36.3	17.6	0.82	1.1	-1.1
<i>Tsuga diversiflora</i>	38.5	25.1	18.1	0.90	0.9	3.7
<i>Cercidiphyllum japonicum</i>	25.7	23.9	12.0	0.62	-2.8	4.1