TITLE:

High contribution of Synechococcus to phytoplankton biomass in the aphotic hypolimnion in a deep freshwater lake (Lake Biwa, Japan)

AUTHOR(S):

Takasu, Hiroyuki; Ushio, Masayuki; LeClair, E.
LeClair; Nakano, Shin-ichi

CITATION:

Takasu, Hiroyuki ...[et al]. High contribution of Synechococcus to phytoplankton biomass in the aphotic hypolimnion in a deep freshwater lake (Lake Biwa, Japan). Aquatic Microbial Ecology 2015, 75(1): 69-79

ISSUE DATE:

2015-05-06

URL:

<http://hdl.handle.net/2433/200804>

RIGHT:

© 2015 Inter-Research; The full-text file will be made open to the public on May 06, 2016 in accordance with publisher's 'Terms and Conditions for Self-Archiving'. ; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。 ; This is not the published version. Please cite only the published version.

Title page:

Original article for *Aquatic Microbial Ecology*

Title: High contribution of *Synechococcus* to phytoplankton biomass in the aphotic hypolimnion in a deep freshwater lake (Lake Biwa, Japan)

Hiroyuki Takasu ^{1,2}, Masayuki Ushio ^{1,3}, Jessica E. LeClair ¹ and Shin-ichi Nakano ^{1*}

Affiliations: ¹ *Center for Ecological Research (CER), Kyoto University, Hirano-cho 2 chome, 509-3, Otsu, Shiga 520-2113, Japan*

² Present address: *Atmosphere and Ocean Research Institute (AORI), The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, Chiba 277-8564, Japan*

³ Present address: Department of Environmental Solution Technology, Faculty of Science and Technology, Ryukoku University, 1-5 Yokoya, Seta Oe-cho, Otsu 520-2194, Japan

*Correspondence to: Shin-ichi Nakano: Mailing address: Center for Ecological Research (CER), Kyoto University, Hirano-cho 2 chome, 509-3, Otsu, Shiga 520-2113, Japan.

Phone and Fax: +81-77-549-8239, +81-77-549-8201

E-mail: nakano@ecology.kyoto-u.ac.jp

Running title: *Synechococcus* in the hypolimnion of Lake Biwa.

Key words: *Synechococcus*, vertical export, hypolimnion, Lake Biwa

27 **Abstract**

28 The effective transport of picophytoplankton to the mesopelagic layer in the ocean by cell
29 aggregation and attachment to large particles has been reported. Those findings suggest that
30 picophytoplankton play important roles in ecological processes in the deep ocean. In
31 contrast, there is no information about vertical transportation of picophytoplankton cells
32 from epilimnion in lakes, though the presence of picophytoplankton cells in hypolimnion
33 have been reported. The present study demonstrated the possible importance of
34 *Synechococcus* (Cyanobacteria) in ecological processes of the hypolimnion in the deep
35 mesotrophic Lake Biwa, Japan. The chlorophyll *a* concentration in the 0.2–2.0- μ m fraction,
36 which is mainly derived from *Synechococcus*, accounted for a large portion (up to 28.8%)
37 of the total chlorophyll *a* concentration in the hypolimnion during the thermal stratification
38 period. We found a significant positive correlation between *Synechococcus* abundances in
39 the epilimnion and hypolimnion during the stratification period. In addition, our incubation
40 experiment revealed that *Synechococcus* did not show remarkable growth during the first 2
41 days in dark conditions. These results suggest the recent delivery of a significant fraction of
42 *Synechococcus* cells from the epilimnion to the hypolimnion. Our results indicate that the
43 abundance of *Synechococcus* makes a greater contribution to ecological processes in the
44 hypolimnion of Lake Biwa than previously hypothesized, and this may also be the case for
45 other deep lakes.

46

Introduction

In pelagic ecosystems, a considerable amount of phytoplankton production is lost through respiration, the release of extracellular organic matter, grazing and lysis mortalities, sedimentation, and physiological death (Bidle & Falkowski 2004, Reynolds 2006). It has been suggested that the mechanisms of production and loss differ between large and small phytoplankton species (Kjørboe 1993), and thus phytoplankton species play different ecological roles in food webs and/or matter cycling (Reynolds 2006). In general, large phytoplankton species ($>20\ \mu\text{m}$) are less vulnerable to zooplankton grazing, and sink faster, than small phytoplankton species (Reynolds 2006). Consequently, large phytoplankton is thought to play important roles in organic matter transportation from the surface to deep aphotic layers in pelagic ecosystems. In contrast, small phytoplankton species ($<20\ \mu\text{m}$) are vulnerable to zooplankton grazing, and previous studies have concluded that most of their production is readily removed from surface layers (Nagata 1988, Nagata et al. 1994, Hirose et al. 2008, Scanlan 2012).

The genus *Synechococcus* (Cyanobacteria) is widely distributed in surface oceans (Stockner 1988). Their cell length is $0.9\ \mu\text{m}$, on average, and they are one of the groups with the smallest size in phytoplankton communities (Kirchman 2008). *Synechococcus* included picophytoplankton sink so slowly (no faster than $0.01\text{--}0.02\ \mu\text{m s}^{-1}$) that the motion of the water is believed to keep them in suspension (Reynolds 2006). Thus, they are too small to sink to the mesopelagic layer of the ocean. However, a recent study using inverse modelling and network analyses suggested that of picophytoplankton carbon biomass, including *Synechococcus*, is transported from the surface to mesopelagic layers (Richardson & Jackson 2007). Recently, the effective transport of *Synechococcus* cells to the mesopelagic layer by cell aggregation (Lomas & Moran 2011) and attachment to large particles (Sohrin et al. 2011) was reported. Those findings suggest that *Synechococcus* play

an important role in ecological processes in the mesopelagic layer of the ocean.

Synechococcus are also distributed in freshwater lakes worldwide (Callieri et al. 2012).

In deep freshwater lakes, the hypolimnion is separated from the epilimnion by the thermic barrier of thermocline. Stockner (1991) pointed out the possibility of transportation of picocyanobacteria from the epilimnion to the hypolimnion. However, unlike studies on vertical transportation of picocyanobacteria in oceans, there is no information about vertical transportation of *Synechococcus* cells to the hypolimnion in lakes. In our previous study, we found picophytoplankton cells in water samples from the hypolimnion (70 m) in deep, mesotrophic Lake Biwa, Japan, during the stratification period (Takasu et al. 2012). Callieri & Pinolini (1995) also reported that picophytoplankton cells were present in the hypolimnion (deeper than 100 m) of the deep Lake Maggiore, Northern Italy. Therefore, vertical transportation of *Synechococcus* from the epilimnion to the hypolimnion in lakes is also possible, similar to vertical transportation in oceans.

In the present study, we hypothesized that *Synechococcus* produced in the epilimnion of Lake Biwa are transported to the hypolimnion. To address this, we assessed the vertical distribution of, and seasonal changes in, *Synechococcus* abundance throughout the water column of Lake Biwa. To verify the major source of the *Synechococcus* population in the hypolimnion of the lake, we also evaluated the growth potential of *Synechococcus* in dark conditions. The results of the present study suggest that a significant fraction of *Synechococcus* cells are transported from the epilimnion to the hypolimnion.

Materials and methods

Sampling

Lake Biwa is a large (surface area: 674 km²), deep (maximum depth: 104 m), monomictic, and mesotrophic lake located in the central part of Honshu Island, Japan. We collected water samples at station Ie-1 (35° 12' 58'' N, 135° 59' 55'' E; ca. 75 m) in the north basin

of the lake.

Samples for assessment of the *Synechococcus* distribution were collected from April through August 2011. Vertical profiles of water temperature were determined using a CTD probe (SBE 911 plus; Sea Bird Electronics, Bellevue, WA, USA). In April, light intensity was measured using an LI-192 underwater quantum meter connected to an LI-1400 data logger (Li-Cor Inc., Lincoln, NE, USA). Secchi disk depth was measured throughout the study period. Samples for chlorophyll *a* analysis were collected at 0, 5, 10, 15, 30, and 70 m using Niskin X bottles and then poured into 500-mL polycarbonate bottles washed with 1.2 M HCl. Samples for *Synechococcus* enumeration were collected at 0, 5, 10, 20, 50, and 70 m, then poured into 100-mL polypropylene bottles and fixed immediately with glutaraldehyde (Wako Pure Chemical Co., Tokyo, Japan) to a final concentration of 1%. In April, samples for *Synechococcus* enumeration were collected at 5 and 70 m. In July, a sample for the incubation experiment was also collected at 5 m using a 10-L acrylic water sampler. For the dilution experiment, approximately 10 L of lake water were poured into acid-washed 10-L polyethylene bags.

Chlorophyll a

To determine chlorophyll *a* concentrations, 100-mL water samples were filtered through 0.2- and 2.0- μ m polycarbonate filters (Whatman International, Ltd., Maidstone, England) and analysed using the *N,N*-dimethylformamide (Wako Pure Chemical Co., Tokyo, Japan) method (Moran & Porath 1980) with a fluorescence spectrometer (RF-5300PC; Shimadzu, Kyoto, Japan). Chlorophyll *a* concentrations in the 0.2–2.0- μ m fraction (hereafter “pico-sized fraction”) were calculated according to the following equations:

$$\text{Chlorophyll } a \text{ in pico-sized fraction} = \text{chlorophyll } a \text{ concentration from 0.2-}\mu\text{m filter} - \text{chlorophyll } a \text{ concentration from 2.0-}\mu\text{m filter}$$

Enumeration of Synechococcus cells

Fixed water samples of 15- to 25-mL were filtered through 0.2- μ m-pore-size black polycarbonate filters (Advantec, Tokyo, Japan), and *Synechococcus* cells retained on the filters were counted by epifluorescence microscopy (BX51, Olympus, Japan) using both blue (460–490-nm excitation by U-MWB2, Olympus) and green (520–550-nm excitation by U-WIG2, Olympus) excitation filter sets. Eukaryotic picophytoplankton exhibited red fluorescence when excited by blue light and weak (red) or no fluorescence under green light (MacIsaac & Stockner 1993). Two types of *Synechococcus* pigments have been described, differing in terms of the phycoerythrin (PE) and phycocyanin (PC) content of phycobiliproteins. PE- and PC-rich *Synechococcus* respectively exhibited orange and dull red fluorescence when excited by blue light, and fluoresced orange and red under green light (MacIsaac & Stockner 1993). These fluorescence characteristics allowed us to separately enumerate the three types of picophytoplankton. We counted at least 300 cells or 100 fields to estimate cell abundance. Images of *Synechococcus* cells were captured at 1000 $\times$ magnification under an epifluorescence microscope equipped with a digital-camera (EOS Kiss X5, Canon, Tokyo, Japan). Digital images were used to determine the length, width, and fluorescence intensity of each cell, and more than 100 *Synechococcus* cells were used for each sample. The image analysis software ImageJ (National Institutes of Health) was used for measurement. Cell volumes were calculated by assuming that the cells were spheres. The cell specific orange fluorescence intensity under green excitation was also determined. PE-rich *Synechococcus* often predominates in Lake Biwa during summer, and the isolated strains exclusively exhibited a strong emission peak of PE (577 nm) under green excitation (546 nm) (Maeda et al. 1992). So, we measured fluorescence intensity of orange cells under the green excitation (520–550-nm excitation) as an indicator of cell specific PE fluorescence.

In addition, we microscopically observed *Synechococcus* microcolonies (from 5 to 50 cells), an aggregation without a clear separation from the single-celled *Synechococcus* (Callieri 2010). So, we also measured the size of the microcolonies in the same manner as individual *Synechococcus* cells.

Incubation experiment in dark conditions

A water sample was gently filtered through 20- μ m mesh to remove mesozooplankton. A 50-L portion of the filtrate was gravity filtered through 0.2- μ m filter cartridges (PALL Acropak Supor membrane capsules, PALL, Co., MI, USA) and collected into tanks. The 0.2- μ m filtrate was then passed through a 30-kDa tangential flow filtration system (PES membrane, Millipore, Co., MA, USA) to prepare a grazer-and-virus-free diluent. To reduce grazing and viral lysis pressure, the 20- μ m filtrate was diluted in 30-kDa diluent to 20% in a 5-L polycarbonate bottle washed with 1.2 M HCl before use. The bottle was then incubated for 48 h at the *in situ* temperature, in the dark. At the beginning (0 h) and end of the incubation (48 h), 50-mL subsamples for the enumeration of *Synechococcus* cells were collected into polypropylene tubes and immediately fixed with glutaraldehyde at a final concentration of 1%. The growth rate (μ , d⁻¹) of *Synechococcus* was calculated from the cell numbers at the beginning and end of the incubation experiment, with the assumption that *Synechococcus* growth would follow an exponential model:

$$\mu = (1/t) \ln (N_t/N_0)$$

where t is the duration of the incubation (days), and N_0 and N_t are *Synechococcus* cells (cells L⁻¹) at the beginning and end of the incubation, respectively.

Statistical analysis

All statistical analyses and visualizing boxplots were performed using the free statistical environment R (R Development Core Team 2013).

Results

Hydrography

The thermal stratification gradually developed from April to August (Fig. 1(a)). The depth of the euphotic zone was estimated on 26 April 2011. Light intensity in the water attenuated exponentially with depth (Fig. 1(b)). The euphotic depth (Z1%), which received 1% of the surface light intensity, was 20 m in April (Fig. 1(b)). The relatively constant Secchi disk depth (5.4 ± 1.5 m, data not shown) indicated that the euphotic depth did not vary markedly during the study period.

Contribution of pico-sized chlorophyll a to total chlorophyll a

Subsurface (0 to 10 m) chlorophyll *a* concentrations of the total ($> 0.2 \mu\text{m}$) and $> 2.0 \mu\text{m}$ fractions were relatively low in July and August (Fig. 2). A single peak of subsurface chlorophyll *a* maximum was detected at 10 or 15 m in the $> 2.0\text{-}\mu\text{m}$ fraction, except in April. In contrast, two peaks of chlorophyll *a* concentration were found in the pico-sized fraction from April and May. The contribution of the pico-sized fraction to total chlorophyll *a* concentration varied in the euphotic layer (Fig. 3). Interestingly, in the hypolimnion, on average, 16.8% of chlorophyll *a* was due to the pico-sized fraction (Fig. 3).

Picophytoplankton abundance

Numbers of *Synechococcus* cells decreased drastically below 10 m in depth (Fig. 4). However, they increased markedly from April to July or August at both the epilimnion (5.7×10^2 to 5.1×10^5 cells mL^{-1} ; average, 1.2×10^5 cells mL^{-1}) and the hypolimnion (7.0×10^2 to 2.4×10^4 cells mL^{-1} ; average, 1.0×10^4 cells mL^{-1}) (Fig. 4). PE-rich *Synechococcus* dominated in the picophytoplankton communities throughout the water column during the

study period (5.7×10^2 to 4.4×10^5 cells mL^{-1} ; average $\pm$ standard deviation, $91.3 \pm 10.3\%$), while PC-rich *Synechococcus* was considerably less abundant (below detection to 7.7×10^4 cells mL^{-1}). In the hypolimnion, PE-rich *Synechococcus* was exclusively found within the picophytoplankton community (average $\pm$ standard deviation, $99\% \pm 2.3\%$). Eukaryotic picophytoplankton cells constituted less than 1% of all picophytoplankton throughout the water column (data not shown).

Picophytoplankton cell volumes and fluorescence intensities

In May and August, the specific cell volume of *Synechococcus* increased with water depth (Fig. 5). In July, the specific cell volume of *Synechococcus* increased with depth until the bottom of the euphotic zone, and then decreased gradually.

In July and August, PE-rich *Synechococcus* microcolonies were found throughout the water column (Fig. 6). The volume of the microcolonies of *Synechococcus* cells typically increased to their maxima (average $\pm$ standard deviation; July, $3.4 \pm 0.9 \mu\text{m}$, $7.2 \pm 5.9 \mu\text{m}^3$; August, $4.0 \pm 1.7 \mu\text{m}$, $17.4 \pm 23.8 \mu\text{m}^3$) at the bottom of the euphotic zone (20 m), and then decreased gradually with depth towards the bottom of the hypolimnion. The cell-specific orange fluorescence intensity showed a scattered distribution (Fig. 7). However, cell-specific orange fluorescence intensities clearly showed that *Synechococcus* in the hypolimnion contained PE in relatively high (May and July), or at least equivalent (April, June, and August), amounts relative to the epilimnion (Fig. 7).

Incubation experiment

During the 48-h incubation in dark conditions, *Synechococcus* abundance did not change significantly (Table 1); indeed, the initial cell abundance was maintained for at least 2 days. The cell-specific orange fluorescence intensity increased during the incubation.

Discussion

The possibility of Synechococcus transportation from the epilimnion to the hypolimnion

Chlorophyll *a* size fractionation method has been used to estimate picophytoplankton contribution to phytoplankton biomass (e.g. Tremblay & Legendre 1994, Marañón et al. 2001). Tremblay & Legendre (1994) concluded that there was no significant differences between the carbon to chlorophyll *a* ratio of small and large phytoplankton. It therefore is appropriate to assume that the chlorophyll *a* distribution in different size classes accounts for the biomass size structure of the phytoplankton assemblages. The most interesting finding of the present study was the constant and relatively high contribution (16.8% on average) of the pico-sized chlorophyll *a* to the total chlorophyll *a* in the hypolimnion throughout the study period (Fig. 3). Although the information about the distribution of *Synechococcus* in the hypolimnion is limited, several studies on the distribution of *Synechococcus* in the hypolimnion of freshwater lakes have been conducted (Padisák et al. 1997, Winder 2009, Callieri et al. 2012). However, to our knowledge, the contribution of *Synechococcus* to the hypolimnetic total phytoplankton biomass has not yet been reported for a large freshwater lake. The present study is the first to report the unexpectedly high contribution of *Synechococcus* biomass to the total phytoplankton biomass in the hypolimnion during the thermal stratification period. The contributions of picophytoplankton in freshwater systems are highly variable (Bell & Kalff 2001). The average of 16.8% in the present study falls into the ranges previously reported from epilimnetic phytoplankton communities (0.2 to 43%; Stockner 1988, Bell & Kalff 2001). The highest contribution in the present study was 28.8% (Fig. 3), suggesting occasional importance of *Synechococcus* biomass in the hypolimnion. In addition, because chlorophyll *a* derived from microcolonies was included in the > 2.0- μ m fraction, our estimation of the *Synechococcus* contribution to phytoplankton biomass may be conservative. *Synechococcus* may be a food source for hypolimnetic nanoflagellates. Therefore, we

believe that our results underscore the importance of *Synechococcus* in the food web and/or matter cycling of the hypolimnion.

Several experimental studies have revealed that cyanobacteria can grow on organic substrates in dark conditions (Rippka 1972, Mannan & Pakrasi 1993). However, our incubation experiment revealed that *Synechococcus* maintained the initial cell abundance for at least 2 days in dark conditions, without significant growth (Table 1). We did not estimate growth rate of *Synechococcus* under the light condition using same manner with the dark incubation experiment. Thus, growth of *Synechococcus* under light condition still remains unclear in the present study. However, we simultaneously measured *Synechococcus* growth rate in diluted lake water with 0.2- μ m filtrate to 20% at the *in situ* light condition in shore of the lake, and *Synechococcus* had positive growth rate (0.22 d^{-1}) in the diluted lake water (our unpublished data). Thus, it is likely that *Synechococcus* in Lake Biwa do not proliferate in dark condition, though they have ability to grow at the *in situ* light condition. Some laboratory experiments have demonstrated *Synechococcus* growth on high concentrations of labile organic substrates under optimal-temperature conditions in dark (Rippka 1972, Mannan & Pakrasi 1993). However, in the hypolimnion of Lake Biwa, labile organic matter concentration is limited, and water temperatures are low (Maki et al. 2010). Thus, *Synechococcus* growth may be limited in the hypolimnion of the lake. Indeed, oceanic *Synechococcus* cannot grow in natural seawater under dark conditions (Sohrin et al. 2011, Timmermans et al. 2005), though they can maintain their populations at a certain level (Sohrin et al. 2011). Hence, *in situ* growth of *Synechococcus* has a minor contribution to changes in their abundance in the hypolimnion of the lake.

The present study is the first to demonstrate the distribution of *Synechococcus* in the entire water column of Lake Biwa. A significant positive correlation between *Synechococcus* abundance in the euphotic zone (10 m) and in the bottom of the hypolimnion (70 m) was found during the stratification period ($r = 0.94$, $p = 0.019$; Fig. 8).

In addition, the incubation experiment suggests that *Synechococcus* cannot grow in dark conditions (Table 1). These results suggest the recent delivery of a significant fraction of *Synechococcus* cells from the epilimnion to the hypolimnion. Thus, the *Synechococcus* population in the hypolimnion may be supplied primarily by the sinking of populations of their epilimnetic counterparts during the stratification period. In deep lakes, PE-rich *Synechococcus* typically dominate at the bottom of euphotic zone and form a deep chlorophyll maximum (Callieri 2007). The deep chlorophyll maximum is quite unstable and suddenly disappear, depending on both abiotic and biotic interactions (Callieri 2012). Thus, PE-rich *Synechococcus* cells may be supplied from deep chlorophyll maximum to hypolimnion of deep lakes, though fate of *Synechococcus* in the deep chlorophyll maximum has not yet been clarified.

It has previously been demonstrated that *Synechococcus* forms aggregates with sinking particles (Waite *et al.*, 2000), zooplankton faecal pellets (Waite *et al.* 2000, Stukel *et al.* 2013), and other *Synechococcus* cells (Waite *et al.* 2000, Lomas & Moran 2011), all of which are thought to be major processes that accelerate the sinking flux of *Synechococcus* in the ocean (Richardson & Jackson 2007, Lomas & Moran 2011, Sohrin *et al.* 2011, Stukel *et al.* 2013). In contrast, in freshwater lakes, co-aggregation of *Synechococcus* with large particles (Klut & Stockner 1991), zooplankton faecal pellets (Callieri 2007) and other *Synechococcus* cells (Callieri 2010, Callieri *et al.* 2012) has been reported, but no association between *Synechococcus*-containing aggregates and the vertical transportation of *Synechococcus* cells to the hypolimnion in lakes has been reported. In addition, there is no information about the hypolimnetic distribution of *Synechococcus*. Because nanoflagellates are the major consumers of *Synechococcus* cells in the epilimnion of Lake Biwa (Nagata 1988), zooplankton faecal pellets are likely not the major transportation carrier of *Synechococcus* in the lake. The formation of microcolonies may accelerate their sinking velocity (Fig. 6). The decrease in microcolony volume from the lower layers of the

euphotic zone to that of the hypolimnion suggests the supply of *Synechococcus* cells removed from the microcolonies in the hypolimnion (Fig. 6). Further study is required to elucidate the mechanisms of *Synechococcus* transportation from the epilimnion to the hypolimnion.

The physiological state of Synechococcus in the hypolimnion

In the present study, the contribution of PE-rich cells to total *Synechococcus* cells in the hypolimnion (average, 98.7%) was higher than that of the epilimnion (average, 86.4%; $p<0.0001$). Previous physiological studies have reported that PE-rich cells have advantageous at low light conditions (Callieri et al. 2012), and the result in the present study supports those in the previous studies. The relatively high fluorescence intensity of *Synechococcus* cells in the hypolimnion than in the epilimnion ($p<0.0001$, Fig. 7) was consistent well with the result of dark incubation experiment (Table 1). It has been reported that cyanobacteria accumulate PE in dark, and their cells immediately initiate photosynthesis when transferred to the light condition (Allen 1984). Thus, increase in cell-specific fluorescence intensity of *Synechococcus* in the hypolimnion may be their natural response to darkness.

Catabolism of cellular stores of endogenous carbon sources has been hypothesized as the mechanism of prolonged cyanobacterial survival in the dark (Montechiaro et al., 2006; Jiao et al., 2014). In the present study, a marked decrease in cell volume below the thermocline was observed in April (Fig. 5), possibly due to the consumption of stored carbon sources to facilitate long-term survival in the dark (Montechiaro et al., 2006). Proteins (amino acids) comprise about half of the carbon in a *Synechococcus* cell (Kaiser & Benner 2008), and phycobiliproteins are the most abundant proteins in the cyanobacterial cells (Allen 1984). Thus, accumulation and catabolism of PE may be one of the reasons of their prolonged survival in the hypolimnion. However, no such decrease in cell volume was

observed in other months (Fig. 5). Therefore, our results do not support the notion that catabolism of cellular stores of endogenous carbon sources is a major survival mechanism in the hypolimnion. Another hypothesis is the direct uptake of dissolved organic matter by *Synechococcus* in the hypolimnion. It is well known that most of the dissolved organic matter in the deep sea is refractory (Hansell 2013). So, it is unlikely that cyanobacteria utilize the dissolved organic matter in the deep ocean (Jiao et al., 2014). In contrast, semi-labile organic matter is supplied to the hypolimnion of Lake Biwa during the mixing period, although a large fraction of the dissolved organic matter in the hypolimnion is also refractory (Maki et al. 2010). Thus, one possible explanation for the survival of *Synechococcus* in the hypolimnion of Lake Biwa is its utilization of semi-labile dissolved organic matter. The permanent oxygenated hypolimnion may also support relatively high abundance of *Synechococcus* at the hypolimnion in Lake Biwa. In general, aerobic catabolism of organic matter results in higher energy gain (Søballe & Pool 1999), though some cyanobacteria can utilize organic matter and survive in dark under anaerobic conditions (Richardson & Castenholz 1987).

Synechococcus abundances below 20 m were nearly constant (Fig. 4), although the cell volume tended to increase with depth, except in April (Fig. 5). This result supports the hypothesis that *Synechococcus* would utilize dissolved organic matter to maintain their cell abundance without significant cell division during the sinking process. Similar results were also reported from oceanic studies (Albertano et al. 1997, Sohrin et al. 2011). In addition, bacterial cell volume in the hypolimnion was larger than that in the epilimnion of Lake Biwa (Takasu et al. 2013). The slow growth rate likely favours the enlargement of *Synechococcus* cells in the absence of cell division (Albertano et al. 1997). Further analyses of *in situ* organic matter utilization ability and the carbon and nitrogen contents of hypolimnetic *Synechococcus* will enhance our understanding of their trophic status and survival mechanisms in the hypolimnion of Lake Biwa.

358

359 *Conclusion*

360 The biomass of *Synechococcus* significantly contributed to that of the total phytoplankton
361 community in the hypolimnion of Lake Biwa during the thermal stratification period,
362 accounting for up to 28.8% (16.8% on average) of the total phytoplankton biomass. The
363 roles of *Synechococcus* in food web and/or matter cycling of the hypolimnion may be more
364 important than previously hypothesized.

365

366 **Acknowledgements**

367 We would like to thank Tadatoshi Koitabashi and Yukiko Goda for their assistance during
368 field sampling. We are grateful to the handling editor, Dr. Rutger de Wit, and anonymous
369 reviewers whose comments greatly improved the manuscript. This study was supported in
370 part by Japan Society for the Promotion of Science (JSPS) KAKENHI Grant Number
371 23370010 and by the Japan Science and Technology Strategic International Research
372 Cooperative Program (Japan-China) on Science and Technology for Environmental
373 Conservation and Construction of a Society with Less Environmental Burden, "Fate of
374 dissolved organic matter in lakes with special reference to loading and pollution", which
375 supported S.N. (Grant Number 21710081). H.T. was supported by JSPS KAKENHI Grant
376 Number 11J00658.

References

- Albertano P, Somma DD, Capucci E (1997) Cyanobacterial picoplankton from the Central Baltic Sea: cell size classification by image-analyzed fluorescence microscopy. *J Plankton Res* 19:1405–1416
- Allen MM (1984) Cyanobacterial cell inclusion. *Ann Rev Microbiol* 38:1–25
- Bell T, Kalff J (2001) The contribution of picophytoplankton in marine and freshwater systems of different trophic status and depth. *Limnol Oceanogr* 46:1243–1248
- Bidle KD, Falkowski PG (2004) Cell death in planktonic, photosynthetic microorganisms. *Nature Rev Microbiol* 2:643–655
- Callieri C, Pinolini ML (1995) Picoplankton in Lake Maggiore, Italy. *Int Revue ges Hydrobiol* 80:491–501
- Callieri C (2007) Picophytoplankton in freshwater ecosystems: The importance of small sized phototrophs. *Freshwater Rev* 1:1–28
- Callieri C (2010) Single cells and microcolonies of freshwater picocyanobacteria: a common ecology. *J Limnol* 69:257–277
- Callieri C, Cronberg G, Stockner JG (2012) Freshwater picocyanobacteria: single cells, microcolonies and colonial forms. In: Whitton B (eds) *Ecology of Cyanobacteria II: Their Diversity in Time and Space*. 2nd edn. Springer Publisher, New York, NY, p 229–271
- Hansell DA (2013) Recalcitrant dissolved organic carbon fractions. *Annu Rev Mar Sci* 5:421–445
- Hirose M, Katano T, Nakano S (2008) Growth and grazing mortality rates of *Prochlorococcus*, *Synechococcus* and eukaryotic picophytoplankton in a bay of the Uwa Sea, Japan. *J Plankton Res* 30:241–250
- Jiao N, Luo T, Zhang R, Yan W, Lin Y, Johnson ZI, Tian J, Yuan D, Yang Q, Sun J, Hu D, Wang P (2014) Presence of *Prochlorococcus* in the aphotic waters of the western

- 403 Pacific Ocean. *Biogeosciences* 11:2391–2400
- 404 Kaiser K, Benner R (2008) Major bacterial contribution to the ocean reservoir of detrital
405 organic carbon and nitrogen. *Limnol Oceanogr* 53:99–112
- 406 Kiørboe T (1993) Turbulence, phytoplankton cell size, and the structure of pelagic food
407 webs. *Adv Mar Biol* 29:1–72
- 408 Kirchman DL (2008) Introduction and overview. In: Kirchman DL (eds) *Microbial*
409 *Ecology of the Oceans*. 2nd edn. John Wiley, New York, NY, p 1–26
- 410 Klut ME, Stockner JG (1991) Picoplankton associations in an ultraoligotrophic lake on
411 Vancouver island, British Columbia. *Can J Fish Aquat Sci* 48:1092–1099
- 412 Lomas MW, Moran SB (2011) Evidence for aggregation and export of cyanobacteria and
413 nano-eukaryotes from the Sargasso Sea euphotic zone. *Biogeosciences* 8:203–216
- 414 MacIsaac EA, Stockner JG (1993) Enumeration of phototrophic picoplankton by
415 autofluorescence microscopy. In: Kemp PF, Sherr BF, Sherr EB, Cole JJ (eds.),
416 *Handbook of Methods in Aquatic Microbial Ecology*. Lewis, Fla, p 187–197
- 417 Maeda H, Kawai A, Tilzer MM (1992) The water bloom of Cyanobacterial plankton in
418 Lake Biwa, Japan. *Hydrobiologia* 248:93–103
- 419 Maki K, Kim C, Yoshimizu C, Tayasu I, Miyajima T, Nagata T (2010) Autochthonous
420 origin of semi-labile dissolved organic carbon in a large monomictic lake (Lake Biwa):
421 carbon stable isotopic evidence. *Limnology* 11:143–153
- 422 Mannan RM, Pakrasi HB (1993) Dark heterotrophic growth conditions result in an increase
423 in the content of photosystem II units in the filamentous cyanobacterium *Anabaena*
424 *variabilis* ATCC 29413. *Plant Physiol* 103:971–977
- 425 Marañón E, Holligan PM, Barciela R, González N, Mouriño B, Pazó MJ, Varela M (2001)
426 Patterns of phytoplankton size structure and productivity in contrasting open-ocean
427 environments. *Mar Ecol Prog Ser* 216:43–56
- 428 Moran R, Porath D (1980) Chlorophyll determination in intact tissues using

- 429 *N,N*-Dimethylformamide. *Plant Physiol* 65:478–479
- 430 Montecchiario F, Hirschmugl CJ, Raven JA, Giordano M (2006) Homeostasis of cell
431 composition during prolonged darkness. *Plant Cell Environ* 19:2198–2204
- 432 Nagata T (1988) The microflagellate–picoplankton food linkage in the water column of
433 Lake Biwa. *Limnol Oceanogr* 33:504–517
- 434 Nagata T, Takai K, Kawanobe K, Kim D-S, Nakazato R, Guselnikova N, Bondarenko N,
435 Mologawayaya O, Kostrova T, Drucker V, Satoh Y, Watanabe Y (1994) Autotrophic
436 picoplankton in southern Lake Baikal: abundance, growth and grazing mortality during
437 summer. *J Plankton Res* 16:945–959
- 438 Padisák J, Krienitz L, Koschel R, Nedoma J (1997) Deep-layer autotrophic picoplankton
439 maximum in the oligotrophic Lake Stechlin, Germany: origin, activity, development
440 and erosion. *Eur J Phycol* 32:403–416
- 441 R Core Team (2013) R: A Language and Environment for Statistical Computing.
- 442 Reynolds C (2006) Mortality and loss processes in phytoplankton. In: Reynolds C (eds)
443 Ecology of phytoplankton. Cambridge University Press, New York p 239–301
- 444 Richardson LL, Castenholz RW (1987) Enhanced survival of the cyanobacterium
445 *Oscillatoria terebriformis* in darkness under anaerobic conditions. *Appl Environ*
446 *Microbiol* 53:2151–2158
- 447 Richardson TL, Jackson GA (2007) Small phytoplankton and carbon export from the
448 surface ocean. *Science* 315:838–840
- 449 Rippka R (1972) Photoheterotrophy and chemoheterotrophy among unicellular blue-green
450 algae. *Arch Microbiol* 87:93–98
- 451 Scanlan DJ (2012) Marine Picocyanobacteria. In: Whitton B (eds) Ecology of
452 Cyanobacteria II: Their Diversity in Time and Space. 2nd edn. Springer Publisher, New
453 York, NY, p 503–533
- 454 Søballe B, Pool KP (1999) Microbial ubiquinones: multiple roles in respiration, gene

- 455 regulation and oxidative stress management. *Microbiology* 145:1817–1830
- 456 Sohrin R, Isaji M, Obara Y, Agostini S, Suzuki Y, Hiroe Y, Ichikawa T, Hidaka K (2011)
- 457 Distribution of *Synechococcus* in the dark ocean. *Aquat Microb Ecol* 64:1–14
- 458 Stockner JG (1988) Phototrophic picoplankton: An overview from marine and freshwater
- 459 ecosystems. *Limnol Oceanogr* 33:765–775
- 460 Stockner JG (1991) Autotrophic picoplankton in freshwater ecosystems: The view from
- 461 summit. *Int Revue ges Hydrobiol* 76:483–492
- 462 Stukel MR, Décima M, Selph KE, Taniguchi DAA, Landry MR (2013) The role of
- 463 *Synechococcus* in vertical flux in the Costa Rica upwelling dome. *Prog Oceanogr* 112–
- 464 113:49–59
- 465 Takasu, H., Kunihiro, T. and Nakano, S. (2012) Vertical community structure of bacterial
- 466 and phytoplankton in Lake Biwa using respiratory quinone and pigment analysis. In:
- 467 Kawaguchi, M., Misaki, K., Sato, H., Yokokawa, T., Itai, T., Nguyen, T.M., Ono, J. and
- 468 Tanabe S. (eds.), *Interdisciplinary Studies on Environmental Chemistry—Advanced*
- 469 *Environmental Studies by Young Scientists*, Terrapub, Tokyo, pp. 377–385.
- 470 Takasu H, Kunihiro T, Nakano S-I (2013) Estimation of carbon biomass and community
- 471 structure of planktonic bacteria in Lake Biwa using respiratory quinone analysis.
- 472 *Limnology* 14:247–256
- 473 Timmermans KR, van der Wagt B, Veldhuis MJW, Maatman A, de Baar HJW (2005)
- 474 Physiological responses of three species of marine pico-phytoplankton to ammonium,
- 475 phosphate, iron and light limitation. *J Sea Res* 53:109–120
- 476 Tremblay JE, Legendre L (1994) A model for the size-fractionated biomass and production
- 477 of marine phytoplankton. *Limnol Oceanogr* 39:2004–2014
- 478 Waite AM, Safi KA, Hall JA, Nodder SD (2000) Mass sedimentation of picoplankton
- 479 embedded in organic aggregates. *Limnol Oceanogr* 45:87–97
- 480 Winder M (2009) Photosynthetic picoplankton dynamics in Lake Tahoe: temporal and

481 spatial niche partitioning among prokaryotic and eukaryotic cells. J Plankton Res
482 31:1307–1320

483 Table 1. Growth rates and cell-specific fluorescence intensity of *Synechococcus* in dark
484 conditions.

Parameter	PE-rich	PC-rich	Total
Growth rate (day ⁻¹)	-0.04	-0.07	-0.04
Cell-specific orange fluorescence intensity at the end of the incubation (relative to initial, %)	119	nd	nd

485 Nd, not determin

Figure legends

Fig. 1. Depth profiles of (a) water temperature and (b) light intensity. The depth of the euphotic zone was estimated on 26 April 2011.

Fig. 2. Depth profiles of chlorophyll *a* concentrations in the (a) > 0.2 -, (b) > 2.0 -, and (c) 0.2 – 2.0 - μm fractions.

Fig. 3. Contribution of the 0.2 – 2.0 - μm fraction to the total chlorophyll *a* concentration ($> 0.2 \mu\text{m}$) shown by a box plot. Boxes and bars indicate the quartile (Q1 and 3) and median values, respectively. Whisker represents the value of the most extreme data point that is no more than 1.5 times the inter-quartile. Outliers are shown by open circles. Data are compiled over the study period.

Fig. 4. Depth profiles of (a) total, (b) phycoerythrin-rich, and (c) phycocyanin-rich *Synechococcus* cells.

Fig. 5. Depth profile of *Synechococcus* cell volumes shown by boxplots. Boxes and bars indicate the quartile (Q1 and 3) and median values, respectively. Whisker represents the value of the most extreme data point that is no more than 1.5 times the inter-quartile. Outliers are shown by open circles.

Fig. 6. Vertical size distribution of phycoerythrin-rich *Synechococcus* microcolonies shown by boxplots. Boxes and bars indicate the quartile (Q1 and 3) and median values, respectively. Whisker represents the value of the most extreme data point that is no more than 1.5 times the inter-quartile. Outliers are shown by open circles. The inserts represent phycoerythrin-rich *Synechococcus* microcolonies under green excitation of epifluorescence microscope. The sample was taken at 70 m in August 2011.

Fig. 7. Depth profile of cell-specific orange fluorescence (phycoerythrin) intensity shown by boxplots. Boxes and bars indicate the quartile (Q1 and 3) and median values, respectively. Whisker represents the value of the most extreme data point that is no more than 1.5 times the inter-quartile. Outliers are shown by open circles.

512 **Fig. 8.** Relationship between *Synechococcus* cell abundances at 10 and 70 m in depth. The
513 *Synechococcus* abundance of April used the data from 5 m rather than 10 m.

Fig.1.Takasu *et al.*

514
515
516
517
518
519

Fig.2.Takasu *et al.*

520

521

522

523

Fig. 3. Takasu *et al.*

524
525
526
527

Fig.4. Takasu *et al.*

528

529

530

Fig.5. Takasu *et al.*

531
532
533
534

Fig. 6. Takasu *et al.*

Fig. 7. Takasu *et al.*

Fig. 8. Takasu *et al.*