

1 **Distribution and geochemical behavior of anionic surfactants determined**
2 **as ethyl violet active substances in Lake Biwa, Japan**

3

4 Hiroo Yoshida, Shunsuke Kudari, Toshitaka Hori & Masahito Sugiyama

5

6 Graduate School of Human and Environmental Studies, Kyoto University, Kyoto,

7 Japan

8

9 Correspondence to: Masahito Sugiyama, Yoshida-nihonmatsu-cho, Sakyo-ku,

10 Kyoto 606-8501, Japan. E-mail: sugiyama.masahito.5s@kyoto-u.ac.jp

11 Telephone: +81-75-753-6836; fax: +81-75-753-7879

12

13 **Abstract**

14 The distribution and geochemical behavior of anionic surfactants (AS)
15 determined as ethyl violet active substances (EVAS) were studied in Lake Biwa,
16 Japan. We determined the vertical distributions of dissolved EVAS (DEVAS),
17 suspended particulate EVAS (PEVAS), particulate organic carbon (POC), and
18 the PEVAS/POC ratio. Statistical analyses conducted using plots of PEVAS vs.
19 POC and PEVAS/ Chlorophyll-*a* (Chl-*a*) vs. POC/Chl-*a* revealed that the
20 distributions of DEVAS and PEVAS were affected by the concentrations of
21 freshly produced POC and aged POC and by the pH of the lake water.
22 Adsorption and desorption of EVAS onto the POC phase were found to be
23 important mechanisms regulating the distribution and
24 environmental/geochemical behavior of EVAS in the lake. The organic
25 carbon-normalized particle–water partitioning coefficient, K_{OC} , for EVAS was also
26 estimated and compared to that for nonylphenol.

27 **Keywords**

28 anionic surfactants; ethyl violet; distribution, geochemical behavior, Lake Biwa

29

30 **Introduction**

31 Linear alkylbenzene sulfonate (LAS) is the primary anionic surfactant found in

32 Japan. This compound is listed as Class I in the Pollutant Release and Transfer

33 Register Law of Japan. A Class I Designated Chemical Substance is defined as

34 continuously occurring throughout much of the environment based on its

35 physical and chemical properties, manufactured volume, and importation and

36 usage, and having risks to human health and the ecosystem (Japan Ministry of

37 the Environment 2013). Production of LAS was 72 kt in 2002 and the amount of

38 LAS discharged to the sewage in Japan was 64 kt in 2001 (National Institute of

39 Technology and Evaluation 2005). The degree of discharge is related to the

40 environmental risk.

41 Research on anionic surfactants (AS) in the aquatic environment has been

42 carried out globally. In Japan, AS concentrations have been reported to range

43 between 50–8500 $\mu\text{g/L}$ in river water (Kobayashii and Yamauchi 1979; Kobuke
44 1981; Yoshimura *et al.* 1984; Hirayama *et al.* 2006), 0–264 $\mu\text{g/g}$ in river
45 sediments (Ambe 1972; Uchiyama 1982; Yoshimura *et al.* 1984), and 10–50
46 $\mu\text{g/g}$ in lake sediments (Ambe 1972). LAS concentrations ranged between
47 0.6–79 $\mu\text{g/L}$ in river water in Japan (Kobuke 2004; Miura *et al.* 2008), 0–342 $\mu\text{g/L}$
48 in South Africa (Gordon *et al.* 2009), and 2–126 $\mu\text{g/L}$ in the Philippines (Dyer *et*
49 *al.* 2003). LAS concentrations also ranged between 60–2300 $\mu\text{g/g}$ in suspended
50 particles in rivers in Japan (Takada and Ishiwatari 1987) and between
51 0.03–17.76 $\mu\text{g/g}$ and 0.101–0.186 $\mu\text{g/g}$ in river sediments in Portugal (Hampel *et*
52 *al.* 2009) and South Africa (Gordon *et al.* 2009), respectively.

53 As noted above, most limnological and environmental studies of AS have been
54 focused on rivers, but few studies have been conducted of AS in lakes. The
55 distribution of AS in rivers is important in determining anthropogenic pollution
56 levels and discharges to the natural environment of AS by human activities.
57 However, these studies alone are not sufficient to fully delineate the

58 environmental and geochemical processes of AS, because water masses are
59 constantly renewed by flows from upstream. On the other hand, water masses
60 may stagnate in lakes and thus, studies on partitioning of AS between lake water
61 and suspended and sediment particles are more feasible.

62 Several authors have studied removal processes for AS in laboratory
63 experiments (Ou *et al.* 1996; Perales *et al.* 1999; Doi *et al.* 2002; Garcia *et al.*
64 2002). These compounds are generally known to be degraded by
65 microorganisms (Perales *et al.* 1999) and adsorbed by solid phases (Ou *et al.*
66 1996; Doi *et al.* 2002; Garcia *et al.* 2002). The former depends on the water
67 temperature, while the latter is related to the organic carbon content of the solid
68 phase, known as the particulate organic carbon (POC) content. To date, there
69 have been few studies of adsorption and partitioning of AS to POC in lakes.
70 Although some authors have addressed POC, they treated it as a homogeneous
71 material despite being a mixture of various organic materials (Yoshida *et al.*
72 2009). In our previous study (Yoshida *et al.* 2009), we clarified that POC in lakes

73 could be classified into two fractions, freshly produced POC (f-POC) and aged
74 POC (a-POC), based on differences in the affinity of nonylphenol (NP) to POC;
75 its affinity to a-POC was approximately 4 times higher than that to f-POC. A
76 similar evaluation is also needed for AS.

77 Colorimetric methods have often been used for analysis of AS. Among these
78 methods, methylene blue, which offers the advantage of being an easy-to-use
79 and convenient technique, is widely used. In recent years, the methylene blue
80 reagent has been replaced by an ethyl violet reagent (Soejima 1982). The ethyl
81 violet colorimetric method is more useful than methylene blue as it has: (1) high
82 sensitivity, (2) a very simple and less time-consuming procedure, (3) little
83 interference by coexisting ions, and (4) good reproducibility and quantification of
84 trace amounts of anionic surfactants (Motomizu *et al.* 1982). It is also used as a
85 standard method in Japan (Japan Standards Association 1998). Therefore, we
86 quantified AS as ethyl violet active substances (EVAS).

87 In this study, we determined the distribution and geochemical behavior of AS in

88 Lake Biwa as EVAS. Using plots of suspended particulate EVAS (PEVAS) vs.
89 POC and PEVAS/Chlorophyll-*a* (Chl-*a*) vs. POC/Chl-*a*, we evaluated adsorption
90 and partitioning of EVAS to POC in Lake Biwa. The average PEVAS/POC ratio
91 and organic carbon-normalized particle–water partitioning coefficient (K_{oc}) of
92 EVAS to f-POC and a-POC were estimated and compared to those of NP.

93

94 **Materials and methods**

95 **Solvents and standards**

96 Sodium sulfate and sodium hydroxide were purchased from Nacalai Tesque
97 (Kyoto, Japan). Acetic acid, toluene, disodium ethylenediaminetetraacetate
98 (EDTA), and sodium dodecyl sulfate (SDS) were obtained from Wako Chemicals
99 (Osaka, Japan). Ethyl violet was obtained from Tokyo Chemical Industry (Tokyo,
100 Japan).

101

102 **Sampling site and sample collection**

103 The sampling sites for this study were located in Lake Biwa (35°15' N, 136°05' E)
104 and the Yasu River. Lake Biwa, which is the largest lake in Japan, is divided into
105 two parts, the northern and southern basins. The former has a volume of 27.3
106 km³ with an average depth of 44 m. Thermal stratification typically occurs from
107 May to January and vertical convection of the entire water column in the lake
108 occurs regularly from February to April. The southern basin of the lake has a
109 volume of 0.2 km³ with an average depth of 3.5 m. The water in this basin is not
110 thermally stratified and is more or less saturated with atmospheric oxygen
111 throughout the year (Sugiyama *et al.* 2005). Field surveys were conducted at
112 Station Ie-1 (35°12.58' N, 135°59.55' E) between September 2000 and August
113 2001.

114 The Yasu River is a large river that flows near Station Ie-1 in Lake Biwa. It has
115 the largest drainage area and the second highest water discharge among the
116 more than 110 rivers flowing into Lake Biwa (Lake Biwa Environmental
117 Research Institute 1986; Kunimatsu 1988). Several towns are located along this

118 river with a total population of 220,000. The river water is generally used for
119 industrial water, domestic supply, and irrigation, and then the discharge is
120 released to the river (Kimaro *et al.* 2002). Field surveys of the river were
121 conducted at Stations YS-1 (35°09.70' N, 135°99.30' E) and YS-2 (35°02.37' N,
122 136°01.12' E) on 5 September 2000 and 9 March 2001.

123 Lake water samples were collected vertically with a Van Dorn water sampler
124 and river water samples were collected with a glass bottle. The samples were
125 filtered through a Whatman GF/F filter (diameter 47 mm, pore size 0.7 μm) to
126 separate suspended particles from the water. The water samples were analyzed
127 within 3 days of sampling. Suspended particle samples collected on the filter
128 were kept frozen until they were analyzed.

129

130 **Analytical procedures**

131 Sample treatment for AS determination was carried out as follows. Forty
132 milliliters of a water sample was transferred into a glass centrifuge tube. Two

133 milliliters 1 mol/L sodium sulfate solution, 2 mL acetic acid–EDTA buffer (pH 5.0),
134 0.8 mL 1 mmol/L ethyl violet solution, and 2 mL toluene were added to the tube.
135 The tube was shaken for 60 min and then centrifuged for 5 min. After phase
136 separation, the absorbance of the toluene phase was measured at 611 nm
137 (UV-1600; Shimadzu, Kyoto, Japan) and calibrated against an SDS solution.

138 Suspended particle and sediment samples were dried at 60 °C and
139 ultrasonicated twice with 10 mL methanol for 45 min at room temperature. The
140 obtained extracts were evaporated until they were dry using a centrifugal
141 vaporizer (Model CVE 2wd; Tokyo Rika Kikai, Tokyo, Japan). The residue was
142 redissolved in 40 mL water and the solution was then treated as a water sample.

143 POC was quantified with a CHN analyzer (Model 2400; Perkin Elmer, Tokyo,
144 Japan). The temperatures of the combustion and reduction chambers were set
145 to 925 °C and 640 °C, respectively.

146 Chl-*a* was quantified spectrophotometrically according to the methods of
147 UNESCO and Lorenzen (Lorenzen 1967).

148

149 **Results and discussion**

150 **Method validation**

151 In the analyses using the ethyl violet colorimetric method, sodium sulfate was
152 used to accelerate phase separation and EDTA was used to reduce
153 interferences from multivalent metal ions. AS were determined as EVAS.

154 Method validation was carried out to evaluate the suitability of this method. SDS
155 was used as a standard material for analysis of AS. The linearity of its calibration
156 curve was evaluated by taking measurements at four different concentrations of
157 SDS. The correlation coefficient r of the curve was >0.999 from 0 to $38 \mu\text{g/L}$ SDS.

158 The detection limit was determined as the concentration of SDS that gave a
159 signal three times the background noise, which was $0.27 \mu\text{g/L}$.

160 Accuracy and precision were calculated based on recovery of SDS from
161 sediments, since sediments have a more complex matrix. For this evaluation,
162 $101 \mu\text{g/g}$ SDS was spiked into a sediment sample containing $235 \mu\text{g/g}$ of EVAS.

163 The recovery rate and precision were 115% and 3.3% ($n = 3$). Thus, this method
164 was found to be suitable for analysis of AS in environmental and geochemical
165 studies.

166

167 **Distribution and geochemical behavior of EVAS in Lake Biwa**

168 Figure 1 shows the vertical distributions for water temperature, pH, and EVAS
169 concentration in Lake Biwa. During the circulation period (2 April 2001), the
170 vertical distribution of dissolved EVAS (DEVAS) was relatively constant,
171 because the lake water was well mixed due to the vertical convection that occurs
172 in early spring. Conversely, the DEVAS concentration during the stagnation
173 period (12 September and 5 October 2000, 11 April 2001) had higher values at
174 the surface and decreased at the thermocline. The concentration in the
175 hypolimnion was nearly constant except near the bottom and was similar to that
176 during the circulation period. The high DEVAS concentrations in the epilimnion
177 are believed to be due to river water with high DEVAS concentrations flowing

178 only into the epilimnion and/or to desorption of AS from the POC phase due to
179 the increase in pH in the surface water during the stagnation period. Average
180 DEVAS concentrations in the Yasu River were 18.2 $\mu\text{g/L}$ on 5 September 2000
181 and 55.7 $\mu\text{g/L}$ on 9 March 2001. These values were higher than those in the
182 epilimnion during the stagnation period and in the entire water column during the
183 circulation period. Weighted average concentrations in the epilimnion were 12.2
184 $\mu\text{g/L}$ on 12 September 2000, 14.6 $\mu\text{g/L}$ on 5 October 2000, and 12.2 $\mu\text{g/L}$ on 11
185 April 2001. During the circulation period (2 April 2001), the weighted average
186 concentration over the entire water column was 10.5 $\mu\text{g/L}$. These results indicate
187 that the DEVAS concentration in the surface water is increased by the Yasu and
188 other rivers flowing into the epilimnion during the stagnation period.

189 In addition, the pH in the epilimnion increases with active biological
190 productivity in the summer in many lakes, including Lake Biwa (Lake Biwa
191 Research Institute and National Institute for Research Advancement 1984;
192 Sugiyama *et al.* 2005). It has been reported that AS desorb from the suspended

193 POC phase when the pH of the aqueous phase increases (Yoshimura and
194 Nakae 1982). Therefore, the DEVAS concentration in the surface water may
195 have increased by desorption of AS from the POC phase in the suspended
196 particulate matter and the sediments in shallow and littoral area of the lake when
197 the pH of the water increased. This hypothesis was supported by the results for
198 PEVAS vs. POC (see below).

199 The concentration of PEVAS decreased from the surface to the thermocline or
200 the middle layer. However, at any given sampling time in Fig. 1, the vertical
201 distribution of PEVAS/POC, which represents the organic carbon-normalized
202 concentration of EVAS in the suspended particles, was neither uniform nor
203 similar to those of PEVAS or POC. Figure 2 plots PEVAS vs. POC. The
204 correlation coefficient r was relatively low at 0.619. There are believed to be two
205 reasons for this result. The first is the treatment of POC as a homogeneous
206 material despite actually being a mixture of various organic materials. The
207 second is not considering changes in the pH of the lake water, although AS is

208 known to desorb from the suspended POC phase when the pH of the aqueous
209 phase increases.

210 The first issue can be explained further based on the results of our previous
211 environmental and geochemical study on NP in Lake Biwa (Yoshida *et al.* 2009).

212 In that study, POC was mathematically classified into two fractions, f-POC and
213 a-POC, based on the affinity of NP to these fractions. The differences between
214 f-POC and a-POC are likely associated with their chemical structure and the
215 nature of the organic matter. Therefore, in the present study, we examined the
216 distribution of PEVAS and POC in a similar manner. The total concentration of
217 POC, [POC], can be expressed as:

218
$$[\text{POC}] = [\text{POC}]_f + [\text{POC}]_a$$

219 where $[\text{POC}]_f$ and $[\text{POC}]_a$ are the concentrations of f-POC and a-POC,
220 respectively. If the EVAS concentration accumulated by f-POC and a-POC are
221 defined as $[\text{PEVAS}]_f$, and $[\text{PEVAS}]_a$, respectively, then the total PEVAS
222 concentration, [PEVAS], can be expressed as:

223
$$[\text{PEVAS}] = [\text{PEVAS}]_f + [\text{PEVAS}]_a$$

224
$$= R_f[\text{POC}]_f + R_a[\text{POC}]_a$$

225
$$= R_f[\text{POC}]_f + R_a([\text{POC}] - [\text{POC}]_f)$$

226 where R_f and R_a are the average PEVAS/POC ratios for f-POC and a-POC,

227 respectively.

228 Phytoplankton is typical of freshly produced organic matter in a lake and the

229 Chl-*a* concentration is an index of phytoplankton abundance. Therefore, if Chl-*a*

230 is regarded as an index of the f-POC concentration, $[\text{POC}]_f$ can be expressed as:

231
$$[\text{POC}]_f = \alpha[\text{Chl-}a]$$

232 where α is the average f-POC/Chl-*a* ratio. The following relationship can be

233 obtained from the above equations:

234
$$[\text{PEVAS}] = R_f\alpha[\text{Chl-}a] + R_a([\text{POC}] - \alpha[\text{Chl-}a])$$

235
$$= R_a[\text{POC}] + \alpha(R_f - R_a)[\text{Chl-}a]$$

236
$$[\text{PEVAS}]/[\text{Chl-}a] = R_a[\text{POC}]/[\text{Chl-}a] + \alpha(R_f - R_a)$$

237 The obtained equation can be interpreted as follows. The plot of PEVAS/Chl-*a*

238 vs. POC/Chl-*a* is predicted to be linear, with a slope and intercept of R_a and $\alpha(R_f$
239 $- R_a)$, respectively. If AS accumulates in a-POC more than in f-POC (i.e., $R_a >$
240 R_f), the intercept will be negative. Figure 3 plots PEVAS/Chl-*a* vs. POC/Chl-*a*.
241 The plot showed good linearity and its correlation coefficient ($r = 0.916$) was
242 higher than that of the PEVAS vs. POC plot ($r = 0.619$). These results suggest
243 that classification of POC into f-POC and a-POC is possible through this method
244 and is important for understanding the behavior of AS in the lake. As expected,
245 the intercept $\alpha(R_f - R_a)$ of the plot was negative (-1.27×10^{-2} g/g). Therefore, the
246 affinity of AS to a-POC was higher than to f-POC. These results are consistent
247 with our previous study showing that NP also has stronger affinity to a-POC than
248 to f-POC (Yoshida *et al.* 2009).

249 The slope R_a and intercept $\alpha(R_f - R_a)$ of the linear regression line in Fig. 3 were
250 5.5×10^{-3} g/g and -1.27×10^{-2} g/g, respectively. Using 87.8 g/g as the average α
251 value (Yoshida *et al.* 2009) based on the average POC/Chl-*a* value in culture
252 experiments of phytoplankton from Lake Biwa (Tezuka 1985), R_f can be

253 estimated as 5.4×10^{-3} g/g. According to the definition given above, R_f and R_a are
254 the average EVAS/POC ratios for f-POC and a-POC, respectively. Therefore,
255 these values indicate that the affinity of AS to a-POC is somewhat higher than
256 that to f-POC.

257 The second reason for the poor correlation coefficient ($r = 0.619$) between
258 PEVAS and POC in Fig. 2 can be explained by the PEVAS vs. POC plots shown
259 in Fig. 4. Based on the pH of the lake water, the data can be classified into 3
260 groups: Group A, samples of the epilimnion in September and October 2000 (pH
261 7.6–7.8; 0–13 m in September and 0–17 m in October); Group B, samples of the
262 hypolimnion in September and October 2000 (pH 6.9–7.1; 20–72 m); and Group
263 C, samples at all depths on 2 and 11 April 2001 (pH 7.1–7.4). The correlation
264 coefficients for the three groups were 0.690, 0.926, and 0.908, respectively. The
265 PEVAS/POC value (3.38) for the epilimnion with high pH values was smaller
266 than those (5.57 and 7.84) for the hypolimnion and at all depths with low pH
267 values. These results are consistent with the increase in the DEVAS

268 concentration in the epilimnion with high pH values during the stagnation period,
269 and also indicates that PEVAS concentrations in the POC decreased with an
270 increase in the pH of the lake water. In other words, EVAS is desorbed from the
271 suspended POC phase when the pH of the aqueous phase increases.

272 We previously reported that some oxyanions are also released from solid
273 phases such as sediments and suspended particulate matter with an increase in
274 the pH of lake water (Harita *et al.* 2005), similar to the present results. Therefore,
275 adsorption and desorption of chemical components onto solid phases with
276 changes in the pH of lake water are considered to be important mechanisms
277 regulating their distribution and environmental/geochemical behavior in lakes.

278 However, such a trend was not observed in our previous study of NP. This is
279 likely because NP, as a non-ionic compound, is not affected by changes in pH.

280 As shown in Fig. 4, the PEVAS/POC value (5.57) in the hypolimnion in
281 September and October was smaller than that (7.84) at all depths in April. This is
282 believed to be due to degradation of PEVAS in the hypolimnion during the

283 stagnation period.

284

285 **Comparison of K_{oc} values for EVAS and NP**

286 The partition coefficient K_{oc} is calculated using the following equation, which

287 uses the DEVAS concentration and the PEVAS/POC ratio (R_f and R_a):

$$288 \quad K_{oc}(\text{mL/g} - \text{POC}) = [\text{PEVAS/POC}(\text{g/g} - \text{POC})]/[\text{DEVAS}(\text{g/mL})]$$

289 The average DEVAS concentrations were 15.9×10^{-9} g/mL and 9.1×10^{-9} g/mL in

290 the epilimnion and hypolimnion, respectively. Using these concentrations, log

291 K_{oc} was determined to be 5.5 and 5.8 for R_f and R_a , respectively. These values

292 are within those (2.5–5.9) obtained in a previous study (Vincent and Glenwood

293 1987).

294 The log K_{oc} values for EVAS (5.5–5.8) were higher than those for NP (4.8 and

295 5.4). The high K_{oc} values for EVAS are likely due to formation of a bulky ionic

296 pair with cationic organic compounds such as quaternary ammonium ions as

297 well as its high affinity to POC. Cationic surfactants, which typically contain

298 quaternary ammonium ions, are widely used in industrial and household
299 detergents and are discharged into lake and river waters. Their production in
300 Japan was 70 kt in 1997 (Japan Surfactant Industry Association 1997). Cationic
301 surfactants easily form a stable ion pair with coexisting anionic surfactants in
302 environmental waters (Utsunomiya *et al.* 1998) and this ion pair is quickly
303 absorbed onto sewage sludge and sediments with a high partitioning coefficient
304 (Sun *et al.* 2003; Sütterlin *et al.* 2008).

305

306 **Conclusions**

307 We studied the distribution and geochemical behavior of EVAS in Lake Biwa,
308 Japan. The distributions of DEVAS and PEVAS were clearly affected by the
309 concentrations of the two POC fractions, f-POC and a-POC, and by the pH of the
310 lake water. Thus, adsorption and desorption of EVAS onto the POC phase were
311 found to be important mechanisms regulating their distribution and
312 environmental/geochemical behavior in lakes.

313

314 **Acknowledgements**

315 This study was partly supported by grants from the Japan Society for the
316 Promotion of Science (12480150 and 15651028) and from the Nippon Life
317 Insurance Foundation. The authors thank Mr. T. Koitabashi and Mr. T. Miyano
318 from the Center for Ecological Research at Kyoto University for their assistance
319 with collection of samples in Lake Biwa.

320

321

322

References

323 Ambe, Y. (1972) Distribution of Alkylbenzenesulfonate in the Bottom Muds of
324 Lake Suwa. *Jpn. J. Limnol.*, **33**, 44–50 (in Japanese).

325 Doi, J., Marks, K. H., DeCarvalho, A. J., McAvoy, D. C., Nielsen, A. M., Kravetz,
326 L. and Cano, M. L. (2002) Investigation of an Onsite Wastewater Treatment
327 System in Sandy Soil: Sorption and Biodegradation of Linear Alkylbenzene

- 328 Sulfonate. *Environ. Toxicol. Chem.*, **21** (12), 2617–2622.
- 329 Dyer, S. D., Peng, C., McAvoy, D. C., Fendinger, N. J., Masscheleyn, P., Castillo,
330 L. V. and Lim, J. M. (2003) The Influence of Untreated Wastewater to
331 Aquatic Communities in the Balatuin River, The Philippines. *Chemosphere*,
332 **52** (1), 43–53.
- 333 Garcia, M. T., Campos, E., Dalmau, M., Ribosa, I. and Sánchez-Leal, J. (2002)
334 Structure-activity Relationships for Association of Linear Alkylbenzene
335 Sulfonates with Activated Sludge. *Chemosphere*, **49** (3), 279–286.
- 336 Gordon, A. K., Muller, W. J., Gysman, N., Marshall, S. J., Sparham, C. J.,
337 O'Connor, S. M. and Whelan, M. J. (2009) Effect of Laundry Activities on
338 In-stream Concentrations of Linear Alkylbenzene Sulfonate in a Small Rural
339 South African River. *Sci. Total. Environ.*, **407** (15), 4465–4471.
- 340 Hampel, M., Canário, J., Branco, V., Vale, C. and Blasco, J. (2009)
341 Environmental Levels of Linear Alkylbenzene Sulfonates (LAS) in Sediments
342 from the Tagus Estuary (Portugal): Environmental Implications. *Environ.*

- 343 *Monit. Assess.*, **149** (1–4), 151–161.
- 344 Harita, Y., Hori, T. and Sugiyama, M. (2005) Release of Trace Oxyanions from
345 Littoral Sediments and Suspended Particles Induced by pH Increase in the
346 Epilimnion of Lakes. *Limnol. Oceanogr.*, **50** (2), 636–645.
- 347 Hirayama, Y., Ikegami, H., Machida, M. and Tatsumoto, H. (2006) Simple and
348 Rapid Determination of Linear Alkylbenzene Sulfonates by In-tube
349 Solid-phase Microextraction Coupled with Liquid Chromatography. *J. Health.*
350 *Sci.*, **52** (3), 228–236.
- 351 Japan Ministry of the Environment (2014) PRTR Substances.
352 <http://www.env.go.jp/en/chemi/prtr/substances/index.html> [accessed 23
353 January 2014]
- 354 Japan Standards Association. (1998) Japanese Industrial Standards K0101.
355 Japan Standards Association, Tokyo.
- 356 Japan Surfactant Industry Association. (1997) Annual Report of Japan. Japan
357 Surfactant Industry Association, Tokyo.

- 358 Kimaro, T., Tachikawa, Y. and Takara, K. (2002) Development of a Hydrological
359 Model for Predicting the Effects of Land Use Changes at Yasu River Basin.
360 *Proceedings of the International Conference on Urban Hydrology for the 21st*
361 *Century*, pp. 696–705.
- 362 Kobayashi, S. and Yamauchi, K. (1979) LAS (MBAS) and Phosphate in Nagara
363 River Water. *Jpn. J. Limnol.*, **40** (1), 29–39 (in Japanese).
- 364 Kobuke, Y. (1981) Anionic Surfactant Concentration in the River and Sea Waters
365 of Hyogo Prefecture and its Relationship to Phosphorus Concentration. *Jpn.*
366 *J. Limnol.*, **42** (4), 189–200 (in Japanese).
- 367 Kobuke, Y. (2004) LAS in Urban Rivers and Factors Contributing to Reduction of
368 their Concentrations. *Water. Sci. Technol.*, **50** (5), 355–361.
- 369 Kunimatsu, T. (1988) Mechanism of Pollution Discharged from River and
370 Presumption of Pollution Load into Lake Biwa. *Biwako Kenkyusyo*
371 *Go-syunen Kinenshi*, 49–63 (in Japanese).
- 372 Lake Biwa Research Institute and National Institute for Research Advancement.

- 373 (1984) Data Book of World Lakes. 513 LECS '84, Otsu, Japan.
- 374 Lake Biwa Environmental Research Institute. (1986) Data Book of Shiga
375 Prefecture Environmental Atlas. Lake Biwa Environmental Research
376 Institute, Shiga, Japan.
- 377 Lorenzen, C. J. (1967) Determination of Chlorophyll and Pheopigments,
378 Spectrophotometric Equations. *Limnol. Oceanogr.*, **12** (2), 343–346.
- 379 Miura, K., Nishiyama, N. and Yamamoto, A. (2008) Aquatic Environmental
380 Monitoring of Detergent Surfactants. *J. Oleo. Sci.*, **57** (3), 161–170.
- 381 Motomizu, S., Fujiwara, S., Fujiwara, A. and Toei, K. (1982) Solvent
382 Extraction-Spectrophotometric Determination of Anionic sSurfactants with
383 Ethyl Violet. *Anal. Chem.*, **54** (3), 392–397.
- 384 National Institute of Technology and Evaluation (2005) Linear Alkylbenzene
385 Sulfonic Acid and its Salt. *Initial Risk Assessment Report*, **1**, 4–7.
- 386 Ou, Z. A., Yediler, Y., He, Y., Jia, L., Kettrup, A. and Sun, T. (1996) Adsorption of
387 Linear Alkylbenzene Sulfonate (LAS) on Soils. *Chemosphere*, **32** (5),

388 827–839.

389 Perales, J. A., Manzano, M. A., Sales, D. and Quiroga, J. A. (1999)

390 Biodegradation Kinetics of LAS in River Water. *Int. Biodeterior.*

391 *Biodegradation*, **43** (4), 155–160.

392 Soejima, K. (1996) Studies on Spectrophotometric Determination of Anionic

393 Surfactants with Ethyl Violet. *Saga-ken Kankyo Senta Shoho*, **11**, 56–59 (in

394 Japanese).

395 Sugiyama, M., Hori, T., Kihara, S. and Matsui, M. (2005) Geochemical Behavior

396 of Trace Elements in Lake Biwa. *Limnology*, **6** (2), 117–130.

397 Sun, H. F., Takata, A., Hata, N., Kasahara, I. and Taguchi S (2003)

398 Transportation and Fate of Cationic Surfactant in River Water. *J. Environ.*

399 *Monit.*, **5**, 891–895.

400 Sütterlin, H., Alexy, R., Coker, A. and Kümmerer, K. (2008) Mixtures of

401 Quaternary Ammonium Compounds and Anionic Organic Compounds in the

402 Aquatic Environment: Elimination and Biodegradability in the Closed Bottle

- 403 Test Monitored by LC–MS/MS. *Chemosphere*, **72** (3), 479–484.
- 404 Takada, H. and Ishiwatari, R. (1987) Linear Alkylbenzenes in Urban Riverine
405 Environments in Tokyo: Distribution, Source, and Behavior. *Environ. Sci.*
406 *Technol.*, **21** (9), 875–883.
- 407 Tezuka, Y. (1985) C:N:P Ratios of Seston in Lake Biwa as Indicators of Nutrient
408 Deficiency in Phytoplankton and Decomposition Process of Hypolimnetic
409 Particulate Matter. *Jpn. J. Limnol.*, **46** (4), 239–246 (in Japanese).
- 410 Uchiyama, M. (1982) The Distribution of Methylene Blue Active Substances
411 (MBAS) in Lake Oze. *Jpn. J. Limnol.*, **43** (1), 39–43 (in Japanese).
- 412 Utsunomiya, A., Mori, Y. and Hasegawa, K. (1998) Adsorption of Linear
413 Alkylbenzenesulfonates and their Complexes with Cationic Surfactants on
414 River Sediment, and their Biodegradation in River Water. *Jpn. J. Toxicol.*
415 *Environ. Health*, **44** (4), 264–276 (in Japanese).
- 416 Vincent, C. H. and Glenwood, K. W. (1987) Structure-Activity Relationships for
417 Sorption of Linear Alkylbenzenesulfonates. *Environ. Sci. Technol.*, **21** (4),

Distribution and Geochemical Behavior of Anionic Surfactants Determined as Ethyl Violet
Active Substances in Lake Biwa, Japan
Hiroo Yoshida, Shunsuke Kudari, Toshitaka Hori, Masahito Sugiyama

418 370–373.

419 Yoshida, H., Kudari, S., Hori, T. and Sugiyama, M. (2009) Distribution of

420 Particulate Nonylphenol in Lake Biwa, Japan. *Water. Air. Soil. Pollut.*, **200**

421 (1–4), 267–276.

422 Yoshimura, K. and Nakae, A. (1982) Studies on the Biodegradation of Linear

423 Alkylbenzene Sulfonate by Activated Sludge (I). *Jpn. J. Water. Pollut. Res.*, **5**

424 (2), 19–25 (in Japanese).

425 Yoshimura, K., Hayashi, K., Kawase, J. and Tsuji, K. (1984) Existence of Anionic

426 Surfactants in River. *Jpn. J. Limnol.*, **45** (1), 51–60 (in Japanese).

427

428 **Figure captions**

429

430 **Fig. 1.** Vertical distributions of WT, pH, DEVAS, PEVAS, POC, PEVAS/POC,
431 and Chl-*a* at Station Ie-1 in Lake Biwa. WT: water temperature.

432

433 **Fig. 2.** Plot of PEVAS vs. POC. The regression line for the plot of PEVAS vs.
434 POC was constrained to pass through the origin to estimate an average
435 PEVAS/POC ratio. The correlation coefficient r was estimated from the PEVAS
436 and POC data.

437


438 **Fig. 3.** Plot of PEVAS/Chl-*a* vs. POC/Chl-*a*. The correlation coefficient r was
439 estimated from the PEVAS/Chl-*a* and POC/Chl-*a* data.

440

441 **Fig. 4.** Plot of PEVAS vs. POC. The data were classified into 3 groups: Group A,
442 samples of the epilimnion (squares) in September and October 2000 (pH


443 7.6–7.8; 0–13 m in September and 0–17 m in October); Group B, samples of the
444 hypolimnion (triangles) in September and October 2000 (pH 6.9–7.1; 20–72 m);
445 and Group C, samples from all depths (circles) on 2 and 11 April 2001 (pH
446 7.1–7.4). The regression line for the plot of PEVAS vs. POC was constrained to
447 pass through the origin to estimate an average PEVAS/POC ratio. The
448 correlation coefficient r was estimated from the PEVAS and POC data.
449

Distribution and Geochemical Behavior of Anionic Surfactants Determined as Ethyl Violet Active Substances in Lake Biwa, Japan
 Hiroo Yoshida, Shunsuke Kudari, Toshitaka Hori, Masahito Sugiyama



450


451 **Fig. 1.**



452

453 **Fig. 2.**

454



455


456 **Fig. 3.**

457

458

459

460



461

462

463

464 **Fig. 4.**