TITLE:

Spectrocolourimetry visualized differences in sexual skin colouration in macaques.

AUTHOR(S):

Ono, Eiri; Suzuki, Juri; Ishida, Takafumi

CITATION:

Ono, Eiri ...[et al]. Spectrocolourimetry visualized differences in sexual skin colouration in macaques.. *Folia primatologica* 2015, 86(3): 178-186

ISSUE DATE:

2015-05

URL:

<http://hdl.handle.net/2433/201386>

RIGHT:

© 2015 S. Karger AG, Basel; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

1 **Spectrocolorimetry visualized differences in sexual skin coloration in macaques**

2 Eiri Ono¹, Juri Suzuki² and Takafumi Ishida¹

4 ¹Department of Biological Sciences, Graduate School of Science, University of Tokyo, 7-3-1

5 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

6 ²Primate Research Institute, Kyoto University, Inuyama, Aichi, Japan

8 **The number of words**

9 4,106 words

11 **Corresponding author**

12 Eiri Ono, Unit of Human Biology and Genetics, Department of Biological Sciences, Graduate

13 School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

14 E-mail: onoeiri@bs.s.u-tokyo.ac.jp

15 Phone number: +81-3-5841-4496

16 Fax number: +81-3-3818-7547

18 **Running Head**

19 Spectrocolorimetry of macaque sexual skin

21 **Keywords**

22 *Macaca fuscata*, *Macaca mulatta*, sexual skin, spectrocolorimetry, CIELAB space

Abstract

Females of some catarrhines develop conspicuous sexual skin transformation in their hindlimbs. Among the macaques, one of the radiated and adapted catarrhine groups with diversified sexual skin transformation, differences in sexual skin coloration between the Japanese macaques *Macaca fuscata* and rhesus macaques *Macaca mulatta* have not been quantitatively analyzed. In this study, sexual skin coloration of these macaques was spectrophotometrically measured in the non-mating season (NMS) and mating season (MS) and represented in the CIELAB space with the variables of L^* , a^* and b^* . In the Japanese macaques, the average $\pm$ SD of L^* , a^* , and b^* was 53.61 ± 3.31 , 11.51 ± 4.57 and 6.66 ± 2.25 in the NMS, and 46.60 ± 2.78 , 19.97 ± 2.99 and 8.80 ± 1.34 in the MS, respectively; while in the rhesus macaques, the average of L^* , a^* , b^* was 60.09 ± 3.96 , 5.99 ± 4.59 and 5.83 ± 2.37 in the NMS, and 52.70 ± 6.54 , 13.62 ± 6.86 and 8.07 ± 1.43 in the MS, respectively. Sexual skin of Japanese macaques was consistently much redder (larger a^*) and darker (smaller L^*) than that of rhesus macaques. The smaller L^* suggested the more dermal melanin content in the Japanese macaques. These closely related macaque species have similar but distinct sexual skin coloration. Spectrophotometry is thus useful to suggest histo-physiological background of coloration.

Introduction

Females of some catarrhines develop conspicuous transformation around the perineal skin in their hindlimbs [Collings, 1926; Dixson, 1983; Anderson and Bielert, 1994; Nunn, 1999; Higham et al., 2010]; this “sexual skin” is thought to be tightly involved in socio-sexual signaling [Wickler, 1967; Gerald et al., 2007; Bradley and Mundy, 2008]. Changes in the sexual skin are simply referred as swelling and coloration. Exaggerated swellings are typical for females in the species with a multimale-multifemale social structure [Nunn, 1999]. Because the swelling reaches its maximum size at around the time of ovulation in the baboons [Higham et al., 2008], mandrills [Setchell and Wickings, 2004], chimpanzees [Deschner et al., 2004] and macaques [Higham et al., 2012], sexual skin swelling is thought to indicate female fertility. On the other hand, sexual skin coloration comprises several facets in the context of fertility and mating behavior [Czaja et al., 1975; Baulu, 1976; Waitt et al., 2006; Higham et al., 2008; Dubuc et al., 2009; Wallner et al., 2011; Watson et al., 2012].

The macaques comprising more than 20 species have radiated and adapted to various environments [Thierry, 2010]. Changes in the sexual skin are also diversified among them. Japanese macaques (*Macaca fuscata*) and rhesus macaques (*M. mulatta*) belong to the *fascicularis* lineage and the former live under various environments from warm to cool temperate zones in Japan, whereas the latter inhabit various environments in the mainland Asia such as India, China and southeast Asia. These two macaques share similar reproductive features [reviewed in Thierry, 2010]; for example, ovarian cycle length (Japanese macaques: 26.5 days; rhesus macaques: 28.3 days), a multimale-multifemale social structure, multi-mounting behavior and seasonal breeding but age at first birth delays in the Japanese macaques (Japanese macaques: five years; rhesus macaques: four years). Sexual swellings were recognized in adolescent females of both species [Anderson and Bielert, 1994], and females' red sexual skin draws male attention [Waitt, et al., 2006; Wallner et al., 2011]. However, sexual skin of the Japanese macaques swells less obviously and is much redder than

that of the rhesus macaques [observed by JS], differences in sexual skin coloration between the two species have not been quantitatively analyzed; these backgrounds prompted us to introduce spectrophotometry into the sexual skin coloration study.

Visual observation, digital photography, and spectrophotometry have been used for skin color measurement [Stevens et al., 2009]. Among them, spectrophotometry obtaining direct color data is applicable to reared animals and the obtained color data are objective without interferences such as stray light. For the evaluation of the obtained color data, there are two major protocols; a model of common color space, such as the CIELAB space that was developed for human color vision and a Receptor-Noise Limited (RNL) model [Vorobyev and Osorio, 1998] that stands species-specific data on the peripheral visual system. Data for species-specific visual system were not available in the Japanese macaques yet but were already obtained for the rhesus macaques [Bowmaker et al., 1978]. The CIELAB space gave great insights into the relationship between skin color and pigmentation, such as by melanin and hemoglobin, in humans [Weatherall and Coombs, 1992; Zonios et al., 2001; Alaluf et al., 2002a; Stephen et al., 2009]; in addition, it was proved in the mandrills (*Mandrillus sphinx*), a member of the papionines, that the CIELAB model gave qualitatively similar results to those in the RNL model [Renoult et al., 2011]. The CIELAB model is thus considered a useful tool to compare skin color in the catarrhines who have a uniform color vision system similar to that of humans [Jacobs and Deegan, 1999; Changizi et al., 2006].

In this study, we aimed to evaluate and distinguish sexual skin coloration objectively in closely related macaques categorized in the *fascicularis* group whose sexual skin appears similar [Nunn, 1999]. We thus obtained color profiles of sexual skin of the Japanese macaques and rhesus macaques with a spectrophotometer and analyzed them in the CIELAB space.

Materials & Methods

96

97 **Subjects**

98 Twelve female Japanese macaques (*Macaca fuscata*) aged from four to six years
99 and 13 female rhesus macaques (*Macaca mulatta*) aged from three to five years reared at
100 Primate Research Institute, Kyoto University, Inuyama, Aichi, Japan were recruited for this
101 study. When we started the data collection in the non-mating season, reddening and swelling
102 of sexual skin of the subjects were not recognized; they were likely to be menarche. We
103 measured their color profiles in July (non-mating season: NMS) and October (mating season:
104 MS) 2010, or in August (NMS) and November (MS) 2011. Two of 12 Japanese macaques
105 were studied in 2010 and the other 10 were in 2011, while six of 13 rhesus macaques were in
106 2010 and the other seven were in 2011. The detailed information of the subjects is in the
107 online supplementary table. This study followed the Guidelines for Care and Use of
108 non-Human Primates, Primate Research Institute, Kyoto University, and was approved by the
109 Ethics committee of animal experiments of the institute.

110

111 **Color measurements**

112 To measure sexual skin coloration, one spot was selected at 2-3 cm under the ischial
113 callosity of each side (two spots for each subject in one season, figure 1). Prior to the
114 measurement, the subjects were anesthetized with ketamine hydrochloride (50 mg/mL
115 Ketalar®, Sankyo-Parke-Davis & Co., Inc., Japan, 2.5 mg/kg) and medetomidine
116 hydrochloride (Domitor® Meiji Seika Kaisha, Ltd. Tokyo, Japan, 0.1 mg/kg), and then the
117 measurement spots were cleaned and shaved. The selected spots were measured in July
118 (NMS) and in October (MS) 2010, also in August (NMS) and in November (MS) 2011. A
119 single color measurement for each spot was conducted with a MINOLTA Spectrocolorimeter
120 CG-411C (for the CIE D65/10° illuminant/observer condition). Three variables in the
121 CIELAB color system, L^* , a^* , and b^* were extracted for each measurement. The average
122 values of L^* , a^* and b^* for each subject were calculated. The variable L^* , a^* and b^* represents

position on the light–dark, red/magenta–green, and yellow–blue axis, respectively. The L^* ranges from 100 to 0, whereas the a^* and b^* , can take any figure theoretically, ranging from about 100 to -100 in general.

Antidote to medetomidine, atipamezole hydrochloride (Domitor® Meiji Seika Kaisha, Ltd. Tokyo, Japan, 0.5 mg/kg), was administered for fully recovering after the measurement.

Data analysis

The data in July 2010 and August 2011 were combined for the NMS, while the data in October 2010 and November 2011 were combined for the MS. Changes in sex skin coloration between NMS and MS were tested with the paired t-test. The differences between the two species in each season were tested with the paired t-test, two-sample t-test and Welch's two-sample t-test if necessary. The relationship between the seasonal differences (MS - NMS) in L^* and a^* was analyzed with a linear regression. P value less than 0.05 ($p < 0.05$) was considered significant.

Results

Sexual skin coloration of Japanese macaques and rhesus macaques in the NMS and MS is visualized in figure 2 with the L^* , a^* , and b^* variables. In the Japanese macaques, the average±SD of L^* , a^* , and b^* was 53.61 ± 3.31 , 11.51 ± 4.57 and 6.66 ± 2.25 in the NMS, and 46.60 ± 2.78 , 19.97 ± 2.99 and 8.80 ± 1.34 in the MS, respectively; while in the rhesus macaques, the average of L^* , a^* , b^* was 60.09 ± 3.96 , 5.99 ± 4.59 and 5.83 ± 2.37 in the NMS, and 52.70 ± 6.54 , 13.62 ± 6.86 and 8.07 ± 1.43 in the MS, respectively. In the MS, decreases in the L^* and increases in the a^* and b^* were consistently observed in each subject of both the Japanese and rhesus macaques (table 1).

Differences in sexual skin coloration by the species and season are also represented in figure 2. In both seasons, the significant species differences were observed for the variable L^* (NMS: $t = -4.22$, $p < 0.001$ by two-sample t-test; MS: $t = -3.08$, $p < 0.01$ by Welch's two-sample t-test) and the variable a^* (NMS: $t = -3.01$, $p < 0.01$ by two-sample t-test; MS: $t = 3.04$, $p < 0.01$ by Welch's two sample t-test) but not for the variable b^* (NMS: $t = 0.89$, $p > 0.05$ by two-sample t-test; MS: $t = 1.31$, $p > 0.05$ by two-sample t-test). In both species, the significant seasonal differences were observed for the variable L^* (Japanese macaques: $t = 6.55$, $p < 0.001$ by paired t-test; rhesus macaques: $t = 6.48$, $p < 0.001$ by paired t-test), the variable a^* (Japanese macaques: $t = -6.20$, $p < 0.001$ by paired t-test; rhesus macaques: $t = -8.41$, $p < 0.001$ by paired t-test) and the variable b^* (Japanese macaques: $t = -3.58$, $p < 0.01$ by paired t-test; rhesus macaques: $t = -3.76$, $p < 0.01$ by paired t-test).

The correlation coefficient between the seasonal differences in L^* and a^* was -0.94 ($r^2 = 0.89$, $p < 0.001$) in the Japanese macaques, and -0.82 ($r^2 = 0.67$, $p < 0.001$) in the rhesus macaques.

Discussion

Sexual skin coloration of female Japanese macaques and rhesus macaques was spectrophotometrically measured and represented in the CIELAB space with which we can biologically interpret sexual skin coloration (table 1 and fig. 2). We analyzed the obtained color values from two aspects: seasonal change and species difference. Seasonal changes in the sexual skin coloration in the Japanese macaques and rhesus macaques were more or less similar; that is, sexual skin became darker, redder and yellower in the MS compared to NMS in both species. In detail, we also found that sexual skin of Japanese macaques was much redder (larger a^*) and darker (smaller L^*) than that of rhesus macaques through the NMS and MS (fig. 2); this is consistent with the general observations that sexual skin of Japanese

macaques appears to be redder than that of rhesus macaques. In terms of CIELAB values, similar dynamic changes in L^* , a^* and b^* were observed and it is thus conceivable that all the color values in the NMS as well as in the MS in both macaques are under the control of the same factor(s). The sexual skin color in the NMS is considered to represent the basic conditions without the surge of reproductive hormones, such as estrogen and progesterone [Nozaki et al., 1995]. Since in the NMS the variable L^* and a^* but not b^* were significantly different between the two species (fig. 2), it is suggested that the presence of different basic conditions that give lighter sexual skin to the rhesus macaques and redder sexual skin to the Japanese macaques. We thus demonstrate that the closely related these macaque species have similar but distinct sexual skin coloration.

Among various factors modifying skin coloration, two major pigments, melanin and hemoglobin, are of our interest. Melanin content modifies all the three color values, in particular, L^* [Takiwaki, 1998; Alaluf et al., 2002b]. In human skin coloration, changes in L^* is mainly due to epidermal melanin content [Alaluf et al., 2002b] that is negatively correlated with the L^* among human population [Alaluf et al., 2002a]. The seasonal decrease in the L^* in the Japanese and rhesus macaques may be attributed to the increase in dermal melanin content. The lower L^* values in the Japanese macaques both in the NMS and MS indicate more dermal melanin contents in the Japanese macaques. Melanin production is induced by ultra-violet (UV) light [Pathak et al., 1962; Jablonski and Chaplin, 2000; Jablonski and Chaplin, 2010] and the epidermal melanin content is subject to adaptive responses to UV light intensity; however, 17- β estradiol for example stimulates melanocyte tyrosinase activity in the human skin [Ranson et al., 1988; McLeod et al., 1994]. We thus cannot rule out possible contributions of other factor(s) to the melanization, and effects of estradiol on the dermal melanin content in macaques should be also elucidated to explain the consistently lower L^* in the Japanese macaques.

The a^* represents the red-green color axis and the higher a^* values were observed in the Japanese macaques in the NMS and MS. In humans, red color of the skin is mainly due to

the superficial vascular plexus in the upper dermis [Takiwaki, 1998; Alaluf et al., 2002b] where reproductive hormones also play certain roles such as improved vascularization by estrogens [Thornton, 2002]. As the reproductive hormones can modulate blood vessels in the sexual skin in the MS [Czaja et al., 1977], the increases in a^* in the MS may be due to vascularization and/or vasodilatation in the Japanese and rhesus macaques. Consistently lower melanin contents in the rhesus macaques were indicated by their higher L^* values both in the NMS and MS. This may correspond to the observation by Montagna et al. (1964) that rhesus macaque has little melanin in the epidermis as well as to the prediction by Collings (1926) that the sexual skin reddening is not due to the pigmentation but to the dynamics of sub-epidermal blood vessels. We conclude that melanin contents support the baseline of the a^* , whereas the increased a^* in the MS is due to blood hemoglobin in both species. The seasonal differences in L^* and a^* were highly correlated in both species (fig. 3). The high correlation indicates L^* and a^* are mostly under the same factors to change, namely, melanin and blood hemoglobin. The r^2 of regression line in the rhesus macaques was lower than that in the Japanese macaques. It indicates other factor(s) of the coloration in the rhesus macaques, for example, sexual swelling. The structural change of sexual skin may also modify the coloration.

It has been reported that adolescent females of both Japanese and rhesus macaques exhibit sexual swellings [Anderson and Bielert, 1994]. However, no Japanese macaques aged from four to six years exhibited sexual swelling, while 11 of the 13 rhesus macaques aged from three to five years did in this study. If the sexual skin coloration and swelling play a role in socio-sexual signaling, possible compensation of their bright red coloration for the lack of sexual swelling should be taken into consideration; it has been reported that hindlimb bright coloration might contain meaningful physiological information in Japanese macaques [Wallner et al., 2011].

For the analyses of spectrophotometric data of mandrills, the RNL model gave similar results as the CIELAB model [Renoult et al., 2011]. We also preliminarily analyzed

data by the RNL model, which were consistent with those in the CIELAB model (data not shown) but the CIELAB model gave higher resolutions. The CIELAB model is suitable for such coloration studies; however, the RNL model that is based on the species-specific visual system may have a potential to approach to understand cognitive consequences in the receiver of color signals. In conclusion, our study has opened objective and quantitative color measurements of sexual skin coloration and demonstrated their similar and dissimilar profiles in the closely related two macaque species. More detailed histological and molecular studies for these macaques' sexual skin will provide evolutionary insights in the mechanism of the sexual skin development in the primates. Spectrocolorimetry should help a further research on species-specific, sex-specific, and age-related coloration in various body parts especially bare skin among primates.

Acknowledgments

We are grateful to Prof. Kenichi Aoki, Department of Biological Sciences, University of Tokyo for his generous permission to use a spectrophotometer. We also thank Prof. Roscoe Stanyon for his comments on this manuscript. This work was supported by the Cooperation Research Program of Primate Research Institute, Kyoto University.

References

Alaluf S, Atkins D, Barrett K, Blout M, Carter N, Heath A (2002). Ethnic variation in melanin content and composition in photoexposed and photoprotected human skin. *Pigment Cell Research* 15:112–118.

Alaluf S, Atkins D, Barrett K, Blount M, Carter N, Heath A (2002). The Impact of

- 258 epidermal melanin on objective measurements of human skin colour. *Pigment Cell Research*
259 15: 119–126.
- 260 Anderson CM, Bielert CF (1994). Adolescent exaggeration in female catarrhine
261 primates. *Primates* 35: 283–300.
- 262 Baulu J (1976). Seasonal sex skin coloration and hormonal fluctuations in
263 free-ranging and captive monkeys. *Hormones and Behavior* 7: 481–494.
- 264 Bowmaker JK, Dartnall HJ, Lythgoe JN, Mollon JD (1978). The visual pigments of
265 rods and cones in the rhesus monkey, *Macaca mulatta*. *Journal of Physiology* 274: 329–348.
- 266 Bradley BJ, Mundy NI (2008). The primate palette: The evolution of primate
267 coloration. *Evolutionary Anthropology: Issues, News, and Reviews* 17: 97–111.
- 268 Brandon-Jones D, Eudey AA, Geissmann T, Groves CP, Melnick DJ, Morales JC,
269 Shekelle M, Stewart CB (2004). Asian primate classification. *International Journal of*
270 *Primatology* 25:97–164.
- 271 Changizi MA, Zhang Q, Shimojo S (2006). Bare skin, blood and the evolution of
272 primate colour vision. *Biology letters* 2: 217–21.
- 273 Collings MR (1926). A study of the cutaneous reddening and swelling about the
274 genitalia of the monkey, *Macacus rhesus*. *The Anatomical Record* 33: 271–287.
- 275 Czaja JA, Eisele SG, Goy RW (1975). Cyclical changes in the sexual skin of female
276 rhesus: relationships to mating behavior and successful artificial insemination. *Federation*
277 *Proceedings* 34: 1680–4.
- 278 Deschner T, Heistermann M, Hodges K, zBoesch C (2004). Female sexual swelling
279 size, timing of ovulation, and male behavior in wild West African chimpanzees. *Hormones*
280 *and Behavior* 46:204–215.
- 281 Dixon A (1983). Observations on the evolution and behavioral significance of
282 ‘sexual skin’ in female primates. In *Advances in the Study of Behavior Vol. 13* (Rosenblatt JS,
283 Hinde RA, Beer C, Busnel MC eds.), pp 63–106.
- 284 Dubuc C, Brent LJN, Accamando AK, Gerald MS, MacLarnon A, Semple S,

- 285 Heistermann M, Engelhardt A (2009). Sexual skin color contains information about the timing
286 of the fertile phase in free-ranging *Macaca mulatta*. *International Journal of Primatology* 30:
287 777–789.
- 288 Gerald MS, Waitt C, Little AC, Kraiselburd E (2007). Females pay attention to
289 female secondary sexual color: an experimental study in *Macaca mulatta*. *International*
290 *Journal of Primatology* 28: 1–7.
- 291 Higham JP, MacLarnon AM, Ross C, Heistermann M, Semple S (2008). Baboon
292 sexual swellings: information content of size and color. *Hormones and Behavior* 53: 452–62.
- 293 Higham JP, Brent LJN, Dubuc C, Accamando AK, Engelhardt A, Gerald MS,
294 Heistermann M, Stevens M (2010). Color signal information content and the eye of the
295 beholder: a case study in the rhesus macaque. *Behavioral ecology* 21: 739–746.
- 296 Higham JP, Heistermann M, Saggau C, Agil, Muhammad PFD, Engelhardt A (2012).
297 Sexual signalling in female crested macaques and the evolution of primate fertility signals.
298 *BMC Evolutionary Biology* 12: 89.
- 299 Jablonski N, Chaplin G (2000). The evolution of human skin coloration. *Journal of*
300 *Human Evolution* 39: 57–106.
- 301 Jablonski N, Chaplin G (2010). Human skin pigmentation as an adaptation to UV
302 radiation. *Proceedings of the National Academy of Sciences of the United States of America*
303 107: 8962–8968.
- 304 Jacobs GH, Deegan JF (1999). Uniformity of colour vision in Old World monkeys.
305 *Proceedings of the Royal Society B: Biological Sciences* 266:2023–8. doi:
306 10.1098/rspb.1999.0881
- 307 McLeod SD, Ranson M, Mason RS (1994). Effects of estrogens on human
308 melanocytes in vitro. *The Journal of Steroid Biochemistry and Molecular Biology* 49: 9–14.
- 309 Nozaki M, Mitsunaga F, Shimizu K (1995). Reproductive senescence in female
310 Japanese monkeys (*Macaca fuscata*): age- and season-related changes in
311 hypothalamic-pituitary-ovarian functions and fecundity rates. *Biology of Reproduction*

- 312 52:1250–1257.
- 313 Pathak MA, Riley FJ, Fitzpatrick TB, Curwen WL (1962). Melanin formation in
314 human skin induced by long-wave ultra-violet and visible light. *Nature* 193:148–150.
- 315 Ranson M, Posen S, Mason RS (1988). Human melanocytes as a target tissue for
316 hormones: in vitro studies with 1 alpha-25, dihydroxyvitamin D3, alpha-melanocyte
317 stimulating hormone, and beta-estradiol. *Journal of Investigative Dermatology* 91:593–8.
- 318 Renoult JP, Schaefer HM, Sallé B, Charpentier MJE (2011). The evolution of the
319 multicoloured face of mandrills: insights from the perceptual space of colour vision. *PloS*
320 *ONE* 6: e29117.
- 321 Setchell JM, Wickings EJ (2004). Sexual swelling in mandrills (*Mandrillus sphinx*):
322 a test of the reliable indicator hypothesis. *Behavioral Ecology* 15:438–445.
- 323 Stephen ID, Law Smith MJ, Stirrat MR, Perrett DI (2009). Facial skin coloration
324 affects perceived health of human faces. *International Journal of Primatology* 30: 845–857.
- 325 Stevens M, Stoddard MC, Higham JP (2009). Studying primate color: towards visual
326 system-dependent methods. *International Journal of Primatology* 30: 893–917.
- 327 Takiwaki H (1998). Measurement of skin color: practical application and theoretical
328 considerations. *The Journal of Medical Investigation* 44:121–126.
- 329 Thierry B (2010). The macaques - a double-layered social organization. In: *Primates*
330 *in Perspective*, 2nd ed. (Campbell C, Fuentes A, MacKinnon K, Bearder S, Stumpf R, eds.),
331 pp 229–241. Oxford University Press.
- 332 Thornton MJ (2002). The biological actions of estrogens on skin. *Experimental*
333 *Dermatology* 11: 487–502.
- 334 Tosi AJ, Morales JC, Melnick DJ (2000). Comparison of Y chromosome and mtDNA
335 phylogenies leads to unique inferences of macaque evolutionary history. *Molecular*
336 *Phylogenetics and Evolution* 17: 133–44.
- 337 Vorobyev M, Osorio D (1998). Receptor noise as a determinant of colour thresholds.
338 *Proceedings of the Royal Society B: Biological Sciences* 265: 351–8.

339 Waitt C, Gerald MS, Little AC, Kraiselburd E (2006). Selective attention toward
340 female secondary sexual color in male rhesus macaques. *American Journal of Primatology*
341 68: 738–44.

342 Wallner B, Aspernig D, Millesi E, Machatschke IH (2011). Non-lactating versus
343 lactating females: a comparison of sex steroids, sexual coloration, and sexual behavior in
344 Japanese macaques. *Primates* 52: 69–75.

345 Watson KK, Ghodasra JH, Furlong MA, Platt ML (2012). Visual preferences for sex
346 and status in female rhesus macaques. *Animal Cognition* 15: 401–7.

347 Weatherall IL, Coombs BD (1992). Skin color measurements in terms of CIELAB
348 color space values. *Journal of Investigative Dermatology* 99: 468–473.

349 Wickler W (1967). Socio-sexual signals and their intra-specific imitation among
350 primates. In *Primate Ethology* (Morris D eds.), pp 69–147. London, Weidenfeld and
351 Nicolson.

352 Zonios G, Bykowski J, Kollias N (2001). Skin melanin, hemoglobin, and light
353 scattering properties can be quantitatively assessed in vivo using diffuse reflectance
354 spectroscopy. *Journal of Investigative Dermatology* 117: 1452–7.

355

356

Table 1. Sexual skin color of the Japanese and rhesus macaques represented with L^* , a^* and b^* variables.

Subject	NMS			MS		
	L^*	a^*	b^*	L^*	a^*	b^*
Mf 1	53.31	9.21	5.64	51.56	12.67	8.63
Mf 2	51.08	10.39	3.72	45.38	18.92	7.39
Mf 3	55.39	15.45	9.88	49.79	20.33	8.78
Mf 4	55.79	10.45	6.49	49.13	17.53	5.66
Mf 5	52.50	17.49	8.63	47.39	22.92	9.18
Mf 6	55.23	5.23	4.13	43.25	20.55	8.18
Mf 7	57.08	10.85	7.50	49.26	18.31	8.97
Mf 8	49.27	19.55	9.42	45.93	23.68	9.77
Mf 9	50.89	13.30	6.31	44.51	19.33	9.43
Mf 10	48.00	12.50	8.05	43.32	20.90	10.85
Mf 11	55.99	9.92	7.30	45.67	21.56	10.16
Mf 12	58.79	3.82	2.84	43.99	22.94	8.62
Mm 1	64.47	2.43	6.44	58.94	4.87	6.13
Mm 2	61.29	4.04	8.40	58.95	8.93	9.28
Mm 3	62.14	0.72	6.35	53.34	9.41	8.91
Mm 4	67.56	-1.29	-0.13	54.76	7.24	6.88
Mm 5	58.88	5.95	6.72	49.26	15.62	7.03
Mm 6	62.62	1.43	3.76	62.44	4.39	8.10
Mm 7	61.40	5.37	5.24	59.79	10.14	8.29
Mm 8	61.49	7.05	6.84	56.83	13.34	9.38
Mm 9	54.53	13.85	8.90	45.49	22.85	8.10
Mm 10	60.21	5.75	5.43	51.72	15.78	8.64
Mm 11	54.51	11.70	7.64	42.61	25.30	11.08
Mm 12	55.26	10.36	3.63	45.38	17.52	6.11
Mm 13	56.87	10.48	6.61	45.69	21.71	7.01

Mf: Japanese macaque; Mm: rhesus macaque; NMS: non-mating season (July, 2010 and Aug, 2011); MS: mating season (October, 2010 and November, 2011)

390 **Legend to figure 1**

391 Measurement spots were selected inside the yellow circles under ischial callosities.

392

393 **Legend to figure 2**

394 Species and seasonal differences in the sexual skin coloration.

395 The dashed lines represent the interspecific comparisons in each season. The black bold lines

396 indicate the means and the black circles indicate the deviated values.

397 *** $P < 0.0001$; ** $P < 0.001$; * $P < 0.05$; NS: not significant.

398

399 **Legend to figure 3**

400 Correlation between the seasonal differences (MS-NMS) in L^* and a^* for the Japanese

401 macaques (Mf) and the rhesus macaques (Mm).

figure 1