

---

原 著

---

## EXPERIMENTAL STUDY OF REFLEX SHOCK

From the Department of Surgery, Mie Medical College,

By

SHIGERU IIDA

[Received for Publication : Aug. 17. 1953]

### I. EXPERIMENTAL STUDY OF PLEURAL SHOCK

The first description of pleural shock was given by ROGER in 1864, who attributed it to a reflex action from the stimulation of nerve-endings in the pleura. Since the artificial pneumothorax was introduced by FORLANINI in 1890 and the thoracic surgery has become a common practice, the number of pleural accidents is gradually increasing.

According to CAPPS, in any operative intervention on the pleural cavity, whether it is a therapeutic pneumothorax, a draining of empyema, a withdrawal of fluid by means of an aspirating needle or merely an exploratory thoracentesis, there lurks the danger of pleural shock. In this state a patient becomes faint, occasionally unconscious and pulse becomes weaker and weaker until it can no longer be detected by the finger. Convulsions and death are rare.

TAKEUCHI (1951) reported that by injecting tincture of iodine into the pleural cavity of rabbits he produced pleural shock, in which blood pressure fell, pulse rate became slower and pupils dilated. There were some evidences suggesting that pleural shock might be related to the vagus nerve.

Some time ago I experienced an accident that aroused my interest in this subject. A man, aged 27, underwent the aspiration of a pleural effusion. He suddenly became faint and then immediately lost consciousness, with the pupils dilated and the pulse almost impalpable and apparently slower. Routine emergency measures were done with no effect; then atropine was injected hypodermically, and the patient recovered in ten minutes. The essential feature in this case is obviously pleural shock caused by the puncture of the pleura. It seemed quite probable that the shock might be related to the vagus nerve, as the patient recovered with atropine. Suggested by this case, the following experimental study has been performed in cats.

#### EXPERIMENTAL TECHNIC

Cats, about 2 kg. in weight were used.

The blood pressure was recorded graphically by inserting a cannula into the right common carotid artery or the femoral artery and in order to measure the pulse rate exactly, an electrocardiograph was employed in some cases. Various stimulants were introduced into the pleural cavity with manometric control. At

necropsy, it was assured whether they had been injected exactly into the pleural cavity or not.

### RESULTS

#### Series I: Thermal Stimulation of Pleura.

A) Injection of hot water 80°C. (5 cc) into the pleural cavity.

Table I. (A)

| No. |                | Before injection | After injection |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 3'  | 5'  | 10' |
| 1   | Blood pressure | 120              | 120             | 114 | 114 | 110 |     |
|     | Pulse rate     | 208              | 208             | 236 | 220 | 232 |     |
| 2   | Blood pressure | 140              | 138             | 138 | 136 | 134 | 126 |
|     | Pulse rate     | 200              |                 | 212 | 208 | 204 | 200 |

The blood pressure showed no marked drop and the pulse rate became rather frequent.

B) Injection of cold water 5°C. (5 cc) into the pleural cavity.

Table I. (B)

| No. |                | Before injection | After injection |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 3'  | 5'  | 10' |
| 3   | Blood pressure | 160              | 180             | 164 | 164 | 170 | 174 |
|     | Pulse rate     | 240              |                 | 200 | 228 | 228 | 180 |

The blood pressure raised a little, but the pulse rate decreased soon after injection.

#### Comment.

It was expected that pleural shock might result, if the pleura was stimulated by hot or cold water. However, there occurred no marked fall of blood pressure and no slowness of pulse rate referable to a neurogenic reflex. Namely, pleural shock could not be produced by the application of thermal stimulus to pleura.

#### Series II: The Chemical Stimulation of Pleura.

A) Injection of 5% tincture of iodine (5 cc) into the pleural cavity.

Table II. (A)

| No. |                | Before injection | After injection |     |     |     |               |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|---------------|-----|
|     |                |                  | immediately     | 1'  | 3'  | 5'  | 10'           | 15' |
| 4   | Blood pressure | 200              | 230             | 176 | 164 | 160 | 160           | 160 |
|     | Pulse rate     | 180              |                 |     | 160 | 140 | 148           | 136 |
| 5   | Blood pressure | 140              | 180             | 101 | 76  | 68  | 62            | 58  |
|     | Pulse rate     | 220              | 200             | 180 | 165 | 160 | 150           | 144 |
| 6   | Blood pressure | 124              | 111             | 96  | 90  | 61  | Died 9' later |     |
|     | Pulse rate     | 200              |                 | 180 | 160 | 140 |               |     |

|    |                |     |     |     |     |     |     |
|----|----------------|-----|-----|-----|-----|-----|-----|
| 7. | Blood pressure | 140 | 138 | 120 | 104 | 114 | 116 |
|    | Pulse rate     | 200 |     | 190 | 180 | 180 | 200 |

Table II. (B)

| No. |                | Before injection | After injection |     |     |     |               |               |     |     |  |
|-----|----------------|------------------|-----------------|-----|-----|-----|---------------|---------------|-----|-----|--|
|     |                |                  | immedi-ately    | 1'  | 2'  | 3'  | 4'            | 5'            | 10' | 15' |  |
| 8   | Blood pressure | 130              | 156             | 86  | 64  | 44  | 10            | Died 5' later |     |     |  |
|     | Pulse rate     | 166              | 160             | 128 | 88  | 52  | 32            |               |     |     |  |
| 9   | Blood pressure | 144              | 190             | 96  | 100 | 108 | 104           | 108           | 130 | 140 |  |
|     | Pulse rate     | 220              | 176             | 176 | 172 | 172 | 168           | 168           | 192 | 208 |  |
| 10  | Blood pressure | 98               | 100             | 40  | 44  | 14  | Died 4' later |               |     |     |  |
|     | Pulse rate     | 224              |                 | 72  | 64  |     |               |               |     |     |  |

In all cases, there was a tendency to low blood pressure and bradycardia. In one case (No. 6), blood pressure fell about 28 mm Hg. a few minutes after injection and the animal died after about 9 minutes.

B) Injection of 10% hydrochloric acid (2 cc) into the pleural cavity.

These experiments were performed in May 1951, and in all animals without exception blood pressure fell and pulse rate became slower. In two animals, blood pressure fell about 50 mm Hg. immediately after injection and they died within five minutes. Comment.

CORDIER(1910) injected iodine into the pleural cavity and produced pleural shock. CROIZER (1927) found that pleural shock




Fig. 1. Injection of 10% hydrochloric acid 2 cc into the pleural cavity. (No.10)




Fig. 2. Injection of 10% hydrochloric acid 2 cc into the pleural cavity. (No. 9)

could be induced when iodine was injected directly into the lung. In rabbits, TAKEUCHI (1951) produced pleural shock by injecting 5% tincture of iodine into the pleural cavity.

In my experiments, the injection of 5% tincture of iodine and 10% hydrochloric acid into the pleural cavity of cats always produced pleural shock, in which blood pressure fell and pulse rate became slower. It was confirmed at necropsy that iodine or hydrochloric acid had been exactly injected into the pleural cavity, the pleura showing discolouration as a result of strong stimulation.

**Series III:** Pharmacostimulation of the pleura.

A) Injection of 0.1% adrenaline (0.5 cc) into the pleural cavity.

**Table III. (A)**

| No. |                | Before injection | After injection |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  |
| 11  | Blood pressure | 150              | 190             | 140 | 150 | 160 | 170 | 170 |
|     | Pulse rate     | 152              |                 | 124 | 124 | 124 | 124 | 124 |
| 12  | Blood pressure | 164              | 150             | 136 | 136 | 136 | 136 | 136 |
|     | Pulse rate     | 220              |                 | 260 | 264 | 260 | 260 | 260 |

B) Injection of 0.1% pilocarpine solution (1 cc) into the pleural cavity.

**Table III. (B)**

| No. |                | Before injection | After injection |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  |
| 13  | Blood pressure | 130              | 150             | 120 | 126 | 134 | 134 | 140 |
|     | Pulse rate     | 216              |                 |     | 196 |     |     |     |
| 14  | Blood pressure | 160              | 170             | 148 | 150 | 148 | 148 | 146 |

The blood pressure fell gradually and pulse rate somewhat decreased, but it seemed uncertain that the changes were due to a neurogenic reflex.

C) Injection of acetylcholine (10 mg) into the pleural cavity.

**Table III. (C)**

| No. |                | Before injection | After injection |    |    |     |    |     |
|-----|----------------|------------------|-----------------|----|----|-----|----|-----|
|     |                |                  | immediately     | 1' | 2' | 3'  | 4' | 5'  |
| 15  | Blood pressure | 140              | 160             | 46 | 40 | 66  | 80 | 82  |
|     | Pulse rate     | 172              |                 |    |    |     |    | 120 |
| 16  | Blood pressure | 136              | 130             | 24 | 50 | 54  | 60 | 62  |
|     | Pulse rate     | 200              |                 |    |    | 124 |    |     |

As soon as acetylcholine was injected into the pleural cavity, blood pressure fell rapidly and pulse rate decreased pronouncedly.

Comment.

By the injection of adrenaline and pilocarpine, reflex shock was not evident. By acetylcholine, however, there occurred marked fall of the blood pressure and bradycardia, which seemed likely due to a neurogenic reflex. This is a finding similar to that following the injection of tincture of iodine or hydrochloric acid, but is more pronounced than the latter, although the fall of blood pressure and bradycardia in case of acetylcholine may be the result of its absorption into general circulation.




Fig. 3. Injection of acetylcholine 10mg into the pleural cavity. (No. 15)

**Series IV:** Injection of Various Chemicals into the Pleural Cavity after Preliminary Injection of Atropine.

In the preceding experiments, it was demonstrated that the syndrome of pleural shock is almost the same as that induced by the intrapleural or hypodermic injection of acetylcholine, namely low blood pressure and bradycardia. It may be expected that preliminary administration of atropine will prevent the pleural shock following the injection of various chemicals into the pleural cavity.

Cats were given 0.05% atropine 0.02 cc (0.01 mg) per kilogram of body weight subcutaneously twice at an interval of twenty minutes. Then various chemicals were injected into the pleural cavity.

A) Injection of 5% tincture of iodine (5 cc) into the pleural cavity after preliminary injection of atropine.

Table IV. (A)

| No. |                | Before injection | After injection |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 3'  | 5'  | 10' | 15' |
| 17  | Blood pressure | 160              | 170             | 164 | 162 | 160 | 160 | 164 |
|     | Pulse rate     | 186              | 180             | 192 | 204 | 204 | 192 | 204 |
| 18  | Blood pressure | 170              | 220             | 180 | 180 | 164 | 156 | 160 |
|     | Pulse rate     | 172              | 200             | 192 | 184 | 180 | 184 | 184 |

At the moment when tincture of iodine was injected into the pleural cavity, cats showed manifestations of pain, and the blood pressure temporarily increased, but soon returned to the previous level. Pulse rate increased. Thus pleural shock, which was otherwise expected to occur, failed entirely to occur by the premedication of atropine.

B) Injection of 10% hydrochloric acid (2 cc) into the pleural cavity after preliminary injection of atropine.

Table IV. (B)

| No. |                | Before injection | After injection |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 3'  | 5'  | 10' | 15' |
| 19  | Blood pressure | 100              | 220             | 106 | 100 | 100 | 104 | 104 |
|     | Pulse rate     | 184              | 184             | 176 | 160 | 174 | 180 | 200 |
| 20  | Blood pressure | 144              | 220             | 140 | 140 | 134 | 136 | 140 |
|     | Pulse rate     | 208              | 192             | 192 | 188 | 180 | 176 | 200 |

pulse rate became a little slower, but blood pressure did not drop. Pleural shock was prevented by the premedication of atropine.

C) Injection of acetylcholine (10 mg) into the pleural cavity after preliminary injection of atropine.

Blood pressure fell after injection, presumably due to the direct action of absorbed acetylcholine to the heart. Pulse rate




Fig. 4. Injection of 10% hydrochloric acid 2 cc into the pleural cavity after preliminary injection of atropine. (No. 19)

Table IV. (C)

| No. |                | Before injection | After injection |    |     |    |    |
|-----|----------------|------------------|-----------------|----|-----|----|----|
|     |                |                  | immediately     | 1' | 2'  | 3' | 4' |
| 21  | Blood pressure | 100              | 102             | 90 | 80  | 82 |    |
|     | Pulse rate     | 210              |                 |    | 192 |    |    |
| 22  | Blood pressure | 112              | 114             | 70 | 71  | 84 | 86 |
|     | Pulse rate     | 224              |                 |    | 220 |    |    |

did not become slower.

Comment.

For the development of a pleural shock, one should think of vagal reflex. In this experiment, therefore, I administered atropine subcutaneously before the injection of various chemicals into the pleural cavity. The result was that pleural shock was arrested by atropine. I believe this is a direct evidence for the main

rôle of the vagus-nerve in the development of pleural shock.

**Series V:** Influence of Season and Atmospheric Temperature upon the Experimental Occurrence of Pleural Shock.

A) Injection of 10% hydrochloric acid (2 cc) into the pleural cavity in about 30°C atmospheric temperature.

This experiment was carried out in July 1951. The rectal temperature of cats was about 38°C.

**Table V. (A)**

| No. |                | Before injection | After injection |     |                 |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----------------|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 3'              | 5'  | 7'  | 10' | 15' |
| 23  | Blood pressure | 134              | 184             | 122 | 124             | 106 | 100 | 108 |     |
|     | Pulse rate     | 236              | 126             | 170 | 150             | 178 | 178 | 160 |     |
| 24  | Blood pressure | 140              | 190             | 130 | 142             | 140 | 130 | 130 |     |
|     | Pulse rate     | 188              | 160             | 216 | 192             | 180 | 184 | 188 |     |
| 25  | Blood pressure | 126              | 190             | 110 | 114             | 116 | 116 | 120 | 138 |
|     | Pulse rate     | 224              | 140             | 172 | 180             | 188 | 212 | 224 | 228 |
| 26  | Blood pressure | 108              | 180             | 114 | 124             | 110 | 80  | 78  |     |
|     | Pulse rate     | 212              | 200             | 188 | 192             | 224 | 228 | 224 |     |
| 27  | Blood pressure | 156              | 220             | 164 | 162             | 144 | 138 | 140 | 138 |
|     | Pulse rate     | 248              |                 | 228 | 240             | 232 | 200 | 192 | 180 |
| 28  | Blood pressure | 160              | 220             | 164 | 104             | 86  |     | 94  | 104 |
|     | Pulse rate     | 208              | 148             | 104 | 200 (arhythmia) |     |     | 120 | 176 |

The typical pleural shock, in which blood pressure fell remarkably and pulse rate became slower, was observed only in No. 28. However there was a suggestive pleural shock in animals Nos. 23, 25 and 26. The others did not show any sign of reflex shock.

B) Injection of 10% hydrochloric acid (2 cc) into the pleural cavity after cooling the body with ice about 30° C atmospheric temperature.

**Table V. (B)**

| No. |                | Before injection | After injection |     |     |               |     |               |     |
|-----|----------------|------------------|-----------------|-----|-----|---------------|-----|---------------|-----|
|     |                |                  | immediately     | 1'  | 3'  | 5'            | 7'  | 10'           | 15' |
| 29  | Blood pressure | 142              | 206             | 116 | 114 | 120           | 114 | 120           | 120 |
|     | Pulse rate     | 180              | 180             | 148 | 128 | 128           | 132 | 168           | 180 |
| 30  | Blood pressure | 150              | 200             | 180 | 22  | Died 4' later |     |               |     |
|     | Pulse rate     | 224              |                 | 144 | 48  |               |     |               |     |
| 31  | Blood pressure | 132              | 184             | 90  | 72  | 74            | 46  | Died 9' later |     |
|     | Pulse rate     | 184              |                 | 52  | 60  | 60            | 68  |               |     |

|    |                |     |     |     |     |     |     |     |     |
|----|----------------|-----|-----|-----|-----|-----|-----|-----|-----|
| 32 | Blood pressure | 130 | 172 | 74  | 62  | 62  | 62  | 74  | 90  |
|    | Pulse rate     | 180 | 108 | 136 | 112 | 180 | 180 | 128 | 144 |
| 33 | Blood pressure | 106 | 190 | 88  | 76  | 84  | 90  | 80  | 92  |
|    | Pulse rate     | 216 | 168 | 172 | 176 | 176 | 172 | 160 | 172 |

Cats were cooled until the rectal temperature fell to about 35.5°C, and one hour afterward, hydrochloric acid was injected into the pleural cavity.

Fall of blood pressure and bradycardia were observed in nearly all cases. In No. 30, which died after about 4 minutes, blood pressure dropped alarmingly. Comment.

From the fact that pleural shock was difficult to take place in summer, it was supposed that there might be some relation between the season or the atmospheric temperature and the pleural shock. Experiment Series V. (B) demonstrated that the cooling of the body was a predisposing factor of pleural shock. According to ISENSCHMIDT, descending of body temperature gives rise to the irritative state of the parasympathetic nervous system. This may explain the result in this experiment.

**Series VI:** Influence of the Section of the Bilateral Vagal Nerves upon Pleural Shock.

By cutting the vagal nerves on both sides in the neck, cats become dyspnoeic, but sooner or later recover with artificial respiration in most cases. After recovery to the normal condition, the following experiments were performed.

A) Injection of 10% hydrochloric acid (2 cc) into the pleural cavity after the section of the bilateral vagal nerves in about 30°C atmospheric temperature.

Table VI. (A)

| No. |                | Before injection | After injection |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 3'  | 5'  | 7'  | 10' | 15' |
| 34  | Blood pressure | 80               | 140             | 70  | 56  | 68  | 68  | 68  | 80  |
|     | Pulse rate     | 232              | 216             | 228 | 168 | 172 | 172 | 180 | 180 |
| 35  | Blood pressure | 90               | 120             | 74  | 54  | 64  | 56  | 74  |     |
|     | Pulse rate     | 264              |                 | 272 | 236 | 264 | 248 | 240 |     |
| 36  | Blood pressure | 116              | 200             | 94  | 86  | 90  | 90  | 90  | 100 |
|     | Pulse rate     | 168              |                 | 144 | 120 | 128 | 132 | 136 | 152 |
| 37  | Blood pressure | 144              | 220             | 140 | 128 | 102 | 90  | 90  | 96  |
|     | Pulse rate     | 176              | 160             | 164 | 164 | 160 | 164 | 156 | 160 |
| 38  | Blood pressure | 180              | 250             | 200 | 196 | 180 | 174 | 150 |     |
|     | Pulse rate     | 220              | 248             | 232 | 208 | 212 | 208 | 208 |     |

The typical pleural shock appeared in no case, although a tendency to shock was noticed in Nos. 34 and 36.

In view of the fact that the typical pleural shock did not occur also in Series



V. (A). There is no much significance in the result of this experiment, that the pleural shock was arrested by the section of the vagi in summer. The most one can say, may be that there is no appreciable difference between Series V. (A) and Series VI. (A).

B) Injection of 10% hydrochloric acid (2 cc) into the pleural cavity after cooling the body together with section of the bilateral vagal nerves in about 30°C atmospheric temperature.

Table VI. (B)

| No. |                | Before injection | After injection |     |               |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|---------------|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 3'            | 5'  | 7'  | 10' | 15' |
| 39  | Blood pressure | 144              | 146             | 142 | 138           | 140 | 140 | 140 | 138 |
|     | Pulse rate     | 200              | 196             | 196 | 200           | 208 | 188 | 188 | 192 |
| 40  | Blood pressure | 106              | 144             | 116 | 116           | 104 | 108 | 108 | 106 |
|     | Pulse rate     | 152              | 152             | 160 | 172           | 172 | 192 | 168 | 168 |
| 41  | Blood pressure | 134              | 150             | 104 | 108           | 108 |     |     |     |
|     | Pulse rate     | 140              | 140             | 156 | 180           | 180 |     |     |     |
| 42  | Blood pressure | 156              | 146             | 80  | Died 3' later |     |     |     |     |
|     | Pulse rate     | 192              | 192             | 184 |               |     |     |     |     |
| 43  | Blood pressure | 156              | 164             | 150 | 150           | 150 | 150 | 150 | 146 |
|     | Pulse rate     | 216              | 216             | 224 | 232           | 216 | 216 | 224 | 228 |

In only one case (No. 42) death occurred 3 minutes after injection of hydrochloric acid, but in others, fall in blood pressure and bradycardia were not produced. Comment.

For the reason that fall in blood pressure and bradycardia did not take place after the section of vagi and the preliminary freezing, it is decidedly clear that the pleural shock is arrested by the section of the bilateral vagal nerves. In short, the vagus nerve plays a principal rôle in the development of pleural shock.




Fig. 5. Injection of 10% hydrochloric acid 2 cc into the pleural cavity after section of bilateral vagal nerves (atmospheric temperature about 30°C). (No. 40)

#### SUMMARY

(1) The pleural shock did not result from the injection of hot or cold water into the pleural cavity.

(2) It developed by the injection of iodine or hydrochloric acid into the pleural cavity. In this state, there were low blood pressure and bradycardia.

(3) Similarly it developed by the injection of acetylcholine into the pleural cavity, but not by adrenaline or pilocarpine.

(4) The pleural shock was arrested by the subcutaneous administration of atropine before injection of various shock-producing chemicals into the pleural cavity.

(5) The freezing of the body makes a predisposing state of pleural shock.

(6) After cutting the both vagi, pleural shock did not occur. Therefore, pleural shock in cats seems to appear by a reflex through the vagus nerve.

(7) The typical manifestations of pleural shock, i. e. low blood pressure and bradycardia are analogous to those of the initial shock after head injuries, as Prof. Araki stated.

## II. EXPERIMENTAL STUDY OF REFLEX SHOCK IN ABDOMEN

The primary shock is believed to be a cardiovascular reflex phenomenon appearing immediately after injury, but the symptomatology is not well established. It seems to consist of low blood pressure, slow pulse, cold skin, pallor face, loss of consciousness and perspiration, whereas in the secondary shock pulse rate becomes frequent and blood pressure falls, as a result of progressive reduction in circulating blood volume. Almost the same features as in the primary shock appear in the case of fainting spells due to cerebral anemia following trivial stimuli, and initial shock after head injuries

GOLTZ observed that on repeated strokes on the frog's abdomen the heart beats become slower or even cease, and he attributed it to reflex inhibition of the heart through the vagus nerve. It has been described that the circulatory disturbance of the primary shock type are occasionally experienced in abdominal diseases accompanied by severe pain, for instance, Japanese "ATEMI", acute pancreatic necrosis, perforation of a gastric ulcer or cancer, and acute intestinal strangulation.

To clarify whether a primary shock with bradycardia and low blood pressure may be induced experimentally from the abdomen, especially from the upper abdomen, the present experiments in cats have been performed.

### EXPERIMENTAL TECHNIC

The method of experiments is the same as that in the pleural shock. Anesthesia was not done in all animals.

### RESULTS

#### Series I: Chemical Stimulation of Abdominal Cavity.

In case of acute perforation of the stomach or intestine, pulse is usually feeble and frequent. However, bradycardia and low blood pressure are occasionally seen, especially immediately after the perforation, presumably as the results of intra-abdominal vagal stimulation. To reveal whether the latter type of shock could be induced by the injection of stimulant chemicals into abdomen, the following experiments were performed.

- 1) Injection of 5% tincture of iodine 10cc into the abdominal cavity.  
 A) Injection into the upper abdominal cavity.

Table VII. (A)

| No. |                | Before injection | After injection |     |     |     |     | Remarks        |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|----------------|
|     |                |                  | immediately     | 3'  | 5'  | 7'  | 10' |                |
| 44  | Blood pressure | 150              | 200             |     | 152 |     | 144 |                |
|     | Pulse rate     | 180              | 210             |     | 200 |     | 200 |                |
| 45  | Blood pressure | 186              | 260             | 170 | 160 | 160 | 158 |                |
|     | Pulse rate     | 200              | 200             | 200 | 200 | 200 | 208 |                |
| 46  | Blood pressure | 160              | 210             | 148 | 140 | 140 | 130 |                |
|     | Pulse rate     | 220              | 220             | 220 | 220 | 200 | 192 |                |
| 47  | Blood pressure | 150              | 176             | 140 | 130 | 130 | 100 | Died 30' later |
|     | Pulse rate     | 240              | 240             | 232 | 220 | 208 | 120 |                |

- B) Injection into the lower abdominal cavity.

Table VII. (B)

| No. |                | Before injection | After injection |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|
|     |                |                  | immediately     | 3'  | 5'  | 7'  | 10' |
| 48  | Blood pressure | 170              | 184             | 164 | 164 | 160 | 160 |
|     | Pulse rate     | 168              | 200             | 200 | 200 | 220 | 220 |

- 2) Injection of hydrochloric acid into the upper abdominal cavity.  
 A) Injection of 10% hydrochloric acid 10 cc.

Table VIII. (A)

| No. |                | Before injection | After injection |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|
|     |                |                  | immediately     | 3'  | 5'  | 7'  | 10' |
| 49  | Blood pressure | 120              | 186             | 110 | 100 | 100 | 104 |
|     | Pulse rate     | 184              |                 | 220 | 200 | 192 | 188 |




Fig. 6. Injection of 10% hydrochloric acid 10 cc into the upper abdominal cavity. (No. 49)

## B) Injection of 20% hydrochloric acid 7.5 cc.

Table VIII. (B)

| No. |                | Before injection | After injection |     |     |    |     |
|-----|----------------|------------------|-----------------|-----|-----|----|-----|
|     |                |                  | immediately     | 3'  | 5'  | 7' | 10' |
| 50  | Blood pressure | 130              | 210             | 110 | 74  | 68 | 44  |
|     | Pulse rate     | 208              | 224             | 278 | 250 |    | 232 |

## C) Injection of 30% hydrochloric acid 2.0 cc.

Table VIII. (C)

| No. |                | Before injection | After injection |     |     |    |     |
|-----|----------------|------------------|-----------------|-----|-----|----|-----|
|     |                |                  | immediately     | 3'  | 5'  | 7' | 10' |
| 51  | Blood pressure | 120              | 190             | 102 | 90  | 80 | 46  |
|     | Pulse rate     | 248              |                 | 248 | 256 |    | 256 |

## D) Injection of 50% hydrochloric acid 10 cc.

Table VIII. (D)

| No. |                | Before injection | After injection |     |     | Remarks       |
|-----|----------------|------------------|-----------------|-----|-----|---------------|
|     |                |                  | immediately     | 3'  | 5'  |               |
| 52  | Blood pressure | 110              | 182             | 60  | 38  | Died 7' later |
|     | Pulse rate     | 172              | 204             | 228 | 200 |               |

## Comment.

In all these experiments, blood pressure and pulse rate increased immediately following injection probably due to a severe pain, and then gradually fell. WEGNER (1876) stated that the main cause of death in acute peritonitis was the absorption of toxic substances produced in the abdominal cavity. Similarly, the death of animals Nos. 47 and 52 might be caused by the absorption of iodine or hydrochloric acid. There was no evidence of reflex shock.

**Series II:** Injection of 10% Hydrochloric Acid 5 cc into the Abdominal Cavity of the Animals with Dysfunction of the Liver.

FINE (1949) and SHORR (1950) report that the hepatic anoxia is an important change in the shock-like state. It has been known that a major operation on a patient with hepatic dysfunction is liable to produce secondary shock. Therefore, it was expected that a primary shock might be more easily produced, when the function of the liver was disturbed. Carbon tetrachloride was given or the common bile duct was ligated, in order to impair the hepatic function of cats irreversibly.

1) The deterioration of the liver by carbon tetrachloride.

Cats had been given 20% carbon tetrachloride olive oil solution 1 cc per kilogram of body weight subcutaneously from 3 to 5 times every other day. The following experiments were carried out two days after the final injection.

A) Injection of 10% hydrochloric acid 5 cc into the abdominal cavity after

treatment with carbon tetrachloride 3 times every other day.

Table IX. (A)

| No. |                | Before injection | After injection |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|
|     |                |                  | immediately     | 3'  | 5'  | 7'  | 10' |
| 53  | Blood pressure | 114              | 170             | 110 | 110 | 114 | 114 |
|     | Pulse rate     | 200              |                 | 220 | 212 | 210 | 210 |

B) The same injection after treatment with carbon tetrachloride 4 times every other day.

Table IX. (B)

| No. |                | Before injection | After injection |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|
|     |                |                  | immediately     | 3'  | 5'  | 7'  | 10' |
| 54  | Blood pressure | 90               | 146             | 80  | 74  | 72  | 72  |
|     | Pulse rate     | 232              | 240             | 210 | 236 | 236 | 242 |
| 55  | Blood pressure | 112              | 168             | 106 | 100 | 94  | 94  |
|     | Pulse rate     | 200              | 228             | 228 | 240 | 228 | 228 |

2) The disturbance of hepatic function due to ligation of the common bile duct.

The cats did not eat anything after the ligation of common bile duct. 3 to 5 days later the liver showed spotted surface and fat-degeneration. Subcutaneous fat-tissues exhibited icteric colour. Injection of hydrochloric acid into the abdominal cavity was performed 3 resp. 5 days following the ligation of common bile duct.




Fig. 7. Injection of 10% hydrochloric acid 5 cc into the upper abdominal cavity of cats, treated with carbon tetrachloride 4 times every other day. (No. 54)

Table X.

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 56  | Blood pressure | 130              | 210             | 116 | 84  | 70  | 86  | 94  | 104 | 104 |
|     | Pulse rate     | 118              |                 | 196 | 188 | 161 | 110 | 128 |     | 152 |
| 57  | Blood pressure | 130              | 161             | 132 | 130 | 130 | 130 | 130 | 130 | 130 |
|     | Pulse rate     | 200              |                 | 232 | 228 | 228 | 224 | 224 | 224 | 224 |

## Comment.

In every case, blood pressure rose a little immediately after injection of hydrochloric acid and soon fell. However, pulse rate somewhat increased.

In short, the primary shock showing bradycardia and low blood pressure did not occur even in the cases of irreversible damage of the liver.

**Series III:** Injection of 10% Hydrochloric Acid 5 cc into the Upper Abdominal Cavity in Splanchnectomized Cats.


It is known that cerebral anemia with bradycardia and lower blood pressure is liable to occur in vagotonics, and that the electric stimulation in the proximal portion of cervical vagus nerve exerts an inhibitory action on the heart, resulting in low blood pressure. Therefore, it should be expected that after cutting sympathetic nerves on both sides, the parasympathetic nerves in the abdomen become predominant, and that reflex phenomenon through the parasympathetic nerve would be more easily produced. Thus injection of hydrochloric acid into the abdominal cavity was performed about one week after the section of upper lumbar sympathetic chains on both sides.

Table XI

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 58  | Blood pressure | 100              | 151             | 96  | 84  | 86  | 81  | 80  | 76  | 76  |
|     | Pulse rate     | 184              | 128             | 160 | 200 | 200 | 196 | 196 | 192 | 192 |
| 59  | Blood pressure | 150              | 200             | 160 | 150 | 146 | 146 | 144 | 144 | 144 |
|     | Pulse rate     | 160              | 128             | 144 | 148 | 148 | 148 | 148 | 150 | 150 |
| 60  | Blood pressure | 120              | 190             | 120 | 121 | 110 | 110 | 106 | 106 | 106 |
|     | Pulse rate     | 212              |                 | 208 | 228 | 200 | 188 | 188 | 188 | 188 |

## Comment.

In every case blood pressure fell gradually but remarkably, and pulse rate became slower immediately following injection and gradually increased later. In Nos. 58 and 59, the blood pressure fell temporarily and pulse rate decreased about one minute after injection, but soon returned to the former level. This response was supposed to be a vagal reflex, but not so strong as to be called a primary shock.



**Fig. 8.** Injection of 10% hydrochloric acid 5 cc into the upper abdominal cavity of splanchnectomized cat. (No.60)

**Series IV:** Injection of 10% Hydrochloric Acid 5 cc into the Upper Abdominal

Cavity after Fasting.

POPOW (1885), HAYASHI (1924), and TABUCHI (1943) observed that the hepatic function is impaired in the state of starvation. SUZUKI and his associates (1951) reported that the secondary shock due to a nervous stimulation is liable to occur in rabbits fasting for 3 to 5 days. For a long time, it has been believed that starvation often produces shock-like symptoms, i. e., starvation is a predisposing state of secondary shock. Thus the following experiments were performed in cats, whose liver had been reversibly damaged by fasting.

- 1) Injection of 10% hydrochloric acid 5 cc into the upper abdominal cavity of cats fasting for 2 days.

Table XII.

| No. |                | Before injection | After injection |    |    |    |     | Remarks |                |
|-----|----------------|------------------|-----------------|----|----|----|-----|---------|----------------|
|     |                |                  | immediately     | 1' | 2' | 3' | 4'  |         | 5'             |
| 61  | Blood pressure | 76               | 84              | 54 | 44 | 50 | 40  | 34      | Died 6' later. |
|     | Pulse rate     | 220              |                 | 80 | 80 | 80 | 100 | 100     |                |

- 2) The same injection in cats fasting for 3 days.

Table XIII

| No. |                | Before injection | After injection |     |     |     |     |     |     | 7'  | 10' |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  |     |     |     |
| 62  | Blood pressure | 110              | 156             | 82  | 120 | 80  | 80  | 80  | 74  | 60  |     |
|     | Pulse rate     | 180              |                 | 92  | 160 | 200 | 200 | 200 | 192 | 180 |     |
| 63  | Blood pressure | 110              | 172             | 110 | 104 | 104 | 100 | 102 | 98  | 96  |     |
|     | Pulse rate     | 220              |                 | 228 | 200 | 192 | 180 | 184 | 200 | 208 |     |
| 64  | Blood pressure | 130              | 200             | 144 | 106 | 96  | 94  | 94  | 92  | 100 |     |
|     | Pulse rate     | 168              |                 | 132 | 148 | 162 | 172 | 176 | 176 | 172 |     |




Fig. 9. Injection of 10% hydrochloric acid 5 cc into the upper abdominal cavity of cats fasting for 3 days. (No. 62)

## 3) The same injection in cats fasting for 7 days.

Table XIV

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 65  | Blood pressure | 130              | 198             | 114 | 112 | 100 | 96  | 92  | 96  | 100 |
|     | Pulse rate     | 220              |                 | 228 | 212 | 208 | 220 | 212 | 216 | 208 |

## 4) The same injection in cats fasting for 12 days.

Table XV.

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 66  | Blood pressure | 110              | 162             | 126 | 114 | 110 | 108 | 102 | 94  | 100 |
|     | Pulse rate     | 192              |                 | 204 | 192 | 188 | 184 | 180 | 190 | 192 |

## Comment.

In No. 61, pulse rate became slow, blood pressure fell rapidly and death occurred after about 6 minutes. However, as blood pressure was considerably low before injection of hydrochloric acid, the symptoms should not be taken for those of a reflex shock, but they were merely agonal signs. Furthermore, in Nos. 63, 65 and 66 blood pressure fell gradually but not remarkably, and pulse rate showed no changes or became frequent. However, in animals Nos. 62 and 64, bradycardia and low blood pressure were observed temporarily a few minutes following injection, and soon blood pressure returned to the former level and pulse rate increased. This response is a temporary vagal reflex and not so heavy as to be called a primary shock. The typical primary shock, such as observed in the thoracic cavity, did not take place from the abdominal cavity. In short, when a severe pain was given by injection of hydrochloric acid into the abdominal cavity after fasting for about 3 days in cats, there seems to occur at first a vagal reflex and then a secondary shock-like state. However, when fasting lasts for a longer period e. g. 7 to 12 days, such a temporary vagal reflex is hardly produced.

**Series V:** Cooling of Body in the State of Starvation and Primary Shock.

It was observed that the shock-like state occurred more frequently about 25 per cent in winter than in summer. WAKIN and GATCH (1943) experienced that animals in shock state are liable to die in cold circumstance. I have reported in this paper that pleural shock in cats is difficult to occur in summer, but when cooled in ice-box until the rectal temperature falls to about 35.5°C, cats can be brought into shock-state even in summer. In such cooled condition of the fasting animals the following experiments were carried out.

## 1) Injection of 10% hydrochloric acid 5 cc into the upper abdominal cavity of cats fasting for 4 days.



**Table XVI.**

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 67  | Blood pressure | 130              | 170             | 140 | 130 | 124 | 122 | 118 | 110 | 108 |
|     | Pulse rate     | 192              |                 | 172 | 172 | 188 | 176 | 200 | 200 | 196 |
| 68  | Blood pressure | 80               | 104             | 80  | 76  | 70  | 68  | 66  | 64  | 64  |
|     | Pulse rate     | 184              |                 | 132 | 132 | 136 | 132 | 132 | 144 | 152 |

2) The same injection in cats fasting for 7 days.

**Table XVII.**

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 69  | Blood pressure | 64               | 74              | 68  | 66  | 66  | 60  | 64  | 64  | 60  |
|     | Pulse rate     | 160              |                 | 200 | 192 | 192 | 192 | 192 | 200 | 200 |

3) The same injection in cats fasting for 10 days.

**Table XVIII.**

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 70  | Blood pressure | 120              | 142             | 90  | 106 | 100 | 104 | 106 | 106 | 106 |
|     | Pulse rate     | 196              |                 | 160 | 200 | 181 | 180 | 180 | 180 | 180 |


4) The same injection in cats fasting for 14 days.

**Table XIX.**

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 71  | Blood pressure | 80               | 132             | 78  | 84  | 82  | 80  | 78  | 80  | 82  |
|     | Pulse rate     | 164              |                 | 168 | 144 | 152 | 160 | 164 | 164 | 164 |

**Comment.**

When starving cats were cooled in ice-box, rectal temperature sometimes fell suddenly, and even death occurred, and a neurogenic shock was liable to develop during the manipulation of the femoral artery for the purpose of measuring blood pressure, just as in Takeuchi's experiment, in



**Fig. 10.** Injection of 10% hydrochloric acid 5 cc into the upper abdominal cavity of cats cooled after fasting for 14 days.(No.71)

which he observed a neurogenic shock to occur immediately following the release of an arterial clamp placed on the carotid artery. The reflex shock of vagal type, however, did not occur in any case in this series of experiment. An initial vagal reflex as observed in previous Series (IV) was evoked in no case but one (No. 70).

Even in the state of instability and disorder of the autonomic nervous function due to freezing and hunger, it seems that a primary shock hardly occurs by the injection of hydrochloric acid into the upper abdominal portion of cats. If a reflex shock would be induced by stimulation of abdominal cavity in man, some additional factors, such as astonishment, fear and other emotional excitement, might play a more important rôle in its development.

**Series V: Injection of Vagostigmine in Fasting Animals and Primary Shock.**

10% hydrochloric acid 5 cc was injected into the upper abdominal cavity of fasting cats about 20 to 30 minutes following the chemical stimulation of parasympathetic nerves by the administration of vagostigmine.

1) Injection of 10% hydrochloric acid 5cc into the upper abdominal cavity after preliminary injection of 0.05% vagostigmine 0.1cc per kilogram of body weight.

A) In cats fasting for 2 days.

**Tablea XX. (A)**

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 72  | Blood pressure | 120              | 180             | 110 | 100 | 100 | 100 | 100 | 96  | 96  |
|     | Pulse rate     | 160              |                 | 152 | 164 | 172 | 160 | 160 | 164 | 160 |
| 73  | Blood pressure | 120              | 196             | 128 | 120 | 102 | 100 | 104 | 108 | 114 |
|     | Pulse rate     | 156              | 80              | 148 | 111 | 141 | 148 | 148 | 160 | 160 |

B) In cats fasting for 3 days.

**Table XX. (B)**

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 74  | Blood pressure | 150              | 178             | 148 | 126 | 140 | 154 | 114 | 110 | 136 |
|     | Pulse rate     | 192              |                 | 88  | 136 | 176 | 160 | 172 | 184 | 212 |

C) In cats fasting for 5 days.

**Table XX. (C)**

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 75  | Blood pressure | 124              | 174             | 110 | 100 | 94  | 100 | 100 | 100 | 100 |
|     | Pulse rate     | 200              |                 | 200 | 196 | 176 | 176 | 176 | 176 | 160 |

|    |                |     |     |     |     |     |     |     |     |     |
|----|----------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 76 | Blood pressure | 130 | 176 | 130 | 120 | 108 | 104 | 106 | 108 | 100 |
|    | Pulse rate     | 152 |     | 104 | 148 | 148 |     | 152 | 156 | 156 |
| 77 | Blood pressure | 110 | 154 | 84  | 100 | 90  | 90  | 88  | 86  |     |
|    | Pulse rate     | 200 |     | 152 | 228 | 216 | 192 | 192 | 200 |     |

D) In cats fasting for 6 days.

Table XX. (D)

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 78  | Blood pressure | 90               | 120             | 70  | 66  | 66  | 66  | 66  | 64  | 62  |
|     | Pulse rate     | 184              |                 | 200 | 200 | 200 | 200 | 196 | 196 | 196 |
| 79  | Blood pressure | 140              | 160             | 130 | 130 | 130 | 126 | 122 | 116 | 120 |
|     | Pulse rate     | 224              |                 | 208 | 208 | 216 | 196 | 208 | 208 | 212 |
| 80  | Blood pressure | 100              | 164             | 114 | 92  | 84  | 82  | 80  | 80  | 78  |
|     | Pulse rate     | 168              |                 | 104 | 180 | 180 | 176 | 168 | 152 | 160 |




Fig. 11. Injection of 10% hydrochloric acid 5cc into the upper abdominal cavity of cats fasting for 6 days and injected preliminarily with 0.05% vagostigmine 0.1cc per kilogram of body weight. (No. 80)

E) In cats fasting for 3 weeks.

Table XX. (E)

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 81  | Blood pressure | 134              | 192             | 120 | 122 | 112 | 110 | 110 | 112 | 118 |
|     | Pulse rate     | 208              |                 | 192 | 172 | 172 | 168 | 176 | 176 | 160 |

A reflex shock of vagal type was not produced by injection of hydrochloric acid into the abdominal cavity following the subcutaneous injection of a small quantity of vagostigmine. However, temporary low blood pressure and bradycardia

which should be taken for a vagal reflex, was observed for a minutes following the injection of hydrochloric acid in animals Nos. 73, 74, 75 and 80.

2) Injection of 10% hydrochloric acid 5 cc into the upper abdominal cavity following preliminary injection of 0.05% vagostigmine 0.2 cc per kilogram of body weight.

A) In non-fasting cats.

Table XXI. (A)

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 7'  | 10' |     |
| 82  | Blood pressure | 136              | 166             | 76  | 70  | 66  | 62  | 52  | 54  | 48  |
|     | Pulse rate     | 196              |                 | 224 | 196 | 192 | 192 | 200 | 176 | 176 |
| 83  | Blood pressure | 90               | 124             | 86  | 86  | 90  | 80  | 74  | 94  | 104 |
|     | Pulse rate     | 160              |                 | 170 | 172 | 172 | 160 | 152 | 164 | 168 |

B) In cats fasting for 3 day.

Table XXI. (B)

| No. |                | Before injection | After injection |     |     |     |     |     |     |     | Remarks        |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|----------------|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |                |
| 84  | Blood pressure | 90               | 110             | 90  | 80  | 70  | 70  | 61  | 42  | 30  | Died 13' later |
|     | Pulse rate     | 192              |                 | 180 | 176 | 160 | 160 | 160 | 150 | 80  |                |

C) In cats fasting for 7 days.

Table XXI. (C)

| No. |                | Before injection | After injection |    |     |    |    |    | Remarks       |
|-----|----------------|------------------|-----------------|----|-----|----|----|----|---------------|
|     |                |                  | immediately     | 1' | 2'  | 3' | 4' | 5' |               |
| 85  | Blood pressure | 150              | 120             | 90 | 66  | 44 | 30 | 20 | Died 6' later |
|     | Pulse rate     | 180              |                 |    | 120 | 80 | 64 | 54 |               |

D) In cats fasting for 8 days.

Table XXI. (D)

| No. |                | Before injection | After injection |     |     |     |     |     |     |                |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|----------------|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10'            |
| 86  | Blood pressure | 86               | 94              | 54  | 36  | 34  | 30  | 20  | 18  | Died 10' later |
|     | Pulse rate     | 172              |                 |     | 80  | 60  | 60  | 40  | 40  |                |
| 87  | Blood pressure | 120              | 160             | 118 | 102 | 92  | 86  | 92  | 90  | 98             |
|     | Pulse rate     | 160              |                 | 148 | 148 | 144 | 140 | 128 | 112 | 112            |

In considering that a large quantity of vagostigmine exerts a toxic effect on respiratory and cardiovascular function, it seems to be the effect of vagostigmine rather than a neurogenic shock, that most of the cats in this series died

within 6 to 13 minutes after injection of hydrochloric acid.

Comment.

Even when the activity of cholinesterase is arrested by vagostigmine and therefore the vagus becomes highly sensitive, the reflex shock does not result from the injection of hydrochloric acid into the abdominal cavity, whereas a moderate vagal reflex develops temporarily. Many cats died within several minutes after injection of hydrochloric acid as a result of toxicity of vagostigmine.

**Series VII :** Iniection of 10% Hydrochloric Acid 5 cc into the Upper Abdominal Cavity of Cats Immediately after Food-Intake.

**Table XXII.**

| No. |                | Before injection | After injection |     |     |     |     |     |     |     |
|-----|----------------|------------------|-----------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                  | immediately     | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 88  | Blood pressure | 100              | 130             | 100 | 90  | 86  | 84  | 84  | 86  | 94  |
|     | Pulse rate     | 240              |                 | 210 | 224 | 232 | 232 | 220 | 208 | 208 |
| 89  | Blood pressure | 98               | 130             | 80  | 90  | 100 | 100 | 104 | 110 | 110 |
|     | Pulse rate     | 208              |                 | 180 | 200 | 200 | 208 | 200 | 192 | 200 |
| 90  | Blood pressure | 120              | 150             | 106 | 104 | 102 | 98  | 102 | 100 | 94  |
|     | Pulse rate     | 200              |                 | 196 | 200 | 192 | 188 | 192 | 200 | 200 |

Comment.

In No. 89. bradycardia and low blood pressure occurred temporarily one to two minutes after injection, however, the changes in blood pressure and pulse rate were not so remarkable as in a primary shock. They mean only a temporary vagal reflex.


**Series VIII :** Injection of 10% Hydrochloric Acid into the Abdominal Cavity of Toads.

1) Experimental Technic.

15 hibernating toads were used. Following the so-called Engelman's hanging method, toads were fixed gently in spine position with the thorax opened and the heart exposed, the heart point being pulled up with a serphine and connected to a lever. Heart beat was recorded by means of a kymograph. All toads recieved no anaesthetics.

2) Result.

Abdomen of toads was opened. Aafter the effects of laparotomy had disappeared, 1 cc of 10% hydrochloric acid was introduced into the upper abdomen. Immediately thereafter, they wriggled, showing pain reflex. In 3 cases there occurred tachycardia, in 7 no change, and in 5 bradycardia. In 2 cases of the last group the typical temporary cardiac arrest follo-



**Fig. 12.** Injection of 10% hydrochloric acid into the abdominal cavity of toads.

wed by marked bradycardia were observed.

### 3) Comment.

Even in toads the injection of hydrochloric acid into the abdominal cavity caused a marked bradycardia rather infrequently. Such bradycardia was observed only in so-called vagotonic animals and tended to be preceded by a temporary cardiac arrest. In this series heart beat of toads returned to preinjection state after about 2 minutes. In considering that in the experiment of pleural shock of cats, the decrease of pulse rate was parallel to the lowering of blood pressure, the latter change seems to have taken place also in toads. Such changes may be taken for a primary shock-like response.

### DISCUSSION

Both in the experiment in rabbits by TAKEUCHI and that in cats by me, the primary shock showing bradycardia and low blood pressure was not produced by the stimulation of the upper abdomen, where parasympathetic nerve seems to be predominant. However, we have occasionally experienced in man a vagal reflex effecting temporary low blood pressure and bradycardia in surgical operations on upper abdominal viscera, such as stomach, bile duct, and pancreas. A reflex phenomenon of this sort is observed not infrequently at the time of traction of the stomach or mechanical stimulation on the common bile duct by the dissection of adhesion, or even at the moment of opening the peritoneum. Clinical manifestations of the reflex are; patients complain of unpleasant sensation in the upper abdominal portion, and become pale, sweaty, but consciousness is maintained. Shortly after cessation of the stimulation, the blood pressure gradually increases and pulse rate returns to normal. Such a response is usually called "traction reflex"; and should not be called "shock". It is also evoked in the state of light anesthesia or in the initial stage of gastric ulcer perforation. The reflex took place in this study following injection of hydrochloric acid into the upper abdominal cavity of cats fasting for 2 or 3 days, but it was not so serious as to be called a neurogenic shock. However, the fact that the injection of hydrochloric acid into the abdominal cavity of toads brought about a marked bradycardia, especially in toads in vagotonic state, should indicate that the instability of autonomic nervous system would be an important factor of a primary shock in human being.

In the experiments with dogs and cats, MARTIN and BURSTEIN (1942) reported that the heart of the animals was arrested by electrical stimulation of the thoracic vagi, while the same stimulation of the abdominal vagi caused lowering of blood pressure. And many cases have been reported of sudden death due to cardiovascular and respiratory disorders during various manipulation of the vagus and its branches in the neck and thorax. In consideration of the fact, that the primary shock, which is produced in rabbits and cats by the stimulation of pleural cavity, is undoubtedly a vagal response, a primary shock of this sort seems to take place by the vagal stimulation 1) in the neck most frequently, 2) in the thorax relatively frequently, and 3) in the abdomen least frequently.

## SUMMARY

1) The primary shock, in which low blood pressure and bradycardia are characteristic, did not result from the injection of irritative chemicals, such as 5% tincture of iodine or 10% hydrochloric acid, into the abdominal cavity of normal cats, whereas the secondary shock, in which low blood pressure and tachycardia are characteristic, was liable to occur.

2) The primary shock did not develop by the injection of hydrochloric acid into the abdominal cavity even in the cats with an irreversible damage of the liver caused by carbon tetrachloride or by ligation of common bile duct. Therefore the disturbance of hepatic function does not seem to be an important predisposing factor of the primary shock.

3) Similarly, following division of bilateral abdominal sympathetics, the primary shock was not evoked by the injection of 10% hydrochloric acid (5 cc) into the abdominal cavity. Only in one case the injection of hydrochloric acid caused a temporary response of bradycardia with low blood pressure.

4) Immediately after the injection of hydrochloric acid into the abdominal cavity of cats fasting for 2 to 3 days, initial low blood pressure was induced and then the secondary shock followed. However, this initial response should not be taken for a primary shock, although it may be surely a vagal reflex. And such a vagal reflex is difficult to occur in cats fasting for a longer period (7 to 12 days). Therefore it appears that fasting for 2 to 3 days may provide an adequate predisposition to evoke a vagal reflex. Taking the primary shock for a serious manifestation of a vagal reflex, the primary shock in human being would be induced by the stimulation of the abdomen in a certain predisposed condition.

5) The shock of vagal type did not develop also in the cats which were cooled after fasting.

6) Immediately following injection of hydrochloric acid into the abdominal cavity of the fasting cats in vagotonic state induced by injection of vagostigmine, a temporary vagal reflex was observed, but it was not so serious as to be called a primary shock.

7) Injection of hydrochloric acid into the abdominal cavity of hibernating toads gave rise to a vagal reflex effecting the inhibition of cardiac function, sometimes the arrest of the heart, especially in those toads in which a parasympathotonic state was induced by a proper preliminary treatment.

8) If the primary shock would occur in man by the stimulation of abdominal viscera, it appears, from the experimental evidences in cats and others, that besides predisposing conditions above mentioned some other factors such as psychic excitement would be necessary to aggravate the instability of the autonomic nervous system. Also the cerebral anemia might serve as a supplementary factor.

## III. REPEATED STROKES ON CAT'S ABDOMEN

Goltz reported that repeated strokes on abdomen of frogs caused reflex inhibition of the heart through the vagus nerve, i. e. heart beat became slow or even

stopped. And also, it is known in Japan that a similar state occurs by an intense stroke on "MIZUOCHI" (epigastrium). Following experiments were carried out to examine, whether a primary shock-like response results from the strokes on abdomen of cats. Cats were fixed tightly in spine position without anesthesia, and repeated strokes on the upper abdominal wall were given from 30 to 50 times in about 15 seconds.

Table XXIII.

| No. |                | Before strokes | After strokes |     |     |     |     |     |     |
|-----|----------------|----------------|---------------|-----|-----|-----|-----|-----|-----|
|     |                |                | immediately   | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  |
| 91  | Blood pressure | 136            | 130           | 140 | 120 | 126 | 120 | 124 | 128 |
|     | Pulse rate     | 188            |               | 216 | 200 | 208 | 200 | 200 | 200 |
| 92  | Blood pressure | 110            | 86            | 100 | 86  | 90  | 94  | 96  | 100 |
|     | Pulse rate     | 220            |               | 240 | 220 | 218 | 200 | 200 | 200 |
| 93  | Blood pressure | 110            | 140           | 114 | 108 | 91  | 94  | 96  | 96  |
|     | Pulse rate     | 200            |               | 120 | 188 | 188 | 188 | 182 | 182 |

In No. 91, blood pressure and pulse rate showed no marked change. In No. 92, blood pressure dropped markedly immediately following cessation of strokes, but soon returned to an almost normal level. Pulse rate became rather frequent. In No. 93, pulse rate decreased immediately after strokes, but blood pressure was raised. In summary typical low blood pressure and bradycardia occurred in no case by stroking the abdomen of normal cats.

Next, the same test was made under local anesthesia of the abdominal wall with 0.5% procaine solution.

Table XXIV

| No. |                | Before strokes | After strokes |     |     |     |     |     |     |
|-----|----------------|----------------|---------------|-----|-----|-----|-----|-----|-----|
|     |                |                | immediately   | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  |
| 94  | Blood pressure | 126            | 104           | 76  | 72  | 70  | 71  | 76  | 76  |
|     | Pulse rate     | 118            |               | 212 | 200 | 188 | 176 | 176 | 180 |
| 95  | Blood pressure | 104            | 66            | 64  | 68  | 68  | 70  | 70  | 74  |
|     | Pulse rate     | 180            |               | 192 | 180 | 180 | 180 | 180 | 180 |

As soon as the stroking ceased, blood pressure fell pronouncedly, but pulse rate rather increased. Namely, bradycardia with low blood pressure did not ensue.

Tapping test of Colz was reexamined with 15 hibernating toads. A typical arrest of the heart or a marked bradycardia occurred in only one case, but moderate bradycardia responses resulted frequently from stroking.

Then, the stroking experiments were carried out in cats anesthetized with urethane.

**Series I:** Stroking of the Abdominal wall of cats between Meals.



A) Stroking of the upper abdominal wall.

Table XXV. (A)

| No. |                | Before strokes | After strokes |     |     |     |     |     |     |
|-----|----------------|----------------|---------------|-----|-----|-----|-----|-----|-----|
|     |                |                | immediately   | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  |
| 96  | Blood pressure | 120            | 90            | 108 | 100 | 88  | 90  | 100 | 110 |
|     | Pulse rate     | 172            | 168           | 160 | 160 | 152 | 152 | 148 | 156 |
| 97  | Blood pressure | 114            | 84            | 86  | 94  | 104 | 108 | 110 | 114 |
|     | Pulse rate     | 144            | 136           | 124 | 124 | 144 | 144 | 144 | 144 |
| 98  | Blood pressure | 120            | 88            | 103 | 104 | 107 | 117 | 120 |     |
|     | Pulse rate     | 200            | 200           | 200 | 200 | 200 | 204 | 204 |     |
| 99  | Blood pressure | 106            | 58            | 63  | 63  | 66  | 56  | 50  | 66  |
|     | Pulse rate     | 176            | 160           | 160 | 168 | 160 | 160 | 160 | 160 |




Fig. 13. Stroking of the upper abdominal wall. (No. 96)

In all cases, blood pressure fell pronouncedly following cessation of strokes and pulse rate somewhat decreased.

B) Stroking of the middle abdominal wall.

Table XXV. (B)

| No. |                | Before strokes | After strokes |     |     |     |     |     |     |
|-----|----------------|----------------|---------------|-----|-----|-----|-----|-----|-----|
|     |                |                | immediately   | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  |
| 96  | Blood pressure | 126            | 116           | 106 | 111 | 118 | 118 | 118 | 118 |
|     | Pulse rate     | 160            |               | 140 | 156 | 156 | 156 | 160 | 160 |
| 97  | Blood pressure | 116            | 96            | 100 | 96  | 98  | 100 | 100 |     |
|     | Pulse rate     | 140            | 140           | 136 | 136 | 140 | 140 | 140 |     |
| 98  | Blood pressure | 120            | 112           | 117 | 120 | 120 | 120 | 120 |     |
|     | Pulse rate     | 216            |               | 216 | 200 | 200 | 200 | 200 |     |

Blood pressure fell a little, but pulse rate showed no marked change.




Fig. 14. Stroking of the middle abdominal wall. (No. 96)

## C) Stroking of the lower abdominal wall.

Table XXV. (C)

| No. |                | Before strokes | After strokes |     |     |     |     |     |     |
|-----|----------------|----------------|---------------|-----|-----|-----|-----|-----|-----|
|     |                |                | immediately   | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  |
| 96  | Blood pressure | 120            | 110           | 126 | 120 | 120 | 122 | 120 | 122 |
|     | Pulse rate     | 160            |               | 168 | 168 | 164 | 160 | 160 | 160 |
| 97  | Blood pressure | 116            | 118           | 118 | 116 | 118 | 116 | 118 |     |
|     | Pulse rate     | 144            | 132           | 136 | 136 | 136 | 136 | 136 |     |
| 98  | Blood pressure | 120            | 118           | 120 | 125 | 126 | 126 | 126 |     |
|     | Pulse rate     | 200            | 200           | 220 | 216 | 216 | 216 | 216 |     |
| 99  | Blood pressure | 100            | 98            | 99  | 106 | 110 | 110 | 112 | 114 |
|     | Pulse rate     | 176            | 160           | 164 | 160 | 172 | 170 | 176 | 176 |

Blood pressure was raised a little and pulse rate became rather frequent.

## Comment.

Low blood pressure responses were evoked markedly by stroking of the abdominal wall in its upper part, moderately in the middle part, and nearly none in the lower part. Pulse rate decreased a little in stroking of the upper abdominal wall, but in Series I (B) and (C), it showed no change or became rather frequent.

Thus, decrease in both blood pressure and pulse rate was obtained by the strokes on the upper abdominal wall only, in which the parasy-




Fig. 15. Stroking of the lower abdominal wall. (No. 96)

mpathetics are predominant. However, it was not so serious as to be called a shock.

**Series II :** Stroking of the Upper Abdominal Wall after Food-Intake.

**Table XXVI.**

| No. |                | Before strokes | After strokes |     |     |     |     |     |     |
|-----|----------------|----------------|---------------|-----|-----|-----|-----|-----|-----|
|     |                |                | immediately   | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  |
| 100 | Blood pressure | 126            | 92            | 56  | 50  | 96  | 108 | 116 | 116 |
|     | Pulse rate     | 240            | 220           | 220 | 120 | 220 | 220 | 220 | 224 |
| 101 | Blood pressure | 102            | 63            | 80  | 61  | 49  | 46  | 52  | 61  |
|     | Pulse rate     | 216            |               | 200 | 208 | 200 | 200 | 183 | 188 |
| 102 | Blood pressure | 100            | 41            | 51  | 46  | 28  | 40  | 40  | 36  |
|     | Pulse rate     | 160            |               | 100 | 140 | 160 | 140 | 110 | 160 |

In all cases, pulse rate more or less decreased. Nausea followed by a remarkable bradycardia was noticed in Nos. 100 and 102, and arrhythmia for 3 minutes in No. 102. Blood pressure fell pronouncedly in all cases for 10 minutes or more.

**Comment.**

In this series of cats, in which the stomach was filled with food, remarkable lowering of blood pressure and slight decrease in pulse rate were observed.


Nausea and arrhythmia were noticed in some cases. Thus, a cardiovascular response effecting low blood pressure and slow pulse rate was evoked in cats after feeding and under anesthesia with urethane. Though it was not such a typical bradycardia as seen in pleural stimulation in cats or in tapping of the abdomen in toads, it may be called a primary shock-like symptom.

**Series III :** Stroking of the Upper Abdominal wall Following Sectioning of Bilateral Vagal Nerves.

Following bilateral division of vagi in the neck dyspnoea appeared and tracheotomy was necessary. Then the cats were stroked on the upper abdomen.

**Comment.**

Blood pressure fell pronouncedly, but pulse rate rather increased. Thus it was shown that bradycardia did not result after division of bilateral vagal nerves.



**Fig. 16.** Stroking of the upper abdominal wall after food-intake. (No. 100)

Table XXVII.

| No. |                | Before strokes | After strokes |     |     |     |     |     |     |     |
|-----|----------------|----------------|---------------|-----|-----|-----|-----|-----|-----|-----|
|     |                |                | immediately   | 1'  | 2'  | 3'  | 4'  | 5'  | 7'  | 10' |
| 103 | Blood pressure | 94             | 64            | 68  | 70  | 74  | 76  | 76  | 76  | 76  |
|     | Pulse rate     | 192            |               | 200 | 208 | 200 | 192 | 192 | 192 | 192 |
| 104 | Blood pressure | 102            | 74            | 76  | 68  | 64  | 66  | 68  | 70  | 70  |
|     | Pulse rate     | 200            | 200           | 208 | 208 | 212 | 216 | 208 | 200 | 200 |
| 105 | Blood pressure | 144            | 130           | 126 | 120 | 100 | 90  | 90  | 90  | 98  |
|     | Pulse rate     | 200            |               | 208 | 192 | 184 | 176 | 172 | 168 | 171 |

## SUMMARY

(1) No marked change was obtained in both blood pressure and pulse rate by repeated strokes on the abdominal wall of cats without anesthesia.

(2) Giving repeated strokes on the abdominal wall under local anesthesia with 0.5% procaine solution, blood pressure fell pronouncedly, and pulse rate became rather frequent.




Fig 17. Stroking of the upper abdominal wall after section of bilateral vagal nerves. (No. 103)

(3) Cats anesthetized with urethane responded with bradycardia and low blood pressure most pronouncedly to repeated strokes on the upper abdominal wall, moderately to that on the middle abdominal wall, but nearly none to that on the lower abdominal wall. This response seems to be a temporary vagal reflex.

(4) Decrease in both pulse rate and blood pressure was produced in cats by stroking of the upper abdominal wall immediately after food-intake under anesthesia with urethane. However, it was not such a typical bradycardia as seen in pleural shock of cats or in a critical state of toads following tapping of the abdominal wall or injecting of 10% hydrochloric acid into the abdominal cavity. It should be called a primary shock-like symptom.

(5) If the same experiment was done after bilateral sectioning of vagal nerves in the neck, blood pressure fell remarkably but pulse rate rather increased. Thus, sectioning of bilateral vagal nerves proved to inhibit the bradycardia response. (Acknowledgement should be made to Prof. Chisato Araki of Kyoto University, Prof. Toskisuhe Yamamoto and Assist. Prof. Tōkichi Takeuchi of this College for their kind suggestions. This study has been supported by a grant from the Educational Department Science Research Foundation.)

## References

1) Araki, Ch. : Pain and Shock, Rinsho no Shinpo, 1,76, Nagai-Shoten, Osaka, 1949. 2) Takeuchi, T. : Experimental Study of Reflex Shock, J. Mie Medical College, 11,55-72, 1951. J. Jap. Surg. Soc., 53,57-63, 1952. 3) Yamamoto, T., Takeuchi, T., and Iida, S. : Reflex Shock, Geka, 14, 135-138, 1952. 4) Miwa, S. : Accidents and Complications of the Artificial Pneumothorax Treatment, Jap. J. Thor. Surg., 3, 22, 1950. 5) Sakamoto, H. : Pleural Shock and Pleural Reflex, Jinko Kikyō Ryōhō, 125, Nihon-Rinsho-Sha, Osaka, 1950. 6) Shibusawa, K. : Shock no Rinsho, Igaku-Shoin, Tokyo, 1950. 7) Blalock, A., and Cressman, R. D. : Experimental Traumatic Shock, Further Studies with Particular Reference to the Role of the Nervous System, Surg. Gynec. & Obst., 68, 278-287, 1939. 8) Gapps, J.A. : Air Embolism Versus Pleural Reflex as the Cause of Pleural Shock, J. A. M. A., 109, 852, 1937. 9) Hamilton, C. E., and Rothstein, E. : Air Embolism, J. A. M. A., 104, 2226, 1936. 10) Andrew Morland : Pleural Shock, Lancet, 6588, 1021-1023, 1949. 11) Suda, I. : Shock, Geka, 12, 9-20, 1950. 12) Tabuchi, Y. : Experimental and Clinical Studies of Secondary Shock, J. Jap. Surg. Soc., 43, 1196-1267, 1942. 13) Shorr,

E., Zweifach, B. W., and Furchgott, R.F. : The Role of Hepato-Renal Vasotropic Principles in Experimental Shock, Research in Medical Science, 301, 1950. 14) Fine, J., Seligman, A.M., and Frank, H. A. : On the Specific Role of the Liver in Hemorrhagic Shock, Ann. Surg., 126, 1002-1009, 1947. 15) Suzuki, H., Shiino, E., Ohi, Y., and Nakamura, R. : Statistical Study on the Incidence of Experimental Nervous Shock, J. Kyoto Pref. Med. Univ., 49, 311-314, 1951. 16) Mine, M., and Tanigawa, H. : Significance of Tonic State of Autonomic Nervous System in Laparotomy, Rinsho Geka 5, 49-52, 1950. 17) Collins, V. J. : Abdominal Reflexes, Anesthesiology, 302-304, Lea & Febiger, Philadelphia, Pa., 1952. 18) Gullikson, M. J., McRae, J. H., and Campbell D.A. : Vagovagal Reflexes. Electrocardiographic Changes during Vagotomy, Surg. Gynec. & Obst. 89, 153, 1949. 19) Weeks, C., Ryan, B. J., and Van Hoy, J. H. : Two Deaths associated with Supradiaphragmatic Vagotomy, J. A. M. A., 132, 988-990, 1946. 20) Chester, S. T., Naffziger, H. C., Fisher, C., Rothenberg, S., and McCorkle, H.J. : The Effect of Stimulation of the Lower Intrathoracic Portion of the Vagus Nerve on the Cardiovascular System of Dogs, Surg. Gynec. & Obst., 94, 23-30, 1952.

## 和文抄録

## 神経反射性ショックの実験的研究

三重県立大学医学部外科学教室 (主任 山本俊介教授)

飯 田 茂

徐脈、血圧下降を呈する一次性ショックを猫の胸部及び腹部に於て検討した処、次の結論を得た。

## I. 肋膜性ショックの実験的研究

1) 肋膜性ショックは胸腔内に熱湯及び冷水を注入しても起し得ないが、沃度チンキ及び塩酸の注入により之を起し得た。

2) 同様に、胸腔内にアセチルコリンを注入することにより肋膜性ショックを起し得たが、アドレナリン及びピロカルピンの注入によりては起し得なかつた。

3) 夏期時には肋膜性ショックは発生し難いが、氷片にて体温を下降せしめると肋膜性ショックは発生し易くなる。即ち、寒冷刺激は肋膜性ショック準備状態

と云い得る。

4) 肋膜性ショックはアトロピンの術前注射又は頸部に於ける両側迷走神経切断により阻止される処から、迷走神経は肋膜性ショックの主役を演じるものであろう。

5) 以上の肋膜性ショックは荒木教授による頭部外傷後発生する一次性ショックの型と類似し、且つ、徐脈、血圧下降型を呈する。

## II. 腹部に於ける神経反射性ショックの実験的研究

1) 正常猫の腹腔内に5%沃度チンキ及び10%塩酸を注入しても一次性ショックはみられず、寧ろ頻脈、血圧下降を呈する二次性ショックが生ずる。

2) 更に術前四塩化炭素の投与及び総輸胆管の結紮

により肝機能障害を生ぜしめた後、或は両側腹部交感神経を切除した後、腹腔内に塩酸を注入しても一次性ショックは生じない。

3) 絶食2~3日後、腹腔内に塩酸を注入すれば、注入直後、徐脈、血圧下降を示し、やがて二次性ショック状態に移行する。この注入直後の変化は一時的に迷走神経反射現象が現われたものではあるが、一次性ショックと呼ぶ程のものではない。しかし、この現象は絶食期間が7~12日の長期に亘れば起り難いところから、2~3日間の絶食時には自律神経の不安状態が高いのであろう。即ち、迷走神経性反射現象には絶食2~3日間の自律神経不安状態が大いに関係する。かかる迷走神経反射の強く起つた場合が一次性ショックであるとすれば、以上のことから、人間に於ける腹腔よりの一次性ショックは自律神経不安状態に於て起り易いであろうことが想像される。

4) 飢餓と寒冷の条件下に於て、猫の腹腔内に塩酸を注入した実験では、徐脈、血圧下降型のショックは起らなかつた。

5) 飢餓時にワゴスチグミンを注射して、副交感神経を興奮せしめた後、塩酸を腹腔内に注入した処、注入直後、一時的迷走神経反射現象を示したが一次性ショックという程のものではなかつた。

6) 冬眠中の鼠に於て、腹腔内塩酸注入試験を行つ

た処、心搏停止、徐脈を呈する迷走神経性反射現象が起つた。しかし、特に副交感神経の興奮状態の強い鼠に於て、かかる反応が著明であつた。

7) 人に於て、腹部から一次性ショックが起り得るとすれば、それは自律神経不安定の状態の上に、更に他の因子、例へば、精神的興奮が高調し、それに疼痛及び臓器の不快感が刺激となり、脳血行が障碍せられて脳貧血の状態となり、一次性ショックを惹起するのではなからうか。

### Ⅲ. 腹壁叩打試験

1) 無麻酔時に於ける猫の腹壁を強打しても血圧、脈搏には変化を生じなかつた。

2) プロカインで局所麻酔された猫の腹壁を強打すれば血圧は著明に下降したが、脈搏はむしろ増加した。

3) ウレタンで麻酔された猫の腹壁を強打すれば徐脈及び血圧反応は上腹部に於て強く、中腹部に於て中等度、下腹部に於ては殆んど認められなかつた。即ち、この反応は一時的迷走神経反応であろうと思われる。

4) 摂食後、麻酔猫の腹部を強打すれば血圧、脈搏の減少を求したが、猫の肋膜性ショックや、鼠の腹壁叩打及び腹腔内塩酸注入によりみられた定型的な徐脈は得られなかつた。即ち、これは一次性ショック類似の症状であらう。

## 脾臓機能亢進症に於ける脾臓

The Spleen in Hypersplenism

R. J. Leffer

Am. J. Pathol. vol. 28 p. 303, 1952

脾臓機能亢進症、即ちバンテュー氏病、先天性溶血性黄疸、特発性血小板減少性紫斑病、更に最近では脾性白血球減少症、脾性貧血等に於て、脾臓は重要な役割をせしめるものであると考えられて来たが、先天性溶血性黄疸を除いては、殆んどこれらの組織病理学的変化は明らかにされていない。私は、末梢血球系減少、骨髓増殖、スプレノメガリー、脾摘出術後の貧血回復等臨床的特色を示す脾臓機能亢進症84例と、対照例221例の脾組織を顕微鏡的に検した結果、脾摘出術前にレ

ントゲン治療を受けた1例のバンテュー氏病を除く脾臓機能亢進症例のことに於て、マルピギー氏小体周縁層の巨大なことを発見した。同様な変化は224対照例中13例即ち5.8%に於てもみとめられたのであるが、脾臓機能亢進症例のそれ程明確ではなかつた。脾臓マルピギー小体の巨大な周縁層は、脾臓機能亢進症の診断的特色として価値のあるものかも知れない。

(戸部隆吉抄訳)