

ERRATUM

Open Access



Erratum to: Impact of medication adherence on renal function in comorbid patients with type 2 diabetes and depression: protocol for a cohort study

Hiroto Ito^{1*}, Tosiya Sato², Noriko Satoh-Asahara³ and Mitsuhiro Noda⁴

Erratum

The authors of this study protocol [1] would like to amend it and clarify that one of the medication adherence scales, the Morisky Medication Adherence Scale 8 (MMAS-8) [2–4], will not be used. This amendment will not cause any changes to the substantive contents of the article since the medication adherence will be assessed by the use of other measures. The Institutional Review Board of the National Center of Neurology and Psychiatry approved this change to the study protocol.

Author details

¹Department of Social Psychiatry, National Institute of Mental Health, National Center of Neurology and Psychiatry, Tokyo, Japan. ²Department of Biostatistics, Kyoto University School of Public Health, Kyoto, Japan. ³Division of Diabetic Research, Clinical Research Institute, National Hospital Organization Kyoto Medical Center, Kyoto, Japan. ⁴Department of Diabetes and Metabolic Medicine, National Centre for Global Health and Medicine, Tokyo, Japan.

Published online: 10 May 2017

References

1. Ito H, Sato T, Satoh-Asahara N, Noda M. Impact of medication adherence on renal function in comorbid patients with type 2 diabetes and depression: protocol for a cohort study. *BMC Fam Pract*. 2015;16:124. doi:10.1186/s12875-015-0339-1.
2. Morisky DE, Ang A, Krousel-Wood M, Ward H. Predictive validity of a medication adherence measure for hypertension control. *J Clin Hypertens*. 2008;10(5):348–54.
3. Krousel-Wood MA, Islam T, Webber LS, Morisky DE, Muntner P. New medication adherence scale versus pharmacy fill rates in seniors with hypertension. *Am J Managed Care*. 2009;15(1):59–66.
4. Morisky DE, DiMatteo MR. Improving the measurement of self-reported medication nonadherence: final response. *J Clin Epidemio*. 2011;64:258–63.

* Correspondence: ItoHiroto@ncnp.go.jp

¹Department of Social Psychiatry, National Institute of Mental Health, National Center of Neurology and Psychiatry, Tokyo, Japan

