

On an inequality of Chern numbers coming from anomaly calculations associated to ABJM(Aharony-Bergman-Jafferis-Maldacena) models

Tomohiro IWAMI (Kyushu Sangyo Univ.)

The Symposium of Algebraic Geometry, Kinosaki, Oct.21-Oct.25, 2013

1 Introduction

Whether there exists "expected" Miyaoka-Yau type inequality, especially including $c_i(X) (i > 2)$, on higher dimensional algebraic variety X of general type, have been posed by Y.Kawamata and N.Nakayama, since the proofs of flip theorem and the existence of the minimal models for 3-folds were established by S.Mori, because of the abundance conjecture on higher dimensional case. Recently, A.Bayer et al [3] showed the similar inequality as Bogomolov-Gieseker type one on projective 3-folds, based on the calculations of black-hole entropy on the context of the recent developments of AdS/CFT (or, gauge/gravity) correspondences. Considering these correspondences, M-theory are expected to have maximal supersymmetry as gravity theory in the area of low-energy. The ABJM (Aharony-Bergman-Jafferis-Maldacena) model [2] in the title, which is expected to be one of such theories having these properties, is a relativistic model with 2-dimensional space and time as a M2 brane in 11-dimensional model having 8-dimensional freeness, which gives explicit values of free energies on gauge theories by scattering amplitudes from the side of bulk (gravity theory) based on AdS/CFT correspondences.

2 Nambu brackets as descriptions of interactions of M2 branes

Considering Lagrangians for M2 branes as dynamics or statistics of them, the Chern-Simons theory as its action $S_{CS} = \frac{k}{4\pi} \int \text{Tr}(A \wedge dA + A^3)$ under a 'tHooft limit of large N with a fixed ratio N/k , where k an integer after quantization, where A is gauge field as 1-form, is required to give description of interactions of M2 branes without field strengths, because the induced equation of motion $F_{\mu\nu} = 0$. However, when the covariant derivatives for scalar fields X^I are introduced as $D_\mu X^I = \partial_\mu X^I + iA_{\mu ab}[T^a, T^b, X^I]$, based on the gauge invariances, where $[T^a, T^b, T^c] = if_d^{abc}T^d$ is called Nambu bracket [7] (this bracket is similar to the one in [8]), the Lie 3-algebra induced by such Nambu brackets is difficult to describe $N > 2$ M2 branes but for 2 M2 branes, under the gauge invariances [4, 5]. So, O.Aharony et al [2] posed the classical moduli theory with $U(N) \times U(N)$ theories under the smaller supersymmetry, and with not receiving quantum corrections as Chern-Simons terms $S_{CS} = \frac{k}{4\pi} \int (A_{(1)} \wedge dA_{(1)} - A_{(2)} \wedge A_{(2)})$ for gauge transformations $A_{(i)} \rightarrow A_{(i)} - d\Lambda_{(i)}$ ($i = 1, 2$) and $C_I \rightarrow e^{i(\Lambda_{(1)} - \Lambda_{(2)})} C_I$ for $A_{(i)}$ field variables and C_I superfields (two of them are chiral and the others are anti-chiral). Then, taking boundary contributions to Chern-Simons terms on \mathbb{C}^4 , $\delta S_{CS} = \frac{k}{2\pi} \int_{\text{boundary}} (\Lambda_{(1)} \wedge F_{(1)} - \Lambda_{(2)} \wedge F_{(2)})$ by Stokes theorem, where $F_{(i)}$ ($i = 1, 2$) are field strengths as 2-forms, the gauge

field strengths are quantized as $\int F_{(i)} \in 2\pi\mathbb{Z}$ on any closed 2-manifold. So, $\Lambda_{(i)} = 2\pi n/k$ for some integer n for the action to transform by 2π , then the moduli space is $\mathbb{C}^4/\mathbb{Z}_k$, where \mathbb{Z}_k -symmetry acts as $C_I \rightarrow e^{2\pi i/k} C_I$.

3 Axial anomaly for Chern-Simons terms in the deformation of NS5-branes and D3-branes

Main Results We consider axial anomaly associated to the brane construction via type IIB strings for Chern-Simons terms with the moduli $\mathbb{C}^4/\mathbb{Z}_k$, associated to probing $(1, k)$ D5-branes on the brane configurations of 2 NS5-branes and N D3-branes with supersymmetry fixing $(1, k)$ D5-branes in the 59-planes in (0123456789) -space [2] as non-relativistic case, motivated by [2, pp.17] with regards to the applications to elliptic fibered Calabi-Yau 4-folds $(\mathbb{T}^2)^4$ with the moduli $\mathbb{C}^4/\mathbb{Z}_k$ as geometric assumptions to have an inequality of Chern classes induced from the anomaly polynomials. So, for (p, q) D5-branes, the Chern-Simons action is $\frac{a}{g_4^2} \frac{p}{q} \int d^3x \epsilon_{\mu\nu\lambda} A^\mu \partial^\nu A^\lambda$ [6] where a is constant determined to be $g_4^2/4\pi$, then the anomaly polynomial [1] $I_{10}(F) = \frac{i^5}{(2\pi)^{45!}} \text{Tr} F^5$ where $F := \epsilon_{\mu\nu\lambda} A^\mu \partial^\nu A^\lambda$ is the field strength with the signature $\epsilon_{\mu\nu\lambda}$. For covariant derivatives $D_\mu X^I = \partial_\mu X^I + iA_{\mu ab}[T^a, T^b, X^I]$ with $[A, B, C] := AC^\dagger B - BC^\dagger A$ [2] weaker than [4, 5], because that the axial anomaly is given by $\partial_k \text{Tr}(\gamma^k \gamma^5 \mathbb{D})$, where γ^k, γ^5 Dirac gamma matrices, for Hermitian operator $\mathbb{D} := \gamma^\mu D_\mu = \gamma^\mu (\partial_\mu - \sum_a (iV_\mu^a + iA_\mu^a T^a \gamma_5)) =: \gamma^\mu (\partial_\mu - iV_\mu - iA_\mu \gamma_5)$ where V_μ vector fields and A_μ^a axial-vector fields, then the resulting anomaly polynomial is $\tilde{I}_{10}(\tilde{F}) = \text{Tr}\{(\epsilon_{\mu\nu\lambda} \gamma^k \gamma^5 \mathbb{D} A^\mu \partial^\nu \gamma^k \gamma^5 \mathbb{D} A^\lambda)^5\}$ up to constant multiplication for the axial gauge transformed \tilde{F} .

References

- [1] L.Alvarez-Gaumi, E.Witten, Phys.B **234**(1983) 269.
- [2] O.Aharony et al: $\mathcal{N} = 6$ superconformal Chern-Simons-matter theories, M2 branes and their gravity duals, [hep-th]0806.1218.
- [3] A.Bayer et al, [math.AG]1103.5010, 1106.3430.
- [4] J.Bagger, N.Lambert, Phys.Rev.D **77**(2008) 065008.
- [5] A.Gustavsson, Nucl.Phys.B **814**(2009) 53.
- [6] O.Bergman et al, [hep-th]9908075.
- [7] Y.Nambu, Phys.Rev D **7**(1973) 2405.
- [8] C.Simpson, Behavior at ..., RIMS symposium "Developments of Moduli: Conference" (June, 2013).