

# Moduli of Bridgeland semistable objects on the projective plane

Tokyo Institute of Technology Ryo Ohkawa

## Outlook

•  $X$  : a smooth projective surface  
 $\mathcal{D}(X)$  : the bounded derived category of Coh( $X$ )  
 Fix  $\alpha \in K(X)$  and  $\sigma$  : a Bridgeland stability condition on  $\mathcal{D}(X)$

Then we consider the moduli functor  $\mathcal{M}_{\mathcal{D}(X)}(\alpha, \sigma)$  of  $\sigma$ -semistable objects with class  $\alpha$  in  $K(X)$

•  $\sigma$  : geometric  $\implies \mathcal{M}_{\mathcal{D}(X)}(\alpha, \sigma) \cong \mathcal{M}_X(\alpha, \omega)$

$\mathcal{M}_X(\alpha, \omega)$  : the moduli functor of  $\omega$ -semistable coherent sheaves on  $X$ ,  
 $\omega$  : an ample divisor class in NS( $X$ )

•  $\sigma$  : algebraic  $\implies \mathcal{M}_{\mathcal{D}(X)}(\alpha, \sigma) \cong \mathcal{M}_B(\alpha_B, \theta_B)$

$\mathcal{M}_B(\alpha_B, \theta_B)$  : the moduli functor of  $\theta_B$ -semistable modules over a finite dimensional C-algebra  $B$   
 where  $\alpha_B \in K(\text{Mod } B)$ ,  $\text{Mod } B$  : an abelian category of finitely generated right  $B$ -modules

• As an application in the case of  $X = \mathbb{P}^2$  we show

$$\mathcal{M}_X(\alpha, \omega) \cong \mathcal{M}_B(\alpha_B, \theta_B)$$

**A Bridgeland stability condition  $\sigma$**

$\sigma$  consists of data  $(Z, \mathcal{A})$ ,

$$Z : K(X) \rightarrow \mathbb{C}, \quad \mathcal{A} \subset \mathcal{D}(X),$$

$Z$  : a group homomorphism,  $\mathcal{A}$  : a full abelian subcategory  
 These data satisfy some axioms

(For example)  
 $0 \neq E \in \mathcal{A} \implies Z(E) \in \mathbb{R}_{>0} \exp(\sqrt{-1}\pi\phi(E))$  with  $0 < \phi(E) \leq 1$

**Definition**  
 A nonzero object  $E \in \mathcal{A}$  : semistable  
 $\iff 0 \neq A \subset E \implies \phi(A) \leq \phi(E)$

**Moduli Functors  $\mathcal{M}_{\mathcal{D}(X)}(\alpha, \sigma)$**

$\mathcal{M}_{\mathcal{D}(\mathbb{P}^2)}(\alpha, \sigma) : (\text{scheme}/\mathbb{C}) \rightarrow (\text{sets})$

For a C-scheme  $S$ ,

$\mathcal{M}_{\mathcal{D}(\mathbb{P}^2)}(\alpha, \sigma)(S)$   
 $= \{E : \text{a family of } \sigma\text{-semistable objects in } \mathcal{A} \text{ with class } \alpha \in K(X)\} \subset \mathcal{D}(X \times S)$

## Main Theorem

In the case of  $X = \mathbb{P}^2$ , we find  $\sigma = (Z, \mathcal{A}) \in \text{Stab}(\mathbb{P}^2)$ ,  
 which is both geometric and algebraic.

**Main Theorem 0.1 (O)**

For any  $\alpha \in K(\mathbb{P}^2)$  with  $0 < c_1(\alpha) \cdot H \leq r(\alpha)$ ,  
 $\Phi_{\mathcal{E}} : E \mapsto \text{R Hom}_{\mathbb{P}^2}(\oplus E_i, E[1])$  gives the isomorphism

$$\mathcal{M}_{\mathbb{P}^2}(\alpha, H) \cong \mathcal{M}_{B_{\mathcal{E}}}(-\alpha_{B_{\mathcal{E}}}, \theta_Z^{\alpha}),$$

where  $H$  is the hyperplane class on  $\mathbb{P}^2$  and

(1)  $\mathcal{E} = \{\mathcal{O}_{\mathbb{P}^2}(1), \Omega_{\mathbb{P}^2}^1(3), \mathcal{O}_{\mathbb{P}^2}(2)\}$   
 or (2)  $\mathcal{E} = \{\mathcal{O}_{\mathbb{P}^2}(1), \mathcal{O}_{\mathbb{P}^2}(2), \mathcal{O}_{\mathbb{P}^2}(3)\}$  (Le Potier)  
 or (3)  $\mathcal{E} = \{\mathcal{O}_{\mathbb{P}^2}(2), \Omega_{\mathbb{P}^2}^1(4), \mathcal{O}_{\mathbb{P}^2}(3)\}$ .

## Geometric Stability

**Theorem 0.3 (O)**

$\alpha \in K(X)$  "normalized",  $\mu_{\omega}(\alpha)$  : the slope of  $\alpha$

We take  $\beta \uparrow \mu_{\omega}(\alpha)$  in NS( $X$ ) $_{\mathbb{R}}$ . Then

$$\mathcal{M}_{\mathcal{D}(X)}(\alpha, \sigma_{(\beta, \omega)}) \cong \mathcal{M}_X(\alpha, \frac{1}{2}K_X, \omega).$$

**Definition of  $\sigma_{(\beta, \omega)}$**  for  $\beta, \omega \in \text{NS}(X)_{\mathbb{R}} = \text{NS}(X) \otimes \mathbb{R}$  with  $\omega$  ample

$\exists \mathcal{A}_{(\beta, \omega)} \subset \mathcal{D}(X)$  : tilting of Coh( $X$ ) by  $\beta \cdot \omega$

$\exists Z_{(\beta, \omega)} : K(X) \rightarrow \mathbb{C} : E \mapsto -\int \exp(-\beta + \sqrt{-1}\omega) \text{ch}(E)$

$\sigma_{(\beta, \omega)} = (Z_{(\beta, \omega)}, \mathcal{A}_{(\beta, \omega)})$  defines a Bridgeland stability condition on  $X$

## Algebraic Stability

If  $\exists \mathcal{E} = \{E_0, \dots, E_n\}$  : a full strong exceptional collection on  $X$

Put  $B_{\mathcal{E}} = \text{End}_X(\oplus_i E_i)$

Then Bondal showed that  $\Phi_{\mathcal{E}} = \text{R Hom}_X(\oplus E_i, -)$  gives the equivalence

$$\Phi_{\mathcal{E}} : \mathcal{D}(X) \cong \mathcal{D}(\text{Mod } B_{\mathcal{E}}).$$

Put  $\mathcal{A}_{\mathcal{E}} = \Phi_{\mathcal{E}}^{-1}(\text{Mod } B_{\mathcal{E}}) \subset \mathcal{D}(X)$  defined by pulling back  
 $\text{Mod } B_{\mathcal{E}} \subset \mathcal{D}(\text{Mod } B)$  by  $\Phi_{\mathcal{E}}$ .

**Proposition 0.2 (O)**

For  $\sigma = (Z, \mathcal{A}_{\mathcal{E}})$ ,

$$\mathcal{M}_{\mathcal{D}(X)}(\alpha, \sigma) \cong \mathcal{M}_{B_{\mathcal{E}}}(\alpha_{B_{\mathcal{E}}}, \theta_Z^{\alpha})$$

where  $\alpha_{B_{\mathcal{E}}} = \Phi_{\mathcal{E}}(\alpha) \in K(\text{Mod } B_{\mathcal{E}})$  and

$\theta_Z^{\alpha}$  is the  $\theta$ -stability of  $B_{\mathcal{E}}$ -modules depending on  $\alpha$  and  $Z$  (and  $\mathcal{E}$ ).



## Conclusion and Further Studies

**Conclusion**

Main Theorem (2) gives another proof of the result by Le Potier (1994).  
 (Similar results are obtained by Barth in the case of  $r(\alpha) = 2$ )  
 $\implies$  The Bridgeland stability condition is the useful new concept to study the usual problem.

**Further Studies**

- Analysis of the wall-crossing phenomena of  $\mathcal{M}_{B_{\mathcal{E}}}(-\alpha_{B_{\mathcal{E}}}, \theta_{B_{\mathcal{E}}})$  when  $\theta_{B_{\mathcal{E}}}$  varies.  
 (the wall-crossing phenomena of  $\mathcal{M}_{\mathbb{P}^2}(\alpha, H)$  never occur because  $\text{NS}(\mathbb{P}^2)_{\mathbb{R}} = \mathbb{R}H$ )
- Our method is applicable for any surface  $X$  with a full strong exceptional collection (Generalization).