TITLE:

Rational curves on hypersurfaces

AUTHOR(S):

Furukawa, Katsuhisa

CITATION:

Furukawa, Katsuhisa. Rational curves on hypersurfaces. 代数幾何学シンポジウム記録 2008, 2008: 138-138

ISSUE DATE:

2008

URL:

<http://hdl.handle.net/2433/215036>

RIGHT:

Rational Curves on Hypersurfaces

古川 勝久 (FURUKAWA, Katsuhisa)

Waseda University
katu@toki.waseda.jp

We study the family of rational curves on hypersurfaces $/k = \bar{k}$, $\text{ch} \geq 0$.

Family of Rational Curves

Let $e \geq 1$, $d \geq 1$, and $n \geq 3$.

Def. For $X = X_d \subset \mathbb{P}^n$: hypersurf of deg d , set

$$R_e(X) := \left\{ \begin{array}{l} \text{smooth rat curves } C \text{ of deg } e \text{ in } \mathbb{P}^n \\ \text{lying in } X \end{array} \right\} \subset \text{Hilb}^{e+1}(X/k).$$

Expected Dimension

For $\mu := (n+1-d)e + n - 4 = \chi(N_{C/X})$,

$$R_e(X) \neq \emptyset \Rightarrow \dim R_e(X) \geq \mu.$$

Starting Point (Lines, $e = 1$)

Thm A (W. Barth-A. Van de Ven '78/79 (C), J. Kollár '96 ($\text{ch} \geq 0$)).

(a) $R_1(X) = \emptyset$ for gen X if $\mu < 0$.

(b) $R_1(X)$: smooth of dim μ for gen X if $\mu \geq 0$.

(c) $R_1(X)$: connected for any X if $\mu \geq 1$, except when $X = X_2 \subset \mathbb{P}^3$.

What can we say about the family $R_e(X)$ for deg $e \geq 2$?

Thm B (J. Harris-M. Roth-J. Starr '04 (C)).

$e \geq 1$, $d < (n+1)/2 \Rightarrow R_e(X)$: integral, loc comp int sch of dim μ for gen X .

Main Thm (Furukawa '07 ($\text{ch} \geq 0$)).

(a) $d \geq \max(e-2, 1) \Rightarrow R_e(X) = \emptyset$ for gen X if $\mu < 0$.

(b) $\left. \begin{array}{l} 1 \leq e \leq 3, d \geq 1, \text{ or} \\ e \geq 4, d \geq 2e-3 \end{array} \right\} \Rightarrow R_e(X)$: smooth of dim μ for gen X if $\mu \geq 0$.

(c) $R_2(X)$: connected for gen X if $\mu \geq 1$, except when $X = X_3 \subset \mathbb{P}^3$.

Exceptional Case of (c)

Prop. $X = X_3 \subset \mathbb{P}^3$: smooth cubic
 $\Rightarrow R_2(X)$ has 27 connected comp.

$\therefore$ For the lines $L_1, \dots, L_{27} \subset X$,

$$R_2(X) \hookrightarrow \bigcup L_i^* \subset (\mathbb{P}^3)^\vee$$

$$C \mapsto (2\text{-pl spanned by } C),$$

where $L_i^* := \{H \subset \mathbb{P}^3 : 2\text{-pl}, L_i \subset H\}$. $\square$

Key of the Proof

Fix $f : \mathbb{P}^1 \xrightarrow{\sim} C \subset \mathbb{P}^n$: deg e .

$$I_C^\circ := \{X = X_d \subset \mathbb{P}^n : \text{smooth}, C \subset X\},$$

$$Z_C^\circ := \{X \in I_C^\circ \mid h^1(N_{C/X}) \neq 0\}.$$

Tools

- ◆ Castelnuovo-Mumford regularity of I_C^2 .
- ◆ Theory of catalecticant matrices.
- ◆ Bounds of a_i for $f^*N_{C/\mathbb{P}^n} \simeq \bigoplus \mathcal{O}_{\mathbb{P}^1}(a_i)$.

Prop. assumption of (b)

$$\Rightarrow \text{codim}(Z_C^\circ, I_C^\circ) \geq \mu + 1.$$

$$\therefore 0 \rightarrow I_C^2 \rightarrow I_C \rightarrow N_{C/\mathbb{P}^n}^\vee \rightarrow 0 \text{ induces}$$

$$H^0(\mathbb{P}^n, I_C(d)) \xrightarrow{\delta_C} \text{Hom}_{\mathcal{O}_{\mathbb{P}^1}}(f^*N_{C/\mathbb{P}^n}, f^*(\mathcal{O}_{\mathbb{P}^n}(d)))$$

$$\cup \mathfrak{b} \quad \cup \mathfrak{f}$$

$$\hat{Z}_C = \delta_C^{-1}(Z) \twoheadrightarrow Z := \{\alpha \mid H^0(\alpha) : \text{not surj}\}.$$

$$(\text{Prop}) \leftarrow \text{codim}(\mathfrak{b}) \leftarrow \text{codim}(\mathfrak{f}). \quad \square$$