TITLE:

Synthesis of type 2 Lewis antigens via novel regioselective glycosylation of an orthogonally protected lactosamine diol derivative

AUTHOR(S):

Yamazaki, Yuji; Sezukuri, Kyohei; Takada, Junko; Obata, Hiroaki; Kimura, Shunsaku; Ohmae, Masashi

CITATION:

Yamazaki, Yuji ...[et al]. Synthesis of type 2 Lewis antigens via novel regioselective glycosylation of an orthogonally protected lactosamine diol derivative. Carbohydrate Research 2016, 422: 34-44

ISSUE DATE:

2016-03

URL:

<http://hdl.handle.net/2433/216657>

RIGHT:

© 2016. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>; The full-text file will be made open to the public on 1 March 2018 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

Carbohydrate Research, Full Paper

Synthesis of type 2 Lewis antigens via novel regioselective glycosylation of an orthogonally protected
lactosamine diol derivative

Yuji Yamazaki, Kyohei Sezukuri, Junko Takada, Hiroaki Obata, Shunsaku Kimura, Masashi Ohmae*

Department of Material Chemistry, Graduate School of Engineering, Kyoto University, Kyoto-
daigaku-katsura, Nishikyo-ku, Kyoto 615-8510, Japan

Tel.: +81-75-383-2403; fax: +81-75-383-2401

E-mail address: ohmae@peptide.polym.kyoto-u.ac.jp

*Corresponding author

Abstract

The novel and efficient synthesis of type 2 Lewis antigens is reported in this study. The rationally designed lactosamine-3,2'-diol derivative with an orthogonal set of protecting groups is efficiently glycosylated with a benzyl protected 1-thio-L-fucoside donor in a unique regioselective manner to produce Lewis x (Le^x) and Lewis y (Le^y) derivatives in good yields. These derivatives can be prepared not only exclusively but also synchronously by choosing the appropriate reaction temperature and donor-acceptor molar ratio. The Le^x derivatives are easily converted into sulfated or non-sulfated Le^x bearing a terminal azido functionalized oligo-(ethyleneoxide) linker; the Le^y derivative having the same linker can also be prepared, all of which can be further used for the chemical modification of other compounds and materials.

Keywords: Type 2 Lewis antigens; Sulfated Lewis x; Lewis y; Regioselective glycosylation; The Heyns rearrangement

1. Introduction

Lewis antigens are well-known as glycan-based blood group antigens,¹ which are classified as type 1 and type 2, depending on the disaccharide core structure. Type 1 Lewis antigens, Lewis a and Lewis b, have a common backbone [$\rightarrow 3\text{Gal}\beta(1\rightarrow 3)\text{GlcNAc}\beta 1\rightarrow$] and exist widely in the membrane of erythrocytes. The type 2 core structure [$\rightarrow 3\text{Gal}\beta(1\rightarrow 4)\text{GlcNAc}\beta 1\rightarrow$] is also found

in numerous glycoconjugates; however, the distribution of its fucosylated derivatives, Lewis x (Le^x) and Lewis y (Le^y), classified as type 2 Lewis antigens (T2-LAs), is limited to some epithelial cells and leukocytes. Notably, T2-LAs are overexpressed in various tumor cells,² and are thus frequently used as biomarkers for the diagnosis of cancer. Furthermore, these T2-LAs are sometimes found in their sulfated form, which includes a 6-*O*-sulfo-GlcNAc residue.³ The Le^x determinant also plays a critical role in various biological events such as inflammation, lymphocyte homing, and infection of pathogens.⁴ Thus, T2-LAs have attracted much attention as promising target compounds for cancer therapy as well as for the treatment of inflammation, infectious diseases, etc.

In order to utilize T2-LAs and their derivatives as bioactive compounds, it is essential to establish versatile, widely applicable synthetic methods. A large number of reports on the synthesis of Lewis antigens have been published to date;⁵ however, none of them have reported the use of common key intermediates to construct T2-LAs, including their sulfated form. In the present study, we report the successful and rapid assembly of T2-LAs (**3–5**) via a refined disaccharide key intermediate **2**, which can be readily prepared from lactulose **1** through the Heyns rearrangement method⁶ (Figure 1). This method is very convenient to obtain useful 2-amino-2-deoxy sugars, particularly lactosamine derivatives.⁷

Figure 1. Efficient synthesis of T2-LAs (**3–5**) via the refined orthogonally protected T2 disaccharide derivative **2** derived from lactulose **1**.

2. Results and discussion

2.1 Refinement of the molecular design of key intermediate **2**

In a previous paper,⁸ we reported the synthesis of T2-LAs having a set of orthogonal protecting groups via the useful disaccharide intermediate **2'** (Figure 2). Compound **2'** was found to be an excellent intermediate for the synthesis of T2-LAs, but had a major drawback concerning the

removal of the anomeric 4-methoxyphenyl (PMP) group: the 6-*O*-*tert*-butyldimethylsilyl (TBDMS) group was found to be labile under oxidation conditions by cerium(IV) ammonium nitrate (CAN). It is acceptable to produce neutral, non-sulfated T2-LAs; however, in order to synthesize structurally complicated and highly bioactive sulfated T2-LAs, the protecting group at glucosamine C6 must be stable under oxidation conditions. Thus, we refined the molecular design from **2'** to **2**: the stability of the *tert*-butyldiphenylsilyl (TBDPS) group is several hundred times higher than that of TBDMS under acidic conditions, whereas both show similar susceptibility to tetra-*n*-butylammonium fluoride (TBAF).⁹ Therefore, TBDPS was selected as a more suitable protecting group for the C6 position of the glucosamine residue. Further, the 4,6-*O*-benzylidene protecting group for the Gal residue was replaced with the *p*-methoxybenzylidene group, which is more rapidly removed by hydrogenolysis.

Figure 2. Refinement of the molecular design of key intermediate **2**.

According to our previous report,⁸ the lactosamine derivative of **6** was readily prepared from lactulose **1** via the Heyns rearrangement (Scheme 1). After removal of the acetyl protecting groups in **6**, the 4'- and 6'-hydroxy moieties were protected with a *p*-methoxybenzylidene group to afford **7** in 69% yield via a two-step procedure. The 6-OH moiety in **7** reacted selectively with TBDPS-Cl

in pyridine, giving **8** in moderate yield (59%). We originally controlled the selectivity in the regioselective 3'-*O*-benzoylation by lowering the reaction temperature (−50 °C). However, the unfavorable 3,3'-di-*O*-benzoyl product was formed even under the optimized conditions designed to obtain the target 3'-mono-*O*-benzoyl product. In the present study, we exclusively obtained the target product **2** at ambient temperature in 73% yield by employing a metal-coordinated regio- and chemoselective nucleophilic substitution method.¹⁰

Scheme 1. Reagents and conditions: (a) 1) MeONa / MeOH, 2) *p*-anisaldehyde dimethylacetal, (±)-10-camphorsulfonic acid/DMF, 30 °C, 24 h, 69% (2 steps); (b) TBDPS-Cl/pyridine, rt, 64 h, 59%; (c) Bu₂SnCl₂, PEMP, BzCl/THF, rt, 48 h, 73%. NPhth: phthalimido, PMP: 4-methoxyphenyl, PEMP: 1,2,2,6,6-pentamethylpiperidine.

2.2 On demand synthesis of Le^x and Le^y derivatives by regioselective α -fucosylation of **2**

Glycosylation of **2** with benzyl-protected phenyl 1-thio-L-fucopyranoside (**9**)¹¹ is the extremely unique reaction throughout the synthesis of T2-LAs (Scheme 2). In order to obtain the Le^x derivative **10**, the glycosylation was carried out at lower temperature (−78 °C) with a slight excess of **9** over **2**; under these conditions, **10** was isolated as the sole product in 76% yield (Table 1, entry 1). Le^y derivative **11** was exclusively formed in 83% yield when more than twice the amount of **9** (2.4 eq) relative to **2** was used at a higher temperature of −40 °C (entry 2). The synchronous synthesis of **10** and **11** using **9** and **2** (entries 3 and 4) is worth mentioning. Compound **11** appeared at a higher temperatures (−40 °C or −50 °C) than that in entry 1. Furthermore, both **10** and **11** were obtained efficiently in 49% and 45% yields, respectively, using an excess amount of **9** (1.8 eq) at −40 °C (entry 4). These results indicate that the synthesis of **10** and **11** can be finely controlled by varying the reaction temperature and the feed ratio. Notably, compounds **10** and **11** can be easily separated by conventional silica gel column chromatography: the *R_f* values for **10** and **11** in an *n*-hexane–EtOAc 2:1 mixture are 0.28 and 0.44, respectively.

Scheme 2. On demand and synchronous syntheses of Le^x and Le^y derivatives via glycosylation of **2** with **9** under the reaction conditions summarized in Table 1.

Table 1. One-pot synthesis of **10** and **11** under different reaction conditions.^a

entry	path	2/eq	9/eq	NIS/eq	TfOH/eq	T/°C	time/h ^b	yield/%	
								10	11
1	a	1.0	1.2	2.5	0.2	−78	1.0	76	n.d. ^c
2	b	1.0	2.4	5.0	0.4	−40	3.0	n.d. ^c	83
3	c	1.0	1.2	2.5	0.2	−50	2.0	25	16
4	c	1.0	1.8	2.5	0.2	−40	1.0	49	45

^aReaction was carried out under the indicated conditions, in CH_2Cl_2 – Et_2O using the NIS–TfOH

activation system. ^bTime for complete consumption of **9**. ^cNot detected.

It is very intriguing that the reactivity of the two hydroxy groups in **2** is strictly fixed as 3-OH > 2'-OH: a mono-fucosylated product at 2'-OH, that is, a type 2H [Fuc α (1 $\rightarrow$ 2)Gal β (1 $\rightarrow$ 4)GlcNAc β 1 $\rightarrow$] derivative is not formed at all in this series of reactions. These results are consistent with our previous report,⁸ although the molecular design of acceptor **2** is slightly different from that of **2'**. Thus, the combination of the newly designed diol acceptor **2** and donor **9** has proved to be highly effective for not only the selective synthesis of Le^x or Le^y derivatives but also the synchronous synthesis of both derivatives in a one-pot reaction. Although there are a few reports on lactosamine diol derivatives for the synthesis of T2-LAs,¹² the order of reactivity of the two hydroxy groups in these derivatives is 2'-OH > 3-OH without exception, which is opposite to that for **2** and renders the preparation of Le^x derivatives difficult. Therefore, compound **2** is the most effective acceptor capable of providing both Le^x and Le^y derivatives when using **9** as the donor.

2.3 Synthesis of 6-O-sulfo-Le^x **3 and non-sulfated Le^x **4** bearing a terminal azido functionalized oligo-ethyleneoxide linker**

For future applications of T2-LAs, we introduced a terminal azido functionalized aglycon moiety. A highly hydrophilic oligo-(ethyleneoxide) structure is advantageous for conjugation with

all kinds of bioactive compounds such as proteins, lipids, polysaccharides, and synthetic polymers.

Furthermore, azides are the first choice in current biochemical and materials sciences as they allow conjugation with a range of substances having alkyne groups via a Huisgen cycloaddition (“click chemistry”).^{13,14} Hence, we selected a 2-(2-(2-(2-azidoethoxy)ethoxy)ethoxy)ethyl group as an efficient linker for T2-LAs, as shown in Fig. 1.

Two types of Le^x derivatives (**3** and **4**) were synthesized following the reactions outlined in Scheme 3. In order to avoid damaging the α -L-fucoside linkage and azido group, hydrogenation of **10** was first carried out with Pd(OH)₂ on activated carbon (Pd(OH)₂-C) under H₂ atmosphere. This reaction normally proceeds to completion within 10 h, and the reaction mixture must be immediately worked-up, as prolonged reaction accelerates the undesirable cleavage of the α -L-fucoside linkage due to the acidity of the reagents. After acetylation, compound **12** was obtained in 39% yield, in two steps. Considering the low yield and our observations by TLC monitoring during the hydrogenation, removal of the benzyl protection in **11** without cleaving the α -L-fucoside linkage seems difficult. The anomeric PMP group of **12** was smoothly removed by CAN oxidation to produce **13** in 61% yield. Compound **13** was converted into the activated glycosyl donor of trichloroacetimidate **14** through the reaction of trichloroacetonitrile with DBU in 83% yield. The linker moiety was introduced through the glycosylation of 2-(2-(2-(2-azidoethoxy)ethoxy)ethoxy)-1-ethanol with **14**, promoted by the addition of TMSOTf at –50 °C, affording **15** in a modest yield of 48%. The acyl groups in **15** were removed successively by treatment with MeONa and hydrazine

monohydrate, followed by acetylation in pyridine and the removal of the TBDPS group with TBAF, affording **16** in 56% yield (4 steps). Sulfation at the 6-OH group in **16** was carried out by the addition of $\text{SO}_3 \cdot \text{NMe}_3$ to produce **17** in excellent yield (91%). Finally, all of the *O*-acetyl groups in **17** were removed with MeONa, which resulted in the target 6-*O*-sulfo- Le^x **3** in 63% yield. The non-sulfated Le^x derivative **4** was obtained from **15** through the reactions in steps e and h in 15% yield (5 steps). Thus, the sulfated and non-sulfated forms of the Le^x derivatives were efficiently synthesized from a common intermediate, **10**, which could be readily prepared through the regioselective glycosylation described above.

Scheme 3. Reagents and conditions: (a) (1) Pd(OH)₂-C, H₂/MeOH, rt, 10 h, (2) Ac₂O/pyridine, rt, overnight, 39% (2 steps); (b) CAN/CH₃CN-H₂O, 0 °C, 6 h, 61%; (c) CCl₃CN, DBU/CH₂Cl₂, 0 °C, 5

h, 83%; (d) HO(CH₂CH₂O)₃C₂H₄N₃, TMSOTf, MS4A/CH₂Cl₂, −50 °C, 8 h, 48%; (e) (1)

MeONa/MeOH, rt, overnight, (2) NH₂NH₂·H₂O/EtOH, 90 °C, 7 h, (3) Ac₂O/pyridine, rt, 48 h, (4)

TBAF–AcOH/THF, rt, 72 h, 56% (4 steps); (f) SO₃·NMe₃/DMF, 55 °C, 72 h, 91%; (g)

MeONa/MeOH, rt, overnight, 63%; (h) MeONa/MeOH, rt, overnight, 15% (5 steps from **15**).

2.4 Synthesis of Le^y bearing a terminal azido functionalized oligo-ethyleneoxide linker **5**

The Le^y derivative bearing a terminal azido functionalized oligo-ethyleneoxide linker **5** was also prepared according to the reactions outlined in Scheme 4. Compound **11** was treated with Pd(OH)₂–C in THF–MeOH (1:1) mixture under H₂ atmosphere as described for the synthesis of **12**. The obtained mixture was subjected to acetylation to provide pure **18**. The anomeric PMP group in **18** was removed by CAN oxidation, followed by trichloroacetimidation to give **19**. Glycosidation of **19** with 2-(2-(2-(2-azidoethoxy)ethoxy)ethoxy)-1-ethanol, which is the same acceptor employed in the synthesis of **15**, proceeded very smoothly with the addition of a catalytic amount of TMSOTf at −50 °C and gave **20** within 30 min in a very good yield of 80%. Compound **20** was converted into **21** through a three-step reaction, i.e., removal of the acetyl and benzoyl groups by MeONa, removal of the phthaloyl group by hydrazine monohydrate and acetylation, which gave **21** in 64% yield. All the protecting groups in **21** were removed by successive treatment with TBAF in THF and MeONa in MeOH, which led to the target Le^y derivative **5** in 57% yield via two steps. Thus, the Le^y derivative can also be synthesized easily from **11**, which in turn can be obtained by the glycosylation

described above.

Scheme 4. Reagents and conditions: (a) (1) $Pd(OH)_2 \cdot C$, $H_2/THF-MeOH$, rt, 10 h, (2) Ac_2O , DMAP/pyridine, rt, 24 h, 68% (2 steps); (b) (1) CAN/CH_3CN-H_2O , rt, 2 h, (2) CCl_3CN , DBU/ CH_2Cl_2 , 0 °C, 4 h, 56% (2 steps); (c) $HO(CH_2CH_2O)_3C_2H_4N_3$, TMSOTf, MS4A/ CH_2Cl_2 , -50 °C, 0.5 h, 80%; (d) (1) $MeONa/MeOH$, rt, 2 h, (2) $NH_2NH_2 \cdot H_2O/EtOH$, 90 °C, 13 h, (3) Ac_2O /pyridine, rt, overnight, 64% (3 steps); (e) (1) TBAF/ THF , rt, 72 h, (2) $MeONa/MeOH$, rt, overnight, 57% (2 steps).

3. Conclusion

In the present study, we have demonstrated for the first time the feasibility of the on-demand synthesis of Le^x and Le^y derivatives, in addition to their synchronous synthesis in one-pot through the combined use of diol acceptor **2** and benzyl-protected thiophenyl fucoside donor **9**. The

selectivity was easily controlled by varying the reaction temperature and the ratio of **2** and **9**. The derivatives of Le^x and Le^y were further functionalized by introducing an oligo-ethyleneoxide-azide linker through glycosylation. The versatile set of orthogonal protecting groups of the obtained Le^x derivative **15** enabled the facile and regioselective synthesis of both sulfated Le^x **3** and non-sulfated Le^x **4**. Le^y derivative **5** was also easily prepared from **11**. Thus, our method is highly efficient for the synthesis of T2-LAs and will be further applied to the preparation of a variety of bioactive materials.

4. Experimental

4.1 General methods

Anhydrous solvents were purchased from Wako Pure Chemical Industries, Ltd., and stored under Ar atmosphere prior to use. Other chemicals were used without further purification unless otherwise stated. Molecular sieves (MS) AW300 and 4A were powdered and activated over 100 °C under reduced pressure with P_2O_5 as desiccant prior to use. Silica gel flash column chromatography was performed on Silica Gel 60, spherical, neutrality (Nacalai Tesque), or with a CombiFlash Rf 75 Var (Teledyne Isco) on RediSep Rf Gold Normal Phase Silica columns. The reactions were monitored by TLC (silica gel 60 F254, Merck) visualized by sprayed with a mixture of $\text{H}_3(\text{PMo}_{12}\text{O}_{40}) \cdot n \text{H}_2\text{O}$ (12.5 g) and $\text{Ce}(\text{SO}_4)_2 \cdot n \text{H}_2\text{O}$ (5 g) in 10% H_2SO_4 (500 mL) and colored by heating at 140 °C. Glycosylation reactions at −50 °C or lower were performed on UCR-150

(Techno-sigma). Optical rotations were measured with a P-1010 polarimeter (Jasco). ^1H and ^{13}C NMR were recorded on a DPX-400 spectrometer (Bruker). Assignments were based on homo- and heteronuclear correlation measurements, and DEPT measurements. High resolution mass spectrometry was carried out with JMS-HX110A spectrometer (Jeol, for FAB-MS) or Exactive spectrometer (Thermo Fisher Scientific, for ESI-MS). Melting points were determined with a MP-500P (Yanaco).

4.2. 4-Methoxyphenyl 4,6-*O*-(4-methoxybenzylidene)- β -D-galactopyranosyl-(1 $\rightarrow$ 4)-2-deoxy-2-phthalimido- β -D-glucopyranoside (7).

Compound **6**⁸ (13.5 g, 16.2 mmol) in dry MeOH (250 mL) was treated with MeONa in MeOH (ca. 28wt%, 2.4 mL) at 0 °C under dry atmosphere for 23 h. The formed precipitate was filtered through filter paper, and washed with MeOH. The filtrate was neutralized by addition of Dowex 50W-X8 (H^+ form), filtered through a cotton bed, and concentrated to dryness by vacuum pump overnight. The former precipitate and the latter residue were combined and dissolved in dry DMF (120 mL). To a solution of the mixture was added *p*-anisaldehyde dimethylacetal (3.16 mL, 18.6 mmol) under acidic conditions in the presence of catalytic amount of ($\pm$)-10-camphorsulfonic acid. After kept stirring at rt for 7 h, excess amount of Et_3N was added to neutralize the reaction system. The mixture was concentrated under diminished pressure, and coevaporated with toluene. The residue was purified by silica gel column chromatography ($\text{CHCl}_3/\text{MeOH}$, 20:1, v/v, containing 0.5% Et_3N) to provide **7**

(7.75 g, 11.1 mmol, 69 %) as a white solid.

$[\alpha]_D^{23} -20.8$ (c 0.64, CHCl_3); mp 128–130 °C ; R_f 0.30 ($\text{CHCl}_3/\text{MeOH}$, 10:1); ^1H NMR (400 MHz, CD_3OD , TMS): δ (ppm) 7.96–7.78 (m, 4H, *NPhth*), 7.47–6.72 (m, 8H, $-\text{C}_6\text{H}_4\text{-OMe}\times 2$), 5.70 (d, 1H, $J_{1,2}$ 8.5 Hz, H-1^{I}), 5.56 (s, 1H, *CH* of *p*-methoxybenzylidene), 4.55 (d, 1H, $J_{1,2}$ 7.5 Hz, H-1^{II}), 4.51 (dd, 1H, $J_{2,3}$ 11.0, $J_{3,4}$ 8.5 Hz, H-3^{I}), 4.28 (dd, 1H, $J_{1,2}$ 8.6, $J_{2,3}$ 11.0 Hz, H-2^{I}), 4.24–4.06 (m, 3H, H-4^{II} , H-6^{IIa} , H-6^{Ia}), 4.06–3.93 (m, 2H, H-6^{IIb} , H-6^{Ib}), 3.84 (t, 1H, $J_{3,4}=J_{4,5}=9.6$ Hz, H-4^{I}), 3.77 (s, 3H, *OMe*), 3.73–3.63 (m, 7H, H-5^{I} , H-2^{II} , H-3^{II} , H-5^{II} , *OMe*); ^{13}C NMR (100 MHz, CDCl_3): δ 168.23, 167.93 (C=O), 159.89, 155.21, 150.79, 134.03, 131.58, 130.38, 127.76, 123.32, 118.32, 114.32, 113.30 (aromatic), 103.99 (C1^{II}), 100.92 (*CH* of *p*-methoxybenzylidene), 97.45 (C1^{I}), 81.92 (C4^{I}), 75.53 (C4^{II}), 75.05 (C5^{I}), 72.08 (C3^{II}), 70.58 (C2^{II}), 69.70 (C3^{I}), 68.73 (C6^{II}), 66.82 (C5^{II}), 61.54 (C6^{I}), 56.07 (C2^{I}), 55.47, 55.17 (*OMe*); HRMS (FAB, positive ion mode, NBA) m/z = 718.2103 [$\text{M} + \text{Na}$] $^+$, calcd for $\text{C}_{35}\text{H}_{37}\text{NO}_{14}\text{Na}$, 718.2112.

4.3. 4-Methoxyphenyl 4,6-*O*-(4-methoxybenzylidene)- β -D-galactopyranosyl-(1 $\rightarrow$ 4)-6-*O*-*tert*-butyldipheylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranoside (8).

To a solution of **7** (7.75 g, 11.1 mmol) in anhydrous pyridine (120 mL) was added *tert*-butyldiphenylchlorosilane (5.36 mL, 20.9 mmol) at rt under dry atmosphere. After stirring for 64 h, MeOH was added to quench excess reagent, and then the mixture was concentrated under reduced pressure. The residue was coevaporated with toluene and extracted with CHCl_3 , washed

successively with satd aq NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered through a Celite bed and concentrated under diminished pressure. The residue was subjected to silica gel column chromatography eluting with CHCl₃/EtOAc (2:1, v/v, containing 0.1% Et₃N) to provide pure **8** (6.12 g, 6.55 mmol, 59 %) as an amorphous powder.

[α]_D²³ −20.6 (*c* 1.0, CHCl₃) ; *R*_f 0.33 (CHCl₃/EtOAc, 1:1), ¹H NMR (400 MHz, CDCl₃, TMS): δ 7.96–6.68 (m, 22H, *NPhth*, −OSiPh₂CMe₃, −C₆H₄OMe×2), 5.75 (d, 1H, *J*_{1,2} 8.0 Hz, H-1^I), 5.45 (s, 1H, CH of 4-methoxybenzylidene), 4.55 (dd, 1H, *J*_{2,3} 10.5, *J*_{3,4} 8.04 Hz, H-3^I), 4.50 (d, 1H, *J*_{1,2} 8.0 Hz, H-1^{II}), 4.44 (dd, 1H, *J*_{1,2} 8.5, *J*_{2,3} 11.0 Hz, H-2^I), 4.30–4.22 (m, 2H, H-6^{II}a, 3^I-OH), 4.17 (d, 1H, *J*_{3,4} 3.0 Hz, H-4^{II}), 4.14–4.08 (m, 1H, H-6^Ia), 4.06–3.97 (m, 2H, H-6^Ib, H-6^{II}b), 3.86 (t, 1H, *J*_{3,4} = *J*_{4,5} = 9.5 Hz, H-4^I), 3.79 (s, 3H, *OMe*), 3.76–3.67 (m, 5H, H-5^I, H-2^{II}, *OMe*), 3.59 (ddd, 1H, *J*_{2,3} 9.5, *J*_{3,OH} 9.5, *J*_{3,4} 3.5 Hz, H-3^{II}), 3.49 (bs, 1H, H-5^{II}), 2.44 (d, 1H, *J*_{3,OH} 9.5 Hz, 3^{II}-OH), 2.34 (d, 1H, *J*_{2,OH} 2.5 Hz, 2^{II}-OH), 1.08 (s, 9H, *Me*₃ of *tert*-Bu); ¹³C NMR (100 MHz, CDCl₃): δ 168.51, 168.04 (C=O), 160.11, 155.33, 150.98, 149.54, 136.20, 135.84, 135.65, 134.09, 133.49, 132.78, 131.70, 130.04, 129.68, 127.76, 127.66, 123.81, 118.80, 114.39, 113.49 (aromatic), 103.73 (C1^{II}), 101.17 (CH of 4-methoxybenzylidene), 97.42 (C1^I), 81.06 (C4^I), 75.32 (C3^{II}), 75.09 (C4^{II}), 72.81 (C5^I), 71.19 (C2^{II}), 69.75 (C3^I), 68.67 (C6^{II}), 66.93 (C5^{II}), 62.41 (C6^I), 56.37 (C2^I), 55.61, 55.22 (*OMe*), 26.81 (*CMe*₃ of *tert*-Bu), 19.36 (*CMe*₃ of *tert*-Bu); HRMS (FAB, positive ion mode, NBA) *m/z* = 956.3308 [*M* + Na]⁺, calcd for C₅₁H₅₅NO₁₄SiNa, 956.3290.

4.4. 4-Methoxyphenyl 3-*O*-benzoyl-4,6-*O*-(4-methoxybenzylidene)- β -D-galactopyranosyl-(1 $\rightarrow$ 4)-6-*O*-*tert*-butyldiphenylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranoside (**2**).

To a solution of compound **8** (2.40 g, 2.57 mmol) in dry THF (50 mL) was added Bu₂SnCl₂ (78.1 mg, 0.26 mmol) and 1,2,2,6,6-pentamethylpiperidine (0.92 mL, 5.14 mmol), followed by kept stirring for 10 min at rt under Ar atmosphere. Benzoyl chloride (0.36 mL, 2.57 mmol) was added dropwise to the mixture at rt under Ar atmosphere. After stirring for 48 h, MeOH was added to quench excess reagent, and the mixture was evaporated under reduced pressure. The residue was diluted with CHCl₃, and washed with satd aq NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered through a Celite bed, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (CHCl₃/EtOAc, 1:0 to 0:1, v/v, linear gradient) to afford **2** (1.95 g, 1.88 mmol, 73%) as colorless amorphous.

$[\alpha]_D^{23} +23.9$ (*c* 1.0, CHCl₃); *R*_f 0.41 (*n*-hexane/EtOAc, 1:1); ¹H NMR (400 MHz, CDCl₃, TMS): δ 8.13–6.69 (m, 27H, aromatic), 5.75 (d, 1H, *J*_{1,2} 8.0 Hz, H-1^I), 5.41 (s, 1H, *CH* of 4-methoxybenzylidene), 5.01 (dd, 1H, *J*_{2,3} 10.0, *J*_{3,4} 3.5 Hz, H-3^{II}), 4.60 (d, 1H, *J*_{1,2} 7.6 Hz, H-1^{II}), 4.54 (t, 1H, *J*_{2,3} = *J*_{3,4} = 8.6 Hz, H-3^I), 4.50–4.41 (m, 2H, H-2^I, H-4^{II}), 4.27 (m, 1H, H-6^{IIa}), 4.23 (s, 1H, 3^I-OH), 4.14–3.98 (m, 4H, H-2^{II}, H-6^{Ia}, H-6^{Ib}, H-6^{IIb}), 3.86 (t, 1H, *J*_{3,4} = *J*_{4,5} = 8.5 Hz, H-4^I) 3.82–3.70 (m, 7H, H-5^I, *OMe*×2), 3.59 (s, 1H, H-5^{II}), 2.17 (d, 1H, *J*_{2,OH} 4.0 Hz, 2^{II}-OH), 1.09 (s, 9H, *Me*₃ of *tert*-Bu); ¹³C NMR (100 MHz, CDCl₃): δ 168.55, 168.13, 166.45 (C=O), 160.03, 155.39, 151.03, 135.94, 135.69, 134.16, 133.48, 133.35, 132.85, 131.79, 130.16, 129.98, 129.83, 129.77, 129.65,

128.53, 127.87, 127.73, 127.53, 123.67, 123.41, 118.80, 114.44, 113.49 (aromatic), 104.03 (C1^{II}), 100.73 (CH of 4-methoxybenzylidene), 97.48 (C1^I), 81.94 (C4^I), 75.40 (C5^I), 74.25 (C3^{II}), 73.31 (C4^{II}), 69.87 (C3^I), 68.69 (C2^{II}), 68.56 (C6^{II}), 66.81 (C5^{II}), 62.79 (C6^I), 56.24 (C2^I), 55.66, 55.29 (OMe), 26.88 (CMe₃ of *tert*-Bu), 19.38 (CMe₃ of *tert*-Bu); HRMS (FAB, positive ion mode, NBA) $m/z = 1037.3693$ [M]⁺, calcd for C₅₈H₅₉NO₁₅Si, 1037.3654.

4.5. 4-Methoxyphenyl 3-*O*-benzoyl-4,6-*O*-(4-methoxybenzylidene)- β -D-galactopyranosyl-(1 $\rightarrow$ 4)-[2,3,4-tri-*O*-benzyl- α -L-fucopyranosyl-(1 $\rightarrow$ 3)]-6-*O*-*tert*-butyldiphenylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranoside (10).

Compound **2** (817 mg, 787 μ mol) was added to a solution of **9** (497 mg, 944 μ mol) in anhydrous CH₂Cl₂ (10 mL), and then diluted with anhydrous Et₂O (20 mL). The mixture was kept stirring at rt for 30 min under Ar atmosphere in the presence of activated powdered molecular sieves (MS) AW 300 (1.00 g). *N*-Iodosuccinimide (443 mg, 1.97 mmol) was added to the mixture, followed by cooling down to -78 °C under Ar atmosphere. Triflic acid (13.8 μ L, 141 μ mol) in anhydrous Et₂O (124 μ L) was added dropwise to the mixture. After stirring for 1 h, excess amount of Et₃N was added to terminate the reaction. After kept stirring for 15 min, the mixture was filtered through a bed of Celite, diluted with CHCl₃, washed successively with 5 wt% aq Na₂S₂O₃, satd aq NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered through a bed of Celite, and concentrated under reduced pressure. The residue was subjected to silica gel column

chromatography (*n*-hexane/EtOAc, 2:1, v/v, containing 0.1% Et₃N), providing pure **10** (867 mg, 596 μmol, 76%) as a white solid.

[α]_D²³ −10.2 (*c* 0.64, CHCl₃); mp 106–107 °C; *R*_f 0.28 (*n*-hexane/EtOAc, 2:1); ¹H NMR (400 MHz, CDCl₃, TMS): δ 8.20–6.64 (m, 42H, aromatic), 5.55 (d, 1H, *J*_{1,2} 8.5 Hz, H-1^I), 5.52 (s, 1H, CH of 4-methoxybenzylidene), 5.13–5.07 (m, 2H, H-1^{III}, H-3^{III}), 4.91–4.84 (m, 2H, H-3^I, H-5^{II}), 4.76–4.69 (m, 2H, H-2^I, H-1^{II}), 4.61 (s, 2H, -CH₂Ph), 4.53–4.25 (m, 6H, H-4^I, H-6^{Ia}, H-4^{III}, H-6^{IIIa}, -CH₂Ph), 4.19–4.10 (m, 2H, H-2^{III}, -CH₂Ph), 4.10–3.96 (m, 3H, H-6^{Ib}, H-6^{IIIb}, H-3^{II}), 3.71 (s, 3H, OMe), 3.69–3.62 (m, 2H, H-5^I, H-2^{II}), 3.59–3.48 (m, 4H, OMe, -CH₂Ph), 3.42 (s, 1H, H-5^{III}), 3.21 (s, 1H, H-4^{II}), 2.38 (d, 1H, *J*_{2,OH} 3.0 Hz, 2^{III}-OH), 1.13 (s, 9H, Me₃ of *tert*-Bu), 1.08 (d, 3H, H-6^{II}, *J*_{5,6} 6.5 Hz); ¹³C NMR (100 MHz, CDCl₃): δ 171.28, 166.35 (C=O), 159.96, 155.37, 151.18, 139.56, 139.48, 138.16, 136.11, 135.44, 134.29, 133.88, 133.53, 132.29, 130.28, 130.00, 129.96, 129.80, 128.63, 128.57, 128.39, 128.29, 128.23, 128.17, 128.05, 127.98, 127.93, 127.85, 127.78, 127.66, 127.46, 127.39, 127.25, 127.12, 127.07, 127.02, 126.78, 123.76, 118.83, 114.44, 113.39 (aromatic), 101.78 (C1^{III}), 99.72 (CH of 4-methoxybenzylidene), 98.19 (C1^{II}), 97.89 (C1^I), 79.14 (C3^{II}), 78.77 (C4^{II}), 76.00 (C2^{II}), 74.90 (CH₂Ph), 74.61 (C4^I), 74.36 (C3^{III}), 73.64 (C4^{III}), 73.42 (C5^I), 73.01 (CH₂Ph), 72.41 (C3^I), 71.47 (CH₂Ph), 69.63 (C2^{III}), 69.08 (C6^{III}), 66.63 (C5^{III}), 66.45 (C5^{II}), 61.79 (C6^I), 56.81 (C2^I), 55.70, 55.06 (OMe), 26.99 (Me₃ of *tert*-Bu), 19.74 (CMe₃ of *tert*-Bu), 16.63 (C6^{II}); HRMS (FAB, positive ion mode, NBA) *m/z* = 1476.5502 [M + Na]⁺, calcd for C₈₅H₈₇NO₁₉SiNa, 1476.5539.

4.6. 4-Methoxyphenyl 2,3,4-tri-*O*-benzyl- α -L-fucopyranosyl-(1 $\rightarrow$ 2)-3-*O*-benzoyl-4,6-*O*-benzylidene- β -D-galactopyranosyl-(1 $\rightarrow$ 4)-[2,3,4-tri-*O*-benzyl- α -L-fucopyranosyl-(1 $\rightarrow$ 3)]-6-*O*-*tert*-butyldiphenylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranoside (11).

Compound **2** (2.01 g, 1.94 mmol) was added to a solution of **9** (2.45 g, 4.66 mmol) in anhydrous CH₂Cl₂ (15 mL), and then diluted with anhydrous Et₂O (30 mL). The mixture was kept stirring at rt for 30 min under Ar atmosphere in the presence of activated powdered MS AW 300 (2.0 g). *N*-Iodosuccinimide (2.18 g, 9.70 mmol) was added to the mixture, and then it was cooled down to $-40\text{ }^{\circ}\text{C}$ under Ar atmosphere. Triflic acid (68 μL , 776 μmol) in anhydrous Et₂O was injected to the mixture. After stirring for 3 h, excess amount of Et₃N was added to terminate the reaction. The mixture was filtered through a bed of Celite, diluted with CHCl₃, washed successively with 5 wt% aq Na₂S₂O₃, satd aq NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered through a Celite bed, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography (*n*-hexane/EtOAc, 1:0 to 0:1, v/v, linear gradient), providing pure **11** (3.03 g, 1.61 mmol, 83%) as a white solid.

$[\alpha]_{\text{D}} -28.2$ (*c* 0.54, CHCl₃); mp 90–91 $^{\circ}\text{C}$; R_{f} 0.79 (*n*-hexane/EtOAc, 1:1); ¹H NMR (400 MHz, CDCl₃, TMS): δ 8.10–6.68 (57H, m, aromatic), 5.57 (1H, d, $J_{1,2}$ 3.6 Hz, H-1^{III}), 5.46 (1H, s, -CH of *p*-methoxybenzylidene), 5.42 (1H, d, $J_{1,2}$ 8.0 Hz, H-1^I), 5.18 (1H, d, $J_{1,2}$ 8.0 Hz, H-1^{II}), 5.13 (1H, dd, $J_{2,3}$ 10.0, $J_{3,4}$ 3.8 Hz, H-3^{II}), 4.98 (1H, d, J_{gem} 11.8 Hz, -CH₂Ph), 4.91 (1H, m, H-5^{IV}), 4.75–4.73 (2H, m, H-2^I, H-3^I), 4.67 (1H, d, $J_{1,2}$ 4.0 Hz, H-1^{IV}), 4.62 (1H, d, J_{gem} 11.6 Hz, -CH₂Ph), 4.57–4.49 (5H,

m, $-\text{CH}_2\text{Ph} \times 4$, H-4^I), 4.45–4.29 (7H, m, H-4^{II}, H-2^{II}, H-6^{IIb}, $-\text{CH}_2\text{Ph} \times 4$), 4.21 (1H, m, H-5^{III}), 4.16 (1H, m, H-6^{Ia}), 4.05–3.98 (4H, m, H-6^{IIa}, H-2^{III}, H-3^{IV}, $-\text{CH}_2\text{Ph}$), 3.76 (3H, s, *OMe*), 3.71–3.65 (2H, m, H-6^{Ib}, H-2^{IV}), 3.55–3.52 (4H, m, *OMe*, H-3^{III}), 3.45 (1H, d, $J_{3,4}$ 3.2 Hz, H-4^{III}), 3.43 (1H, d, J_{gem} 12.4 Hz, $-\text{CH}_2\text{Ph}$), 3.33 (1H, m, H-5^{II}), 3.20 (1H, m, H-5^I), 3.09 (1H, s, H-4^{IV}), 1.30 (1H, d, $J_{5,6}$ 7.2 Hz, H-6^{III}), 1.19 (1H, d, $J_{5,6}$ 6.4 Hz, H-6^{IV}), 1.11 (9H, s, Me_3 of *tert*-Bu); ¹³C NMR (100 MHz, CDCl_3): δ 165.61, 159.97, 155.59, 151.32, 139.74, 139.57, 138.95, 138.78, 138.45, 138.19, 136.06, 135.15, 133.093, 133.71, 132.32, 130.18, 130.09, 129.94, 129.89, 129.70, 128.88, 128.46, 128.39, 128.33, 128.30, 128.27, 128.21, 128.18, 128.11, 128.04, 127.93, 127.82, 127.71, 127.57, 127.44, 127.37, 127.34, 127.10, 127.08, 127.05, 126.97, 126.65, 119.18, 114.45, 113.43 (C=O, aromatic), 99.86 (C1^{II}), 99.83 (CH of *p*-methoxybenzylidene), 98.75 (C1^{IV}), 98.41 (C1^I), 98.36 (C1^{III}), 79.48 (C4^{III}), 79.38 (C3^{IV}), 78.73 (C4^{IV}), 77.96 (C3^{III}), 77.55 (C4^I), 76.61 (C2^{III}), 76.16 (C3^{II}), 75.99 (C5^I), 75.07, 74.73 (CH_2Ph), 73.30 (C4^{II}), 73.28 (CH_2Ph), 73.09 (CH_2Ph), 72.94 (C3^I, CH_2Ph), 72.83 (C2^{IV}), 72.31 (C2^{II}), 71.38 (CH_2Ph), 69.14 (C6^{II}), 67.51 (C5^{III}), 66.84 (C5^{IV}), 66.33 (C5^{II}), 61.47 (C6^I), 56.92 (C2^I), 55.84, 55.10 (*OMe*), 26.87 (Me_3 of *tert*-Bu), 19.74 (CMe_3 of *tert*-Bu), 16.39 (C6^{III}), 16.31 (C6^{IV}); HRMS (FAB, positive ion mode, NBA) m/z = 1892.7506 $[\text{M} + \text{Na}]^+$, calcd for $\text{C}_{112}\text{H}_{115}\text{NO}_{23}\text{SiNa}$, 1892.7527.

4.7. 4-Methoxyphenyl 2,4,6-tri-*O*-acetyl-3-*O*-benzoyl- β -D-galactopyranosyl-(1 $\rightarrow$ 4)-[2,3,4-tri-*O*-acetyl- α -L-fucopyranosyl-(1 $\rightarrow$ 3)]-6-*O*-*tert*-butyldiphenylsilyl-2-deoxy-2-phthalimido- β -D-

glucopyranoside (**12**).

Compound **10** (0.43 g, 294 μ mol) was dissolved in the mixture of dry MeOH (4.0 mL) and dry THF (4.0 mL) followed by addition of Pd(OH)₂ on activated carbon (20%, 200 mg). After stirring at rt under H₂ atmosphere for 10 h, the mixture was filtered through filter paper, followed by concentration under reduced pressure. To a solution of the residue in pyridine (8 mL) was added Ac₂O (350 μ L, 3.70 mmol) under dry atmosphere at rt overnight, and methanol was added to quench excess reagents. The mixture was concentrated and coevaporated with toluene under reduced pressure. The residue was dissolved in CHCl₃, and washed with satd aq NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered through a Celite bed, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (Rf 75 system, *n*-hexane/EtOAc, 1:0 to 0:1, v/v, linear gradient) to afford **12** (153 mg, 116 μ mol, 39%) as colorless amorphous.

$[\alpha]_D^{30} -42.7$ (*c* 0.05, CHCl₃); *R*_f 0.32 (*n*-hexane/EtOAc, 1:1); ¹H NMR (CDCl₃, 400 MHz, TMS): δ 7.95–6.69 (23H, m, aromatic), 5.60 (d, 1H, *J*_{3,4} 3.2 Hz, H-4^{III}), 5.55 (d, 1H, *J*_{1,2} 8.4 Hz, H-1^I), 5.42 (s, 1H, H-4^{III}), 5.29–5.02 (m, 5H, H-3^{II}, H-5^{II}, H-1^{III}, H-2^{III}, H-3^{III}), 4.98 (d, 1H, *J*_{1,2} 4.0 Hz, H-1^{II}), 4.87–4.82 (m, 2H, H-3^I, H-2^{II}), 4.58–4.51 (m, 2H, H-2^I, H-6^{IIIa}), 4.37–4.30 (m, 2H, H-4^I, H-6^{IIIb}), 3.91–3.84 (m, 1H, H-5^I), 3.73 (s, 3H, *OMe*), 3.53 (bd, 1H, *J*_{5,6} 10.0 Hz, H-5^{III}), 2.14–1.80 (m, 18H, *Ac*), 1.27 (d, 3H, *J*_{5,6} 6.0 Hz, H-6^{II}), 1.12 (s, 9H, *Me*₃ of *tert*-Bu); ¹³C NMR (CDCl₃, 100 MHz): δ 170.87, 170.73, 170.48, 170.42, 169.80, 169.05, 165.32 (C=O), 155.57, 150.98, 136.13, 135.89, 134.56, 133.60, 130.25, 130.04, 129.94, 128.70, 128.23, 128.15, 127.84, 123.81, 118.83, 114.54

(aromatic), 99.92 (C1^{III}), 97.74 (C1^I), 95.45 (C1^{II}), 75.56 (C5^I), 74.17 (C4^I), 72.00 (C3^{III}), 71.60 (C4^{II}), 71.48 (C3^I), 71.36 (C5^{III}), 69.35 (C2^{III}), 68.30 (C3^{II}), 67.94 (C2^{II}), 67.07 (C4^{III}), 64.37 (C5^{II}), 61.19 (C6^I), 61.10 (C6^{III}), 56.60 (C2^I), 55.74 (OMe), 27.01 (*Me*₃ of *tert*-Bu), 20.95, 20.87, 20.81, 20.67, 20.62 (Ac), 19.46 (CMe₃ of *tert*-Bu), 16.10 (C6^{II}); HRMS (ESI, positive ion mode) *m/z* = 1340.4334 [M + Na]⁺, calcd for C₆₈H₇₅NO₂₄SiNa, 1340.4360.

4.8. 2,4,6-Tri-*O*-acetyl-3-*O*-benzoyl-β-D-galactopyranosyl-(1→4)-[2,3,4-tri-*O*-acetyl-α-L-fucopyranosyl-(1→3)]-6-*O*-*tert*-butyldiphenylsilyl-2-deoxy-2-phthalimido-D-glucopyranose (13).

Compound **12** (153 mg, 116 μmol) was dissolved in a mixed solution of CH₃CN-H₂O (8.0 mL – 2.0 mL) followed by addition of cerium(IV) ammonium nitrate (CAN) (190 mg, 348 μmol). After stirring at rt for 6 h, the mixture was concentrated under reduced pressure to remove CH₃CN. The residue was dissolved in CHCl₃, washed successively with satd aq NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered through a Celite bed, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography eluting with CH₃Cl/EtOAc (1:0 to 0:1, v/v, linear gradient, R_f 75 system) to afford **13** (86 mg, 70.9 μmol, 61%) as a yellowish amorphous powder.

[α]_D²⁸ –22.2 (*c* 0.1, CHCl₃); R_f 0.56 (CH₃Cl/EtOAc, 2:1); ¹H NMR (CDCl₃, 400 MHz, TMS): δ 7.94–7.41 (19H, m, aromatic), 5.58 (d, 1H, *J*_{3,4} 3.2 Hz, H-4^{III}), 5.39 (d, 1H, *J*_{3,4} 3.2 Hz, H-4^{II}), 5.29–5.16 (m, 3H, H-1^I, H-3^{II}, H-2^{III}), 5.13–5.00 (m, 3H, H-1^{III}, H-5^{II},), 5.03 (dd, 1H, *J*_{2,3} 10.0, *J*_{3,4} 10.0 Hz, H-3^{III}), 4.95 (d, 1H, *J*_{1,2} 4.0 Hz, H-1^{II}), 4.84–4.77 (m, 2H, H-3^I, H-2^{II}), 4.54 (dd, 1H, *J*_{5,6a} 6.5,

$J_{6a,6b}$ 11.2 Hz, H-6^{IIIa}), 4.33–4.10 (m, 3H, H-4^I, H-6^{IIIb}, H-2^I), 4.04 (m, 2H, H-6^{Ia}, H-6^{Ib}), 3.84–3.79 (m, 1H, H-5^{III}), 3.47 (m, 1H, H-5^I), 2.66 (d, 1H, $J_{1,OH}$ 8.2 Hz, 1-OH), 2.13–1.85 (m, 18H, Ac), 1.25 (d, 3H, $J_{5,6}$ 6.0 Hz, H-6^{II}), 11.5 (s, 9H, CMe_3 of *tert*-Bu); ^{13}C NMR ($CDCl_3$, 100 MHz): δ 170.89, 170.77, 170.48, 170.27, 169.82, 169.12, 165.35 (C=O), 136.26, 135.61, 134.55, 133.62, 133.45, 132.46, 130.34, 130.12, 129.77, 129.15, 128.73, 128.21, 127.89, 123.77 (aromatic), 99.86 (C1^{III}), 95.30 (C1^{II}), 92.86 (C1^I), 75.79 (C5^I), 74.19 (C4^I), 72.04 (C3^{III}), 71.64 (C4^{II}), 71.26 (C5^{III}, C3^I), 69.36 (C2^{III}), 68.28 (C3^{II}), 68.05 (C2^{II}), 67.02 (C4^{III}), 64.37 (C5^{II}), 61.37 (C6^I), 61.09 (C6^{III}), 58.58 (C2^I), 27.12 (CMe_3 of *tert*-Bu), 20.99, 20.92, 20.84, 20.73, 20.71, 20.65 (Ac), 19.55 (CMe_3 of *tert*-Bu), 16.12 (C6^{II}); HRMS (ESI, positive ion mode) m/z = 1234.3911 $[M + Na]^+$, calcd for $C_{61}H_{69}NO_{23}SiNa$, 1234.3927.

4.9. 2,4,6-Tri-*O*-acetyl-3-*O*-benzoyl- β -D-galactopyranosyl-(1 $\rightarrow$ 4)-[2,3,4-tri-*O*-acetyl- α -L-fucopyranosyl-(1 $\rightarrow$ 3)]-6-*O*-*tert*-butyldiphenylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl trichloroacetimidate (14).

To a solution of compound **13** (32 mg, 26.4 μ mol) in dry CH_2Cl_2 (5.0 mL) was added trichloroacetonitrile (53 μ L, 527 μ mol). After stirring at 0 °C under Ar atmosphere for 30 min, DBU (1 μ L, 7.9 μ mol) was added, then the reaction mixture was kept stirring at 0 °C under Ar atmosphere for 5 h. The mixture was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (*n*-hexane-ethyl acetate containing 0.5% Et_3N , 1:0 to 0:1, v/v, linear gradient,

Rf 75 system) to afford **14** (30 mg, 22.1 μ mol, 83%) as colorless amorphous.

$[\alpha]_D^{27} -42.7$ (C 0.1, CHCl_3); R_f 0.41 (n -hexane/EtOAc 1:1 containing with Et_3N); ^1H NMR (CDCl_3 , 400 MHz, TMS): δ 8.55 (s, 1H, $-\text{CNHCCl}_3$), 7.96–7.30 (m, 19H, aromatic), 6.36 (d, 1H, $J_{1,2}$ 8.8 Hz, H-1^I), 5.60 (d, 1H, $J_{3,4}$ 3.2 Hz, H-4^{III}), 5.42 (d, 1H, $J_{3,4}$ 2.8 Hz, H-4^{II}), 5.31 (dd, 1H, $J_{1,2}$ 10, $J_{2,3}$ 10 Hz, H-2^{III}), 5.24–5.19 (m, 2H, H-1^{III}, H-3^{II}), 5.16–5.12 (m, 2H, H-3^{III}, H-5^{II}), 5.01 (d, 1H, $J_{1,2}$ 4.0 Hz, H-1^{II}), 4.91 (t, 1H, $J_{2,3}=J_{3,4}=8.7$ Hz, H-3^I), 4.82 (dd, 1H, $J_{1,2}$ 3.9, $J_{2,3}$ 10.9 Hz, H-2^{II}), 4.65–4.52 (m, 2H, H-2^I, H-6^{IIIa}), 4.40–4.31 (m, 2H, H-4^I, H-6^{IIIb}), 4.08 (bs, 2H, H-6^{Ia}, H-6^{Ib}), 3.86 (m, 1H, H-5^{III}), 3.66–3.60 (m, 1H, H-5^I), 2.13, 2.12, 2.09, 2.08, 1.93, 1.84 (s $\times$ 6, 18H, Ac), 1.28 (d, 3H $J_{5,6}$ 6.4 Hz, H-6^{II}), 1.15 (s, 9H, Me_3 of *tert*-Bu); ^{13}C NMR (CDCl_3 , 100 MHz): δ 170.86, 170.82, 170.42, 170.21, 169.77, 169.03, 165.30, 160.73 (C=O, $-\text{CNHCCl}_3$), 136.14, 135.38, 134.59, 133.60, 133.47, 131.95, 130.24, 130.02, 129.75, 128.69, 128.20, 127.80, 123.71 (aromatic), 99.95 (C1^{III}), 95.39 (C1^{II}), 93.62 (C1^I), 90.47 ($-\text{CCl}_3$), 76.06 (C5^I), 73.95 (C4^I), 71.98 (C3^{III}), 71.56 (C4^{II}), 71.43 (C5^{III}), 71.20 (C3^I), 69.31 (C2^{III}), 68.22 (C3^{II}), 68.09 (C2^{II}), 67.10 (C4^{III}), 64.38 (C5^{II}), 61.18 (C6^{III}), 60.09 (C6^I), 55.58 (C2^I), 26.98 ($\text{C}(\text{CH}_3)_3$ of *tert*-Bu), 20.95, 20.89, 20.81, 20.65, 20.61 (Ac), 19.55 ($\text{C}(\text{CH}_3)_3$ of *tert*-Bu), 16.09 (C6^{II}); HRMS (ESI, positive ion mode) $m/z = 1377.2974$ [$\text{M} + \text{Na}$]⁺, calcd for $\text{C}_{63}\text{H}_{69}\text{N}_2\text{O}_{23}\text{SiCl}_3\text{Na}$, 1377.3024.

4.10. **2-(2-(2-(2-Azidoethoxy)ethoxy)ethoxy)ethyl** **2,4,6-tri-*O*-acetyl-3-*O*-benzoyl- β -D-galactopyranosyl-(1 $\rightarrow$ 4)-[2,3,4-tri-*O*-acetyl- α -L-fucopyranosyl-(1 $\rightarrow$ 3)]-6-*O*-tert-**

butyldiphenylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranoside (**15**).

To a solution of compound **14** (40 mg, 29.8 μ mol) in dry CH_2Cl_2 (5.0 mL) was added 2-(2-(2-(2-azidoethoxy)ethoxy)ethoxy)ethanol (19 mg, 88.5 μ mol) and activated 4Å molecular sieves (MS4A, 80 mg). After stirring at -50°C under Ar atmosphere for 30 min, TMSOTf (2 μ L, 8.85 μ mol) was added to the mixture, followed by stirring at -50°C under Ar atmosphere for 8 h. After the reaction was completed, the reaction mixture was neutralized by addition of Et_3N , filtered through a Celite bed, and the filtrate was concentrated under reduced pressure. The residue was dissolved in CHCl_3 , and washed with satd aq NaHCO_3 and brine. The organic layer was dried over MgSO_4 , filtered through a Celite bed, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (*n*-hexane/EtOAc 1:0 to 0:1, v/v, R_f 75 system, linear gradient) to afford **15** (20 mg, 14.2 μ mol, 48%) as colorless amorphous.

$[\alpha]_{\text{D}}^{29} -44.4$ (*c* 0.1, CHCl_3); R_f 0.31 (*n*-hexane/EtOAc 2:3); ^1H NMR (CDCl_3 , 400 MHz, TMS): δ 7.95–7.36 (m, 19H, aromatic), 5.58 (d, 1H, $J_{3,4}$ 3.2 Hz, H-4^{III}), 5.40 (d, 1H, $J_{3,4}$ 2.4 Hz, H-4^{II}), 5.28 (dd, 1H, $J_{1,2}$ 8.4, $J_{2,3}$ 10.0 Hz, H-2^{III}), 5.22–5.17 (m, 2H, H-1^{III}, H-3^{II}), 5.13–5.07 (m, 3H, H-1^I, H-3^{III}, H-5^{II}), 4.95 (d, 1H, $J_{1,2}$ 4.0 Hz, H-1^{II}), 4.82 (dd, 1H, $J_{1,2}$ 4.0, $J_{2,3}$ 10.9 Hz, H-2^{II}), 4.77 (t, 1H, $J_{2,3}$ = $J_{3,4}$ = 9.9 Hz, H-3^I), 4.55 (dd, 1H, $J_{5,6a}$ 6.7, $J_{6a,6b}$ 11.6 Hz, H-6^{IIIa}), 4.35–4.22 (m, 3H, H-2^I, H-4^I, H-6^{IIIb}), 4.07–3.98 (m, 2H, H-6^{Ia}, H-6^{Ib}), 3.89–3.83 (m, 2H, H-5^{III}, PEG), 3.65–3.25 (m, 16H, H-5^I, PEG), 2.12, 2.10, 2.06, 1.91, 1.82 (s $\times$ 6, 18H, Ac), 1.25 (d, 3H, $J_{5,6}$ 6.4 Hz, H-6^{II}), 1.15 (s, 9H, *Me*₃ of *tert*-Bu); ^{13}C NMR (CDCl_3 , 100 MHz): δ 170.91, 170.89, 170.51, 170.29, 169.83, 169.12, 165.36

(C=O), 136.23, 135.50, 134.48, 133.62, 133.46, 132.25, 130.29, 130.07, 129.79, 129.16, 128.73, 128.24, 127.85, 123.68 (aromatic), 99.93 (C1^{III}), 98.18 (C1^I), 95.36 (C1^{II}), 75.44 (C5^I), 74.38 (C4^I), 72.06 (C3^{III}), 71.68 (C4^{II}), 71.54 (C3^I), 71.31 (C5^{III}), 70.72, 70.64, 70.62, 70.54, 70.20, 70.16 (PEG), 69.45 (C2^{III}), 68.53 (PEG), 68.32 (C3^{II}), 68.04 (C2^{II}), 67.11 (C4^{III}), 64.33 (C5^{II}), 61.18 (C6^I, C6^{III}), 56.71 (C2^I), 50.79 (PEG), 27.07 (CMe₃ of *tert*-Bu), 20.98, 20.95, 20.85, 20.71, 20.67 (Ac), 19.54 (CMe₃ of *tert*-Bu), 16.14 (C6^{II}); HRMS (ESI, positive ion mode) m/z = 1435.5026 [M + Na]⁺, calcd for C₆₉H₈₄N₄O₂₆SiNa, 1435.5041

4.11. 2-(2-(2-(2-azidoethoxy)ethoxy)ethoxy)ethyl 2,3,4,6-tetra-*O*-acetyl- β -D-galactopyranosyl-(1 $\rightarrow$ 4)-[2,3,4-tri-*O*-acetyl- α -L-fucopyranosyl-(1 $\rightarrow$ 3)]-2-acetamido-2-deoxy- β -D-glucopyranoside (16).

Compound **15** (51 mg, 36.8 μ mol) was suspended in dry MeOH (5.0 mL) followed by the addition of *ca.* 28 wt-% MeONa in MeOH (3 μ L, 19.7 μ mol). After stirring at rt under dry atmosphere overnight, the reaction mixture was neutralized by the addition of Dowex 50W-X4 (H⁺ form), filtered through cotton, and concentrated under reduced pressure. The residue dissolved in EtOH (5.0 mL) was added NH₂NH₂·H₂O (10 μ L, 205.8 μ mol). After stirring at 90 °C for 7 h, reaction mixture was concentrated to dryness under reduced pressure. The residue was dissolved in pyridine (5.0 mL) followed by addition of Ac₂O (65 μ L, 687.6 μ mol) and DMAP (3 mg, 24.6 μ mol). After stirring at rt under dry atmosphere for 36 h, MeOH was added to quench excess reagents, followed by

concentration under reduced pressure. The residue was dissolved in CHCl_3 , and washed with satd aq NaHCO_3 and brine. The organic layer was dried over MgSO_4 , filtered through a Celite bed, and concentrated under reduced pressure. The residue was roughly purified by silica gel column chromatography eluting with $\text{CH}_3\text{Cl}/\text{MeOH}$ (1:0 to 6:1, v/v, Rf 75 system, linear gradient). The obtained compound was dissolved in dry THF (3.0 mL) followed by the addition of AcOH (5 μL , 91.8 μmol) and 1M TBAF in THF (207 μL , 207 μmol). After stirring at rt under Ar atmosphere for 72 h, the reaction mixture was concentrated under reduced pressure and extracted with CHCl_3 , washed with satd aq NaHCO_3 and brine. The organic layer was dried over MgSO_4 , filtered through a Celite bed, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography eluting with $\text{CHCl}_3/\text{MeOH}$ (1:0 to 6:1, v/v, linear gradient, Rf 75 system) to afford the compound **16** (21 mg, 20.5 μmol , 56%, 4 steps) as colorless amorphous.

$[\alpha]_{\text{D}}^{26} -71.4$ (c 0.2, CHCl_3); R_f 0.24 ($\text{CHCl}_3/\text{MeOH}$, 10:1); ^1H NMR (CDCl_3 , 400 MHz, TMS): δ 6.31 (d, 1H, $J_{2,\text{NHAc}}$ 9.4 Hz, NHAc), 5.42–5.38 (m, 3H, H-1^{II}, H-4^{II}, H-4^{III}), 5.22 (dd, 1H $J_{2,3}$ 11.0, $J_{3,4}$ 3.4 Hz, H-3^{II}), 5.10–4.98 (m, 4H, H-2^{II}, H-5^{II}, H-2^{III}, H-3^{III}), 4.73 (d, 1H, $J_{1,2}$ 7.28 Hz, H-1^{III}), 4.67 (d, 1H, $J_{1,2}$ 8.40 Hz, H-1^I), 4.50 (dd, 1H, $J_{5,6a}$ 6.2, $J_{6a,6b}$ 11.4 Hz, H-6a^{III}), 4.32 (dd, 1H, $J_{5,6b}$ 7.9, $J_{6a,6b}$ 11.4 Hz, H-6b^{III}), 4.03–3.90 (m, 4H, H-2^I, H-4^I, H-6^{Ia}, H-5^{III}), 3.85–3.54 (m, 16H, H-3^I, H-6^{Ib}, PEG), 3.45–3.39 (m, 2H, PEG), 3.28–3.24 (m, 1H, H-5^I), 2.30 (dd, 1H, $J_{6a,\text{OH}}$ 3.96, $J_{6Ib,\text{OH}}$ 9.6 Hz, ¹⁶-OH), 2.19, 2.14, 2.13, 2.08, 2.05, 1.98, 1.97, 1.95 (s $\times$ 8, 24H, Ac), 1.19 (d, 3H $J_{5,6}$ 6.5 Hz, H-6^{II}); ^{13}C NMR (CDCl_3 , 100 MHz): δ 171.70, 171.15, 170.87, 170.67, 170.64, 170.14, 169.79, 169.26 (C=O),

102.11 (C1^I), 100.40 (C1^{III}), 95.58 (C1^{II}), 75.45 (C5^I), 74.55 (C4^I), 73.76 (C5^{III}), 71.73 (C4^{III}, PEG), 71.09, 71.05 (C3^I, C3^{III}), 70.99, 70.69, 70.67, 70.52, 70.10 (PEG), 69.47 (C2^{III}), 68.59 (PEG), 68.11 (C2^{II}, C3^{II}), 67.05 (C4^{II}), 64.14 (C5^{II}), 60.87 (C6^I), 60.59 (C6^{III}), 55.97 (C2^I), 50.76 (PEG), 23.35, 21.26, 20.96, 20.86, 20.83, 20.81, 20.76, 20.70 (Ac), 15.97 (C6^{II}); HRMS (ESI, positive ion mode) $m/z = 1047.3740$ [M + Na]⁺, calcd for C₄₂H₆₄N₄O₂₅Na, 1047.3757.

4.12. Triethylammonium {2-(2-(2-(2-azidoethoxy)ethoxy)ethoxy)ethyl 2,3,4,6-tetra-*O*-acetyl-β-D-galactopyranosyl-(1→4)-[2,3,4-tri-*O*-acetyl-α-L-fucopyranosyl-(1→3)]-2-acetamido-2-deoxy-6-*O*-sulfonato-β-D-glucopyranoside} (17).

Compound **16** (28 mg, 27.3 μmol) was dissolved in dry DMF (3.0 mL) and Et₃N (600 μL). After stirring at 55 °C under Ar atmosphere for 20 min, the reaction mixture was added to SO₃·NMe₃ (145 mg, 934 μmol). After stirring at 55 °C for 72 h, MeOH (1 mL) was added to quench excess reagents, and evaporated under reduced pressure. The residue was purified by LH-20 size exclusion column chromatography eluting with MeOH to afford **17** (30 mg, 24.9 μmol, 91%) as colorless amorphous.

[α]_D²³ −83.8 (*c* 0.07, MeOH); *R*_f 0.17 (CHCl₃/MeOH 10:1); ¹H NMR (400 MHz, CDCl₃, TMS): δ 9.46 (1H, bs, SO₃H), 6.43 (d, 1H, *J*_{2,NHAc} 9.6 Hz, NHAc), 5.45 (bs, 1H, H-4^{III}), 5.39 (d, 1H, *J*_{1,2} 3.8 Hz, H-1^{II}), 5.36 (bd, 1H, *J*_{3,4} 3.0 Hz, H-4^{II}), 5.26 (dd, 1H, *J*_{2,3} 10.9, *J*_{3,4} 3.3 Hz, H-3^{II}), 5.10–4.97 (m, 5H, H-2^{II}, H-3^{III}, H-5^{II}, H-1^{III}, H-2^{III}), 4.60 (d, 1H, *J*_{1,2} 8.40 Hz, H-1^I), 4.44 (dd, 1H, *J*_{5,6a} 5.8, *J*_{6a,6b} 11.2 Hz, H-6^{IIIa}), 4.32 (bs, 2H, H-6^{Ia}, H-6^{Ib}), 4.26 (dd, 1H, *J*_{5,6b} 8.7, *J*_{6a,6b} 11.1 Hz, H-6^{IIIb}), 4.05–

3.93 (m, 3H, H-2^I, H-5^{III}, PEG), 3.80–3.54 (m, 14H, PEG), 3.47–3.42 (m, 4H, H-3^I, H-4^I, H-5^I, PEG), 3.20 (6H, m, N(CH₂CH₃)₃), 2.17, 2.14, 2.12, 2.08, 2.06, 1.96, 1.95, 1.94 (s×8, 24H, Ac), 1.40 (t, 9H, J 7.3 Hz, N(CH₂CH₃)₃), 1.20 (d, 3H, *J*_{5,6} 6.5 Hz, H-6^{II}); ¹³C NMR (CDCl₃, 100 MHz): δ 171.60, 171.26, 170.83, 170.61, 169.83, 169.80, 169.79, 169.76 (C=O), 102.04 (C1^I), 99.66 (C1^{III}), 95.53 (C1^{II}), 74.63 (PEG), 73.79 (C3^I, C5^I), 73.43 (C5^{III}), 71.83 (C4^{II}), 71.65 (PEG), 71.29 (C3^{III}), 70.88 (PEG), 70.72 (C4^I), 70.65, 70.61, 70.46, 70.06 (PEG), 69.44 (C2^{III}), 68.47 (PEG), 68.20 (C2^{II}), 67.94 (C3^{II}), 67.25 (C4^{III}), 64.91 (C6^I), 64.30 (C5^{II}), 60.64 (C6^{III}), 55.46 (C2^I), 50.78 (PEG), 46.71 (N(CH₂CH₃)₃), 23.30, 21.22, 21.02×2, 20.88, 20.83, 20.77, 20.69 (Ac), 15.97 (C6^{II}), 8.83 (N(CH₂CH₃)₃): HRMS (ESI, negative ion mode) *m/z* = 1103.3360 [M – HNEt₃][–], calcd for C₄₈H₆₃N₄O₂₈S, 1103.3350.

4.13. 4-Methoxyphenyl 2,3,4-tri-*O*-acetyl- α -L-fucopyranosyl-(1→2)-4,6-di-*O*-acetyl-3-*O*-benzoyl- β -D-galactopyranosyl-(1→4)-[2,3,4-tri-*O*-acetyl- α -L-fucopyranosyl-(1→3)]-6-*O*-tert-butylidiphenylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranoside (18).

To a solution of compound **11** (1.00 g, 0.53 mmol) in 20.0 mL of THF–MeOH (1:1, v/v) was added Pd(OH)₂-C (20%, 0.25 g, 1.78 mmol). After stirring at rt under H₂ for 10 h, the reaction mixture was filtered through a Celite bed, and the filtrate was concentrated under reduced pressure. The residue was dissolved in dry pyridine (10.0 mL), followed by addition of DMAP (64 mg 0.53 mmol) and Ac₂O (1.0 mL, 10.6 mmol). After stirring at rt under dry atmosphere for 24 h, MeOH was added to quench

excess reagents, and then concentrated and coevaporated with toluene under reduced pressure. The residue was dissolved in CHCl_3 , and washed with satd aq NaHCO_3 and brine. The organic layer was dried over MgSO_4 , filtered through a Celite bed, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography eluting with *n*-hexane/EtOAc (1:2, v/v, containing 0.5% Et_3N) to afford **18** (562 mg, 0.36 mmol, 68%) as colorless amorphous.

$[\alpha]_{\text{D}}^{27} -110.1$ (*c* 0.03, CHCl_3); R_f 0.63 (*n*-hexane/EtOAc, 1:2); ^1H NMR (400 MHz, CDCl_3 , TMS): δ 7.89–7.32, 7.23–6.74 (23H, m, aromatic), 5.51 (bd, 1H, $J_{3,4}$ 3.4 Hz, H-4^{II}), 5.45 (d, 1H, $J_{1,2}$ 8.5 Hz, H-1^I), 5.40 (bd, 1H, $J_{3,4}$ 2.7 Hz, H-4^{III}), 5.26–5.09 (7H, m, H-1^{IV}, H-4^{IV}, H-3^{III}, H-3^{II}, H-1^{II}, H-2^{IV}, H-5^{II}), 4.97–4.81 (m, 4H, H-1^{III}, H-3^{IV}, H-2^{III}, H-3^I), 4.59 (dd, 1H, $J_{1,2}$ 8.6, $J_{2,3}$ 10.0 Hz, H-2^I), 4.53–4.40 (m, 3H, H-4^I, H-6^{IIa}, H-5^{IV}), 4.30 (dd, 1H, $J_{5,6}$ 7.4, $J_{6a,6b}$ 11.4 Hz, H-6^{IIb}), 4.23–4.15 (m, 2H, H-6^{Ia}, H-6^{Ib}), 3.92 (bt, 1H, $J_{1,2}=J_{2,3}=9.0$ Hz, H-2^{II}), 3.85–3.80 (m, 1H, H-5^{II}), 3.75 (s, 3H, OMe), 3.47–3.43 (m, 1H, H-5^I), 2.15, 2.11, 2.09, 2.08, 2.07, 1.92, 1.86, 1.72 (s×8, 24H, Ac), 1.27–1.24 (m, 6H, H-6^{III}, H-6^{IV}), 1.16 (s, 9H, Me_3 of *tert*-Bu); ^{13}C NMR (CDCl_3 , 100 MHz): δ 171.00, 170.88, 170.67, 170.40, 170.15, 169.88, 169.85, 169.61, 169.15 (C=O), 155.69, 1150.94, 135.99, 135.41, 134.55, 133.90, 133.46, 132.04, 130.04, 129.99, 129.93, 129.96, 128.90, 128.27, 128.15, 127.89, 127.86 (aromatic), 99.86 (C1^{II}), 98.16 (C1^I), 97.72 (C1^{IV}), 95.92 (C1^{III}), 75.42 (C5^I), 74.21 (C2^{II}), 74.07 (C3^{II}), 73.04 (C4^I), 72.47 (C3^I), 71.70 (C4^{III}), 71.20 (C5^{II}), 71.06 (C4^{IV}), 68.23 (C3^{III}), 67.98 (C3^{IV}), 67.67 (C2^{III}), 67.27 (C4^{II}), 66.88 (C2^{IV}), 65.41 (C5^{IV}), 64.41 (C5^{III}), 61.14, 61.08 (C6^I, C6^{II}), 56.83 (C2^I), 55.80 (OMe), 27.16 (Me_3 of *tert*-Bu), 21.06, 20.97, 20.82, 20.73, 20.69×2, 20.67, 20.34 (Ac),

19.43 (CMe₃ of *tert*-Bu), 15.59, 15.52 (C6^{III}, C6^{IV}); HRMS (ESI, positive ion mode) m/z =

1570.5151 [M + Na]⁺, calcd for C₇₈H₈₉NO₃₀SiNa, 1570.5136.

4.14. 2,3,4-Tri-*O*-acetyl- α -L-fucopyranosyl-(1 $\rightarrow$ 2)-4,6-di-*O*-acetyl-3-*O*-benzoyl- β -D-galactopyranosyl-(1 $\rightarrow$ 4)-[2,3,4-tri-*O*-acetyl- α -L-fucopyranosyl-(1 $\rightarrow$ 3)]-6-*O*-*tert*-butyldiphenylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl trichloroacetimidate (19).

Compound **18** (562 mg, 0.36 mmol) was dissolved in a mixed solution of CH₃CN (8.0 mL)–H₂O (2.0 mL) followed by addition of CAN (592 mg, 1.08 mmol). After stirring at rt for 2 h, the reaction mixture was extracted with CHCl₃, washed successively with satd aq NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered through a Celite bed, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (CHCl₃/MeOH, 30:1, containing 0.5% Et₃N) to afford the corresponding anomer-free compound (440 mg). To a solution of this compound (440 mg) in dry CH₂Cl₂ (10 mL) was added CCl₃CN (290 μ L, 2.90 mmol). After stirring at 0 °C under Ar for 15 min, DBU (13 μ L, 87 μ mol) was added to the mixture. After kept stirring for 4 h, the mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (CHCl₃/EtOAc, 3:1, containing 0.5% Et₃N) to afford **19** (314 mg, 0.20 mmol, 2 steps, 56%) as colorless amorphous.

$[\alpha]_D^{27}$ –77.4 (c 0.5, CHCl₃); R_f 0.57 (CHCl₃/EtOAc 2:1); ¹H NMR (400 MHz, CDCl₃, TMS): δ 8.58 (s, 1H, *NH*), 7.90–7.32 (19H, m, aromatic), 6.32 (d, 1H, $J_{1,2}$ 8.8 Hz, H-1^I), 5.51 (bd, 1H, $J_{3,4}$ 3.6 Hz,

H-4^{II}), 5.39 (bd, 1H, $J_{3,4}$ 2.8 Hz, H-4^{III}), 5.26–5.24 (m, 2H, H-1^{IV}, H-4^{IV}), 5.23–5.16 (m, 2H, H-3^{II}, H-3^{III}), 5.14–5.07 (m, 3H, H-5^{III}, H-1^{II}, H-2^{IV}), 5.02–4.95 (m, 2H, H-3^{IV}, H-1^{III}), 4.94–4.84 (m, 2H, H-3^I, H-2^{III}), 4.63 (dd, 1H, $J_{1,2}$ 8.9, $J_{2,3}$ 10.2 Hz, H-2^I), 4.53–4.44 (m, 2H, H-4^I, H-6^{IIa}), 4.42–4.36 (m, 1H, H-5^{IV}), 4.33–4.18 (m, 3H, H-6^{Ia}, H-6^{Ib}, H-6^{IIb}), 3.92 (dd, 1H, $J_{1,2}$ 8.1, $J_{2,3}$ 9.9 Hz, H-2^{II}), 3.80–3.75 (m, 1H, H-5^{II}), 3.62–3.58 (m, 1H, H-5^I), 2.12, 2.11, 2.09, 2.08, 2.07, 1.93, 1.86, 1.71 (s×8, 24H, Ac), 1.28–1.24 (m, 6H, H-6^{III}, H-6^{IV}), 1.16 (s, 9H, CMe_3 of *tert*-Bu); ¹³C NMR (CDCl₃, 100 MHz): δ 170.97, 170.86, 170.53, 170.38, 170.13, 169.86, 169.61, 168.05, 165.23, 160.74 (C=O), 136.03, 135.40, 134.65, 133.87, 133.57, 132.19, 131.45, 130.09, 130.01, 129.57, 128.95, 128.72, 128.20, 127.88, 123.72 (aromatic, C=NH), 100.01 (C1^{II}), 97.80 (C1^{IV}), 95.89 (C1^{III}), 94.08 (C1^I), 90.48 (CCl₃), 76.03 (C5^I), 74.39 (C2^{II}), 73.97 (C3^{II}), 72.99 (C4^I), 72.12 (C3^I), 71.65 (C4^{III}), 71.25 (C5^{II}), 71.15 (C4^{IV}), 68.14 (C3^{III}), 67.86 (C3^{IV}), 67.83 (C2^{III}), 67.33 (C4^{II}), 67.02 (C2^{IV}), 65.57 (C5^{IV}), 64.42 (C5^{III}), 61.14, 61.08 (C6^I, C6^{III}), 55.73 (C2^I), 27.14 (CMe_3 of *tert*-Bu), 21.00, 20.98, 20.82, 20.70, 20.68, 20.65, 20.64, 20.34 (Ac), 19.66 (CMe_3 of *tert*-Bu), 16.14, 15.73 (C6^{III}, C6^{IV}); HRMS (ESI) m/z = 1340.4334 [M + Na]⁺, calcd for C₆₈H₇₅NO₂₄SiNa, 1340.4360.

4.15. 2-(2-(2-(2-azidoethoxy)ethoxy)ethoxy)ethyl 2,3,4-tri-*O*-acetyl- α -L-fucopyranosyl-(1→2)-4,6-di-*O*-acetyl-3-*O*-benzoyl- β -D-galactopyranosyl-(1→4)-[2,3,4-tri-*O*-acetyl- α -L-fucopyranosyl-(1→3)]-6-*O*-*tert*-butyldiphenylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranoside (20).

To a solution of compound **19** (314 mg, 200 μ mol) and 2-(2-(2-azidoethoxy)ethoxy)ethoxy)ethanol (65 mg, 0.30 mmol) in dry CH_2Cl_2 (5.0 mL) was added activated MS4A. The mixture was kept stirring at -50°C for 15 min under Ar atmosphere, followed by addition of TMSOTf (11 μ L, 59 μ mol). After stirring at -50°C for 30 min, Et_3N was added to terminate the reaction. The mixture was filtered through a Celite bed, and the filtrate was washed successively with satd aq NaHCO_3 and brine. The organic layer was dried over MgSO_4 , filtered through a Celite bed, and concentrated under reduced pressure. The residue was purified by silica gel chromatography (*n*-hexane/ EtOAc 1:2, v/v, containing 0.5% Et_3N) to afford **20** (262 mg, 159 μ mol, 80%) as colorless amorphous.

$[\alpha]_{\text{D}}^{25} -73.6$ (*c* 0.1, CHCl_3); R_f 0.39 (*n*-hexane/ EtOAc 2:3); ^1H NMR (400 MHz, CDCl_3 , TMS): δ 7.90–7.36 (19H, m, aromatic), 5.50 (bd, 1H, $J_{3,4}$ 3.8 Hz, H-4^{II}), 5.38 (bd, 1H, $J_{3,4}$ 3.0 Hz, H-4^{III}), 5.26–5.08 (m, 7H, H-1^{IV}, H-4^{IV}, H-3^{III}, H-3^{II}, H-1^I, H-5^{III}, H-3^{IV}), 5.02 (d, 1H, $J_{1,2}$ 8.5 Hz, H-1^I), 4.98–4.93 (m, 2H, H-2^{IV}, H-1^{III}), 4.87 (dd, 1H, $J_{1,2}$ 4.0, $J_{2,3}$ 11.0 Hz, H-2^{III}), 4.78 (t, 1H, $J_{2,3}=J_{3,4}=9.7$ Hz, H-3^I), 4.50–4.38 (m, 3H, H-5^{IV}, H-6^{IIa}, H-4^I), 4.36–4.26 (m, 2H, H-2^I, H-6^{IIb}), 4.20 (bs, 2H, H-6^{Ia}, H-6^{Ib}), 3.95–3.88 (m, 2H, H-2^{II}, PEG), 3.82–3.78 (m, 1H, H-5^{II}), 3.70–3.23 (m, 16H, H-5^I, PEG), 2.11, 2.10, 2.08 $\times$ 2, 2.07, 1.92, 1.87, 1.73 (s $\times$ 8, 24H, Ac), 1.27–1.24 (m, 6H, C6^{III}, C6^{IV}), 1.16 (s, 9H, Me_3 of *tert*-Bu); ^{13}C NMR (CDCl_3 , 100 MHz): δ 170.96 $\times$ 2, 170.88, 170.67, 170.38, 170.13, 169.86 $\times$ 2, 169.63, 165.22 $\times$ 2 (C=O), 136.06, 135.47, 134.46, 133.88, 133.52, 132.40, 130.06, 130.01, 129.56, 128.94, 128.73, 128.22, 127.87, 123.61 (aromatic), 99.88 (C1^{II}), 98.50 (C1^I), 97.63 (C1^{IV}), 95.81 (C1^{III}), 75.30

(C5^I), 74.07 (C2^{II}, C3^{II}), 73.28 (C4^I), 72.51 (C3^I), 71.72 (C4^{III}), 71.14 (C5^{II}), 71.07(C4^{IV}), 770.87, 70.83, 70.76, 70.72, 70.66, 70.61, 70.55, 70.15, 70.10, 68.88 (PEG), 68.19 (C3^{III}), 68.03 (C2^{IV}), 67.75 (C2^{III}), 67.30 (C4^{II}), 66.93 (C3^{IV}), 65.41 (C5^{IV}), 64.34 (C5^{III}), 61.27 (C6^I), 61.04 (C6^{II}), 56.87 (C2^I), 50.80 (PEG), 27.15 (CMe₃ of *tert*-Bu), 21.01, 20.96, 20.83, 20.70×2, 20.68×2, 20.37 (Ac), 19.61 (CMe₃ of *tert*-Bu), 16.14, 15.65 (C6^{III},C6^{IV}); HRMS (ESI) m/z = 1665.5807 [M + Na]⁺, calcd for C₇₉H₉₈N₄O₃₂SiNa, 1665.5831.

4.16. 2-(2-(2-(2-azidoethoxy)ethoxy)ethoxy)ethyl 2,3,4-tri-*O*-acetyl- α -L-fucopyranosyl-(1→2)-3,4,6-tri-*O*-acetyl- β -D-galactopyranosyl-(1→4)-[2,3,4-tri-*O*-acetyl- α -L-fucopyranosyl-(1→3)]-6-*O*-*tert*-butyldiphenylsilyl-2-acetamido-2-deoxy- β -D-glucopyranoside (21).

To a solution of compound **20** (262 mg, 159 μ mol) in MeOH (5.0 mL) was added MeONa in MeOH solution (*ca.* 28 wt%; 32 μ L, 159 μ mol). After stirring at rt for 2 h, DOWEX 50W-X8 (H⁺ form) was added to neutralize the reaction system, and then filtered through cotton, concentrated under reduced pressure to obtain the crude mixture (203 mg). To a solution of the mixture (164 mg, 136 μ mol) in EtOH (4.0 mL) was added NH₂NH₂·H₂O (33 μ L, 681 μ mol). After stirring at 90 °C for 13 h, the reaction mixture was concentrated under reduced pressure. The residue was dissolved in pyridine (5.0 mL), followed by addition of Ac₂O (192 μ L, 2.04 mmol) and DMAP (8 mg, 68.0 μ mol). After stirring at rt under dry atmosphere overnight, MeOH (1 mL) was added to quench excess reagents, and the mixture was concentrated with toluene under reduced pressure. The

residue was dissolved in CHCl_3 , washed with satd aq NaHCO_3 and brine. The organic layer was dried over MgSO_4 , filtered through a Celite bed, and concentrated under reduced pressure. The residue was purified by silica gel chromatography (MeOH/EtOAc , 1:30, v/v, containing 1% Et_3N) to afford **21** (151 mg, 101 μmol , 64%) as colorless amorphous.

$[\alpha]_{\text{D}}^{25} -115.5$ (c 0.24, CHCl_3); R_f 0.51 ($\text{CHCl}_3/\text{MeOH}$, 10:1); ^1H NMR (400 MHz, CDCl_3 , TMS): δ 7.76–7.70, 7.42–7.32 (m, 10H, aromatic), 6.11 (d, 1H, $J_{2,\text{NH}}$ 8.6 Hz, NH), 5.39–5.37 (m, 2H, H-1^{IV}, H-4^{III}), 5.33–5.28 (m, 2H, H-4^I, H-1^{III}), 5.24 (dd, 1H, $J_{2,3}$ 11.0, $J_{3,4}$ 3.0 Hz, H-3^{III}), 5.17 (bs, 1H, H-4^{IV}), 5.07–4.95 (m, 5H, H-1^{II}, H-2^{III}, H-3^{IV}, H-2^{IV}, H-5^{III}), 4.92 (dd, 1H, $J_{2,3}$ 10.0, $J_{3,4}$ 3.5 Hz, H-3^{II}), 4.61 (d, 1H, $J_{1,2}$ 7.4 Hz, H-1^I), 4.45 (dd, 1H, $J_{5,6a}$ 6.4, $J_{6a,6b}$ 11.5 Hz, H-6^{IIa}), 4.32–4.20 (m, 3H, H-4^I, H-6^{IIb}, H-5^{IV}), 4.15–4.04 (m, 2H, H-6^{Ia}, H-6^{Ib}), 3.93–3.86 (m, 4H, H-2^I, H-3^I, PEG $\times$ 1), 3.78–3.59 (m, 14H, H-2^{II}, H-5^{II}, PEG $\times$ 6), 3.42–3.39 (m, 2H, PEG $\times$ 1), 3.16–3.12 (m, 1H, H-5^I), 2.18, 2.14, 2.13, 2.11, 2.05 $\times$ 2, 1.99, 1.96, 1.95, 1.91 (s $\times$ 10, 3H $\times$ 10, Ac $\times$ 10), 1.17–1.06 (m, 15H, H-6^{IV}, H-6^{III}, Me_3 of *tert*-Bu); ^{13}C NMR (100 MHz, CDCl_3): δ 171.64, 171.37, 170.90, 170.66, 170.60 $\times$ 2, 170.39, 170.24, 169.86, 169.80 (C=O), 136.05, 135.46, 133.61, 132.47, 129.89, 128.07, 127.81 (aromatic), 101.48 (C1^I), 100.56 (C1^{II}), 96.64 (C1^{III}), 96.09 (C1^{IV}), 75.37 (C5^I), 73.61 (C2^{II}), 73.40 (C3^{II}), 73.08 (C4^I), 71.79 (C4^{III}), 71.36, 71.07 (PEG), 70.97 (C4^{IV}), 70.94 (C5^{II}), 70.68, 70.67, 70.55, 70.07 (PEG), 68.19 (C3^{III}), 68.10 (C3^I, C2^{IV}), 68.04 (C2^I), 67.96 (C2^{III}), 67.72 (C3^{IV}), 67.33 (C4^{II}), 65.03 (C5^{IV}), 64.06 (C5^{III}), 61.36 (C6^I), 61.04 (C6^{II}), 50.78 (PEG), 27.10 (Me_3 of *tert*-Bu), 23.48, 21.33, 20.96, 20.84 $\times$ 2, 20.83 $\times$ 3, 20.76, 20.71, (Ac), 19.57 (CMe_3 of *tert*-Bu) 16.14, 15.71 (C6^{III}, C6^{IV});

HRMS (ESI, positive ion mode) $m/z = 1510.6196$ [$M + NH_4$] $^+$, calcd for $C_{68}H_{100}N_5O_{31}Si_1$,
1510.6172.

4.17. Sodium {2-(2-(2-(2-azidoethoxy)ethoxy)ethoxy)ethyl β -D-galactopyranosyl-(1 $\rightarrow$ 4)-[α -L-fucopyranosyl-(1 $\rightarrow$ 3)]-2-acetamido-2-deoxy-6-*O*-sulfonato- β -D-glucopyranoside} (3).

To a solution of the compound **17** (30 mg, 24.9 μ mol) in MeOH (3.0 mL) was added MeONa (*ca.* 28 wt-%) in MeOH (2 μ L, 12.4 μ mol). After stirring at rt overnight, the reaction mixture was concentrated under reduced pressure. The residue was purified by Biogel P-2 gel column eluting with H₂O to afford **3** (13 mg, 15.6 μ mol, 63%) as colorless amorphous.

$[\alpha]_D^{25} -37.5$ (*c* 0.2, MeOH); R_f 0.51(CHCl₃/MeOH/H₂O, 5:4:1); 1H NMR (400 MHz, CD₃OD): δ 5.04 (d, 1H, $J_{1,2}$ 3.9 Hz, H-1^{II}), 4.83–4.76 (m, 1H, H-5^{II}), 4.59 (d, 1H, $J_{1,2}$ 7.6 Hz, H-1^{III}), 4.55 (d, 1H, $J_{1,2}$ 7.7 Hz, H-1^I), 4.44 (dd, 1H, $J_{5,6a}$ 3.4, $J_{6a,6b}$ 10.8 Hz, H-6^{IIIa}), 4.31 (dd, 1H, $J_{5,6b}$ 2.5, $J_{6a,6b}$ 10.9 Hz, H-6^{IIIb}), 3.98–3.84 (m, 5H, H-2^I, H-3^{II}, H-4^I, PEG $\times$ 1), 3.82 (d, 1H, $J_{3,4}$ 2.9 Hz, H-4^{III}), 3.79–3.61 (m, 18H, H-6^{Ia}, H-6^{Ib}, H-2^{II}, H-3^I, H-4^{II}, H-5^{III}, PEG $\times$ 6), 3.56–3.1 (m, 2H, H-3^{III}, H-5^I), 3.47 (dd, 1H, $J_{1,2}$ 7.6, $J_{2,3}$ 9.6 Hz, H-2^{III}), 3.42–3.38 (m, 2H, PEG $\times$ 1), 1.97 (s, 3H, Ac), 1.17 (d, 3H, $J_{5,6}$ 6.5 Hz, H-6^{II}); ^{13}C NMR (CD₃OD, 100 MHz) δ 173.83 (C=O), 103.61 (C1^{III}), 102.40 (C1^I), 100.15 (C1^{II}), 76.40 (C5^I), 76.27, 75.17, 75.07, 74.80, 73.72 (C4^{II}, C3^{III}, C5^{III}, C3^I, C4^I), 72.97 (C2^{III}), 71.51, 71.48, 71.44, 71.40, 71.35 (PEG), 71.15 (C3^{II}), 71.04 (PEG), 70.11 (C4^{III}), 69.98 (C2^{II}), 69.94 (PEG), 67.72 (C5^{II}), 66.96 (C6^{III}), 62.66 (C6^I), 56.71 (C2^I), 51.77 (PEG), 23.08 (Ac), 16.61 (C6^{II}); HRMS (ESI,

negative ion mode) $m/z = 809.2626$ $[M - Na]^-$, calcd for $C_{28}H_{49}N_4O_{21}S$, 809.2610.

4.18. 2-(2-(2-(2-azidoethoxy)ethoxy)ethoxy)ethyl β -D-galactopyranosyl-(1 $\rightarrow$ 4)-[α -L-fucopyranosyl-(1 $\rightarrow$ 3)]-2-acetamido-2-deoxy- β -D-glucopyranoside (4).

Compound **15** (51 mg, 36.1 μ mol) was suspended in dry MeOH (3.0 mL) followed by the addition of *ca.* 28 wt-% MeONa in MeOH (5 μ L, 32.9 μ mol). After stirring at rt under dry atmosphere overnight, the reaction mixture was neutralized by the addition of Dowex 50W-X4 (H^+ form), filtered through cotton, and concentrated under reduced pressure. The residue dissolved in EtOH (4.0 mL) was added $NH_2NH_2 \cdot H_2O$ (10 μ L, 212.8 μ mol). After stirring at 90 $^{\circ}C$ overnight, the reaction mixture was concentrated under reduced pressure. To a solution of the residue in pyridine (3.0 mL) was added Ac_2O (50 μ L, 530.7 μ mol). After stirring at rt under dry atmosphere for 36 h, MeOH (1 mL) was added to quench excess reagents, then concentrated with toluene under reduced pressure. The residue was purified by silica gel column chromatography ($CH_3Cl/MeOH$, 1:0 to 6:1, v/v, linear gradient, Rf 75 system). The obtained compound was dissolved in dry THF (3.0 mL) followed by the addition of $AcOH$ (3 μ L, 58.6 μ mol) and 1M TBAF in THF (550 μ L, 550 μ mol). After stirring at rt under Ar atmosphere for 3 days, the reaction mixture was concentrated under reduced pressure and extracted with $CHCl_3$, washed with satd aq $NaHCO_3$ and brine. The organic layer was dried over $MgSO_4$, filtered through a Celite bed, and concentrated under reduced pressure. The residue was roughly purified by silica gel column chromatography ($CHCl_3/Methnol$, 1:0 to 6:1, v/v, linear gradient, Rf 75

system) to afford **16**. To a solution of **16** in MeOH (4.0 mL) was added MeONa in MeOH (*ca.* 28 wt-%, 6 μ L, 40.9 μ mol). After stirring at rt overnight, the reaction mixture was concentrated under reduced pressure. The residue was subjected to LH-20 size-exclusion column chromatography eluting with H₂O, and then to reversed phase column chromatography eluting with H₂O/MeOH (1:0 to 0:100, v/v, linear gradient, R_f 75 system) to afford **4** (4 mg, 5.47 μ mol, 15%, 5 steps) as colorless amorphous.

$[\alpha]_D^{24} -56.0$ (*c* 0.04, MeOH); R_f 0.58 (CHCl₃/MeOH/H₂O, 5:4:1); ¹H NMR (400 MHz, CD₃OD): δ 5.03 (d, 1H, *J*_{1,2} 3.9 Hz, H-1^{II}), 4.86–4.81 (m, 1H, H-5^{II}), 4.54 (d, 1H, *J*_{1,2} 7.5 Hz, H-1^I), 4.44 (d, 1H, *J*_{1,2} 7.3 Hz, H-1^{III}), 3.95–3.84 (m, 6H, H-6^{III}a, H-4^I, H-2^I, H-3^{II}, PEG $\times$ 1), 3.82–3.60 (m, 19H, H-4^{III}, H-6^Ia, H-4^{II}, H-6^{III}b, H-6^Ib, H-5^{III}, H-2^{II}, PEG $\times$ 6), 3.54–3.35 (m, 6H, H-2^{III}, H-3^{III}, H-3^I, H-5^I, PEG $\times$ 1), 1.97 (s, 3H, Ac), 1.18 (d, 3H, *J*_{5,6} 6.6 Hz, H-6^{II}); ¹³C NMR (CD₃OD, 100 MHz): δ 173.90 (C=O), 103.89 (C1^{III}), 102.40 (C1^I), 100.35 (C1^{II}), 77.42 (C5^I), 76.66 (C3^I, C5^{III}), 75.17 (C3^{II}), 74.90 (C3^{III}), 73.69 (C4^{II}), 72.75 (C2^{III}), 71.69, 71.62, 71.54, 71.51 (PEG), 71.22 (C4^I), 71.13 (PEG), 70.00 (C4^{III}), 69.94 (C2^{II}), 69.90 (PEG), 67.67 (C5^{II}), 62.77 (C6^I), 61.39 (C6^{III}), 57.38 (C2^I), 51.77 (PEG), 23.12 (Ac), 16.61 (C6^{II}); HRMS (ESI, positive ion mode) *m/z* = 753.3000 [M + Na]⁺, calcd for C₂₈H₅₀N₄O₁₈Na, 753.3018.

4.19. 2-(2-(2-(2-azidoethoxy)ethoxy)ethoxy)ethyl α -L-fucopyranosyl-(1 $\rightarrow$ 2)- β -D-galactopyranosyl-(1 $\rightarrow$ 4)-[α -L-fucopyranosyl-(1 $\rightarrow$ 3)]-2-acetamido-2-deoxy- β -D-glucopyranoside

(5).

Compound **21** (24 mg, 16.1 μmol) was dissolved in THF (3 mL) followed by addition of AcOH (3 μL , 64.3 μmol) and TBAF (156 μL , 156 μmol). After stirring at rt under Ar atmosphere for 72 h, the reaction mixture was concentrated under reduced pressure and extracted with CHCl_3 , washed successively with satd aq NaHCO_3 and brine. The organic layer was dried over Na_2SO_4 , filtered through a Celite bed, and concentrated under reduced pressure. The residue was roughly purified by silica gel column chromatography eluting with $\text{CH}_3\text{Cl}/\text{MeOH}$ (1:0 to 6:1, v/v, linear gradient, R_f 75 system). To a solution of the obtained product in dry MeOH (3 mL) was added MeONa in MeOH (*ca.* 28 wt-%, 10 μL , 56.8 μmol). After stirring at rt overnight, the reaction mixture was added DOWEX 50W-X4 (H^+ form) to neutralize the reaction system, and then filtered and concentrated under reduced pressure. The residue was purified by LH20 column chromatography eluting with MeOH to afford **5** (8 mg, 9.1 μmol , 57%, 2 steps) as colorless amorphous.

$[\alpha]_{\text{D}}^{22} -104.4$ (*c* 0.01, MeOH); R_f 0.53 ($\text{CHCl}_3/\text{MeOH}/\text{water}$, 5:4:1); ^1H NMR (400 MHz, CD_3OD): δ 5.17 (d, 1H, $J_{1,2}$ 3.2 Hz, H-1^{IV}), 5.03 (d, 1H, $J_{1,2}$ 3.9 Hz, H-1^{III}), 4.88–4.79 (m, 1H, H-5^{IV}), 4.54–4.51 (m, 2H, H-1^I, H-1^{II}), 4.22–4.16 (m, 1H, H-5^{III}), 3.95–3.61 (m, 29H, PEG $\times$ 6, H-2^I, H-3^I, H-4^I, H-6^{Ia}, H-6^{Ib}, H-2^{II}, H-3^{II}, H-4^{II}, H-5^{II}, H-6^{IIa}, H-6^{IIb}, H-2^{III}, H-3^{III}, H-4^{III}, H-2^{IV}, H-3^{IV}, H-4^{IV}), 3.47–3.43 (m, 1H, H-5^I), 3.39–3.35 (m, 2H, PEG $\times$ 1), 1.97 (s, 3H, Ac), 1.26–1.21 (m, 6H, H-6^{III}, H-6^{IV}); ^{13}C NMR (CDCl_3 , 100 MHz): δ 173.89 (C=O), 102.53, 102.21 (C1^I, C1^{II}), 102.14 (C1^{IV}), 100.36 (C1^{III}), 79.45, 77.45, 76.72, 76.64, 76.52, 75.30, 74.46, 73.71, 73.67, 71.87, 71.68, 71.60, 71.53,

71.25, 71.13, 70.79, 70.13, 69.96, 69.93 (C5^I, C4^I, C5^{II}, C3^I, C4^{II}, C2^{II}, C3^{II}, C4^{II}, C2^{III}, C3^{III}, C2^{IV}, C3^{IV}, C4^{IV}, PEG), 68.28 (C5^{III}), 67.65 (C5^{IV}), 62.94, 62.71 (C6^I, C6^{II}), 57.32 (C2^I), 51.78 (PEG), 23.12 (Ac), 16.86, 16.80 (C6^{III}, C6^{IV}); HRMS (ESI, positive ion mode) m/z = 899.3592 [M + Na]⁺, calcd for C₃₄H₆₀N₄O₂₂Na, 899.3597.

4.20. A typical procedure for synchronous synthesis of **10** and **11**.

Compound **2** (100 mg, 96.3 μmol) was added to a solution of **9** (90 mg, 170.9 μmol) in anhydrous CH₂Cl₂ (1.0 mL), and then diluted with anhydrous Et₂O (2.0 mL). The mixture was kept stirring at rt for 30 min under Ar atmosphere in the presence of activated MSAW 300 (100 mg). *N*-Iodosuccinimide (55 mg, 244.5 μmol) was added to the mixture, and then it was cooled down to −40 °C under Ar atmosphere. Triflic acid (1.7 μL, 19.3 μmol) in anhydrous Et₂O (100 μL) was injected to the mixture. After stirring for 1 h, excess amount of Et₃N was added to terminate the reaction. The mixture was filtered through a Celite bed, diluted with CHCl₃, washed successively with 5% aq Na₂S₂O₃, satd aq NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered through a Celite bed, and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography (*n*-hexane/EtOAc, 1:0 to 0:1, v/v, linear gradient, Rf75 system), providing **10** (69 mg, 48.4 μmol, 49%) and **11** (73 mg, 43.3 μmol, 45%).

Acknowledgments

This work was partially supported by JSPS KAKENHI Grant Number 24550134.

Supplementary data

^1H , H,H-COSY and ^{13}C NMR spectra of compounds **2–5**, **7–8**, **10–21** are provided as

Supplementary data online version.

References

1. Morgan, W. T. J.; Watkins, W. M. *Glycoconjugate J.* **2000**, *17*, 501–530.
2. a) Sakamoto, J.; Furukawa, K.; Cordon-Cardo, C. *Cancer Res.* **1986**, *46*, 1553–1561, b) Murata, K.; Egami, H.; Shibata, Y.; Sakamoto, K.; Misumi, A.; Ogawa, M. *Am. J. Clin. Pathol.* **1992**, *98*, 6775, c) Yin, B. W. T.; Finstad, C. L.; Kitamura, K.; Federici, M. G.; Welshinger, M.; Kudryashov, V.; Hoskins, W. J.; Welt, S.; Lloyd, K. O. *Int. J. Cancer* **1996**, *65*, 406–412.
3. Shida, K.; Misonou, Y.; Korekane, H.; Seki, Y.; Noura, S.; Ohue, M.; Honke, K.; Miyamoto, Y. *Glycobiology* **2009**, *19*, 1018–1033.
4. Rosen, S. D. *Annu. Rev. Immunol.* **2004**, *22*, 129–156.
5. For example, see the following literatures and the references cited therein: a) Miermont, A.; Zeng, Y.; Jing, Y.; Ye, X.-s.; Huang, X. *J. Org. Chem.* **2007**, *72*, 8958–8961, b) Tsai, B.-L.; Han, J.-L.; Ren, C.-T.; Wu, C.-Y.; Wong, C.-H. *Tetrahedron Lett.* **2011**, *52*, 2132–2135, c) Mong, K.-K. T.; Wong, C.-H. *Angew. Chem. Int. Ed.* **2002**, *41*, 4087–4090, d) Pratt, M. R.; Bertozzi, C. R. *Org. Lett.* **2004**, *6*,

- 2345–2348, e) Santra, A.; Yu, H.; Tasnima, N.; Muthana, M. M.; Li, Y.; Zeng, J.; Kenyon, N. J.; Louie, A. Y.; Chen, X. *Chem. Sci.* **2015**, accepted manuscript, DOI 10.1039/C5SC04104J, f) Matsushita, T.; Nagashima, I.; Fumoto, M.; Ohta, T.; Yamada, K.; Shimizu, H.; Hinou, H.; Naruchi, K.; Ito, T.; Kondo, H.; Nishimura, S.-I. *J. Am. Chem. Soc.* **2010**, *132*, 16651–16656.
6. Stütz, A. E.; Dekany, G.; Eder, B.; Illaszewicz, C.; Wrodnigg, T. M. *J. Carbohydr. Chem.* **2003**, *22*, 253–265.
7. a) Shan, Y.; Oulaidi, F.; Lahmann, M. *Tetrahedron Lett.* **2013**, *54*, 3960–3961, b) Hara, A.; Imamura, A.; Ando, H.; Ishida, H.; Kiso, M. *Molecules* **2014**, *19*, 414–437.
8. Ohmae, M.; Takada, J.; Murakami, H.; Kimura, S. *Chem. Lett.* **2011**, *40*, 438–439.
9. Nelson, T. D.; Crouch, R. D. *Synthesis* **1996**, 1031–1069.
10. Muramatsu, W. *J. Org. Chem.* **2012**, *77*, 8083–8091.
11. Komba, S.; Ishida, H.; Kiso, M.; Hasegawa, A. *Bioorg. Med. Chem.* **1996**, *4*, 1833–1847.
12. 1) Windmüller, R.; Schmidt, R. R. *Tetrahedron Lett.* **1994**, *35*, 7927–7930, 2) Depré, D.; Düffels, A.; Green, L. G.; Lenz, R.; Ley, S. V.; Wong C.-H. *Chem. Eur. J.* **1999**, *5*, 3326–3340, 3) Mandal, P. K.; Turnbull, W. B. *Carbohydr. Res.* **2011**, *346*, 2113–2120.
13. Kolb, H. C.; Finn, M. G.; Sharpless, K. B. *Angew. Chem. Int. Ed.* **2001**, *40*, 2004–2021.
14. Hein, C. D.; Liu, X.-M.; Wang, D. *Pharm. Res.* **2008**, *25*, 2216–2230.