TITLE:

Measurement of the Strong Interaction Induced Shift and Width of the 1s State of Kaonic Deuterium at J-PARC

AUTHOR(S):

Zmeskal, J.; Sato, M.; Ajimura, S.; Bazzi, M.; Beer, G.; Berucci, C.; Bhang, H.; ... Yamaga, T.; Yamazaki, T.; Zhang, Q.

CITATION:

Zmeskal, J. ...[et al]. Measurement of the Strong Interaction Induced Shift and Width of the 1s State of Kaonic Deuterium at J-PARC. Acta Physica Polonica B 2015, 46: 101-112

ISSUE DATE:

2015-01

URL:

<http://hdl.handle.net/2433/216662>

RIGHT:

© 2015 ACTA PHYSICA POLONICA B.; 'Acta Physica Polonica B' : Since 1999 all current and past articles are freely available on an Open Access basis and under the conditions of Creative Common License, CC-BY 4.0. Starting from January 1, 2014 we are partially funded by SCOAP3.

MEASUREMENT OF THE STRONG INTERACTION INDUCED SHIFT AND WIDTH OF THE $1s$ STATE OF KAONIC DEUTERIUM AT J-PARC

J. ZMESKAL^a, M. SATO^b, S. AJIMURA^c, M. BAZZI^d, G. BEER^e, C. BERUCCI^a
H. BHANG^f, D. BOSNAR^g, M. BRAGADIREANU^h, P. BUEHLER^a, L. BUSO^{i,j}
M. CARGNELLI^a, S. CHO^f, A. CLOZZA^d, C. CURCEANU^d, A. D'UFFIZI^d
S. ENOMOTO^k, L. FABBETTI^l, D. FASO^{i,j}, C. FIORINI^{m,n}, H. FUJIOKA^o
F. GHIO^p, R. GOLSER^q, C. GUARALDO^d, T. HASHIMOTO^b, R.S. HAYANO^r
T. HIRAIWA^c, M. IIO^k, M. ILIESCU^d, K. INOUE^s, S. ISHIMOTO^k
T. ISHIWATARI^t, K. ITAHASHI^b, M. IWAI^k, M. IWASAKI^{b,u}, S. KAWASAKI^s
J. LACHNER^q, P. LEVI SANDRI^d, Y. MA^b, J. MARTON^a, Y. MATSUDA^v
Y. MIZOI^w, O. MORRAⁱ, P. MOSKAL^x, T. NAGAE^o, H. NOUMI^c, H. OHNISHI^b
S. OKADA^b, H. OUTA^b, D. PIETREANU^h, K. PISCICCHIA^{d,y}, M. POLI LENER^d
A. ROMERO VIDAL^z, Y. SADA^c, A. SAKAGUCHI^s, F. SAKUMA^b
E. SBARDELLA^d, A. SCORDO^d, M. SEKIMOTO^k, H. SHI^d, M. SILARSKI^{d,x}
D. SIRGHI^{d,h}, F. SIRGHI^{d,h}, K. SUZUKI^a, S. SUZUKI^k, T. SUZUKI^r, K. TANIDA^f
H. TATSUNO^k, M. TOKUDA^u, A. TOYODA^k, I. TUCAKOVIC^d, K. TSUKADA^α
O. VAZQUEZ DOCE^l, E. WIDMANN^a, T. YAMAGA^t, T. YAMAZAKI^{b,r}
Q. ZHANG^b

^aStefan Meyer Institute for Subatomic Physics, Vienna, Austria;

^bRIKEN Nishina Center, RIKEN, Wako, Japan; ^cResearch Center for Nuclear Physics (RCNP), Osaka University, Osaka, Japan; ^dLaboratori Nazionali di Frascati dell INFN, Frascati, Italy;

^eDepartment of Physics and Astronomy, University of Victoria, Victoria, Canada; ^fDepartment of Physics, Seoul National University, Seoul, South Korea; ^gPhysics Department, University of Zagreb, Zagreb, Croatia; ^hNational Institute of Physics and Nuclear Engineering — IFIN HH, Romania; ⁱINFN Sezione di Torino, Torino, Italy; ^jDipartimento di Fisica Generale, Università di Torino, Torino, Italy; ^kHigh Energy Accelerator Research Organization (KEK), Tsukuba, Japan; ^lExcellence Cluster Universe (TUM), Garching, Germany; ^mPolitecnico di Milano, Dipartimento di Elettronica, Milano, Italy; ⁿINFN Sezione di Milano, Milano, Italy;

^oDepartment of Physics, Kyoto University, Kyoto, Japan; ^pINFN Sezione di Roma I, Istituto Superiore di Sanita, Roma, Italy; ^qIsotope Research and Nuclear Physics, University of Vienna, Vienna, Austria; ^rDepartment of Physics, University of Tokyo, Tokyo, Japan; ^sDepartment of Physics, Osaka University, Osaka, Japan; ^tDepartment of Physics, University of Vienna, Vienna, Austria; ^uDepartment of Physics, Tokyo Institute of Technology, Tokyo, Japan; ^vGraduate School of Arts and Sciences, University of Tokyo, Tokyo, Japan; ^wLaboratory of Physics, Osaka Electro-Communication University, Osaka, Japan; ^xInstitute of Physics, Jagiellonian University, Kraków, Poland; ^yMuseo Storico della Fisica e Centro Studi e Ricerche Enrico Fermi, Roma, Italy; ^zUniversidade de Santiago de Compostela, Santiago de Compostela, Spain; ^αDepartment of Physics, Tohoku University, Sendai, Japan

(Received December 8, 2014)

The antikaon–nucleon ($\bar{K}N$) interaction close to threshold provides crucial information on the interplay between spontaneous and explicit chiral symmetry breaking in low-energy QCD. In this context, the importance of kaonic deuterium X-ray spectroscopy has been well recognized, but no experimental results have yet been obtained due to the difficulty of the measurement. We propose to measure the shift and width of the kaonic deuterium $1s$ state with an accuracy of 60 eV and 140 eV, respectively, at J-PARC. These results, together with the kaonic hydrogen data (KpX at KEK, DEAR and SIDDHARTA at DAΦNE), will then permit the determination of values of both the isospin $I = 0$ and $I = 1$ antikaon–nucleon scattering lengths and will provide the most stringent constraints on the antikaon–nucleon interaction, promising a breakthrough. Refined Monte Carlo studies were performed, including the investigation of background suppression factors for the described setup. These studies have demonstrated the feasibility of determining the shift and width of the kaonic deuterium atom $1s$ state with the desired accuracy of 60 eV and 140 eV.

DOI:10.5506/APhysPolB.46.101

PACS numbers: 25.80.Nv, 29.30.Kv, 29.40.Wk

1. Introduction

1.1. Physics motivation

The antikaon–nucleon ($\bar{K}N$) interaction close to threshold provides crucial information on the interplay between spontaneous and explicit chiral symmetry breaking in low-energy QCD. Techniques such as scattering experiments and X-ray spectroscopy of kaonic atoms [1] as well as production experiments close to threshold [2–7] have been used over many years. Theoretical investigations based on the experimental results have also been performed [8–19]. These approaches are complicated due to the presence of the $\Lambda(1405)$ resonance located just below the K^-p threshold. At present, there are no lattice QCD calculations of antikaon–nucleon scattering lengths, although a theoretical framework has recently been proposed [20]. In particular, the kaonic atom X-ray data provide the most precise values of the antikaon–nucleon scattering lengths at threshold. The K^-p interaction is now well understood from the recent results of kaonic hydrogen obtained from KpX [21] at KEK, DEAR [22] and finally from SIDDHARTA at DAΦNE [23] along with theoretical calculations based on these results [24, 25]. A first milestone was the KpX experiment which used fiducial volume cuts to suppress background, thus solving the long standing kaonic hydrogen puzzle. Although the importance of kaonic deuterium X-ray spectroscopy has been well recognized for more than 30 years (Dalitz *et al.* [26]), no experimental results have yet been obtained due to the difficulty of the X-ray measurement. “*The necessity to perform measurements of the kaonic*

deuterium ground state observables is justified by the fact that, unlike the case of pionic atoms, the measurement of only the kaonic hydrogen spectrum does not allow — even in principle — to extract independently both s -wave $\bar{K}$ -nucleon scattering lengths a_0 and a_1 ”, quoting from Meißner [27]. The kaonic deuterium X-ray measurement represents the most important experimental information missing in the field of the low-energy antikaon-nucleon interactions today. The kaonic hydrogen and deuterium data will also be indispensable when applying unitarized chiral perturbation theory to account for the $\Lambda(1405)$ resonance in kaon-hadronic systems in nuclear matter. There are several hints of K^- quasi-bound states in few-body nuclear systems by FINUDA [28] at DAΦNE and DISTO [29] at SATURNE and at J-PARC with the E15 experiment [30] a search for a K^-pp bound system is ongoing, but the situation is not at all clear.

2. Present state of the art

Recently, the most precise values of the strong interaction shift and width of the kaonic hydrogen $1s$ state were obtained by the SIDDHARTA experiment [23]. The result of the pioneering KpX experiment (KEK-PS E228) together with those from DEAR and SIDDHARTA are shown in Fig. 1. In the latest SIDDHARTA kaonic hydrogen measurement at DAΦNE, an exploratory kaonic deuterium measurement was also made, in part to clarify the hydrogen background as well as to set a limit on the strength of the deuterium signal [31].

Fig. 1. Comparison of experimental results for the strong interaction shift and width of the kaonic hydrogen $1s$ state.

The experimentally determined shift and width are related to the s -wave scattering lengths at threshold [1]. Because of isospin conservation, only the average value of the isospin $I = 0$ and $I = 1$ scattering lengths (a_0 and a_1) was obtained from the kaonic hydrogen measurement. In order to determine the isospin dependent scattering lengths, a measurement of the shift and width of the kaonic deuterium $1s$ state is definitely needed [24–27, 32–34]. On the theoretical side, there are many recent publications predicting quite consistent values of the shift and width for the kaonic deuterium $1s$ state (see Table I).

TABLE I

Compilation of predicted K^-d scattering lengths a_{k-d} and corresponding experimental quantities ϵ_{1s} and Γ_{1s} (taken from [31], Table 1).

a_{k-d} [fm]	ϵ_{1s} [eV]	Γ_{1s} [eV]	Ref.
$-1.58 + i1.37$	-887	757	Mizutani 2013 [35]
$-1.48 + i1.22$	-787	1011	Schevchenko 2012 [36]
$-1.46 + i1.08$	-779	650	Meissner 2011 [25]
$-1.42 + i1.09$	-769	674	Gal 2007 [33]
$-1.66 + i1.28$	-884	665	Meißner 2006 [27]

3. Proposed experimental method

The proposed experiment will measure the transition X-ray energies to the ground state of kaonic deuterium atoms using recently developed Silicon Drift Detectors (SDDs) and using the experience gained with SDDs at KEK with the E570 [37] experiment. In addition, charged particle tracking for background reduction will be used, as previously developed for the KpX experiment E228 [21] at KEK, as well as the background studies performed for the E17 experiment [38] at J-PARC to measure $K^-^3\text{He}$. Finally, the results obtained from an exploratory measurement of kaonic deuterium with SIDDHARTA [31] were used to confine the values for the X-ray yield, shift and width used in the Monte Carlo studies for the experiment proposed in this article. The experimental challenge is the very small kaonic deuterium X-ray yield and the difficulty to perform X-ray spectroscopy in the high radiation environment of an extracted beam. It is, therefore, crucial to control and improve the signal-to-background ratio for a successful observation of the kaonic deuterium X-rays.

4. The proposed experimental setup

The proposed kaonic deuterium experiment will use the excellent features of the K1.8BR kaon beam line together with the K1.8BR spectrometer. We plan to upgrade the spectrometer with a cryogenic deuterium target surrounded by recently developed SDDs with a total area of 246 cm^2 (380 single elements) for X-ray detection.

4.1. The setup at the K1.8BR spectrometer

The K1.8BR multi-purpose spectrometer [39] (see Fig. 2) has a quite unique features for our needs, namely a large acceptance cylindrical spectrometer system (CDS), consisting of a cylindrical drift chamber (CDC) for charged particle tracking, essential for efficient background reduction for the proposed study of kaonic deuterium X-rays. The cylindrical drift chamber, with a diameter of 1060 mm and length of 950 mm, is surrounded by a cylindrical detector hodoscope (CDH) to trigger on decay particles. These parts of the experimental setup will be utilized for the proposed experiment. They will be upgraded with a cryogenic target system and a novel X-ray detector system with a total area of 246 cm^2 , having excellent energy resolution (better than 150 eV at 6 keV) and improved timing ($\sim 100 \text{ ns}$). The proposed setup (Fig. 3) will allow tracking of the incoming kaons with segmented plastic scintillators — used also as start counters (T1, T0) —

Fig. 2. The J-PARC K1.8BR spectrometer. The setup consists of a beam line spectrometer, a cylindrical spectrometer system (CDS), a beam sweeping dipole magnet, a caved beam dump equipped with beam monitor hodoscopes, a neutron counter made of an array of plastic scintillation counters equipped with charged veto counters and a proton counter hodoscope.

and the beam line chamber (BLC). The charged particles produced due to kaon absorption on the nucleus will be tracked by the large cylindrical drift chamber (CDC), while the cylindrical detector hodoscope (CDH) is used as the trigger in coincidence with the start counters (T1, T0).

Fig. 3. Sketch of the proposed setup for the kaonic deuterium measurement.

4.2. New Silicon Drift Detectors (SDDs)

We plan to use new SDD chips, which were developed in collaboration with Fondazione Bruno Kessler FBK and Politecnico di Milano in Italy.

Monolithic arrays with 9 detector segments with a total area of 5.76 cm^2 (see Fig. 4) are ideally suited for the proposed experiment. For these SDDs, special preamplifier chips (CUBE), were developed. First test measurements were performed successfully. For example, a 72 hour stability test, performed at an SDD temperature of 120 K, achieved an energy resolution of 130 eV at 6 keV (see Fig. 5). The drift time at this working temperature was measured to be around 200 ns.

Fig. 4. Monolithic array of 3×3 SDDs each $8 \times 8 \text{ mm}^2$ (left); backside of the SDD array mounted on a printed circuit board with preamplifier chip (CUBE) for detector segment (right).

Fig. 5. Test of a single cell at the temperature of 100 K, measured for 72 hours using a Fe-55 radioisotope, which decays via electron capture to Mn-55. The Mn K_{α} -line at 5.9 keV, emitted after electron capture, is measured with high resolution, leading to a FWHM of 127 eV.

4.3. Cryogenic deuterium target and SDD mounting structure

The cryogenic target cell will be made of a 75 μm Kapton body with a diameter of 65 mm and a length of 160 mm, with reinforcement structure made out of aluminium. In addition, the cooling and mounting structure of the SDDs is used to reinforce the target cell in the longitudinal direction. The working temperature of the target cell is around 25 K with a maximum pressure of 0.35 MPa. With these parameters, a gas density of 5% liquid deuterium density (LDD) will be achieved. Finally, 48 monolithic SDD arrays will be placed close together around the target, with a total area of 246 cm^2 containing 380 readout channels (see Fig. 6).

Fig. 6. Design of the cryogenic target and X-ray detector system. The target cell, with a diameter of 65 mm and a length of 160 mm, is closely surrounded by SDDs, about 5 mm away from the target wall.

5. Monte Carlo studies

5.1. Degradation optimisation, kaon stopping distribution

The kaon beam properties were taken from a measurement at the K1.8 BR in June 2012 with a kaon momentum of 1000 MeV/c. For use as a kaon degrader, several materials (carbon, polyethylene and iron) were compared in a simulation for kaons with a central momentum of 700 MeV/c. The highest stopping rates were obtained with a carbon degrader of about 40 cm thickness. In Table II, the optimised results for the carbon degrader thicknesses are summarised for gas and liquid targets. An additional prism-shaped degrader was used to compensate for the position dependence of the momentum. The results of the kaon stopping optimisation in gaseous and liquid deuterium targets respectively is shown in Fig. 7. The simulation to optimise the kaon stops in deuterium started with $7 \times 10^6 K^-$ and a kaon momentum of 0.7 GeV/c. All further calculations are performed only for a gaseous target with a density of 5% LDD.

TABLE II

Kaon stopping density in gaseous and liquid targets, optimised for carbon degrader. The target density is given relative to the liquid deuterium density (LDD).

Degradation thickness [cm]	Kaon target stops per beam kaon (10^{-3})	Target density
40	0.31	0.03
40	0.50	0.05
39	9.5	1

Fig. 7. Kaon stopping distribution in the deuterium gas target with a density of 5% LDD (left), and in the liquid deuterium target (right).

5.2. X-ray yields, cascade calculations

The most recent cascade calculations for kaonic deuterium are shown in Fig. 8, where the X-ray yields per stopped kaon for K_α , K_β and K_γ transitions over a wide density range were calculated. As input in our Monte Carlo simulation, we used a K_α yield of 0.1% for the gas targets and 0.01% for the liquid target, which are between the calculated values of Koike [40] and Jensen [41].

Fig. 8. Kaonic deuterium cascade calculations, for the X-ray yield of K_α , K_β , K_γ and K_{tot} ; figure from reference [40] (left) and from [41] (right).

5.3. Feasibility study

We have simulated the kaonic deuterium X-ray spectrum (3×10^8 K^- produced per day) using the new X-ray detector with an active area of 246 cm^2 . The gaseous target density is 5% of LDD. The results of this calculation are summarised in Table III.

The yield ratios of the $K_\alpha:K_\beta:K_{tot}$ transitions were taken from the kaonic hydrogen data, but with an assumed K_α yield of 10^{-3} for a gaseous deuterium target. For the strong interaction induced shift and width, theoretical predictions of shift = -800 eV and width of 750 eV were used (see Table I). The simulated spectrum for the transition energies of kaonic deuterium atoms is shown in Fig. 9, using the vertex cut method (successfully used with E570 at KEK and planned as well for E17 at J-PARC) with a defined volume inside the target (5 mm away from the walls), and a charged particle veto for tracks passing through or nearby to the SDDs. For a gas density of 5%, the estimated signal to background ratio (integral) is 1:4 (see Fig. 9). Fitting a set of simulated data, we extracted the shift and width with a precision of 60 eV and 140 eV, respectively, which is comparable with the precision of the kaonic hydrogen (K^-p) SIDDHARTA result of the shift and width measurement.

TABLE III

Simulated values for synchronous background and the K_α transition X-ray yield. For the asynchronous background, the adapted estimation of E17 is used. Background and signal are calculated for an active SDD area of 246 cm^2 .

Produced K^- per day (80% duty cycle)	3.0×10^8
Numbers of K^- at beam counter (T1)	17.4×10^6
Trigger: $T0 \otimes$ (charged particle in CDH)	11.4×10^6
Number of gas stops	1.5×10^5
Total synchronous BG per keV at 7 keV	3.6×10^4
Synchronous BG per keV at 7 keV with	90
Fiducial cut and charged particle veto	
asynchronous BG per keV at 7 keV	30
Detected K_α events for a yield $Y(K_\alpha) = 0.1\%$	
with fiducial cut and charged particle veto	30

Fig. 9. Simulated kaonic deuterium X-ray spectrum, assuming 10^{10} produced K^- and a detector area of 246 cm^2 . In addition, demanding a vertex cut (fiducial volume 5 mm off the walls) as well as the charged particle veto for tracks passing through the SDDs.

6. Conclusion

Assuming a kaon intensity of about $3 \times 10^8 K^-$ per day with a momentum of $700 \text{ MeV}/c$ and rejecting the kaon correlated background using various cuts, it will be possible to collect 1000 K_α events with a signal to

background ratio of 1:4 in 30 days. The asynchronous background will be efficiently reduced due to the unprecedented timing resolution of the new SDD (~ 100 ns).

This work is supported by RIKEN, KEK, RCNP, a Grant-in-Aid for Scientific Research on Priority Areas [No. 17070005 and No. 20840047], a Grant-in-Aid for Specially Promoted Research [No. 20002003], a Grant-in-Aid for Young Scientists (Start-up) [No. 20028011], a Grant-in-Aid for Scientific Research on Innovative Areas [No. 21105003], the Austrian Science Fund (FWF) [P24756-N20], and the Polish National Science Centre grant No. 2013/11/B/ST2/04245.

REFERENCES

- [1] J. Zmeskal, *Progr. Part. Nucl. Phys.* **61**, 512 (2008).
- [2] P. Winter *et al.*, *Phys. Lett.* **B635**, 23 (2006).
- [3] Y. Maeda *et al.*, *Phys. Rev.* **C77**, 015204 (2008).
- [4] M. Silarski *et al.*, *Phys. Rev.* **C80**, 045202 (2009).
- [5] Q.-J. Ye *et al.*, *Phys. Rev.* **C85**, 035211 (2012).
- [6] M. Silarski, P. Moskal, *Phys. Rev.* **C88**, 025205 (2013).
- [7] G. Agakishiev *et al.*, *Phys. Rev.* **C87**, 025201 (2013).
- [8] U.-G. Meißner, U. Raha, A. Rusetsky, *Eur. Phys. J.* **C35**, 349 (2004).
- [9] B. Borasoy, R. Nisßler, W. Weise, *Phys. Rev. Lett.* **94**, 213401 (2005).
- [10] B. Borasoy, R. Nisßler, W. Weise, *Eur. Phys. J.* **A25**, 79 (2005).
- [11] B. Borasoy, U.-G. Meißner, R. Nisßler, *Phys. Rev.* **C74**, 055201 (2006).
- [12] J.A. Oller, J. Prades, M. Verbeni, *Phys. Rev. Lett.* **95**, 172502 (2005).
- [13] J.A. Oller, *Eur. Phys. J.* **A28**, 63 (2006).
- [14] B. Borasoy, R. Nisßler, W. Weise, *Phys. Rev. Lett.* **96**, 199201 (2006).
- [15] J.A. Oller, J. Prades, M. Verbeni, *Phys. Rev. Lett.* **96**, 199202 (2006).
- [16] J. Révai, N.V. Shevchenko, *Phys. Rev.* **C79**, 035202 (2009).
- [17] A. Cieplý, J. Smejkal, *Eur. Phys. J.* **A34**, 237 (2007).
- [18] E. Oset, A. Ramos, *Nucl. Phys.* **A635**, 99 (1998).
- [19] W. Weise, *Nucl. Phys.* **A835**, 51 (2010).
- [20] M. Lage, U.-G. Meißner, A. Rusetsky, *Phys. Lett.* **B681**, 439 (2009).
- [21] M. Iwasaki *et al.*, *Phys. Rev. Lett.* **78**, 3067 (1997).
- [22] G. Beer *et al.*, *Phys. Rev. Lett.* **94**, 212302 (2005).
- [23] M. Bazzi *et al.*, *Phys. Lett.* **B704**, 113 (2011).
- [24] Y. Ikeda, T. Hyodo, W. Weise, *Phys. Lett.* **B706**, 63 (2011).

- [25] M. Döring, U.-G. Meißner, *Phys. Lett.* **B704**, 663 (2011).
- [26] R.H. Dalitz, J. McGinley, C. Belyea, S. Anthony, Theory of low-energy kaon–nucleon scattering, in: Proc. Int. Conf. Hypernuclear and Kaon Physics, ed. by B. Povh, MPI, Heidelberg 1982, p. 201.
- [27] U.-G. Meißner, U. Raha, A. Rusetsky, *Eur. Phys. J.* **C47**, 473 (2006).
- [28] M. Agnello *et al.*, *Phys. Rev. Lett.* **94**, 212303 (2005).
- [29] T. Yamazaki *et al.*, *Phys. Rev. Lett.* **104**, 132502 (2010).
- [30] M. Iio *et al.*, Proposals for Nuclear and Particle Physics Experiments at J-PARC, the 1st PAC meeting, Fri 30 June–Sun 02 July, 2006.
- [31] M. Bazzi *et al.*, *Nucl. Phys.* **A907**, 69 (2013).
- [32] S. Bianco *et al.*, *Riv. Nuovo Cim.* **22**, 1 (1999).
- [33] A. Gal, *Int. J. Mod. Phys.* **A22**, 226 (2007).
- [34] V. Baru, E. Epelbaum, A. Rusetsky, *Eur. Phys. J.* **A42**, 111 (2009).
- [35] T. Mizutani, C. Fayard, B. Saghai, K. Tsushima, [arXiv:1211.5824](https://arxiv.org/abs/1211.5824) [hep-ph].
- [36] N.V. Shevchenko, *Nucl. Phys.* **A890–891**, 50 (2012).
- [37] S. Okada *et al.*, *Phys. Lett.* **B653**, 387 (2007).
- [38] G. Beer *et al.*, Proposals for Nuclear and Particle Physics Experiments at J-PARC, the 1st PAC meeting, Fri 30 June–Sun 02 July, 2006.
- [39] K. Agari *et al.*, *Prog. Theor. Exp. Phys.*, 02B011 (2012).
- [40] T. Koike, T. Harada, Y. Akaishi, *Phys. Rev.* **C53**, 79 (1996).
- [41] T.S. Jensen, Proceedings of DAΦNE 2004: Physics at meson factories, June 2004.