

Title	Taxonomic Relationships of Two Species of Rhacophorus from Vietnam (Amphibia, Rhacophoridae)
Author(s)	MATSUI, MASAFUMI
Citation	Current Herpetology (2005), 24(2): 91-93
Issue Date	2005-12
URL	http://hdl.handle.net/2433/216878
Right	© 2005 by The Herpetological Society of Japan
Type	Journal Article
Textversion	publisher

Taxonomic Relationships of Two Species of *Rhacophorus* from Vietnam (Amphibia, Rhacophoridae)

Author(s): MASAFUMI MATSUI

Source: Current Herpetology, 24(2):91-93.

Published By: The Herpetological Society of Japan

DOI: [http://dx.doi.org/10.3105/1345-5834\(2005\)24\[91:TROTSO\]2.0.CO;2](http://dx.doi.org/10.3105/1345-5834(2005)24[91:TROTSO]2.0.CO;2)

URL: [http://www.bioone.org/doi/](http://www.bioone.org/doi/full/10.3105/1345-5834%282005%2924%5B91%3ATROTSO%5D2.0.CO%3B2)

[full/10.3105/1345-5834%282005%2924%5B91%3ATROTSO%5D2.0.CO%3B2](http://www.bioone.org/doi/full/10.3105/1345-5834%282005%2924%5B91%3ATROTSO%5D2.0.CO%3B2)

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

Taxonomic Relationships of Two Species of *Rhacophorus* from Vietnam (Amphibia, Rhacophoridae)

MASAFUMI MATSUI

Graduate School of Human and
Environmental Studies, Kyoto University,
Sakyo, Kyoto 606–8501, JAPAN

Abstract: *Rhacophorus notater* Smith, 1924, described from a single juvenile, is regarded as conspecific with *R. annamensis* Smith, 1924. Both species were described from the same locality in Vietnam. Although the name *R. notater* appears before *R. annamensis* in the same publication, *R. annamensis* is more popular in name among herpetologists, and its biological details are better clarified than those of *R. notater*. Thus, I propose to treat *R. notater* as a junior synonym of *R. annamensis*.

Key words: Ontogenetic color change; *Rhacophorus annamensis*; *Rhacophorus notater*; Synonymy; Taxonomy; Vietnam

Rhacophorus annamensis was described from Daban (alt. 200 m), Phan Rang Province, S. Annam (=Vietnam) by Smith (1924). The species was once regarded as a subspecies of *R. pardalis* (Wolf, 1936; Bourret, 1942). *Rhacophorus annamensis*, however, is morphologically quite distinct from *R. pardalis* Günther, 1859, and is now considered a good species (Inger, 1954; Inger et al., 1999). Smith (1924) also described another, more enigmatic rhacophorid, *R. notater*, from the same locality as *R. annamensis* (Daban, alt. 200 m) in the same publication.

My examination of the type of *R. notater* Smith, 1924, deposited in the Natural History Museum, London (BM 1947.2.8.83=1924.1.31.3, originally M. Smith 2797), confirmed that the individual is just metamorphosed with

the trace of a tail stub. The specimen has a snout-vent length (SVL) of 19.0 mm, a tibia length of 10.6 mm, and a head width of 6.7 mm. The body is pale gray brown with discrete dark brown blotches on the back (Fig. 1).

A photograph of a juvenile *R. annamensis* just after metamorphosis (Abb. 6 of Ryboltovsky, 1999: 12) clearly shows that the young of this species have a body color quite different from that of adults. Ryboltovsky (1999) described the ground color of the body as bright gray or whitish, with large black spots. This peculiar color pattern is quite similar to that of *R. notater*. Indeed Wolf (1936) suspected that *R. notater* was a young *R. annamensis*.

Ryboltovsky (1999) reported the size at metamorphosis of reared individuals to range from 12 to 14 mm in SVL, and this size range is smaller than the size of *R. notater*. However, the size at metamorphosis varies greatly in anurans according to the conditions the tadpoles have experienced. In contrast, the body color varies less evidently than the body size (my unpublished data).

Recently, Orlov et al. (2002: 96) noted close similarity of body color between *R. notater* and young *R. annamensis*, and listed only *R. annamensis* in their checklist of Vietnamese amphibians, omitting *R. notater*.

I fully agree with Orlov et al. (2002) in that the two species are very similar, and further consider them to be conspecific. Although Smith (1924) described *R. notater* on page 227 before *R. annamensis* on page 229 of the same paper (Proc. Zool. Soc. London, 1924), *R. annamensis* is more popular in name and its biological details are better established (e.g., Inger et al., 1999) than those of *R. notater*. Therefore, I herewith formally propose to relegate *R. notater* in the synonymy of *R. annamensis*.

ACKNOWLEDGMENTS

I thank C. McCarthy and B. T. Clarke of The Natural History Museum, London, for allowing me to examine specimens under their



FIG. 1. *Rhacophorus notater*. Dorsal (top left), ventral (top right) and lateral (bottom) views of the holotype.

care, and T. Hikida, H. Ota, N. Ananjeva and N. Orlov for providing literature. This study was supported by a grant under The Monbusho International Scientific Research Program (Field Research, No. 10041166).

LITERATURE CITED

- BOURRET, R. 1942. Les Batraciens de l'Indochine. Mem. Inst. Océanogr. l'Indochine, Hanoi 6: x, 1-547, 4 pls.
- INGER, R. F. 1954. Systematics and zoogeography

- of Philippine Amphibia. *Fieldiana: Zool.* 33: 181–531,
- INGER, R. F., N. ORLOV, AND I. DAREVSKY. 1999. Frogs of Vietnam: A report on new collections. *Fieldiana: Zool. N. S.* (92): 1–46.
- ORLOV, N. L., R. W. MURPHY, N. B. ANANJEVA, S. A. RYABOY, AND T.-C. HO. 2002. Herpetofauna of Vietnam, a checklist. Part I. Amphibia. *Russ. J. Herpetol.* 9: 81–104.
- RYBOLTOVSKY, E. 1999. Bemerkungen zur nachzucht des Flugfrosches *Rhacophorus annamensis*. *Elaphe* 4: 9–12.
- SMITH, M. A. 1924. New tree-frogs from Indo-China and the Malay Peninsula. *Proc. Zool. Soc. London* 1924: 225–234, 3 pls.
- WOLF, S. 1936. Revision der untergattung *Rhacophorus* (Ausschliesslich der Madagaskar-Formen). *Bull. Raffles Mus.* 12: 137–217.
-

Accepted: 16 November 2005