TITLE:

A New Species of *Leptolalax* from Sarawak, Western Borneo (Anura: Megophryidae)

AUTHOR(S):

Matsui, Masafumi; Zainudin, Ramlah; Nishikawa, Kanto

CITATION:

Matsui, Masafumi ...[et al]. A New Species of *Leptolalax* from Sarawak, Western Borneo (Anura: Megophryidae). *Zoological Science* 2014, 31(11): 773-779

ISSUE DATE:

2014-11

URL:

<http://hdl.handle.net/2433/216902>

RIGHT:

© 2014 Zoological Society of Japan

A New Species of *Leptolalax* from Sarawak, Western Borneo (Anura: Megophryidae)

Masafumi Matsui^{1*}, Ramlah Zainudin², and Kanto Nishikawa¹

¹Graduate School of Human and Environmental Studies, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan

²Molecular Ecology Laboratory, Faculty of Resource Science and Technology, Universiti Malaysia Sarawak, 94300, Kota Samarahan, Sarawak, Malaysia

A new megophryid species is described from southwestern Sarawak, Malaysian Borneo. In appearance, *Leptolalax marmoratus* sp. nov. is most similar to *L. hamidi* also from southwestern Sarawak, but differs from it by mtDNA sequence, larger body size, and higher dominant frequency of advertisement call. The assumption that more than one species of *Leptolalax* coexist at one locality in Borneo is supported. The finding of the new species raises the species number of *Leptolalax* known from Borneo to nine, and the island is thought to be one of the diversification centers of the genus.

Key words: acoustics, Borneo, *Leptolalax*, new species, Southeast Asia, molecular phylogeny

INTRODUCTION

It is now becoming common to identify cryptic anuran species through molecular phylogenetic methods, but an even more biologically significant method is acoustic analysis. Analyses of call characteristics play an important role in detecting cryptic taxa among frogs, and this is particularly the case in the Southeast Asian megophryid genus *Leptolalax* Dubois, 1980, whose members are otherwise very difficult to identify (Malkmus and Riede, 1993; Matsui, 1997, 2006; Matsui et al., 2009; Matsui and Dehling, 2012; Dehling and Matsui, 2013; Matsui et al., 2014). Eight species of *Leptolalax*, *L. arayai* Matsui, 1997, *L. dringi* Dubois, 1987, *L. fritinniensis* Dehling and Matsui, 2013, *L. gracilis* (Günther, 1872), *L. hamidi* Matsui, 1997, *L. maurus* Inger, Lakim, Biun, and Yambun, 1997, *L. pictus* Malkmus, 1992, and *L. sabahmontanus* Matsui, Nishikawa, and Yambun, 2014, have been recognized from the island of Borneo. Call characteristics of all of the named species have been described and proved to be species-specific (Malkmus and Riede, 1993; Matsui, 1997; Malkmus et al., 2002; Sukumaran et al., 2010; Matsui and Dehling, 2012; Dehling and Matsui, 2013; Matsui et al., 2014).

During our field survey in southwestern Sarawak, Malaysian part of Borneo Island (Fig. 1), we collected several *Leptolalax* specimens that are very similar to *L. hamidi* in color pattern, but which had significantly larger body sizes than that species. Moreover, their calls were higher in frequency than in *L. hamidi* in spite of their larger body size. Detailed analyses of acoustic, morphological, and molecular characteristics have revealed that they represent a distinct species that is different from any other congeners. In this paper, we describe this species as new to science.

Fig. 1. Map of Sarawak, Malaysian Borneo, showing the known distribution of *Leptolalax marmoratus* sp. nov. (filled circles), *L. hamidi* (open circles), and literature record of either of the two species (half-filled circles). The filled star indicates the type locality.

MATERIALS AND METHODS

We collected *Leptolalax* in southwestern Sarawak (Annah Rais, Padawan; Gunung [= Mount] Penrissen, Padawan; and Ranchan, Serian) during our field survey during 2009–2014. We made recordings of calls in the field using digital recorders (Zoom H2 or Olympus LS11). At the time of recording, we made temperature measurements using a quick-recording thermistor thermometer. We analyzed the recorded calls with SoundEdit Pro (MacroMind-Paracom, Inc.) and Raven Lite 1.0 for Mac OS X (<http://www.birds.cornell.edu/raven>) on a Macintosh computer. Temporal data were obtained from the oscillogram and frequency information was obtained from the audiospectrograms using fast Fourier transformation (1024 point Hanning window).

After recording calls, we collected specimens, took tissues for subsequent biochemical analysis, and fixed the specimens for vouchers. Specimens, fixed in 10% formalin and later preserved in 70% ethanol, are stored at the Molecular Ecology Laboratory, Faculty of Resource Science and Technology, Universiti Malaysia Sarawak (UNIMAS), Sarawak Research Collections, Forest Research Center

* Corresponding author. Tel. : +81-75-753-6846;
Fax : +81-75-753-6846;
E-mail: fumi@zoo.zool.kyoto-u.ac.jp
doi:10.2108/zs140137

Sarawak (SRC), and Graduate School of Human and Environmental Studies, Kyoto University (KUHE). Twenty body measurements were taken, mainly following Matsui (1984, 1997): (1) snout-vent length (SVL); (2) head length (HL); (3) snout-nostril length (S-NL); (4) nostril-eyelid length (N-EL); (5) snout length (SL); (6) eye length (EL, including eyelid); (7) tympanum-eye length (T-EL); (8) tympanum diameter (TD); (9) head width (HW); (10) internarial distance (IND); (11) interorbital distance (IOD); (12) upper eyelid width (UEW); (13) forelimb length (FLL); (14) lower arm length (LAL); (15) inner palmar tubercle length (IPTL); (16) hindlimb length (HLL); (17) tibia length (TL); (18) foot length (FL); (19) inner metatarsal tubercle length (IMTL); and (20) first toe length (1TOEL). We made all measurements to the nearest 0.1 mm with dial calipers under a binocular dissecting microscope. We followed Savage (1997) in describing toe-webbing states.

DNA sequence data were obtained from tissue samples preserved in 99% ethanol. We reconstructed phylogenetic trees from approximately 2300 base pairs (bp) of the partial sequences of mitochondrial 12S and 16S rRNA genes to clarify the genetic structure of six specimens of *Leptolalax* sp. from Sarawak (KUHE 53191 and 53192 from Annah Rais, Padawan; KUHE 53925 and 54471 from Gunung Penrissen, Padawan; KUHE 53204 and 53227 from Ranchang, Serian). For comparisons, *L. hamidi* from Bukit Lanjak (KUHE 17545, paratype) was also sequenced. The resultant sequences were deposited in GenBank (AB969287–969292). Published data for each one sample of *L. arayai* (BORNEENSIS 22931 from Liwagu, Kinabalu: AB8475558), *L. dringi* (KUHE 55610 from Camp 4 of Gunung Mulu: AB847553), *L. fritinniensi* (KUHE 55371 from Headquarters, Gunung Mulu: AB847557), *L. gracilis* (KUHE 55624 from Camp 1, Gunung Mulu: AB847560), *L. maurus* (SP 21450 from Kinabalu: AB847559), *L. pictus* (KUHE 39298 from Poring, Kinabalu: AB847556), and *L. sabahmontanus* (BORNEENSIS 12632 from Liwagu, Kinabalu: AB847551), and outgroup species *L. heteropus* (Boulenger, 1900) (KUHE 15490 from Larut, Peninsular Malaysia: AB847561) and *Leptobrachium hasseltii* Tschudi, 1838 (KUHE 42820 from Java: AB530424) were used for comparisons. Methods for DNA extraction and, amplification and sequencing of the mtDNA fragments are the same as those reported by Matsui et al. (2010) and Hamidy et al. (2011). The alignment matrix with 2289 mtDNA nucleotide sites was subjected to estimate phylogenetic relationships using maximum likelihood (ML) and Bayesian inference (BI). Pairwise comparisons of uncorrected sequence divergences (p-distance) were also calculated for 16S rRNA. Details for all these procedures are given in Hamidy et al. (2011).

RESULTS

We obtained 2289 bp of concatenated fragments of mtDNA genes for all samples, including out-groups (Fig. 2), of which sites, 756 were variable, and 398 were parsimoniously informative. The best substitution model was GTR with gamma shape parameter (G) of 0.312 for ML and 0.332 for BI, and the likelihood values (-lnLs) of the ML and BI trees were 11797.418 and 11817.707, respectively.

Phylogenetic analyses employing two different optimality criteria yielded nearly identical relationships. As shown in the ML tree in Fig. 2, six specimens of *Leptolalax* sp. from Sarawak obtained from three localities proved to form a fully supported clade (MLBS = 100, Bayesian posterior probability [BPP] = 1.00). *Leptolalax* sp. from Sarawak, *L. arayai*, and *L. hamidi* also formed a fully supported clade, but within this clade, sister species relationship of *L. arayai* and *L. hamidi* was not resolved (MLBS = 63, BPP = 0.70). The genetic distance (uncorrected p-distance of 16S rRNA) of the clade of *Leptolalax* sp. from Sarawak from *L. hamidi* was substantially large, with the mean of 6.0% (range = 5.9–

Fig. 2. Maximum likelihood (ML) tree of a 2289 bp sequence of mitochondrial 12S and 16S rRNA for samples of *Leptolalax* (for sample details, see text). Numbers above or below branches represent bootstrap supports for ML inference and Bayesian posterior probability (ML-BS/BPP).

Table 1. Uncorrected p-distances (in %) for fragment of 16S rRNA among *Leptolalax* taxa compared.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1 <i>Leptolalax</i> sp. KUHE 53192 Annah Rais															
2 <i>Leptolalax</i> sp. KUHE 53191 Annah Rais	0.3														
3 <i>Leptolalax</i> sp. KUHE 53204 Serian	0.3	0													
4 <i>Leptolalax</i> sp. KUHE 53227 Serian	0.3	0	0												
5 <i>Leptolalax</i> sp. KUHE 53925 Penrissen	0.3	0	0	0											
6 <i>Leptolalax</i> sp. KUHE 54471 Penrissen	0.3	0	0	0	0										
7 <i>L. hamidi</i> KUHE 17545	6.2	5.9	5.9	5.9	5.9	5.9									
8 <i>L. arayai</i> BORN 22931	5.7	5.5	5.5	5.5	5.5	5.5	6.0								
9 <i>L. sabahmontanus</i> BORN 12632	10.9	10.7	10.7	10.7	10.7	10.7	11.1	11.5							
10 <i>L. dringi</i> KUHE 55610	11.1	10.9	10.9	10.9	10.9	10.9	11.6	11.6	5.0						
11 <i>L. fritinniensi</i> KUHE 55371	11.5	11.3	11.3	11.3	11.3	11.3	11.9	11.7	6.8	7.2					
12 <i>L. pictus</i> KUHE 39298	10.1	9.8	9.8	9.8	9.8	9.8	10.9	10.5	6.1	5.9	6.3				
13 <i>L. maurus</i> SP 21450	9.9	9.7	9.7	9.7	9.7	9.7	10.5	9.7	11.1	11.6	12.0	11.2			
14 <i>L. gracilis</i> KUHE 55624	15.2	15.0	15.0	15.0	15.0	15.0	15.6	15.0	14.7	14.7	15.3	13.5	14.4		
15 <i>L. heteropus</i> KUHE 15490	21.0	20.9	20.9	20.9	20.9	20.9	20.5	20.9	20.1	20.1	19.8	19.1	20.9	21.1	
16 <i>Leptobrachium hasseltii</i> KUHE 42820	21.6	21.4	21.4	21.4	21.4	21.4	21.5	21.7	23.1	23.5	22.5	21.9	21.8	23.3	23.2

Fig. 3. Dorsolateral view of a male paratype of *Leptotalax marmoratus* sp. nov. (SRC unnumbered [former KUHE 53192]) from the type locality.

6.2%: Table 1), and was even larger than that from *L. arayai* (mean = 5.5%, range = 5.5–5.7%: Table 1). These values approximated the distances between distinctly different species *L. dringi* and *L. sabahmontanus* (5.0%), and *L. fritinniens* and *L. pictus* (6.3%).

In addition to genetic separation, the specimens of *Leptotalax* sp. from Sarawak are also clearly separated morphologically and acoustically from all the other congeners. We thus describe *Leptotalax* sp. from Sarawak as follows:

SYSTEMATICS

Leptotalax marmoratus sp. nov.

(Figs. 4, 5)

Diagnosis

A large-sized form of the genus *Leptotalax* (males 32.3–38.0 mm, females 41.4–46.8 mm in SVL); tibiotarsal articulation of adpressed limb at least reaching nostril; differing from all the other congeners except for *L. hamidi* and *L. pictus* by unique dorsal coloration, with large dark brown markings on a tan to light brown background of dorsum, and creamy ventrum without dark markings; from very similar *L. hamidi*, it differs in its much larger body size (males 28.7–28.8 mm, females 35.1–37.4 mm SVL in *L. hamidi*), presence of discrete dark spots posterior to sacrum, and higher dominant frequency of advertisement call (7.4 to 8.2 kHz vs. 6.7 to 7.3 kHz in *L. hamidi*) in spite of larger body size; it overlaps *L. pictus* in body size (males 29.8–34.0 mm and females 38.6–42.3 mm) but differs by much more conspicuous, discrete dorsal markings, including a large dark marking on top of snout, and higher dominant frequency of advertisement call (snout marking faint and dominant frequency 6.8–7.2 kHz in *L. pictus*).

Etymology

The specific epithet is an adjective from Greek “marmaro”, meaning marble, in reference to beautifully blotched dorsum of the new species.

Holotype

KUHE 53193, an adult male from Annah Rais, Padawan, Kuching Division, Sarawak, Malaysia (01°08'35" N, 110°15'47" E, 71 m a.s.l.), collected on 28 August 2009

Fig. 4. Dorsal (A) and ventral (B) views of male holotype of *Leptotalax marmoratus* sp. nov. (KUHE 53193). Scale bar = 10 mm.

Fig. 5. Ventral view of right hand (A) and foot (B) of male holotype of *Leptotalax marmoratus* sp. nov. (KUHE 53193). Scale bar = 5 mm.

by M. Matsui.

Paratypes

All from Sarawak. USM GR 006, 095, two females from Sungei (= River) Regu, Padawan (01°12'16" N, 110°16'07" E) collected on 16 and 18 August 2009 by R. Zainudin; SRC unnumbered (former KUHE 48804, one male, collected on 7 March 2014 by M. K. Nishikawa from Ranchan, Serian, Samarahan Division [01°08'28" N, 110°34'59" E, 52 m a.s.l.], and former KUHE 53192, one male, data same as the holotype); KUHE 53191, 53194, 53195, two males and one female, data same as the holotype; KUHE 48805, 48821, 48822, two males and one female, collected on 7 March 2014 by M. Matsui, K. Nishikawa, and K. Eto, KUHE 53204, 53227, 53228, 53229, three males and one female, collected on 29 August 2009 by M. Matsui and K. Nishikawa from Ranchan, Serian; KUHE 48516, 48517, 48521, 48622, two males and two females, collected between 19–21 December 2013 by M. Matsui, K. Nishikawa, and T. Shimada at 01°08'34" N, 110°13'37" E, 432 m a.s.l., KUHE 53924, 53925, two males, collected on 2 September 2010 by M. Matsui and K. Nishikawa at 01°07'31" N, 110°12'49" E, 905 m a.s.l., and KUHE 54471, one male, collected on 21 February 2012 by K. Nishikawa at 01°07'31" N, 110°12'49" E, 905 m a.s.l. on Gunung Penrissen, Kuching Division.

Referred specimens

KUHE 48624, one newly metamorphosed juvenile from Gunung Penrisen (01°08'38" N, 110°13'45" E, 455 m a.s.l.); KUHE 48683, one juvenile from Ranchan, Serian (01°08'28" N, 110°34'59" E, 52 m a.s.l.).

Description of holotype (measurements in mm)

SVL 36.2; habitus slender (Fig. 4); head longer (HL 14.2, 39.2%SVL) than broad (HW 11.4, 31.5%SVL); snout obtusely pointed, rounded in profile, projecting beyond lower jaw; eye large, length (EL 5.6, 15.5%SVL) slightly greater than snout length (SL 5.3, 14.6%SVL); canthus distinct, slightly constricted; lore slightly oblique, concave; nostril lateral, below canthus, distinctly closer to tip of snout than to eye; internarial distance (IND 3.0, 8.3%SVL) subequal to interorbital distance (IOD 2.9, 8.0%), latter much narrower than upper eyelid (UEW 3.9, 10.8%); pineal spot absent; tympanum distinct, diameter (TD 2.8, 7.7%SVL) half that of eye, and separated from eye by one-third of tympanic diameter (T-EL 1.1, 3.0%SVL); vomerine teeth absent; tongue notched, without papillae; vocal sac subgular, bipartite, consisting of two ventrolaterally situated parts; small vocal openings posterior to rictus, close to eustachian tube openings.

Forelimb long (FLL 27.2, 75.1%SVL; LAL 18.9, 52.2%SVL) and slender; fingers slender, unwebbed; finger length formula: $I < II < IV < III$ (Fig. 5A), first finger slightly shorter than second, latter slightly shorter than fourth; finger tips slightly swollen; no fringes of skin along fingers; inner palmar tubercle large (IPTL 2.9, 8.0%SVL), not extending onto first metacarpal and smaller outer palmar tubercle; sub-articular tubercles indistinct, replaced by indistinct, low callous tissue; nuptial pads absent.

Hindlimb long (HLL 62.8, 173.5%SVL); tibia relatively long (TL 20.2, 55.8%SVL), heels overlapping when limbs are held at right angles to body; tibiotarsal articulation of adpressed limb reaching beyond snout; foot (FL 16.9, 46.7%SVL) shorter than tibia; toe length formula $I < II < V < III < IV$; third toe longer than fifth; toe tips similar to those of fingers; webbing confined to bases of toes (Fig. 5B), formula: $I \frac{1}{4} - 2 + II \frac{1}{5} - 3 III \frac{2}{3} - 3 \frac{3}{4} IV 4 - 2 \frac{2}{3} V$; toes without lateral fringes; subarticular tubercles obscure, but elongate, replaced by low callous tissue, a relatively short one on each of second, third, and fifth toes, larger ones on fourth toe; inner metatarsal tubercle low, length (IMTL 2.1, 5.8%SVL) about one-third of first toe (1TOEL 3.0, 8.3%SVL); no outer metatarsal tubercle.

Skin dorsally nearly smooth, scattered with small tubercles of varying sizes; top of snout smooth but eyelid with small tubercles; a low, supratympanic ridge from eye to axilla; sides with larger tubercles; chest and abdomen smooth; supra-axillary gland small, white in color; pectoral gland and ventrolateral glandular ridges absent; femoral gland indistinct, in a large white blotch.

Color

In life tan dorsally on head and body, with large, discrete, cream-edged, dark brown spots, including interorbital, frontal, canthal, and post-sacral spots; pinkish tinges between spots (Fig. 4A); tympanum brown edged black above and white below; lips barred with black; sides with

dark spots; groin with dark spot not extending onto thigh; limbs marked dorsally with alternating light and dark brown crossbars; no light-colored area around the elbow, between upper and lower arms; throat largely unpigmented postero-laterally; chest, and abdomen immaculate white (Fig. 4B); iris clay-white in ventral two-thirds, reddish in dorsal third, with inner edge around pupil ruby-red. In preservative, color and pattern have slightly faded in preservative, but not obviously changed. The pinkish tinges between dorsal dark markings have disappeared, and upper and lower halves of iris have become indistinguishable.

Variation

Individuals of the type series are generally similar to each other in morphology. Variation in size and body proportions is given in Table 2. Specimens from three different localities do not differ in size and proportions of body. Females (SVL 41.4–46.8 mm) are significantly larger than males (SVL 32.3–38.0 mm; t-test, $P < 0.01$), but the sexes do not differ in all morphometric values relative to SVL (Dunn's multiple comparisons test, $P > 0.05$). Tibiotarsal articulation of adpressed limb reaches nostril (in 30.8% of total individuals examined), tip of snout (23.1%), and beyond snout (46.2%) in males, and between anterior corner of eye and nostril (28.6%), nostril (14.3%), nostril to tip of snout (14.3%), and tip of snout (42.9%) in females. Dark brown marking on dorsum slightly varies in the size and shape.

Eggs and larvae

The diameter of 12 ova from ovary of KUHE 53195 ranged from 1.28–1.59 (mean $\pm$ SD = 1.47 ± 0.10) mm. Both animal and vegetal poles were uniformly cream in color. Nothing is known about larvae.

Call characteristics

The temporal and the frequency characteristics of advertisement calls are nearly similar among individuals from three localities. Calls of a male (KUHE 48805), recorded at Ranchan, Serian on 7 March 2014 at an air temperature of 26.4°C, were analyzed in detail. The calls lasted 4.7 ± 2.1 (mean $\pm$ SD; 1.9 – 6.7 , $n = 6$) s with an interval of 1.6 ± 0.1 (1.4 – 1.7 , $n = 5$) s. The call consisted of long series of notes (Fig. 6A), the number of which within a series was 58.8 ± 25.5 (24 – 83 , $n = 15$). Note repetition rate was 12.2 ± 0.1 (12.0 – 12.3 , $n = 6$) per s. Individual notes lasted 27.9 ± 2.1 (24 – 32 , $n = 15$) ms and were separated from each other by an interval of 53.9 ± 3.1 (46 – 59 , $n = 14$) ms. Each note was composed of several indistinct pulses (Fig. 6F). Marked frequency modulation was always observed within a note (Fig. 6E). The dominant frequency at the beginning was 8.03 ± 0.18 (7.83 – 8.55 , $n = 15$) kHz and decreased towards the end of the note to 7.55 ± 0.14 (7.43 – 7.99 , $n = 15$) kHz. The frequency difference between the beginning and the end of the note was 0.48 ± 0.14 (0.16 – 0.64 , $n = 15$) kHz. Usually, the first note of a series was higher in frequency than the subsequent ones (Fig. 6A, C). Irregular intensity modulation was observed within a single call (Fig. 6A).

Comparisons

Leptotalax marmoratus differs from congeneric species occurring outside Borneo in the same manner as in *L.*

Table 2. Measurements in adults of *Leptolax marmoratus* sp. nov. and *L. hamidi*. SVL (mean $\pm$ SD, in mm) and medians of ratios (R) of other characters to SVL, followed by ranges in parenthesis. See text for character abbreviations.

	<i>L. marmoratus</i>		<i>L. hamidi</i>	
	14M	7F	2M	3F
SVL	34.7 $\pm$ 1.63 (32.3–38.0)	44.2 $\pm$ 1.79 (41.4–46.8)	28.8 (28.7–28.8)	36.3 $\pm$ 1.15 (35.1–37.4)
RHL	38.5 (37.2–40.5)	37.3 (35.0–39.7)	40.2 (39.9–40.4)	39.3 (35.0–39.6)
RS-NL	5.8 (5.3–6.6)	5.31 (4.7–6.2)	6.4 (5.6–7.3)	6.0 (5.9–7.7)
RN-EL	8.3 (7.0–9.4)	8.6 (7.5–9.4)	8.7 (8.0–9.4)	8.0 (7.4–9.7)
RSL	13.7 (13.2–15.2)	13.9 (12.8–14.6)	13.9 (12.8–15.0)	14.0 (13.7–14.4)
REL	15.5 (14.3–17.1)	14.4 (13.2–15.5)	17.0 (16.7–17.4)	15.4 (14.6–16.0)
RT-EL	2.7 (2.3–4.2)	3.5 (2.6–4.5)	2.7 (2.6–2.8)	2.7 (2.3–3.2)
RTD	7.2 (6.6–9.0)	6.6 (5.4–7.8)	8.7 (8.3–9.1)	7.2 (6.3–8.5)
RHW	31.5 (30.4–34.2)	32.3 (31.1–33.6)	31.6 (30.7–32.5)	33.4 (33.0–34.5)
RIND	8.4 (7.0–9.4)	7.7 (6.5–8.5)	8.7 (8.7–8.7)	8.2 (7.0–8.5)
RIOD	8.4 (6.0–9.7)	7.5 (6.8–9.1)	10.1 (9.4–10.8)	7.2 (6.3–8.5)
RUEW	11.3 (10.2–13.7)	11.1 (10.4–13.0)	11.8 (11.5–12.2)	11.1 (11.0–11.5)
RFL	70.7 (68.9–75.4)	70 (69.4–76.2)	71.5 (69.7–73.3)	78.3 (71.7–79.1)
RLAL	53.1 (52.1–57.2)	53.2 (51.3–55.8)	52.2 (51.2–53.1)	53.0 (50.8–57.8)
RIPTL	7.5 (6.6–9.9)	7.5 (5.3–10.0)	8.0 (7.7–8.3)	7.1 (7.0–7.7)
RHLL	173.5 (168.5–185.6)	177.2 (163.0–181.7)	172.7 (170.0–175.3)	176.4 (174.3–184.6)
RTL	55.6 (52.4–58.4)	56.2 (51.7–57.6)	55.1 (54.4–55.9)	55.1 (54.1–59.3)
RFL	46.4 (44.3–50.2)	47.8 (43.5–49.3)	45.0 (44.6–45.5)	48.1 (44.9–49.6)
RIMTL	5.7 (4.5–6.3)	5.3 (4.6–6.1)	5.0 (4.9–5.2)	5.5 (5.1–6.1)
R1TOEL	7.5 (7.2–9.2)	7.7 (7.3–8.8)	–	8.7 (8.0–9.4)

fritinniens (Dehling and Matsui, 2013) and *L. sabahmontanus* (Matsui et al., 2014), and comparisons with Bornean species are pertinent. *Leptolax marmoratus* morphologically differs from all the described Bornean species in the following ways. By the absence of ventral markings, it resembles *L. pictus*, *L. hamidi*, and *L. arayai*. Especially, the new species closely resembles *L. hamidi* by having large dark brown dorsal markings with light outlines. However, from *L. hamidi*, *L. marmoratus* differs by having a larger body size, male SVL 32.3–38.0 mm, female 41.4–46.8 mm in SVL, and presence of discrete dark spots posterior to sacrum (vs. 28.7–28.8 mm, female 35.1–37.4 mm, a faint dark spot sometimes

Fig. 6. Sonogram (A, C, E) and wave form (B, D, F) showing advertisement calls of a paratype of *Leptolax marmoratus* sp. nov. (KUHE 48805) from Ranchan, Serian, recorded at an air temperature of 26.4°C.

present between sacrum and cloaca in *L. hamidi*). It differs from *L. arayai* also by the larger body size (male SVL 29.2–32.0 mm in *L. arayai*), lack of strongly shagreened back, presence of dorsal markings, and absence of flush color on ventral side of legs and groin (vs. dorsum tuberculated, without conspicuous markings, venter yellow, and groin and ventral sides of limbs orange in *L. arayai*). *Leptolax marmoratus* resembles *L. pictus* in the body size (male SVL 29.8–34.0 mm SVL in *L. pictus*) and possession of dark brown markings with conspicuous thin light outlines. However, the new species differs from *L. pictus* by having much more conspicuous, discrete dorsal markings, especially a dark marking on top of the snout, which is as large as the tympanum and as distinct as the interorbital marking (vs. dorsal markings less conspicuous, and sometimes not discrete, and a marking on the snout is faint and much less marked than the interorbital marking in *L. pictus*).

The white ventrum without markings clearly differentiates *L. marmoratus* from *L. dringi*, *L. sabahmontanus*, *L. maurus*, *L. gracilis* and *L. fritinniens*, all of which have more or less developed ventral dark marking or a dark brown to black ventrum. Of these, the new species is larger than *L. dringi*, *L. sabahmontanus*, and *L. maurus* (male SVL 32.3–38.0 mm vs., 26.6–31.3 mm in *L. dringi*, 25.4–27.9 mm in *L. sabahmontanus*, and 25.0–26.9 mm in *L. maurus*), but is

similar to *L. gracilis* and *L. fritinniensi* in body size (male SVL 31.1–39.2 mm in *L. gracilis* and 27.4–34.8 mm in *L. fritinniensi*). Furthermore, the new species differs from *L. dringi*, *L. sabahmontanus*, *L. maurus*, and *L. gracilis* by having a bipartite vocal sac in males (vs. medially arranged single vocal sac in *L. dringi*, *L. sabahmontanus*, *L. maurus*, and *L. gracilis*). *Leptolalax fritinniensi* has a bipartite vocal sac like the new species, but the body coloration is completely different from it.

The new species also differs from the known Bornean congeners acoustically. By having clear frequency modulation within a note, the new species differs *L. sabahmontanus*, *L. dringi*, *L. arayai*, *L. maurus*, and *L. gracilis* that have calls lacking frequency modulation and lower dominant frequency than the new species (7.43–8.22 kHz at 26.4°C vs. 6.90–7.35 kHz at 17.2–17.7°C in *L. sabahmontanus*; 6.05–6.40 kHz at unknown temperature in *L. dringi*; 5.40–5.90 kHz at 17.4°C in *L. arayai*; 5.15 kHz at unknown temperature in *L. maurus*; 2.60–2.80 kHz at 20.0–26.2°C in *L. gracilis*). It resembles *L. hamidi*, *L. pictus*, and *L. fritinniensi* in having clear frequency modulation within a note, but the dominant frequency of the call of the new species is higher than that of *L. hamidi* (6.70–7.30 kHz at 22.9–24.1°C) and *L. pictus* (6.80–7.15 kHz at 19–22°C), although it overlaps *L. fritinniensi* (7.25–9.20 kHz at 24.3–24.9°C).

Range

Southwestern region of Sarawak (Fig. 1): Besides the type locality, Annah Rais, Padawan, Kuching Division, the species was recorded from Sungei Regu, Padawan; Ranchan, Serian, Samarahan Division; Gunung Penrisen, Kuching Division. The known localities range 52–905 m in altitude.

Natural history

At all three localities, males of *L. marmoratus* sp. nov. were calling at night on the shrubs 1–50 m from streams (width < 10 m). Breeding season seems to last long, and calls were heard from late February to early March, from late August to early September, and in late December. At Padawan, two females were found on rocks, one 2 m from a stream, and another in a stream in mid August, and one female was observed to approach to a calling male in late August. Species associated with the new species were *Ansonia spinulifer* (Mocquard, 1890), *Phrynomantis asper* (Gravenhorst, 1829), *Leptobrachella mjobergi* Smith, 1925, *Leptolalax gracilis* (Günther, 1872), *Ingerana baluensis* (Boulenger, 1896), *Limnonectes khasianus* (Anderson, 1871), *L. leporinus* Andersson, 1923, *Hylarana raniceps* (Peters, 1871), *Odorrana hosii* (Boulenger, 1891), *Meristogenys jerboa* (Günther, 1872), *Staurois guttatus*, and *Kurixalus appendiculatus* (Günther, 1858).

DISCUSSION

Generally, a larger male frog, with a larger larynx, has a call of lower frequency than a smaller conspecific (e.g., Ryan, 1980), and this holds for interspecifically in some frog groups (e.g., Blair, 1964). However, this seems to be not the case between the new species and *L. hamidi*, which are very similar in general morphology but differ in body size. As shown above, the dominant frequency of 6.70–7.30 kHz in *L. hamidi* (Matsui, 1997) with male body size of 28.7–28.8

mm are lower than those found in the new species (7.43–8.22 kHz), notwithstanding larger body size of the latter species (male SVL 32.3–38.0 mm). Actually, the male SVL and dominant frequency of the mating call do not correlate in the known species of Bornean *Leptolalax* (detailed data shown above). Except for *L. gracilis* (male SVL 31.1–39.2 mm, 2.60–2.80 kHz), larger species like *L. marmoratus* sp. nov. have even higher frequencies than the smaller species like *L. maurus* (25.0–26.9 mm, 5.15 kHz).

In his description of *L. hamidi*, Matsui (1997) erroneously cited Inger et al.'s (1995) data for body size range of the species. *Leptolalax* sp. (morphotype from Sarawak with non-spotted venter) reported by Inger et al. (1995) seems to include both *L. hamidi* and the present new species, since the body size given by Inger et al. (males 28.7–31.3 mm, females 36.1–42.8 mm in SVL) overlaps that of *L. hamidi* (males 28.7–28.8 mm, females 35.1–37.4 mm: Matsui, 1997 and newer data) and also the present new species (males 32.3–38.0 mm, females 41.4–46.8 mm).

Leptolalax sp. (morphotype from Sarawak with non-spotted venter) of Inger et al. (1995) was recorded from headwaters of Sungai Sekerang, Second Division (= Sri Aman Division), and Nanga Tekalit, Kapit District, Seventh Division (= Kapit Division) (Fig. 1). Of these, Sungai Sekerang is close to the type locality of *L. hamidi*, but Nanga Tekalit is further northeast from there. We recently found *L. hamidi* on Gunung Mulu, northern Sarawak, which is far distant from the type locality (Matsui, unpublished data). Thus the range of *L. hamidi* seems wide and may overlap that of the new species.

In all the three known localities, the new species is syntopic with *L. gracilis*, with a similar body size but calls with much lower frequency. This differential acoustic trait would likely prohibit interspecific hybridization between them, working as a pre-mating reproductive isolation mechanism. Therefore, it is possible that the new species and *L. hamidi* occur syntopically because of their different call characteristics.

The finding of *L. marmoratus* sp. nov. raises the species number of *Leptolalax* known from Borneo to nine, and our earlier assumption that the island of Borneo is one of the diversification centers of the genus (Dehling and Matsui, 2013) is further strengthened. Similarly, another hypothesis that coexistence of more than one species of *Leptolalax* at one locality is common in northwestern Borneo (Matsui et al., 2014) can be applied to southwestern region of the island, and also would hold for the entire island, including hitherto little surveyed Indonesian regions of Kalimantan.

ACKNOWLEDGMENTS

The field survey in Sarawak by MM and KN was made under research permissions of Pejabat Jabatan Perhutanan (NPW. 907.4.4(V)-93 and NCCD.907.4.4(Jld.VI)-122). The Sarawak Forestry Corporation and the Sarawak Biodiversity Centre kindly provided facilities for conducting research. We are grateful to the following for their encouragements and/or permission to conduct research and export specimens: Datu Haji Len Talif Salleh, Haji ali Bin Yusop, Marker Nyogin, Mohd. Shabudin Sabki, Uning, Haji Abang Mohamad Mohtar Bin Abang Pawozan, Paulus Ak Meleng, and Takao Ichioka. MM and KN thank Koshiro Eto, Tsutomu Hikida, Tomohiko Shimada, and Siew Teck Yeo for field companionship, and Amir Hamidy, Koshiro Eto, and Norihiro Kuraishi for help in the

laboratory. Field trips by MM and KN were made possible by grants from The Monbusho through the Japan Society for the Promotion of Science (Field Research, Nos. 20405013 and 23405014).

REFERENCES

- Blair WF (1964) Evolution at populational and intrapopulational levels: isolating mechanisms and interspecies interactions in anuran amphibians. *Quart Rev Biol* 39: 333–334
- Dehling JM, Matsui M (2013) A new species of *Leptolalax* (Anura: Megophryidae) from Gunung Mulu National Park, Sarawak, East Malaysia (Borneo). *Zootaxa* 3670: 33–44
- Hamidy A, Matsui M, Shimada T, Nishikawa K, Yambun P, Sudin A, et al. (2011) Morphological and genetic discordance in two species of Bornean *Leptobrachium* (Amphibia, Anura, Megophryidae). *Mol Phyl Evol* 61: 904–913
- Inger RF, Stuebing RB, Tan FL (1995) New species and new records of anurans from Borneo. *Raffles Bull Zool* 43: 115–131
- Malkmus R, Riede K (1993) Nachtrag zu *Leptolalax pictus* Malkmus, 1992 (Anura: Pelobatidae). *Sauria Berlin* 15: 7–9
- Malkmus R, Manthey U, Vogel G, Hoffman P, Kosuch J (2002) Amphibians and Reptiles of Mount Kinabalu (North Borneo). ARG Gantner, Ruggell
- Matsui M (1984) Morphometric variation analyses and revision of the Japanese toads (Genus *Bufo*, Bufonidae). *Contrib Biol Lab Kyoto Univ* 26: 209–428
- Matsui M (1997) Call characteristics of Malaysian *Leptolalax* with a description of two new species (Anura: Pelobatidae). *Copeia* 1997: 158–165
- Matsui M (2006) Three new species of *Leptolalax* from Thailand (Amphibia, Anura, Megophryidae). *Zool Sci* 23: 821–830
- Matsui M, Dehling M (2012) Notes on an enigmatic Bornean megophryid, *Leptolalax dringi* Dubois, 1987 (Amphibia: Anura). *Zootaxa* 3317: 49–58
- Matsui M, Belabut DM, Ahmad N, Yong H-S (2009) A new species of *Leptolalax* (Amphibia, Anura, Megophryidae) from Peninsular Malaysia. *Zool Sci* 26: 243–247
- Matsui M, Hamidy A, Murphy RW, Khonsue W, Yambun P, Shimada T, et al. (2010) Phylogenetic relationships of megophryid frogs of the genus *Leptobrachium* (Amphibia, Anura) as revealed by mtDNA gene sequences. *Mol Phyl Evol* 56: 259–272
- Matsui M, Nishikawa K, Yambun P (2014) A new *Leptolalax* from mountains of Sabah, Borneo (Amphibia, Anura, Megophryidae). *Zootaxa* 3753: 440–452
- Ryan MJ (1980) Female mate choice in a Neotropical frog. *Science* 209: 523–525
- Savage JM (1997) Digital webbing formulae for anurans: a refinement. *Herpetol Rev* 28: 131
- Sukumaran J, Das I, Haas A (2010) Description of the advertisement calls of some Bornean frogs. *Russ J Herpetol* 17: 189–194

(Received June 11, 2014 / Accepted July 21, 2014)