

Title	STRUCTURE OF THE POSITIVE RADIAL SOLUTIONS FOR A SUPERCRITICAL NEUMANN PROBLEM IN A BALL (Progress in Qualitative Theory of Ordinary Differential Equations)
Author(s)	宮本, 安人
Citation	数理解析研究所講究録 (2014), 1901: 92-98
Issue Date	2014-06
URL	http://hdl.handle.net/2433/223022
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

STRUCTURE OF THE POSITIVE RADIAL SOLUTIONS FOR A SUPERCRITICAL NEUMANN PROBLEM IN A BALL

Yasuhito Miyamoto¹
Graduate School of Mathematical Sciences,
University of Tokyo

Let $\Omega \subset \mathbb{R}^N$, $N \geq 3$, be a bounded domain with smooth boundary. We study the positive solution of the Neumann problem

$$(1) \quad \begin{cases} \varepsilon^2 \Delta u - u + u^p = 0 & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\varepsilon > 0$ is a positive parameter. This problem arises in stationary problems of the shadow system of the Gierer-Meinhardt model and the Keller-Segel model with logarithmic sensitivity function. When the domain is the entire space \mathbb{R}^N , the problem (1) also appears in the study of the standing wave of the nonlinear Schrödinger equation. The problem (1) has attracted much attention for more than two decades. Solutions of various shapes have been found in [4, 6, 11, 12]. However, many authors study the case $1 < p < p_S$. Here,

$$p_S := \begin{cases} \frac{N+2}{N-2} & \text{if } N \geq 3, \\ \infty & \text{if } N = 1, 2. \end{cases}$$

When $p > p_S$, the Sobolev embedding $H^1(\Omega) \hookrightarrow L^{p+1}(\Omega)$ does not hold and it is difficult to use variational methods. There are few results about the structure of the positive solutions in the case $p \geq p_S$.

We consider the positive radial solutions when $p > p_S$ and $\Omega = B$. Then (1) can be reduced to the ODE

$$(2) \quad \begin{cases} u_{rr} + \frac{N-1}{r} u_r + \lambda f(u) = 0 & (0 < r < 1), \\ u_r(1) = 0, \\ u(r) > 0 & (0 \leq r \leq 1), \end{cases}$$

¹This work was partially supported by the Japan Society for the Promotion of Science, Grant-in-Aid for Young Scientists (B) (Subject No. 24740100) and by Keio Gijuku Academic Development Funds.




FIGURE 1. Schematic picture of the bifurcation diagram of (2) in the case $p_S < p < p_{JL}$.

where $f(u) = -u + u^p$ and $\lambda = 1/\varepsilon^2 > 0$. Adimurthi and Yadava [1, 2] studied the critical case $p = p_S$ when Ω is a unit ball B . They have shown that if $N \geq 7$, then (2) has a solution for all small $\lambda > 0$, while if $N \in \{4, 5, 6\}$, then (2) has no solution for small $\lambda > 0$. Del Pino et.al. [3] constructed a bubble tower solution when p is slightly greater than p^* .

We study the bifurcation diagram of the radial solutions of (1), using ODE techniques. In this study the existence of the singular solution of (2) plays an important role.

Theorem A. *Suppose that $p > p_S$. The problem (1) has infinitely many singular solutions $(\lambda_n^*, U_n^*(r)) \in \mathbb{R}_+ \times (C^2(0, 1) \cap C^0(0, 1] \cap H^1(B))$ ($n = 1, 2, \dots$ and $\lambda_1^* < \lambda_2^* < \dots \rightarrow \infty$) such that the following assertions hold:*

(i) $U_n^*(r)$ satisfies

$$(3) \quad U_n^*(r) = A(p, N)(\sqrt{\lambda_n^*}r)^{-\theta}(1 + o(1)) \quad \text{as } (r \downarrow 0),$$

where

$$(4) \quad A(p, N) := \{\theta(N - 2 - \theta)\}^{\frac{1}{p-1}}.$$

(ii) $\mathcal{Z}_{(0,1]}[U_n^*(\cdot) - 1] = n$.

(iii) $U_n^*(r) > 0$ ($0 < r \leq 1$).

Moreover, the singular solution (λ_n^*, U_n^*) is unique, i.e., if $(\tilde{\lambda}_n^*, \tilde{U}_n^*)$ is a singular solution such that (i) and (ii) hold, then $(\tilde{\lambda}_n^*, \tilde{U}_n^*) = (\lambda_n^*, U_n^*)$.

The main result is the following:

Theorem B. *Suppose that $p > p_S$. Let \mathcal{S} be the set of the regular solutions. Then*

$$\mathcal{S} = C_0 \cup \bigcup_{n=1}^{\infty} (C_n^+ \cup C_n^-),$$

where C_n^+ (resp. C_n^-) is the branch emanating from the trivial branch $\{(\lambda, 1)\}_{\lambda > 0}$, which we denote by C_0 , such that $u(0) > 1$ (resp. $u(0) < 1$). C_n^\pm is a C^1 -function of $\gamma := u(0)$, hence C_n^\pm can be described as $\{(\lambda_n(\gamma), u_n(r, \gamma))\}$. Moreover, the following hold:

- (i) $\lambda_n(1) = \bar{\lambda}_n$,
- (ii) $\lambda_n(\gamma) \rightarrow \lambda_n^*$ ($\gamma \rightarrow \infty$),
- (iii) if $p_S < p < p_{JL}$, then $\lambda_n(\gamma)$ oscillates infinitely many times around λ_n^* , where

$$p_{JL} := \begin{cases} 1 + \frac{4}{N - 4 - 2\sqrt{N-1}} & \text{if } N \geq 11, \\ \infty & \text{if } 2 \leq N \leq 10, \end{cases}$$

- (iv) $\lambda_n(\gamma) \rightarrow \infty$ ($\gamma \downarrow 0$),
- (v) if $\gamma > 0$ is small, then $u_1(r, \gamma)$ is non-degenerate in the space of radial functions and it concentrates on the boundary,
- (vi) $\lambda_1(\gamma) < \lambda_2(\gamma) < \dots$.

Figure 1 is a schematic picture of the bifurcation diagram of (2) in the case $p_S < p < p_{JL}$. When $p_S < p < p_{JL}$, (2) has infinitely many regular solutions for $\lambda = \lambda_n^*$. Each branch blows up at λ_n^* , while it is unbounded in the positive direction of λ in the subcritical case $1 < p < p_S$.

The following corollary is an immediate consequence of Theorem B.

Corollary C. *Suppose that $p > p_S$. There exist $\underline{\lambda} > 0$ and $\bar{\lambda} (> \underline{\lambda})$ such that a radially decreasing solution of (2), which belongs to C_1^+ , does not exist for $\lambda \in (0, \underline{\lambda}) \cup (\bar{\lambda}, \infty)$.*

The main tool of the proof is an intersection number between the singular solution and a regular solution. Using a scaling argument, one can show that each branch has infinitely many turning points if $p_S < p < p_{JL}$. See [8] for details of the proof. In the case $p \geq p_{JL}$ we do not know the number of the turning points of each branch.

Let us explain the strategy of the proof. Let $u(s) := U(r)$ and $s := \sqrt{\lambda}r$. The equation (1) is transformed to the problem

$$(5) \quad \begin{cases} u_{ss} + \frac{N-1}{s}u_s + f(u) = 0, & 0 < s < \sqrt{\lambda}, \\ u_s(\sqrt{\lambda}) = 0, \\ u > 0, \end{cases} \quad 0 \leq s \leq \sqrt{\lambda}.$$

First we construct the singular solution $u^*(s)$ of the equation in (5) near $s = 0$ and show that $u^*(s) = As^{-\theta}(1 + o(1))$ ($s \downarrow 0$). Here $A := A(p, N)$ and $A(p, N)$ is defined by (4). Second we show that the domain of $u^*(s)$ can be extended to $0 < s < \infty$, that $u^*(s)$ satisfies the equation in (5), and that $u^*(s) > 0$ for $s > 0$. Third we show that $u^*(s)$ oscillates around 1 infinitely many times as $s \rightarrow \infty$ and that $u^*(s)$ has the set of the critical points $\{s_n^*\}_{n=1}^\infty$ of u^* such that $0 < s_1^* < s_2^* < \dots \rightarrow \infty$ and

$$\begin{cases} s_n^* \text{ is a local minimum point of } u^* \text{ and } u^*(s_n^*) < 1 \text{ if } n \in \{1, 3, 5, \dots\}, \\ s_n^* \text{ is a local maximum point of } u^* \text{ and } u^*(s_n^*) > 1 \text{ if } n \in \{2, 4, 6, \dots\}. \end{cases}$$

We set $\lambda_n^* := (s_n^*)^2$ and $U_n^*(r) := u^*(s)$ ($s = \sqrt{\lambda_n^*}r$). Then, (λ_n^*, U_n^*) is a singular solution stated in Theorem A.

Let $(\lambda_n(\gamma), u(s, \gamma))$ denote the solution of (5) such that $u(0, \gamma) = \gamma$ and $u_s(0, \gamma) = 0$. We show that $\lambda_n(\gamma) \rightarrow \lambda_n^*$ as $\gamma \rightarrow \infty$ and that $u(s, \gamma)$ converges to $u^*(s)$ in an appropriate sense. In [7] Merle and Peletier proved a similar convergence result for the Dirichlet problem

$$\begin{cases} U_{rr} + \frac{N-1}{r}U_r + \lambda U + U^p = 0, & 0 < r < 1, \\ U(1) = 0, \\ U > 0, \end{cases} \quad 0 \leq r < 1.$$

when $p > p_S$. We show that $u(s, \gamma) \rightarrow u^*(s)$, following arguments in the proof of [7, Theorem A].

We show that $\lambda_n(\gamma)$ oscillates around λ_n^* if $p_S < p < p_{JL}$. Let $\rho := \gamma^{\frac{p-1}{2}}s$. We define $\tilde{u}(\rho, \gamma) := u(s, \gamma)/\gamma$ and $\tilde{u}^*(\rho) := u^*(s)/\gamma$. We use the intersection number between \tilde{u} and \tilde{u}^* . The function $\tilde{u}(\rho, \gamma)$ satisfies

$$(6) \quad \begin{cases} \tilde{u}_{\rho\rho} + \frac{N-1}{\rho}\tilde{u}_\rho + \tilde{u}^p - \frac{1}{\gamma^{p-1}}\tilde{u} = 0, & 0 < \rho < \infty, \\ \tilde{u}(0) = 1, \tilde{u}_\rho(0) = 0. \end{cases}$$

Let $\bar{u}(\rho, \gamma)$ be the regular solution of

$$(7) \quad \begin{cases} \bar{u}_{\rho\rho} + \frac{N-1}{\rho}\bar{u}_\rho + \bar{u}^p = 0, & 0 < \rho < \infty, \\ \bar{u}(0) = \gamma, \bar{u}_\rho(0) = 0. \end{cases}$$

We show that as $\gamma \rightarrow \infty$,

$$\tilde{u}(\rho, \gamma) \rightarrow \bar{u}(\rho, 1) \quad \text{in} \quad C_{loc}^2(0, \infty) \cap C_{loc}^0[0, \infty)$$

and

$$\tilde{u}^*(\rho) \rightarrow \bar{u}^*(\rho) \quad \text{in} \quad C_{loc}^0(0, \infty),$$

where $\bar{u}^*(\rho)$ a singular solution of the equation in (7). We recall the fact that $\mathcal{Z}_{(0, \infty)}[\bar{u}^*(\cdot) - \bar{u}(\cdot, 1)] = \infty$. Hence, for each $\delta > 0$,

$$(8) \quad \mathcal{Z}_{(0, \delta)}[u^*(\cdot) - u(\cdot, \gamma)] \rightarrow \infty \quad (\gamma \rightarrow \infty),$$

since $s \in (0, \delta)$ is corresponding to $\rho \in (0, \delta\gamma^{\frac{N-1}{2}})$ and $\delta\gamma^{\frac{N-1}{2}} \rightarrow \infty$ ($\gamma \rightarrow \infty$). Since each zero of $u^*(\cdot) - u(\cdot, \gamma)$ is simple, each zero depends continuously on γ . The divergence (8) tells us that a zero which is simple enters the interval $(0, \sqrt{\lambda_n^*}]$ from $s = \sqrt{\lambda_n^*}$ infinitely many times. Therefore, there exists a sequence of large numbers $\{\gamma_j\}_{j=1}^{\infty}$ ($\gamma_1 < \gamma_2 < \dots \rightarrow \infty$) such that $u^*(\sqrt{\lambda_n^*}) = u(\sqrt{\lambda_n^*}, \gamma_j)$ and the following holds: $u_s(\sqrt{\lambda_n^*}, \gamma_j) < 0$ for $j \in \{1, 3, 5, \dots\}$ and $u_s(\sqrt{\lambda_n^*}, \gamma_j) > 0$ for $j \in \{2, 4, 6, \dots\}$. Using the convergence $u(s, \gamma) \rightarrow u^*(s)$, we show that if $n \in \{1, 3, 5, \dots\}$ (resp. $n \in \{2, 4, 6, \dots\}$)

$$(9) \quad \lambda_n(\gamma_j) \begin{cases} > \lambda_n^*, & (j \in \{1, 3, 5, \dots\}), \\ < \lambda_n^*, & (j \in \{2, 4, 6, \dots\}), \end{cases}$$

$$\left(\text{resp. } \lambda_n(\gamma_j) \begin{cases} < \lambda_n^*, & (j \in \{1, 3, 5, \dots\}), \\ > \lambda_n^*, & (j \in \{2, 4, 6, \dots\}). \end{cases} \right)$$

which implies that $\lambda_n(\gamma)$ oscillates around λ_n^* infinitely many times as $\gamma \rightarrow \infty$.

This method can be applied to Dirichlet problems. In [9] the author studies

$$(10) \quad \begin{cases} \Delta u + \lambda f(u) = 0 & \text{in } B, \\ u > 0 & \text{in } B, \\ u = 0 & \text{on } \partial B, \end{cases}$$

where $f(u) = u^p + g(u)$, $p > p^*$, and $|g(u)| < Cu^{p-\varepsilon}$. We assume that $f \in C^1$ and $f(u) > 0$ for $u \geq 0$. If $3 \leq N \leq 10$, then the branch of the solutions of (10) has infinitely many turning points. It is shown that there is a nonlinear term $f(u)$ such that the branch has finitely many turning points.

In [10] he studies (10), where $f(u) = e^u + g(u)$ and $|g(u)| < Ce^{(1-\varepsilon)u}$. In [10] it is shown that if $3 \leq N \leq 9$, then the branch of the solutions

of (10) has infinitely many turning points around some $\lambda^* > 0$ and that if

$$(f1') \quad N \geq 10, \quad -e^u < g'(u) \leq \frac{N-10}{8}e^u \text{ in } (0, \infty),$$

$$\text{and } g''(u) > -e^u \text{ in } (0, \infty),$$

then the branch does not have a turning point and blows up at λ^* . In particular, the branch consists only of the minimal solutions. Thus, when (f1') is satisfied, then the bifurcation diagram is qualitatively the same as the case $f(u) = e^u$.

REFERENCES

- [1] Adimurthi and S. Yadava, *Existence and nonexistence of positive radial solutions of Neumann problems with critical Sobolev exponents*, Arch. Rational Mech. Anal. **115** (1991), 275–296.
- [2] Adimurthi and S. Yadava, *Nonexistence of positive radial solutions of a quasilinear Neumann problem with a critical Sobolev exponent*, Arch. Rational Mech. Anal. **139** (1997), 239–253.
- [3] M. del Pino, and M. Musso, and A. Pistoia, *Super-critical boundary bubbling in a semilinear Neumann problem*, Ann. Inst. H. Poincaré Anal. Non Linéaire **22** (2005), 45–82.
- [4] C. Gui and J. Wei, *On multiple mixed interior and boundary peak solutions for some singularly perturbed Neumann problems*, Canad. J. Math. **52** (2000), 522–538.
- [5] Z. Guo and J. Wei, *Global solution branch and Morse index estimates of a semilinear elliptic equation with super-critical exponent*, Trans. Amer. Math. Soc. **363** (2011), 4777–4799.
- [6] A. Malchiodi and M. Montenegro, *Boundary concentration phenomena for a singularly perturbed elliptic problem*, Comm. Pure Appl. Math. **55** (2002), 1507–1568.
- [7] F. Merle and L. Peletier, *Positive solutions of elliptic equations involving supercritical growth*, Proc. Roy. Soc. Edinburgh Sect. A **118** (1991), 49–62.
- [8] Y. Miyamoto, *Structure of the positive radial solutions for the supercritical Neumann problem $\varepsilon^2 \Delta u - u + u^p = 0$ in a ball*, UTMS Preprint Series 2013, <http://kyokan.ms.u-tokyo.ac.jp/users/preprint/pdf/2013-6.pdf>
- [9] Y. Miyamoto, *Structure of the positive solutions for supercritical elliptic equations in a ball*, to appear in J. Math. Pures Appl.
- [10] Y. Miyamoto, *Classification of bifurcation diagrams for elliptic equations with exponential growth in a ball*, to appear in Ann. Mat. Pura Appl.
- [11] J. Wei, *On the boundary spike layer solutions to a singularly perturbed Neumann problem*, J. Differential Equations **134** (1997), 104–133.
- [12] J. Wei, *On the interior spike layer solutions to a singularly perturbed Neumann problem*, Tohoku Math. J. **50** (1998), 159–178.

Graduate School of Mathematical Sciences
University of Tokyo
3-8-1 Komaba, Meguro-ku, Tokyo 153-8914
JAPAN
E-mail address: miyamoto@ms.u-tokyo.ac.jp

東京大学大学院数理科学研究科 宮本 安人