

Hida distribution construction of $P(\phi)_d$ ($d \geq 4$) indefinite metric quantum field models without BPHZ renormalization

Sergio ALBEVERIO * and Minoru W. YOSHIDA †

December 22, 2013

Abstract

By removing the divergent functions (i.e., the divergent Fynman graphs) a system of "Schwinger functions" which corresponds to a $(\Phi^4)_4$ Euclidean quantum field theory is constructed. The system of "Schwinger functions", which are defined through the Hida distributions, satisfies the property of OS 2) (Euclidean covariance), OS 4) (Symmetry) and OS 5) (Cluster property). It does not satisfy OS 3) (Reflection positivity). Then, for the system of "Schwinger functions" a possibility that it admits an analytic continuation to a system of "Wightman functions" satisfying the modified Wightman axioms is discussed.

1 Explanation of well known results on $d = 2$ by means of probabilistic words

Below, let $d = 4$ or $d = 2$ with an adequate understanding.

Let \dot{W} be the random variable such that $\dot{W}(\omega) \in \mathcal{S}'(\mathbb{R}^d \rightarrow \mathbb{R})$, $P - a.e. \omega \in \Omega$, and for each $\varphi \in \mathcal{S}(\mathbb{R}^d \rightarrow \mathbb{R})$, $\langle \dot{W}, \varphi \rangle_{\mathcal{S}', \mathcal{S}}$ is a real valued Gaussian random variable (*white noise on \mathbb{R}^d*) satisfying

$$E[\langle \dot{W}, \varphi \rangle_{\mathcal{S}', \mathcal{S}}] = 0, \quad (1)$$

$$E[\langle \dot{W}, \varphi_1 \rangle_{\mathcal{S}', \mathcal{S}} \cdot \langle \dot{W}, \varphi_2 \rangle_{\mathcal{S}', \mathcal{S}}] = \int_{\mathbb{R}^d} \varphi_1(\mathbf{x}) \varphi_2(\mathbf{x}) d\mathbf{x}, \quad \forall \varphi_1, \varphi_2 \in \mathcal{S}(\mathbb{R}^d \rightarrow \mathbb{R}). \quad (2)$$

Let $J_{d=2}^{\frac{1}{2}}$ be the integral kernel of the pseudo differential operator on $\mathcal{S}(\mathbb{R}^2)$ such that $(-\Delta + 1)^{-\frac{1}{2}}$ with $\Delta = \Delta_{d=2}$

the Laplace operator on \mathbb{R}^2 . For $\varphi, f_j \in \mathcal{S}(\mathbb{R}^2 \rightarrow \mathbb{R})$, $j = 1, \dots, n$, let

$$\phi(f_j) = \int_{\mathbb{R}^2} \left(\int_{\mathbb{R}^2} f_j(\mathbf{x}) J_{d=2}^{\frac{1}{2}}(\mathbf{x} - \mathbf{y}) d\mathbf{x} \right) \dot{W}(\mathbf{y}) d\mathbf{y}, \quad (3)$$

*Inst. Angewandte Mathematik, Universität Bonn

†Dept. Math. Tokyo City Univ.

and

$$\langle : \phi_{d=2}^4 : , \varphi \rangle \quad (4)$$

$$= \int_{(\mathbb{R}^2)^4} \left\{ \int_{\mathbb{R}^2} \varphi(\mathbf{x}) \prod_{i=1}^4 J_{d=2}^{\frac{1}{2}}(x - y_i) dx \right\} : \dot{W}(\mathbf{y}_1) \cdots \dot{W}(\mathbf{y}_4) : d\mathbf{y}_1 \cdots d\mathbf{y}_4. \quad (5)$$

Then,

$$e^{-\lambda \langle : \phi_{d=2}^4 : , 1_\Lambda \rangle} \in \cap_{p \geq 1} L^p(\Omega; P), \quad (6)$$

and the Schwinger function for $d = 2$

$$S_n(f_1, \dots, f_n) \equiv \frac{1}{Z(\lambda; \Lambda)} E \left[\phi(f_1) \cdots \phi(f_n) e^{-\lambda \langle : \phi_{d=2}^4 : , 1_\Lambda \rangle} \right], \quad (7)$$

is well defined.

2 Formulation for $d = 4$

By changing the 2-dimensional space time Gaussian white noise process by 4-dimensional space time ones in the above discussion, and if we apply the same considerations to $(\Phi^4)_4$ quantum field model, then all the terms in $(\phi(f_1) \cdots \phi(f_n) (\langle : \phi_{d=4}^4 : , 1_\Lambda \rangle)^k)$ will not have the right to be random variables.

Denote

$$\mathbf{x} \equiv (t, \vec{x}) \in \mathbb{R} \times \mathbb{R}^3 = \mathbb{R}^4, \quad \xi \equiv (\tau, \vec{\xi}) \in \mathbb{R} \times \mathbb{R}^3 = \mathbb{R}^4.$$

Let

$$\mathcal{F}[\varphi](\xi) = (2\pi)^{-2} \int_{\mathbb{R}^4} e^{-\sqrt{-1}\mathbf{x} \cdot \xi} \varphi(\mathbf{x}) d\mathbf{x}, \quad \mathcal{F}^{-1}[\varphi](\mathbf{x}) = (2\pi)^{-2} \int_{\mathbb{R}^4} e^{\sqrt{-1}\mathbf{x} \cdot \xi} \varphi(\xi) d\xi$$

For each $\epsilon > 0$, let $j_\epsilon^{\frac{1}{2}}(\xi)$, $j^{\frac{1}{2}}(\xi)$, $j_\epsilon(\xi)$, and $j(\xi)$, resp., be the symbol of the pseudo differential operators, resp., such that

$$\begin{aligned} j_\epsilon^{\frac{1}{2}}(\xi) &\equiv (|\xi|^2 + 1 + \epsilon(|\xi|^2 + 1)^2)^{-\frac{1}{2}}, & j^{\frac{1}{2}}(\xi) &\equiv (|\xi|^2 + 1)^{-\frac{1}{2}}, \\ j_\epsilon(\xi) &\equiv (|\xi|^2 + 1 + \epsilon(|\xi|^2 + 1)^2), & j(\xi) &\equiv (|\xi|^2 + 1). \end{aligned}$$

and define

$$\begin{aligned} (J_\epsilon^{\frac{1}{2}} \varphi)(\mathbf{x}) &= \mathcal{F}^{-1}(j_\epsilon^{\frac{1}{2}} \hat{\varphi})(\mathbf{x}), & (J^{\frac{1}{2}} \varphi)(\mathbf{x}) &= \mathcal{F}^{-1}(j^{\frac{1}{2}} \hat{\varphi})(\mathbf{x}), \\ (J_\epsilon \varphi)(\mathbf{x}) &= \mathcal{F}^{-1}(j_\epsilon \hat{\varphi})(\mathbf{x}), & (J \varphi)(\mathbf{x}) &= \mathcal{F}^{-1}(j \hat{\varphi})(\mathbf{x}). \end{aligned}$$

Symbolically

$$\begin{aligned} J_\epsilon^{\frac{1}{2}} &= (-\Delta_{d=4} + 1 + \epsilon(-\Delta_{d=4} + 1)^2)^{-\frac{1}{2}}, & J^{\frac{1}{2}} &= (-\Delta_{d=4} + 1)^{-\frac{1}{2}}, \\ J_\epsilon &= (-\Delta_{d=4} + 1 + \epsilon(-\Delta_{d=4} + 1)^2)^{-1}, & J &= (-\Delta_{d=4} + 1)^{-1}, \end{aligned}$$

where

$$\Delta_{d=4} \equiv \frac{\partial}{\partial t} + \frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z}, \quad \text{with } \mathbf{x} = (t, x, y, z).$$

Let

$$\langle \phi(\omega), \varphi \rangle_{\mathcal{S}', \mathcal{S}} = \int_{\mathbb{R}^4} \left(\int_{\mathbb{R}^4} \varphi(\mathbf{x}) J^{\frac{1}{2}}(\mathbf{x} - \mathbf{y}) d\mathbf{x} \right) \dot{W}(\mathbf{y}) d\mathbf{y}, \quad \text{for } \varphi \in \mathcal{S}'(\mathbb{R}^4 \rightarrow \mathbb{R}). \quad (8)$$

Also, define an $\mathcal{S}'(\mathbb{R}^4 \rightarrow \mathbb{R})$ -valued random variable $\phi_\epsilon(\omega)$ such that

$$\langle \phi_\epsilon(\omega), \varphi \rangle_{\mathcal{S}', \mathcal{S}} = \int_{\mathbb{R}^4} \left(\int_{\mathbb{R}^4} \varphi(\mathbf{x}) J_\epsilon^{\frac{1}{2}}(\mathbf{x} - \mathbf{y}) d\mathbf{x} \right) \dot{W}(\mathbf{y}) d\mathbf{y}, \quad \text{for } \varphi \in \mathcal{S}'(\mathbb{R}^4 \rightarrow \mathbb{R}), \quad (9)$$

and for $p \in \mathbb{N}$, define

$$\langle : \phi_\epsilon^p : , \varphi \rangle = \int_{(\mathbb{R}^4)^p} \left\{ \int_{\mathbb{R}^4} \varphi(\mathbf{x}) \prod_{i=1}^p J_\epsilon^{\frac{1}{2}}(\mathbf{x} - \mathbf{y}_i) d\mathbf{x} \right\} : \prod_{j=1}^p \dot{W}(\mathbf{y}_j) : \prod_{j=1}^p d\mathbf{y}_j, \quad (10)$$

We then see that (cf. [FelMagRivSé], [CRiv])

$$\langle : \phi_\epsilon^4 : , 1_\Lambda \rangle^k = \sum_{G' \in A'(k; \epsilon, \Lambda)} G' + \sum_{G \in A(k; \epsilon, \Lambda)} G, \quad (11)$$

where $A'(k; \epsilon, \Lambda)$ is a set of random variables such that for $G'(k; \epsilon, \Lambda) \in A'(k; \epsilon, \Lambda)$ there exists an $n \in \mathbb{N} \cup \{0\}$ (in fact $0 \leq n \leq 4k$) and

$$\lim_{\epsilon \downarrow 0, \Lambda \uparrow \mathbb{R}^4} E[(\phi(f_1) \cdots \phi(f_n) G'(k; \epsilon, \Lambda))] \text{ diverges, } \forall f_j \in \mathcal{S}(\mathbb{R}^4), j = 1, \dots, n$$

and $A(k; \epsilon, \Lambda)$ is a set of random variables such that for $G(k; \epsilon, \Lambda) \in A(k; \epsilon, \Lambda)$

$$\lim_{\epsilon \downarrow 0, \Lambda \uparrow \mathbb{R}^4} E[(\phi(f_1) \cdots \phi(f_k) G(k; \epsilon, \Lambda))] \text{ converges,}$$

$$\forall f_j \in \mathcal{S}(\mathbb{R}^4), j = 1, \dots, n, \quad \forall n \in \mathbb{N} \cup \{0\}, \quad (12)$$

By using (12) we can define a *Hida distribution*:

$$G = \lim_{\epsilon \downarrow 0, \Lambda \uparrow \mathbb{R}^4} G(k; \epsilon, \Lambda). \quad (13)$$

Definition 1 [Tensor Product]

Let F and G be the random variables in $\cap_{p \geq 1} L^p(\Omega; P)$ defined by the multiple stochastic integral with respect to \dot{W} such that

$$F = \int_{(\mathbb{R}^4)^n} f(\mathbf{x}_1, \dots, \mathbf{x}_n) : \prod_{k=1}^n \dot{W}(\mathbf{x}_k) : d\mathbf{x}_1 \cdots d\mathbf{x}_n, \quad (14)$$

$$G = \int_{(\mathbb{R}^4)^m} g(\mathbf{x}_1, \dots, \mathbf{x}_m) : \prod_{k=1}^m \dot{W}(\mathbf{x}_k) : d\mathbf{x}_1 \cdots d\mathbf{x}_n. \quad (15)$$

The *tensor product* $F \otimes G$ of F and G is defined by

$$F \otimes G = \int_{(\mathbb{R}^4)^{n+m}} f(\mathbf{x}_1, \dots, \mathbf{x}_n) g(\mathbf{x}_{n+1}, \dots, \mathbf{x}_{n+m}) : \prod_{k=1}^{n+m} \dot{W}(\mathbf{x}_k) : d\mathbf{x}_1 \cdots d\mathbf{x}_{n+m}. \quad (16)$$

As a usual notation by [GrotStreit], the notion expressed by $F \otimes G$ above is equivalent with the Wick product $F \diamond G$. ■

Definition 2 [simplified definition of Hida-distributions]

For $f^{(n)}(\mathbf{x}_1, \dots, \mathbf{x}_n) \in \mathcal{S}((\mathbb{R}^4)^n)$, $n \in \mathbb{N}$ ($f^{(0)}$ is understood as a constant), a random variable φ defined by

$$\varphi = \sum_{n=0}^{\infty} \int_{(\mathbb{R}^4)^n} f^{(n)}(\mathbf{x}_1, \dots, \mathbf{x}_n) : \prod_{k=1}^n \dot{W}(\mathbf{x}_k) : d\mathbf{x}_1 \cdots d\mathbf{x}_n,$$

is said to be in $(\mathcal{S})_r$ for $r \in \mathbb{N} \cup \{0\}$ if

$$\|\varphi\|_{L^2(\Omega, P), r}^2 \equiv \sum_{n=0}^{\infty} n! \left\| \left(\prod_{k=1}^n (-\Delta_{\mathbf{x}_k} + |\mathbf{x}_k|^2 + 1)^r \right) f^{(n)} \right\|_{L^2((\mathbb{R}^4)^n)}^2 < \infty, \quad (17)$$

where $\Delta_{\mathbf{x}_k} \equiv \frac{\partial^2}{\partial^2 t_k} + \frac{\partial^2}{\partial^2 x_k} + \frac{\partial^2}{\partial^2 y_k} + \frac{\partial^2}{\partial^2 z_k}$ for $\mathbf{x}_k = (t_k, x_k, y_k, z_k) \in \mathbb{R}^4$.
 $\|\varphi\|_{L^2(\Omega, P), r}^2$ is equivalent with

$$E \left[\left(\sum_{n=0}^{\infty} \int_{(\mathbb{R}^4)^n} \left(\prod_{k=1}^n (-\Delta_{\mathbf{x}_k} + |\mathbf{x}_k|^2 + 1)^r \right) f^{(n)}(\mathbf{x}_1, \dots, \mathbf{x}_n) : \prod_{k=1}^n \dot{W}(\mathbf{x}_k) : d\mathbf{x}_1 \cdots d\mathbf{x}_n \right)^2 \right].$$

We say that a sequence of random variables $\{G_\epsilon\}_{\epsilon>0}$ defines a Hida distribution $G \in (\mathcal{S})_{-r}$ if there exists a constant $K < \infty$ and

$$|E[G_\epsilon \cdot \varphi]| \leq K \|\varphi\|_{L^2(\Omega, P), r}, \quad \forall \varphi \in (\mathcal{S})_r, \quad (18)$$

and the limit exists for $\forall \varphi \in (\mathcal{S})_r$:

$$\lim_{\epsilon \downarrow 0} E[G_\epsilon \varphi] \quad (19)$$

The following continuous linear functional G on $(\mathcal{S})_r$ is called as a Hida distribution in $(\mathcal{S})_{-r}$:

$$\langle G, \varphi \rangle = \lim_{\epsilon \downarrow 0} E[G_\epsilon \varphi] \quad (20)$$

■

Remark 1 i) If the defining sequence $\{G_\epsilon\}_{\epsilon>0}$ of a *Hida distribution* G is composed by elements of n -times multiple stochastic integrals, then

$$E \left[G_\epsilon \cdot \left(\int_{(\mathbb{R}^4)^m} f(\mathbf{x}_1, \dots, \mathbf{x}_m) : \prod_{k=1}^m \dot{W}(\mathbf{x}_k) : d\mathbf{x}_1 \cdots d\mathbf{x}_m \right) \right] = 0, \quad (21)$$

for any $f \in \mathcal{S}(\mathbb{R}^m)$ with $m \neq n$.

ii) Since,

$$\begin{aligned} & \left\| \left(\prod_{k=1}^n (-\Delta_{\mathbf{x}_k} + |\mathbf{x}_k|^2 + 1)^r \right) f \right\|_{L^2((\mathbb{R}^4)^n)} \\ &= \left\{ \int_{(\mathbb{R}^4)^n} \left(\left(\prod_{k=1}^n (|\mathbf{x}_k|^2 + 1)^{-2} (|\mathbf{x}_k|^2 + 1)^2 (-\Delta_{\mathbf{x}_k} + |\mathbf{x}_k|^2 + 1)^r \right) f \right)^2 d\mathbf{x}_1 \cdots d\mathbf{x}_n \right\}^{\frac{1}{2}} \\ &\leq \left(\sup_{\mathbf{x}_1, \dots, \mathbf{x}_n} \left| \left(\prod_{k=1}^n (|\mathbf{x}_k|^2 + 1)^2 (-\Delta_{\mathbf{x}_k} + |\mathbf{x}_k|^2 + 1)^r \right) f \right| \right) \cdot \left\| \prod_{k=1}^n (|\mathbf{x}_k|^2 + 1)^{-2} \right\|_{L^2((\mathbb{R}^4)^n)} \\ &\leq K \cdot p_{m,k}(f), \quad \forall f(\mathbf{x}_1, \dots, \mathbf{x}_n) \in \mathcal{S}((\mathbb{R}^4)^n), \end{aligned} \quad (22)$$

where $p_{m,k}(\cdot)$ is a semi-norm of $\mathcal{S}((\mathbb{R}^4)^n)$ such that

$$p_{m,k}(f) = \sum_{|\alpha| \leq m} \sup_{\bar{\mathbf{x}}} (1 + |\bar{\mathbf{x}}|^2)^k |D^\alpha f(\bar{\mathbf{x}})|, \quad (23)$$

with

$$\begin{aligned} \bar{\mathbf{x}} &= (\mathbf{x}_1, \dots, \mathbf{x}_n) \in (\mathbb{R}^4)^n, \quad \mathbf{x}_k = (t_k, x_k, y_k, z_k) \in \mathbb{R}^4, \\ \alpha &= (\alpha_1, \dots, \alpha_n), \quad \alpha_k = (\alpha_{k1}, \alpha_{k2}, \alpha_{k3}, \alpha_{k4}), \quad |\alpha| = \sum_{k=1}^n \sum_{i=1}^4 \alpha_{ki}, \\ D^\alpha &= \prod_{k=1}^n \left(\frac{\partial}{\partial t_k} \right)^{\alpha_{k1}} \left(\frac{\partial}{\partial x_k} \right)^{\alpha_{k2}} \left(\frac{\partial}{\partial y_k} \right)^{\alpha_{k3}} \left(\frac{\partial}{\partial z_k} \right)^{\alpha_{k4}}. \end{aligned}$$

By this, if a *Hida distribution* G is defined through a sequence $\{G_\epsilon\}_{\epsilon>0}$ of n -times multiple stochastic integrals, then it can be identified with an element of $\mathcal{S}'((\mathbb{R}^4)^n)$:

$\exists K < \infty$ and $\exists m, k \in \mathbb{N}$ that depend only on r and

$$\begin{aligned} |< G, \varphi >| &\leq K \cdot p_{m,k}(f) \\ \mathcal{S}'((\mathbb{R}^4)^n) \ni G : \mathcal{S}((\mathbb{R}^4)^n) \ni f &\longmapsto < G, \varphi >, \end{aligned} \quad (24)$$

for

$$\varphi = \int_{(\mathbb{R}^4)^n} f(\mathbf{x}_1, \dots, \mathbf{x}_n) : \prod_{k=1}^n \dot{W}(\mathbf{x}_k) : d\mathbf{x}_1 \cdots d\mathbf{x}_n.$$

■

3 Well defined terms for $d = 4$ and the strategy of the consideration

Through the discussions in the previous sections, we can define

$$e^{-\lambda \langle : \phi_\epsilon^4 : , 1_\Lambda \rangle} = \sum_{k=0}^{\infty} \frac{(-\lambda)^k}{(k!)} \left(\langle : \phi_\epsilon^4 : , 1_\Lambda \rangle \right)^k. \quad (25)$$

The equality holds for $P - a.e.$ $\omega \in \Omega$, because both side of the equality are real valued random-variables, regardless they are integrable or not. The number of terms (graphs) of $(\langle : \phi_\epsilon^4 : , 1_\Lambda \rangle)^k$, is estimated by $(k!)^2$:

$$(\langle : \phi_\epsilon^4 : , 1_\Lambda \rangle)^k = \sum_{G \in \tilde{A}(k; \epsilon, \Lambda)} G, \quad (26)$$

where, each $G \in \tilde{A}(k; \epsilon, \Lambda)$ is a *tensor product* of multiple stochastic integrals, and in the sense of Fynman graph it is a graph with k -verticies, also the cardinality of the set $\tilde{A}(k; \epsilon, \Lambda)$ is the order $(k!)^2$. By the notation of (11), $\tilde{A}(k; \epsilon, \Lambda)$ can be expressed by the direct sum

$$\tilde{A}(k; \epsilon, \Lambda) = A'(k; \epsilon, \Lambda) \oplus A(k; \epsilon, \Lambda). \quad (27)$$

For $r \geq 0$, let

$$\Lambda_r \equiv \left\{ (t, x, y, z) \in \mathbb{R}^4 \mid \sqrt{t^2 + x^2 + y^2 + z^2} \leq r \right\}. \quad (28)$$

For each $k \in \mathbb{N} \cup \{0\}$, define $A_r(\epsilon, k)$, a subset of $A(k; \epsilon, \Lambda_r) \subset \tilde{A}(k; \epsilon, \Lambda_r)$ in (28), as follows:

$$\begin{aligned} A_r(\epsilon, k) &= \text{the set with the elements } G \in \tilde{A}(k; \epsilon, \Lambda_r) \text{ (cf. (40)), each of which is} \\ &\quad \text{a tensor product of stochastic integrals, and each component of} \\ &\quad \text{the tensor product is identified with a connected Fynman graph} \\ &\quad \text{who is in type O or type I.} \end{aligned} \quad (29)$$

Where, *type O* resp., and *type I* is defined as follows:

Type O is the set of graphs, each element of which satisfies the following:

- (O-i) each vertex of the graph has at least 1 free (open) leg,
- (O-ii) there exists at least 1 vertex which has more than 2 free legs,
- (O-iii) there exists a connected path passing through every vertex of the graph exactly once,
- (O-iv) in the graph there is no sub graph such that two vertices connect two or three legs each other.

Type I is the set of graphs, each element of which satisfies the following:

- (I) between each vertex and the other vertex in the graph there exists exactly only one (directed) path that connects these two vericies, and each vertex of the graph has at least 1 free (open) leg.

We would like to show that the limit

$$S_n(\phi(f_1), \dots, \phi(f_n)) \equiv \lim_{r \rightarrow \infty} \left\{ \lim_{\epsilon \downarrow 0} E \left[\phi(f_1) \cdots \phi(f_n) \left(\sum_{k=0}^{\infty} \frac{(-\lambda)^k}{k!} \left(\sum_{G \in A_r(\epsilon, k)} G \right) \right) \right] \right\}, \quad (30)$$

for $f_j \in \mathcal{S}(\mathbb{R}^4 \rightarrow \mathbb{R})$, $j = 1, \dots, n$, $n \in \mathbb{N} \cup \{0\}$, defines a system of "Schwinger functions" $\{S_n\}_{n \in \mathbb{N} \cup \{0\}}$ in a "modified sense". Because of (0.21), (O-i), (O-ii) and (I), since the sum is precisely a finite sum, and the expectation is clearly well-defined:

$$\begin{aligned} & E \left[\phi(f_1) \cdots \phi(f_n) \left(\sum_{k=0}^{\infty} \frac{(-\lambda)^k}{k!} \left(\sum_{G \in A_r(\epsilon, k)} G \right) \right) \right] \\ &= E \left[\phi(f_1) \cdots \phi(f_n) \left(\sum_{k=0}^n \frac{(-\lambda)^k}{k!} \left(\sum_{G \in A_r(\epsilon, k)} G \right) \right) \right], \end{aligned} \quad (31)$$

and the limit of (30) can be taken within the framework of finite sum of *Hida distributions* (cf. Definition 2).

In fact the limit exists and we have Theorem's 4.1 and 4.2 below. To give the statements we prepare some notions and notations:

Denote the number of the elements of $A_r(\epsilon, k)$ by $N(A(k))$, and give an index to each $G \in A_r(\epsilon, k)$ to indicate it as $G_j(k; \epsilon, r)$, $j = 1, \dots, N(A(k))$. Then,

$$A_r(\epsilon, k) = \left\{ G_j(k; \epsilon, r) \right\}_{j=1, \dots, N(A(k))}, \quad k \in \mathbb{N}. \quad (32)$$

Recall that for $f_j \in \mathcal{S}(\mathbb{R}^4)$, $j = 1, 2, \dots$,

$$\begin{aligned} \phi(f_1)\phi(f_2) &= : \phi(f_1)\phi(f_2) : + E[\phi(f_1)\phi(f_2)] \\ \phi(f_1)\phi(f_2)\phi(f_3)\phi(f_4) &= : \phi(f_1)\phi(f_2)\phi(f_3)\phi(f_4) : \\ &+ (: \phi(f_1)\phi(f_2) :) E[\phi(f_3)\phi(f_4)] + (: \phi(f_1)\phi(f_3) :) E[\phi(f_2)\phi(f_4)] \\ &+ (: \phi(f_1)\phi(f_4) :) E[\phi(f_2)\phi(f_3)] + (: \phi(f_2)\phi(f_3) :) E[\phi(f_1)\phi(f_4)] \\ &+ (: \phi(f_2)\phi(f_4) :) E[\phi(f_1)\phi(f_3)] + (: \phi(f_3)\phi(f_4) :) E[\phi(f_1)\phi(f_2)] \\ &+ \sum_{\text{distinguished } i_l \text{'s}} E[\phi(f_{i_1})\phi(f_{i_2})] E[\phi(f_{i_3})\phi(f_{i_4})], \end{aligned}$$

and

$$: \phi(f_1) \cdots \phi(f_n) := \int_{(\mathbb{R}^4)^n} \prod_{j=1}^n (J^{\frac{1}{2}} f_j)(\mathbf{x}_j) : \prod_{j=1}^n \dot{W}(\mathbf{x}_j) : d\mathbf{x}_1 \cdots d\mathbf{x}_n. \quad (33)$$

By using the above, through a simple evaluation we have the following:

Lemma 3.1 For each $r \geq 0$, $\epsilon > 0$ and $k \geq 1$, let $A_r(\epsilon, k)$ be the set of Fynman graphs (i.e., set of multiple stochastic integrals) defined by (29). Then, by using the expression (32), the following hold:

$$N(A(k)) \simeq (k!)^{\frac{3}{2}}, \quad (34)$$

and by denoting the number of free legs of $G_j(k; \epsilon, r) \in A_r(\epsilon, k)$ by $N_f(G_j(k; \epsilon, r))$,

$$k + 2 \leq N_f(G_j(k; \epsilon, r)) \leq 4k, \quad j = 1, \dots, N(A(k)), \quad (35)$$

also (cf. (33))

$$E [(: \phi(f_1) \cdots \phi(f_n) :) G_j(k; \epsilon, r)] = 0, \quad \text{if } n \neq N_f(G_j(k; \epsilon, r)), \quad (36)$$

$$E [(\phi(f_1) \cdots \phi(f_n)) G_j(k; \epsilon, r)] = 0, \quad \text{if } n - 2 < k, \quad (37)$$

moreover, there exists $M < \infty$ and

$$\left| E [(: \phi(f_1) \cdots \phi(f_n) :) G_j(k; \epsilon, r)] \right| \leq M^k (n!) \prod_{i=1}^n \|\hat{f}_i\|_{L^1(\mathbb{R}^4)}, \quad (38)$$

for $\forall f_i \in \mathcal{S}(\mathbb{R}^4 \rightarrow \mathbb{R})$, $i = 1, \dots, n$, $\forall n \in \mathbb{N}$; $\forall G_j(k; \epsilon, r) \in A_r(\epsilon, k)$; $\forall k \in \mathbb{N}$, $\forall \epsilon > 0$, and $\forall r > 0$ (if $n \neq N_f(G_j(k; \epsilon, r))$, then "right hand side of (33)" = 0).

□

From Lemma 4.1, we immediately have

Theorem 3.1 For each $k \in \mathbb{N}$ and $j = 1, \dots, N(A(k))$, there exists a Hida distribution $G_j(k)$, which is a Fynman graph, such that

$$\lim_{r \rightarrow \infty} \lim_{\epsilon \downarrow 0} G_j(k; \epsilon, r) = G_j(k) \quad \text{with} \quad N_f(G_j(k)) = N_f(G_j(k; \epsilon, r)) \quad (39)$$

and the set $A_r(\epsilon, k)$ converges to a set of Hida distributions $A(k)$,

$$\lim_{r \rightarrow \infty} \lim_{\epsilon \downarrow 0} A_r(\epsilon, k) = A(k) = \{G_j(k)\}_{j=1, \dots, N(A(k))}. \quad (40)$$

Denote

$$\langle (: \phi(f_1) \cdots \phi(f_n) :), G_j(k) \rangle \equiv \lim_{r \rightarrow \infty} \lim_{\epsilon \downarrow 0} E [(: \phi(f_1) \cdots \phi(f_n) :) G_j(k; \epsilon, r)], \quad (41)$$

then there exists $M < \infty$ and

$$\left| \langle (: \phi(f_1) \cdots \phi(f_n) :), G_j(k) \rangle \right| \leq M^k (n!) \prod_{i=1}^n \|(-\Delta_{d=4} + 1)^2 f_i\|_{L^2(\mathbb{R}^4)}, \quad (42)$$

$$\left| \langle (\phi(f_1) \cdots \phi(f_n)), G_j(k) \rangle \right| \leq M^k (n!) \prod_{i=1}^n p_{4,2}(f_i), \quad (43)$$

for $\forall f_i \in \mathcal{S}(\mathbb{R}^4 \rightarrow \mathbb{R})$, $i = 1, \dots, n$, $\forall n \in \mathbb{N}$; $\forall G_j(k) \in A(k)$; $\forall k \in \mathbb{N}$
(if $n \neq N_f(G_j(k))$, then "right hand side of (42) and (43)" = 0), where $p_{m,k}(f)$ is the semi-norm defined by (23).

□

By the above theorem, we can set the following Definition.

Definition 3 Let $A(0) = \{1\}$, namely, $G_1(0) = 1$ and $N(A(0)) = 1$. For each $\lambda \in \mathbb{Z}$ let $\{S_n^\lambda\}_{n \in \mathbb{N} \cup \{0\}}$, be a system of Schwartz distributions such that

$$S_0^\lambda = S_1^\lambda = 0, \quad (44)$$

$$\begin{aligned} \langle S_n^\lambda, f_1 \otimes \cdots \otimes f_n \rangle &= \sum_{k=0}^{\infty} \frac{(-\lambda)^k}{k!} \left(\sum_{j=1}^{N(A(k))} \langle (\phi(f_1) \cdots \phi(f_n)), G_j(k) \rangle \right) \\ &= \sum_{k=0}^{n-2} \frac{(-\lambda)^k}{k!} \left(\sum_{j=1}^{N(A(k))} \langle (\phi(f_1) \cdots \phi(f_n)), G_j(k) \rangle \right) \\ \text{for } f_i &\in \mathcal{S}(\mathbb{R}^4), i = 1, \dots, n, n \in \mathbb{N}. \end{aligned} \quad (45)$$

□

For the statements of the next theorem, we recall the OS (Osterwalder-Schrader) axioms (cf., e.g., [Si1]):

It is said that a set of functions $\{S_n\}$ is a *system of Schwinger functions* with *Osterwalder-Schrader axioms OS'1-5*, if it satisfies

OS 1') (Temperedness + Analytic continuity)

$$S_n^p \in \mathcal{S}'_\neq(\mathbf{R}^{dn}), \quad \overline{S_n^p(f)} = S_n^p(\theta f^*),$$

and S_n^p is a Laplace transform of some $M_n \in \mathcal{S}'((\mathbf{R}_+^d)^{n-1})$, precisely

$$\begin{aligned} S_n^p(x_1^0, \vec{x}_1; \dots; x_n^0, \vec{x}_n) &= \int_{(\mathbf{R}_+^d)^{n-1}} M(\tau_1, \xi_1; \dots; \tau_{n-1}, \xi_{n-1}) \\ &\times \exp \left\{ \sum_{j=1}^{n-1} (\sqrt{-1} \xi_j \cdot (\vec{x}_{j+1} - \vec{x}_j) - \tau_j (x_{j+1}^0 - x_j^0)) \right\} d\xi_1 d\tau_1 \cdots d\xi_{n-1} d\tau_{n-1}; \end{aligned} \quad (46)$$

OS 2) (Euclidean covariance);

OS 3) (Reflection positivity);

- OS 4) (Symmetry);
- OS 5) (Cluster property).

Theorem 3.2 Let $\{S_n^\lambda\}_{n \in \mathbb{N} \cup \{0\}}$ be the system of Schwartz distributions defined by Definition 3, then it is a system of "modified Schwinger functions" in the sense that it satisfies OS 2), OS 4) and OS 5).

□

Remark 2.

We have an affirmative rigorous result on the analyticity OS 1') for $\{S_n^\lambda\}_{n \in \mathbb{N} \cup \{0\}}$. It will be announced in forthcoming papers. ■

References

- [AFeY] Albeverio, S., Ferrario, B., Yoshida, M.W., On the essential self-adjointness of Wick powers of relativistic fields and of fields unitary equivalent to random fields, *Acta Applicandae Mathematicae* **80** 309-334 (2004).
- [AGW1] Albeverio, S., Gottschalk, H., Wu, J.-L., Convolved generalized white noise, Schwinger functions and their analytic continuation to Wightman functions, *Rev. Math. Phys.* **8** (1996) 763-817.
- [AGW2] Albeverio, S., Gottschalk, H., Wu, J.-L., Models of local relativistic quantum fields with indefinite metric (in all dimensions), *Comm. Math. Phys.* **184** (1997), 509-531.
- [AH-K] Albeverio, S., Høegh-Krohn, R., Uniqueness and the global Markov property for Euclidean fields: The case of trigonometric interactions, *Comm. Math. Phys.* **68** (1979), 95-128.
- [AR] Albeverio, S., Röckner, M., Classical Dirichlet forms on topological vector spaces-closability and a Cameron-Martin formula, *J. Functional Analysis* **88** (1990) 395-43
- [AY1] Albeverio, S., Yoshida, M. W., $H - C^1$ maps and elliptic SPDEs with polynomial and exponential perturbations of Nelson's Euclidean free field, *J. Functional Analysis* **196** (2002) 265-322.
- [AY2] Albeverio, S., Yoshida, M. W., Hida distribution construction of non-Gaussian reflection positive generalized random fields, *Infinite Dimensional Analysis, Quantum Probability and Related Topics.* **12** (2009) 21-49.
- [Araki] Araki, H., On a pathology in indefinite inner product spaces, *Commn. Math. Phys.* **85** (1982), 121-128.

- [BaSeZh] Baez, J.C., Segal, I.E., Zhou, Z., *Introduction to Algebraic and Constructive Quantum Field Theory*, Princeton Univ. Press 1992
- [BogPara] Bogoliubow, N. N., Parasiuk, O. S., *Über die Multiplikation der Kausalfunktionen in der Quantentheorie der Felder*, **97** (1957), 227-266.
- [BogLogT] Bogoliubov, N. N., Logunov, A. A., Todorov, I. T., *Introduction to axiomatic quantum field theory*, Translated from the Russian by Stephen A. Fulling and Ludmila G. Popova. Edited by Stephen A. Fulling. Mathematical Physics Monograph Series, No. 18. W. A. Benjamin, Inc., Reading, Mass.-London-Amsterdam, 1975.
- [Bo] Borchers, H.-J., *Algebraic aspects of Wightman field theory*, in R.N. Sen and C. Weil (eds.), Statistical mechanics and Field Theory, Haifa Lectures 1971; New York: Halstedt Press, 1972
- [BrFSok] Brydges, D. C., Fröhlich, J., Sokal, A. D., A new proof of the existence and nontriviality of the continuum φ_2^4 and φ_3^4 quantum field theory, *Commn. Math. Phys.* **91** (1983) 141-186.
- [CRiv] de Calan, C., Rivasseau, V., Local existence of the Borel transform in Euclidean Φ_4^4 , *Commn. Math. Phys.* **82** (1981) 69-100.
- [DaFelRiv] David, F., Feldman, J., Rivasseau, V., On larde order bahavior of Φ_4^4 , *Commn. Math. Phys.* **116** (1988) 215-233.
- [DüR] Dütsch, M., Rehren, K.-H., A comment on the dual field in the AdS-CFT correspondence, *Lett. Math. Phys.* 62 (2002), no. 2, 171-184
- [Ep] Epstein, H., On the Borchers class of a free field, *Nuovo Cimento* **27** 1963, 886-893.
- [Fel] Feldman, J., The $\lambda\varphi_3^4$ field theory in a finite volume, *Commn. Math. Phys.* **37** (1974) 93-120.
- [FelMagRivSé] Feldman, J., Magnen, J., Rivasseau, V., Séneor, R., Bound on completely convergent Euclidean Fynman graphs, *Commn. Math. Phys.* **98** (1985) 273-288.
- [FernFSok] Fernandez, R., Fröhlich, J., Sokal, A. D., *Random walks, critical phenomena, and triviality in quantum field theory*, Texts and Monographs in Physics. Springer-Verlag, Berlin, 1992.
- [GRSi] Guerra, F., Rosen, L., Simon, B., The $P(\phi)_2$ Euclidean quantum field theory as classical statistical mechanics. I, II, *Ann. of Math.* **101** (1975) 111-189
- [GJ] Glimm, J., Jaffe, A., *Quantum Physics: A Functional Integral Point of View*, 2nd ed., Springer, Berlin, 1987.
- [GrotStreit] Grothaus, M., Streit, L., Construction of relativistic quantum fields in the framework of white noise analysis, *J. Math. Phys.* **40** (1999) 5387-5405.
- [Heg] Hegerfeldt, G. C., From Euclidean to relativistic fields and on the notion of Markoff fields, *Comm. Math. Phys.* **35** (1974) 155-171.

- [Hepp] Hepp, K., Proof of the Bogoliubov-Parasiuk theorem on renormalization, *Comm. Math. Phys.* **2** (1966) 301-326.
- [H1] Hida, T., Generalized multiple Wiener integrals, *Proc. Japan Acad. Ser. A Math. Sci.* **54** (1978) 55-58.
- [H2] Hida, T., *Brownian motion*, Springer-Verlag, New York Heidelberg Berlin 1980.
- [HKPStreit] Hida, T., Kuo, H.-K., Potthoff, J., Streit, L., *White Noise: An Infinite Dimensional Calculus*, Kluwer Academic Publishers, Dordrecht, 1993.
- [HStreit] Hida, T., Streit, L., On quantum theory in terms of white noise, *Nagoya Math. J.* **68** (1977) 21-34.
- [Ho] Hofman, G., On GNS representations on inner product space, *Commun. Math. Phys.* **191**, 299-323 (1998)
- [IkW] Ikeda, N., Watanabe, S., *Stochastic differential equations and diffusion processes*, second edition, North-Holland, 1989.
- [ItoKR] Ito, K. R., *Publ. RIMS Kyoto Univ.* **14**, 503 (1978)
- [Jo] Jost, R., *The general theory of quantized fields*, Ed. Mark Kac, Lectures in Applied Mathematics (Proceedings of the Summer Seminar, Boulder, Colorado, 1960), Vol. IV American Mathematical Society, Providence, R.I. 1965 xv+157 pp.
- [Klein1] Klein, A., Renormalized products of the generalized free field & its derivatives, *Pac. J. Math.* **45** (1973) 275-292.
- [Klein2] Klein, A., Gaussian OS-positive processes, *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **40** (1977) 115-124.
- [MagNicRivSé] Magnen, J., Nicolo, F., Rivasseau, V., Séneor, R., A Lipatov bound for Φ_4^4 Euclidean field theory, *Comm. Math. Phys.* **108** (1987) 257-289.
- [Mizohata] Mizohata, S.: *The theory of partial differential equations*. Cambridge University Press, New York, 1973.
- [MoStro] Morchio, G., Strocchi, F., Infrared singularities, vacuum structure and pure phase in local quantum field theory, *Ann. Inst. H. Poincaré* **A33** (1980) 251-282.
- [NaMu1] Nagamachi, S., Mugibayashi, N., Hyperfunction quantum field theory, *Comm. Math. Phys.* **46** (1976) 119-134.
- [NaMu2] Nagamachi, S., Mugibayashi, N., Hyperfunction quantum field theory. II. Euclidean Green's functions, *Comm. Math. Phys.* **49** (1976) 257-275.
- [NaMu3] Nagamachi, S., Mugibayashi, N., Hyperfunctions and renormalization, *J. Math. Phys.* **27** (1986) 832-839.

- [Ne1] Nelson, E., Construction of quantum fields from Markoff fields, *J. Functional Analysis* **12** (1973) 97–112.
- [Ne2] Nelson, E., The free Markov field, *J. Functional Analysis* **12** (1973) 221-227.
- [Nu] Nualart, D., *The Malliavin calculus and related topics*, Springer-Verlag, New York/Heidelberg/Berlin, 1995.
- [Oj1] Ojima, I., Entropy production and nonequilibrium stationarity in quantum dynamical systems, Physical meaning of van Hove limit., *J. Statist. Phys.* **56** (1989) 203-226.
- [Oj2] Ojima, I.: *How to formulate non-equilibrium local states in QFT? General characterization and extension to curved spacetime.* A garden of quanta, 365-384, World Sci. Publishing, River Edge, NJ, 2003.
- [OS1] Osterwalder,K.,Schrader, R., Axioms for Euclidean Green's functions I, *Comm. Math. Phys.* **31** (1973) 83-112.
- [OS2] Osterwalder,K.,Schrader, R., Axioms for Euclidean Green's functions II, *Comm. Math. Phys.* **42** (1975) 281-305.
- [ReSi] Reed, M., Simon, B., *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness*, Academic Press, 1975.
- [Rehren] Rehren, K-H., Comments on a Recent Solution to Wightman's Axioms, *Comm. Math. Phys.* **178** (1996) 453-465.
- [Si1] Simon, B., *The $P(\Phi)_2$ Euclidean (Quantum) Field Theory*, Princeton Univ. Press, Princeton, NJ., 1974.
- [Si2] Simon, B., Borel summability of the ground-state energy in spacially cutoff $(\varphi^4)_2$, *Phys. Rev. letters* **25** (1970) 1583-1586.
- [SmSol1] Smirnov, A.G., Soloviev, M.A., Spectral properties of Wick power series of a free field with an indefinite metric, *Theoret. and Math. Phys.* **125** (2000) 1349-1362.
- [SmSol2] Smirnov, A.G., Soloviev, M.A., Wick power series that converge to nonlocal fields, *Theoret. and Math. Phys.* **127** (2001) 632-645.
- [Sok] Sokal, A. D., An improvement of Watson's theorem on Borel summability, *J. Math. Phys.* **21** (1980) 261-263.
- [Speer] Speer, E. R., On the structure of analytic renormalization, *Comm. Math. Phys.* **23** (1971) 23-36.
- [StWi] Streater R.F., Wightman A.S., *PCT, Spin and Statistics, and all that*, Princeton Univ. Press 1964

- [Stro] Strocchi F., *Selected topics on the General Properties of Quantum Field Theory*, Lect. Notes in Physics, **51** World Sci., Singapore-New York-London-Hong Kong, 1993.
- [Widder] Widder, D. V., *The Laplace transform*, Princeton University Press, Princeton, Eighth printing 1972.
- [Wi1] Wightman, A.S., *Introduction to new aspects of relativistic dynamics of quantum fields*, Cargèse Lect. Theor. Phys, Ed. M. Lévy, 171-291, Gordon and Breach New York (1967)
- [Wi2] Wightman, A.S., *Recent achievements of axiomatic field theory*, Sem. on Theor. Phys. Trieste 1962 Int. At. En. Ag., 11-58 Vienna (1963)
- [Y1] Yoshida, M.W., Construction of infinite-dimensional interacting diffusion processes through Dirichlet forms, *Probab. Theory Relat. Fields* **106** (1996) 265-297.
- [Y2] Yoshida, M.W., Non-linear continuous maps on abstract Wiener spaces defined on space of tempered distributions, *Bulletin of the Univ. Electro-Commun.* **12** (1999) 101-117.