

# On reflection numbers under large continuum

神戸大学大学院・システム情報学研究科 梶野 昌

Sakaé Fuchino

Graduate School of System Informatics, Kobe University  
Rokko-dai 1-1, Nada, Kobe 657-8501 Japan  
fuchino@diamond.kobe-u.ac.jp

## Abstract

We give a survey of the results on the reflection numbers which are connected with Rado's Conjecture, Galvin's Conjecture, countable chromatic number of graphs, Hamburger's Problem on metrizability or freeness of Boolean algebras. In particular we examine some models of set theory in which the continuum is very large while some of these numbers are less than or equal to the continuum.

## 1 Reflection numbers

Let  $\mathcal{C}$  be a class of structures of some given type or signature. We also allow here classes of structures of types of higher-order; several classes of topological spaces, for example, are also considered under such  $\mathcal{C}$ 's. We often identify  $A \in \mathcal{C}$  with its underlying set. We assume that  $\mathcal{C}$  is provided with a notion of substructures  $\sqsubseteq_{\mathcal{C}}$  with appropriate properties like transitivity. For  $A$  and  $\kappa \in \text{Card} \setminus \omega + 1$ , let

---

*Date:* February 1, 2016    *Last update:* February 15, 2016 (15:45 JST)

*2010 Mathematical Subject Classification:* 03E35, 03E55, 03E65

*Keywords:* large continuum, strongly compact cardinal, reflection principles, Cohen reals

The research is supported by Grant-in-Aid for Exploratory Research No. 26610040 of the Ministry of Education, Culture, Sports, Science and Technology Japan (MEXT). A part of this work was done during the programme: Mathematical, Foundational and Computational Aspects of the Higher Infinite at the Isaac Newton Institute for Mathematical Sciences, Cambridge. The author would like to thank the Isaac Newton Institute for support and hospitality.

An extended version of this paper with more details and proofs is downloadable as:

<http://fuchino.ddo.jp/papers/RIMS15-large-continuum-x.pdf>

$$(1.1) \quad S_{<\kappa}^{\mathcal{C}}(A) = \{B \in \mathcal{C} : B \sqsubseteq_{\mathcal{C}} A, |B| < \kappa\}.$$

We also assume that  $S_{<\kappa}^{\mathcal{C}}(A)$  contains a club in  $[A]^{<\kappa}$  for all  $A \in \mathcal{C}$  and uncountable  $\kappa \leq |A|$ .

For such a class of structures  $\mathcal{C}$  and a property  $P$ , the reflection number  $\mathfrak{Rfl}(\mathcal{C}, P)$  of the property  $P$  in  $\mathcal{C}$  is defined by

$$(1.2) \quad \mathfrak{Rfl}(\mathcal{C}, P) = \begin{cases} \min\{\kappa \in \text{Card} : \text{for all } A \in \mathcal{C} \text{ if } A \not\models P \text{ then there are} \\ \text{club many } B \in S_{<\kappa}^{\mathcal{C}}(A) \text{ with } B \not\models P\}, \\ \text{if } \{\kappa \in \text{Card} : \dots\} \neq \emptyset; \\ \infty, & \text{otherwise.} \end{cases}$$

If  $P$  is hereditary, that is, if  $A \models P$  for  $A \in \mathcal{C}$  and  $B \sqsubseteq_{\mathcal{C}} A$  always imply  $B \models P$ , the reflection number  $\mathfrak{Rfl}(\mathcal{C}, P)$  can be represented more simply by

$$(1.3) \quad \mathfrak{Rfl}(\mathcal{C}, P) = \begin{cases} \min\{\kappa \in \text{Card} : \text{for all } A \in \mathcal{C} \text{ if } A \not\models P \text{ then there is} \\ B \in S_{<\kappa}^{\mathcal{C}}(A) \text{ with } B \not\models P\}, \\ \text{if } \{\kappa \in \text{Card} : \dots\} \neq \emptyset; \\ \infty, & \text{otherwise.} \end{cases}$$

Among the instances (A)  $\sim$  (E) of reflection numbers  $\mathfrak{Rfl}(\mathcal{C}, P)$  we are going to introduce in the next section, all of the properties  $P$  considered there except that in (E) are hereditary.

## 2 Some examples of reflection numbers

We are going to consider the following instances (A)  $\sim$  (E) of  $\mathcal{C}$  and  $\mathcal{P}$  in the general setting of the reflection number  $\mathfrak{Rfl}(\mathcal{C}, P)$  defined in the last section.

(A)  $\mathcal{C} = \text{trees}$ ;  $P = \text{“special”}$ .

Recall that a tree  $T$  is said to be special if  $T$  can be represented as a union of countably many antichains (i.e. pairwise incomparable sets).

The statement  $\mathfrak{Rfl}(\mathcal{C}, P) = \aleph_2$  for these  $\mathcal{C}$  and  $P$  is known as the *Rado Conjecture* and studied extensively, e.g. in [15], [17], [18], [4], [12]. We shall call this reflection number *the reflection number of the Rado Conjecture* and denote it by  $\mathfrak{Rfl}_{\text{Rado}}$ .

We have  $\aleph_1 < \mathfrak{Rfl}_{\text{Rado}} \leq \infty$ . The first inequality is trivial. The equation  $\mathfrak{Rfl}_{\text{Rado}} = \infty$  is possible: This holds for example if we have  $\square_{\kappa}$  for class many  $\kappa \in \text{Card}$  — actually  $\text{ADS}^-(\kappa)$  for class many regular uncountable  $\kappa$  is enough (see [12]).

The Rado Conjecture ( $\mathfrak{Rfl}_{\text{Rado}} = \aleph_2$ ) is known to be consistent: for example, it can be forced when we start from a model of ZFC with a strongly compact cardinal  $\kappa$  and collapse it to  $\aleph_2$  by  $\text{Col}(\aleph_1, \kappa)$  (Todorćević [15], see also Section 3 below).

The Rado Conjecture is known to imply many interesting mathematical consequences:

- $2^{\aleph_0} \leq \aleph_2$  (Todorčević [17]).
- strong forms of Chang’s Conjecture (Todorčević [17], Doebler [4], Fuchino, Sakai, Torres and Usuba [12]).
- $\mathfrak{Rfl}_{\text{Rado}} = \aleph_2$  implies the Fodor-type Reflection Principle (FRP) (Fuchino, Sakai, Torres and Usuba [12]) and hence all consequences of FRP like SCH (Fuchino and Rinot [10] — a direct proof of SCH from the Rado Conjecture was given in Todorčević [17]), stationarity reflection (of sets of ordinals of countable cofinality) etc.
- Semistationary Reflection Principle (SSR) of Sakai [14] (Doebler [4]).

(B)  $\mathcal{C}$  = partial orderings;  $P$  = “countable union of chains”.

The statement  $\mathfrak{Rfl}(\mathcal{C}, P) = \aleph_2$  for these  $\mathcal{C}$  and  $P$  is known as the *Galvin Conjecture*. It is a longstanding open problem if the Galvin Conjecture is consistent with ZFC. We shall call  $\mathfrak{Rfl}(\mathcal{C}, P)$  for these  $\mathcal{C}$  and  $P$  *the reflection number of the Galvin Conjecture* and denote it with  $\mathfrak{Rfl}_{\text{Galvin}}$ .

(C)  $\mathcal{C}$  = graphs;  $P$  = “of countable chromatic number”.

Recall that a graph  $\Gamma$  is of countable chromatic number if and only if it is a union of countably many subsets  $\Gamma_n$ ,  $n \in \omega$  of  $\Gamma$  such that each of  $\Gamma_n$ ’s contains no adjacent pairs. We shall call  $\mathfrak{Rfl}(\mathcal{C}, P)$  for these  $\mathcal{C}$  and  $P$  *the reflection number of countable chromatic number* and denote it with  $\mathfrak{Rfl}_{\text{chr}}$ . In spite of the similar definition to that of  $\mathfrak{Rfl}_{\text{Rado}}$ , it is known that  $\mathfrak{Rfl}_{\text{chr}}$  can never be  $\aleph_2$ :

**Theorem 2.1** (Erdős and Hajnal [5])  $\beth_\omega \leq \mathfrak{Rfl}_{\text{chr}}$ . □

Erdős and Hajnal originally proved this theorem under GCH. A modification of the proof without relying on GCH can be found e.g. in [11].

**Corollary 2.2** *Assuming the consistency of ZFC + “there are at least two strongly compact cardinals”, the inequality  $\mathfrak{Rfl}_{\text{Rado}} < \mathfrak{Rfl}_{\text{chr}} < \infty$  is consistent.*

**Proof.** The model obtained by collapsing the first strongly compact cardinal  $\kappa$  by  $\text{Col}(\aleph_1, \kappa)$  will do. The inequality  $\mathfrak{Rfl}_{\text{chr}} < \infty$  follows from Lemma 2.3 below.

□ (Corollary 2.2)

**Lemma 2.3** (Todorčević [18])

$\mathfrak{Rfl}_{\text{Rado}} \leq \mathfrak{Rfl}_{\text{Galvin}} \leq \mathfrak{Rfl}_{\text{chr}} \leq$  *the  $\omega_1$ -strongly compact cardinal (if it exists).*

**Proof.**  $\mathfrak{Refl}_{\text{Rado}} \leq \mathfrak{Refl}_{\text{Galvin}}$ : Suppose that  $T$  is a tree such that every subtrees of  $T$  of cardinality  $< \kappa = \mathfrak{Refl}_{\text{Galvin}}$  are special. We have to show that  $T$  is special.

Let  $<_w$  be a well-ordering on  $T$  and let  $<_w^T$  be the relation on  $T$  defined by

$$(2.1) \quad t <_w^T t' \Leftrightarrow t \text{ and } t' \text{ are incomparable in } T \text{ and } t \upharpoonright \alpha^* <_w t' \upharpoonright \alpha^* \text{ where} \\ \alpha^* = \min\{\alpha \in \text{On} : t \upharpoonright \alpha \text{ and } t' \upharpoonright \alpha \text{ are incomparable}\}.$$

It is easy to see that  $<_w^T$  is a partial ordering on  $T$ .

For any  $T' \subseteq T$ ,

$$(2.2) \quad T' \text{ is an antichain if and only if } T' \text{ is linearly ordered with respect to } <_w^T.$$

It follows that any  $T' \in [T]^{<\kappa}$  is the union of countably many linearly ordered sets with respect to  $<_w^T$ . Since  $\kappa = \mathfrak{Refl}_{\text{Galvin}}$ , it follows that  $T$  itself is the union of countably many linearly ordered sets with respect to  $\leq_w^T$ . By (2.2), this means that  $T$  is special.

$\mathfrak{Refl}_{\text{Galvin}} \leq \mathfrak{Refl}_{\text{chr}}$ : Suppose that  $P = \langle P, \leq_P \rangle$  is a partial ordering such that every  $Q \in [P]^{<\kappa}$  for  $\kappa = \mathfrak{Refl}_{\text{chr}}$  are the union of countably many linearly ordered sets with respect to  $\leq_P$ .

We have to show that  $P$  itself is also the union of countably many linearly ordered subsets with respect to  $\leq_P$ .

For  $p, q \in P$ , let

$$(2.3) \quad p E q \Leftrightarrow p \text{ and } q \text{ are incomparable with respect to } \leq_P.$$

Then, for any  $Q \subseteq P$ ,

$$(2.4) \quad Q \text{ is linearly ordered with respect to } \leq_P \Leftrightarrow Q \text{ contains no adjacent pair} \\ \text{with respect to } E.$$

Thus, by the assumption on  $P$ , we have that all subsets  $Q$  of  $P$  of cardinality  $< \kappa$  are countable chromatic with respect to  $E$ . By  $\kappa = \mathfrak{Refl}_{\text{chr}}$  it follows that  $(\langle P, E \rangle)$  is also countable chromatic. The latter means by (2.4) that  $P$  is the union of countably many linearly ordered subsets with respect to  $\leq_P$ .

$\mathfrak{Refl}_{\text{chr}} \leq$  the  $\omega_1$ -strongly compact cardinal: Recall that a cardinal  $\kappa$  is the  $\omega_1$ -strongly compact cardinal if and only if it is the smallest cardinal  $\kappa$  such that, for any  $\mathcal{L}_{\omega_1, \omega}$ -theory  $T$ , if all  $T' \in [T]^{<\kappa}$  are satisfiable (that is, they have models) then  $T$  itself is also satisfiable.

Suppose that  $\kappa$  is  $\omega_1$ -strongly compact and let  $\Gamma = \langle \Gamma, R_\Gamma \rangle$  be a graph such that all  $\Gamma' \in [\Gamma]^{<\kappa}$  are countable chromatic.

It is then easy to see that all subsets  $T'$  of cardinality  $< \kappa$  of the following set  $T$  of formulas in  $L_{\omega_1, \omega}$  are satisfiable:

$$\begin{aligned}
T = & \{c_g R_\Gamma c_{g'} : g, g' \in \Gamma, g R_\Gamma g'\} \\
& \cup \{\neg c_g R_\Gamma c_{g'} : g, g' \in \Gamma, g \not R_\Gamma g'\} \\
& \cup \{\text{“}\Gamma_n(\cdot) \text{ is pairwise non adjacent with respect to } R_\Gamma\text{”} : n \in \omega\} \\
& \cup \{\forall x \left( \bigwedge_{n \in \omega} \Gamma_n(x) \right)\}.
\end{aligned}$$

By  $\omega_1$ -strong compactness of  $\kappa$ , it follows that  $T$  has a model  $M$ .

Clearly

$$(2.5) \quad \{\{g \in \Gamma : M \models \Gamma_n(c_g) : n \in \omega\}$$

is a partition of  $\Gamma$  into countably many pairwise non adjacent subsets with respect to  $R_\Gamma$ . □ (Lemma 2.3)

Let us continue with some other examples of reflection numbers:

(D)  $\mathcal{C}$  = first countable topological spaces;  $P$  = “metrizable”.

The first countability is added to avoid some trivial examples of non reflection:

**Example 2.4** (Hajnal and Juhász [13]) *For an uncountable regular cardinal  $\kappa$ , let  $X = \kappa + 1$  be the topological space whose open sets are generated by*

$$(2.6) \quad \{\{\alpha\} : \alpha < \kappa\} \cup \{[\alpha, \kappa + 1) : \alpha < \kappa\}.$$

*All  $Y \in [X]^{<\kappa}$  are discrete and hence metrizable but  $X$  is not since  $\chi(\kappa, X) = \kappa$ . □*

The question about the consistence of  $\mathfrak{Rfl}(\mathcal{C}, P) = \aleph_2$  for these  $\mathcal{C}$  and  $P$  is called *Hamburger’s Problem* and is still open.

We shall call this reflection number *the reflection number of Hamburger’s Problem* and denote it with  $\mathfrak{Rfl}_{\text{HP}}$ .

**Lemma 2.5** (1)  $\aleph_1 < \mathfrak{Rfl}_{\text{HP}} \leq \infty$ .

(2)  $\mathfrak{Rfl}_{\text{HP}} = \infty$  is consistent.

(3)  $\mathfrak{Rfl}_{\text{HP}} \leq$  the  $\omega_1$ -strongly compact cardinal (if it exists).

**Proof.** (1): Clear.

(2): This holds if  $\square_\kappa$  holds for cofinally many  $\kappa$  — actually  $\text{ADS}^-(\kappa)$  for class many regular uncountable  $\kappa$  is enough (see Proposition 6.3 in [7]).

(3): Suppose that  $(X, \mathcal{O})$  is a first countable topological space such that all subspaces  $Y \in [X]^{<\kappa}$  are metrizable. For each  $x \in X$ , let  $\{O_{x,n} : n \in \omega\}$  be an open neighborhood base of  $x$ .

Let  $T$  be the  $\mathcal{L}_{\omega_1, \omega}$  theory in the language with the binary relation symbols  $O_n(x, y)$  for all  $n \in \omega$  coding “ $y \in O_{x,n}$ ” and the binary symbols  $d_q(x, y)$  for all  $q \in \mathbb{Q}_{\geq 0}$  which should code “ $d(x, y) \leq q$ ”:

$$\begin{aligned}
(2.7) \quad T = & \{O_n(c_a, c_b) : a, b \in X, b \in O_{a,n}\} \\
& \cup \{\neg O_n(c_a, c_b) : a, b \in X, b \notin O_{a,n}\} \\
& \cup \{\forall x \forall y (d_q(x, y) \rightarrow d_q(y, x)) : q \in \mathbb{Q}_{\geq 0}\} \\
& \cup \{\forall x \forall y (d_q(x, y) \rightarrow d_{q'}(x, y)) : q, q' \in \mathbb{Q}_{\geq 0}, q \leq q'\} \\
& \cup \{\forall x \forall y (d_0(x, y) \rightarrow x \equiv y)\} \\
& \cup \{\forall x \forall y \forall z (d_q(x, y) \wedge d_{q'}(y, z) \rightarrow d_{q+q'}(x, z)) : q, q' \in \mathbb{Q}_{\geq 0}\} \\
& \cup \{\forall x \bigwedge_{q \in \mathbb{Q}_{> 0}} \forall y (d_q(x, y) \rightarrow O_n(x, y)) : n \in \omega\} \\
& \cup \{\forall x \bigwedge_{q \in \mathbb{Q}_{> 0}} \bigwedge_{n \in \omega} \forall y (O_n(x, y) \rightarrow d_q(x, y))\}
\end{aligned}$$

Clearly all  $T' \in [T]^{<\kappa}$  are satisfiable. Since  $\kappa$  is  $\omega_1$ -strongly compact, it follows that  $T$  is also satisfiable. Let  $M$  be a model of  $T$ . Then  $d : X^2 \rightarrow \mathbb{R}$  defined by

$$(2.8) \quad d(a, b) = \min\{q \in \mathbb{Q} : M \models d_q(c_a, c_b)\} \text{ for } a, b \in X$$

is a metric on  $X$  generating the topology of  $(X, \mathcal{O})$ . □ (Lemma 2.5)

(E)  $\mathcal{C}$  = atomless Boolean algebras;  $P$  = “free”.

We shall denote the corresponding reflection number by  $\mathfrak{Rfl}_{\text{fBa}}$ .

**Lemma 2.6** (1)  $\aleph_1 < \mathfrak{Rfl}_{\text{fBa}} \leq \infty$ .

(2)  $\mathfrak{Rfl}_{\text{fBa}} = \infty$  is consistent.

(3)  $\mathfrak{Rfl}_{\text{fBa}} \leq$  the first supercompact cardinal (if it exists).

**Proof.** (1):  $\aleph_1 < \mathfrak{Rfl}_{\text{fBa}}$  since all countable atomless Boolean algebras are free.

(2): By Corollary 2.5. in [8].

(3): Suppose that  $\kappa$  is a supercompact cardinal and  $B$  is an atomless Boolean algebra such that there is a club  $\mathcal{D} \subseteq S_{<\kappa}^{\mathcal{C}}(B)$  such that all  $A \in \mathcal{D}$  are free where  $\mathcal{C}$  denotes the class of all atomless Boolean algebras. Without loss of generality, we may assume that  $|B| \geq \kappa$ . Let  $j : V \xrightarrow{\kappa} M$  be  $|B|$ -supercompact elementary embedding. Thus we have  $j(\kappa) \geq |B|$  and  ${}^{|B|}M \subseteq M$ . Then  $A^* = j''B \in j(\mathcal{D})$  and  $A^* \in M$ . By the elementarity of  $j$  it follows that  $M \models A^*$  is free. Thus  $A \cong A^*$  is really free. □ (Lemma 2.6)

**Problem 2.7** Does “ $\mathfrak{Rfl}_{\text{fBa}} \leq$  the  $\omega_1$ -strongly compact cardinal” hold?

### 3 A consistency proof of the Rado Conjecture

Todorćević proved the following proposition under the existence of a supercompact cardinal. Almost the same proof will do under a strongly compact cardinal.

**Proposition 3.1** (Todorćević [15]) *Suppose that  $\kappa$  is strongly compact and  $\mathbb{P} = \text{Col}(\aleph_1, \kappa)$ . Then we have*

$$(3.1) \quad \Vdash_{\mathbb{P}} \text{“}\mathfrak{Rfl}_{\text{Rado}} = \aleph_2\text{”}.$$

For the proof of Proposition 3.1, we use the following lemma also due to Todorćević:

**Lemma 3.2** *Suppose that  $T$  is a non-special tree and  $\mathbb{P}$  a  $\sigma$ -closed p.o. Then  $\Vdash_{\mathbb{P}}$  “ $T$  is a non-special tree”.*

**Proof.** We prove the contraposition of the statement. Suppose that, for a tree  $T$  and a  $\sigma$ -closed p.o.  $\mathbb{P}$ , we have  $\Vdash_{\mathbb{P}}$  “ $T$  is a special tree”. Note that it follows that

$$(3.2) \quad ht(T) \leq \omega_1.$$

Let  $\underset{\sim}{f}$  be a  $\mathbb{P}$ -name of a witness of the specialness of  $T$ , that is,

$$(3.3) \quad \Vdash_{\mathbb{P}} \text{“}\underset{\sim}{f} : T \rightarrow \omega \text{ and} \\ \underset{\sim}{f}(t) \neq \underset{\sim}{f}(t') \text{ for all } t, t' \in T \text{ such that } t \text{ and } t' \text{ are comparable”}.$$

For  $t \in T$  we can define  $p_t \in \mathbb{P}$  and  $n_t \in \omega$  by induction on  $ht(t)$  such that

$$(3.4) \quad p_t \text{ decides } \underset{\sim}{f}(t) \text{ to be } n_t; \text{ and}$$

$$(3.5) \quad p_t \leq_{\mathbb{P}} p_{t'} \text{ if } t \leq_T t'.$$

Note that this is possible by the  $\sigma$ -closedness of  $\mathbb{P}$  and (3.2).

The mapping  $f^* : T \rightarrow \omega$  defined by

$$(3.6) \quad f^*(t) = n_t \text{ for } t \in T$$

witnesses the specialness of  $T$  (in the ground model). □ (Lemma 3.2)

**Proof of Proposition 3.1:** Let  $\mathbb{P} = \text{Col}(\aleph_1, \kappa)$  and let  $G$  be a  $(V, \mathbb{P})$ -generic filter. Note that

$$(3.7) \quad V[G] \models \kappa = \omega_2.$$

Suppose that  $T = \langle T, \leq_T \rangle \in V[G]$  is a tree of size  $\lambda > \aleph_1$  in  $V[G]$  (so  $\lambda$  is a cardinal in  $V$  with  $\lambda \geq \kappa$ ) such that

$$(3.8) \quad V[G] \models \text{“}T \text{ is non-special”}.$$

By (1.3), it is enough to show that, in  $V[G]$ , there is a non-special  $T' \in [T]^{\aleph_1}$ .

Without loss of generality, we may assume that (the underlying set of)  $T$  is the set of ordinals  $\lambda$ .

Let  $j : V \xrightarrow{\leq} M$  be a  $\lambda$ -strongly compact embedding. That is,  $j$  is an elementary embedding such that

$$(3.9) \quad \kappa = \text{crit}(j);$$

(3.10)  $\kappa M \subseteq M$ ; and

(3.11) for all  $X \in [M]^{\leq \lambda}$  (in  $V$ ) there is  $Y \in ([M]^{< j(\kappa)})^M$  with  $X \subseteq Y$ .

Let  $\mathbb{P}^* = j(\mathbb{P})$ . Then, by (3.10), we have  $\mathbb{P}^* = \text{Col}(\aleph_1, j(\kappa))$  (in  $V$ ). Let  $G^* \supseteq G$  be a  $(V, \mathbb{P}^*)$ -generic filter.  $j$  can be extended to  $j^* : V[G] \xrightarrow{\cong} M[G^*]$  by declaring  $j^*(\dot{a}^G) = (j(\dot{a}))^{G^*}$  for all  $\mathbb{P}$ -name  $\dot{a}$ .

Since  $|j''\lambda| = \lambda$  (in  $V$ ), there is  $Y \in ([M]^{< j(\kappa)})^M$  such that  $j''\lambda \subseteq Y \subseteq j(\lambda)$  by (3.11). Let  $T^* = \langle Y, j^*(\leq_T) \cap Y^2 \rangle$ . Then  $T^* \in M[G^*]$  and, by (3.7),

(3.12)  $M[G^*] \models$  “ $T^*$  is a subtree of  $j(T)$  of cardinality  $< \aleph_2$ ”.

In  $V[G^*]$ ,  $T$  is embeddable in  $T^*$  and, by Lemma 3.2, we have  $V[G^*] \models$  “ $T$  is non-special”. It follows that  $M[G^*] \models$  “ $T^*$  is non-special”.

Thus

(3.13)  $M[G^*] \models$  “there is a non-special  $T' \in [j^*(T)]^{< \aleph_2}$ ”.

By the elementarity of  $j^*$ , it follows that

(3.14)  $V[G] \models$  “there is a non-special  $T' \in [T]^{< \aleph_2}$ ”.

□ (Proposition 3.1)

## 4 Models with large continuum

The continuum can be consistently very large in many different ways. For example it can be weakly inaccessible (provided that we work under the consistency of ZFC + “there exists an inaccessible cardinal”).

More extreme situations would be:

(4.1) There is an inner model  $M$  with  $\text{Card}^M \cap 2^\omega = \text{Card} \cap 2^\omega +$  and  $2^\omega$  is a fairly large cardinal (e.g. strongly compact, supercompact etc.) in  $M$ ; or

(4.2)  $2^{\aleph_0}$  is a generic large cardinal, that is, there are generic elementary embeddings with the critical point  $2^{\aleph_0}$ .

We can attain the situations like in (4.1) or (4.2) e.g. by starting from a large cardinal, say a strongly compact  $\kappa$ , and adding  $\kappa$  many reals in a coherent manner.

Reflection numbers can be small under large continuum in such sense. The following result is an example of this:

**Theorem 4.1** (Dow, Tall and Weiss [2]) *Suppose that  $\kappa$  is a strongly compact cardinal and  $\mathbb{P} = \text{Fn}(\kappa, 2)$ . Then  $\Vdash_{\mathbb{P}}$  “ $\mathfrak{Refl}_{\text{HP}} \leq 2^{\aleph_0}$ ”.* □



In the following we show that  $\mathfrak{Rfl}_{\text{Rado}}$  behaves similarly in the generic extension of Theorem 4.1.

**Theorem 4.2** *Suppose that  $\kappa$  is a strongly compact cardinal. Then, for any  $\mu \geq \kappa$  and  $\mathbb{P} = \text{Fn}(\mu, 2)$ , we have*

$$(4.3) \quad \Vdash_{\mathbb{P}} \text{“} \mathfrak{Rfl}_{\text{Rado}} \leq 2^{\aleph_0} \text{”}.$$

*In particular, by setting  $\mu > \kappa$ , we obtain the consistency of  $\mathfrak{Rfl}_{\text{Rado}} < 2^{\aleph_0}$  (modulo a strongly compact cardinal).*

Note that, in contrast, Rado’s Conjecture (i.e.  $\mathfrak{Rfl}_{\text{Rado}} = \aleph_2$ ) implies that  $2^{\aleph_0} \leq \aleph_2 = \mathfrak{Rfl}_{\text{Rado}}$  (Todorćević [17]).

Mateo Viale asked if  $\mathfrak{Rfl}_{\text{Rado}} = \aleph_3$  implies  $2^{\aleph_0} \leq \aleph_3$ . This is still open. More generally:

**Problem 4.3** *Is it provable in ZFC that  $\mathfrak{Rfl}_{\text{Rado}} = \aleph_n$  implies  $2^{\aleph_0} \leq \aleph_n$  for all  $n \geq 3$ ?*

For the proof of Theorem 4.2, we need the following lemma. Remember that a p.o.  $\mathbb{P}$  is said to be  $\sigma$ -centered if it can be represented as the union of countably many centered subsets (i.e. filter bases).

**Lemma 4.4** (1) *Suppose that  $T$  is a non-special tree. If  $\mathbb{P}$  is a  $\sigma$ -centered p.o., then  $\Vdash_{\mathbb{P}} \text{“} T \text{ is not special”}$ .*

(2) *Suppose that  $T$  is a non-special tree. Then, for any  $\mu$  and  $\mathbb{P} = \text{Fn}(\mu, 2)$ , we have  $\Vdash_{\mathbb{P}} \text{“} T \text{ is not special”}$ .*

**Proof.** (1): Suppose that  $T$  is a non-special tree and  $\mathbb{P} = \bigcup_{n \in \omega} C_n$  where each  $C_n$  is centered.

If  $\not\Vdash_{\mathbb{P}} \text{“} T \text{ is not special”}$  then there are  $p \in \mathbb{P}$  and  $\mathbb{P}$ -names  $\underset{\sim}{A}_m$ ,  $m \in \omega$  such that

$$(4.4) \quad p \Vdash_{\mathbb{P}} \text{“} T \text{ is a union of pairwise incomparable } \underset{\sim}{A}_m, m \in \omega \text{”}.$$

For each  $m, n \in \omega$

$$(4.5) \quad A_{m,n} = \{t \in T : q \Vdash_{\mathbb{P}} \text{“} t \in \underset{\sim}{A}_m \text{” for some } q \in C_n \text{ such that } q \leq_{\mathbb{P}} p\}$$

is a pairwise incomparable subset of  $T$  and  $T = \bigcup \{A_{m,n} : m, n \in \omega\}$ . This is a contradiction to the assumption that  $T$  is not special.

(2): Note that  $\mathbb{P}$  is  $\sigma$ -centered if and only if  $\mu \leq 2^{\aleph_0}$ . Thus, if  $\mu \leq 2^{\aleph_0}$ , then the claim follows from (1).

Suppose that  $\mu > 2^{\aleph_0}$  and  $\not\Vdash_{\mathbb{P}} \text{“} T \text{ is not special”}$ . By the homogeneity of  $\mathbb{P}$  it follows that

$$(4.6) \quad \Vdash_{\mathbb{P}} \text{“} T \text{ is special”}.$$

Let  $\mathbb{Q} = \text{Col}(\aleph_1, \kappa^+)$ . By (4.6) we have  $\Vdash_{\mathbb{P} * \mathbb{Q}} \text{“} T \text{ is special”}$  where  $\mathbb{Q}$  is a  $\mathbb{P}$ -name of a p.o. such that  $\mathbb{P} * \mathbb{Q} \sim \mathbb{Q} * \mathbb{P}$ . Thus we have

$$(4.7) \quad \Vdash_{\mathbb{Q} * \mathbb{P}} \text{“} T \text{ is special”}.$$

Since  $\Vdash_{\mathbb{Q}} \text{“} \mathbb{P} \text{ is } \sigma\text{-centered”}$ , it follows that  $\Vdash_{\mathbb{Q}} \text{“} T \text{ is special”}$  by (1).

By Lemma 3.2, it follows that  $T$  is special. □ (Lemma 4.4)

**Proof of Theorem 4.2:** Suppose that  $G$  is a  $(V, \mathbb{P})$ -generic filter and  $T = \langle T, <_T \rangle$  a non-special tree in  $V[G]$ . Let  $V[G] \models \lambda = |T|$  and let  $j : V \rightarrow M$  be an elementary embedding such that  $M \subseteq V$  is a transitive class  $\subseteq V$ ,  $\text{crit}(j) = \kappa$ ,  $j(\kappa) > \lambda$  and

$$(4.8) \quad \text{for all } X \in [M]^{\leq \lambda} \text{ (in } V \text{) there is } Y \in ([M]^{< j(\kappa)})^M \text{ with } X \subseteq Y.$$

Let  $\mathbb{P}^* = j(\mathbb{P}) = \text{Fn}(j(\mu), 2)$  and let  $G^*$  be a  $(\mathbb{P}^*, V)$ -generic filter with  $G^* \supseteq G$ .

Let  $j^* : V[G] \xrightarrow{\sim} V[G^*]$  be the extension of  $j$  defined by  $j^*([a]^G) = [j(a)]^{G^*}$  for a  $\mathbb{P}$ -name  $a$ . By the ccc of  $\mathbb{P}^*$  and (4.8), we have

$$(4.9) \quad \text{for all } X \in [M[G^*]]^{\leq \lambda} \text{ (in } V[G^*] \text{) there is } Y \in ([M[G^*]]^{< j(\kappa)})^{M[G^*]} \text{ with } X \subseteq Y.$$

Thus, in  $M[G^*]$ , there is a subtree  $T'$  of  $j^*(T)$  of size  $< j(\kappa)$  containing  $j^*T$ . Since  $V[G^*] \models j^*T \cong T$ , and  $V[G^*] \models \text{“} T \text{ is not special”}$  by Lemma 4.4, (2), we have  $V[G^*] \models \text{“} j^*T \text{ is not special”}$ . Since  $V[G^*] \models j^*T \subseteq T'$ , it follows that  $V[G^*] \models \text{“} T' \text{ is not special”}$  and hence  $M[G^*] \models \text{“} T' \text{ is not special”}$ . Thus

$$(4.10) \quad M[G^*] \models \text{“there is a non-special subtree } T' \text{ of } j^*(T) \text{ of size } < j^*(\kappa)\text{”}.$$

By elementarity of  $j^*$ , it follows that

$$(4.11) \quad V[G] \models \text{“there is a non-special subtree } T' \text{ of } T \text{ of size } < \kappa\text{”}.$$

Since  $\kappa \leq \mu \leq (2^{\aleph_0})^{V[G]}$ , we have

$$(4.12) \quad V[G] \models \text{“there is a non-special subtree } T' \text{ of } T \text{ of size } < 2^{\aleph_0}\text{”}.$$

□ (Theorem 4.2)

In analogy to the common terminology in set-theoretic topology, we say a structure  $A$  is indestructibly  $\neg P$  for a property  $P$ , if  $\Vdash_{\mathbb{P}} \text{“} A \models \neg P \text{”}$  holds for any  $\sigma$ -closed p.o.  $\mathbb{P}$ .

For a class  $\mathcal{C}$  of structures as in Section 1 and a property  $P$ , we define the *indestructible reflection number*  $\aleph_{\text{ref}}^*(\mathcal{C}, P)$  of  $P$  in  $\mathcal{C}$  is defined by:

$$(4.13) \quad \mathfrak{Refl}^*(\mathcal{C}, P) = \begin{cases} \min\{\kappa \in \text{Card} : \text{for all } A \in \mathcal{C} \text{ if } A \text{ is indestructively } \neg P \\ \text{then there are club many } B \in S_{<\kappa}^{\mathcal{C}}(A) \\ \text{with } B \not\equiv P\}, \\ \text{if } \{\kappa \in \text{Card} : \dots\} \neq \emptyset; \\ \infty, \quad \text{otherwise.} \end{cases}$$

Let  $\mathfrak{Refl}_{\text{Rado}}^*$ ,  $\mathfrak{Refl}_{\text{Galvin}}^*$ ,  $\mathfrak{Refl}_{\text{chr}}^*$ ,  $\mathfrak{Refl}_{\text{HP}}^*$ ,  $\mathfrak{Refl}_{\text{fBa}}^*$  be the indestructible version of corresponding reflection numbers in (A)  $\sim$  (E). By Lemma 3.2,  $\mathfrak{Refl}_{\text{Rado}}^* = \mathfrak{Refl}_{\text{Rado}}$ . Since the proof of Lemma 2.3 also holds for indestructible version we have:

$$\begin{array}{ccc} \mathfrak{Refl}_{\text{Galvin}} & \leq & \mathfrak{Refl}_{\text{chr}} \\ \vee & & \vee \\ \mathfrak{Refl}_{\text{Rado}} & \leq & \mathfrak{Refl}_{\text{Galvin}}^* \leq \mathfrak{Refl}_{\text{chr}}^* \end{array}$$

The following Theorems can be proved similarly to Proposition 3.1 and Theorem 4.2:

**Theorem 4.5** *Suppose that  $\kappa$  is a strongly compact cardinal and  $\mathbb{P} = \text{Col}(\aleph_1, \mu)$  for some  $\mu \geq \kappa$ . Then we have*

$$(4.14) \quad \Vdash_{\mathbb{P}} \text{“} \mathfrak{Refl}_{\text{Galvin}}^* = \mathfrak{Refl}_{\text{chr}}^* = \mathfrak{Refl}_{\text{HP}}^* = \aleph_2 \text{”}.$$

**Theorem 4.6** *Suppose that  $\kappa$  is strongly compact. Then, for any  $\mu \geq \kappa$  and  $\mathbb{Q} = \text{Fn}(\mu, 2)$ , we have*

$$(4.15) \quad \Vdash_{\mathbb{Q}} \text{“} \mathfrak{Refl}_{\text{Galvin}}^* = \mathfrak{Refl}_{\text{chr}} \leq 2^{\aleph_0} \text{”}.$$

Similarly to the theorems above  $\Vdash_{\mathbb{P}} \text{“} \mathfrak{Refl}_{\text{fBa}}^* = \aleph_2 \text{”}$  and  $\Vdash_{\mathbb{Q}} \text{“} \mathfrak{Refl}_{\text{fBa}}^* \leq 2^{\aleph_0} \text{”}$  can be obtained when we start from a supercompact  $\kappa$ .

## 5 Martin's axiom and large continuum

Let us remind first the following classical theorem by Kurepa, just for completeness:

**Lemma 5.1** (Đ. Kurepa 1940) *For a tree, the following are equivalent:*

- (a) *T is special.*
- (b) *There is a strictly order preserving  $f : T \rightarrow \mathbb{Q}$ .*

**Proof.** (b)  $\Rightarrow$  (a): Suppose that  $f : T \rightarrow \mathbb{Q}$  is strictly order preserving. Then  $f^{-1}''\{q\}$ ,  $q \in \mathbb{Q}$  is a countable partition of  $T$  into pairwise incomparable sets.

(a)  $\Rightarrow$  (b): Suppose that  $h : T \rightarrow \omega$  is such that

(5.1)  $h^{-1}''\{n\}$  is pairwise incomparable for all  $n \in \omega$ .

Let  $f : T \rightarrow \mathbb{Q}$  be defined by

$$(5.2) \quad f(t) = \sum \{2^{-(k+1)} : k \leq h(t), \text{ there is } t' \leq_T t \text{ such that } h(t') = k\}.$$

To show that this  $f$  is strictly order preserving, assume that  $t <_T u$ . By (5.1), we have  $h(t) \neq h(u)$ .

If  $h(t) > h(u)$ , then

$$\begin{aligned} f(t) &= \sum \{2^{-(k+1)} : k \leq h(u), \text{ there is } t' \leq_T t \text{ such that } h(t') = k\} \\ &\quad + \sum \{2^{-(k+1)} : h(u) < k \leq h(t), \text{ there is } t' \leq_T t \text{ such that } h(t') = k\} \\ &< \sum \{2^{-(k+1)} : k \leq h(u), \text{ there is } t' <_T u \text{ such that } h(t') = k\} + 2^{-(h(u)+1)} \\ &= f(u). \end{aligned}$$

If  $h(t) < h(u)$ , then

$$\begin{aligned} f(t) &= \sum \{2^{-(k+1)} : k \leq h(t), \text{ there is } t' \leq_T t \text{ such that } h(t') = k\} \\ &< \sum \{2^{-(k+1)} : k \leq h(t), \text{ there is } t' \leq_T u \text{ such that } h(t') = k\} + 2^{-(h(u)+1)} \\ &\leq f(u). \end{aligned}$$

□ (Lemma 5.1)

Let  $\mathfrak{ma}$  denote the first cardinal  $\kappa$  for which Martin's Axiom for a family of  $\kappa$  many dense sets does not hold. Note that  $\mathbf{MA}_\kappa$  is equivalent to  $\mathfrak{ma} \geq \kappa^+$ .

**Theorem 5.2** (Baumgartner-Malitz-Reinhardt [1]) *For any tree  $T$  of size  $< \mathfrak{ma}$  without uncountable branches, there is a mapping  $f : T \rightarrow \mathbb{Q}$  such that, for any  $t, t' \in T$ ,  $t <_T t'$  implies  $f(t) < f(t')$  (Note that, by Lemma 5.1, there is such a mapping  $f$  if and only if  $T$  is special).*

The Theorem 5.2 follows from Lemma 5.4 below. The following combinatorial Lemma 5.3 is the key to the Lemma 5.4:

**Lemma 5.3** *Suppose that  $T$  is a tree without uncountable branches. If  $S \subseteq [T]^{<\aleph_0}$  is uncountable and pairwise disjoint, then there are  $s, s' \in S$  such that any  $t \in s$  and  $t' \in s'$  are incomparable.*

**Proof.** Without loss of generality we may assume that  $S \subseteq [T]^n$  for some  $n \in \omega$ .

Assume toward a contradiction that, for all  $s, s' \in S$ , there are  $t \in s$  and  $t' \in s'$  such that  $t$  and  $t'$  are compatible.

We assume that there is some canonical linear ordering on  $T$  and with this ordering we can talk about “the  $k$ th element of  $s$ ” for  $s \in S$  and  $k < n$ . Let  $D$  be a ultrafilter over  $S$  with  $\{S \setminus u : u \in [S]^{\leq \aleph_0}\} \subseteq D$ . Thus, all elements of  $D$  are uncountable.

For  $s \in S$ ,  $t \in s$  and  $k < n$ , let

$$(5.3) \quad F_t^k = \{s' \in S : t \text{ is compatible with the } k\text{th element of } s'\}.$$

By assumption we have  $S = \bigcup_{k < n, t \in s} F_t^k$ . Thus, for each  $s \in S$ , there is  $t_s \in s$  and  $k_s < n$  such that  $F_{t_s}^{k_s} \in D$ .

Let  $k^* < n$  be such that  $S' = \{s \in S : k_s = k^*\}$  is uncountable.

The following claim is a contradiction to the assumption on  $T$  as desired:

**Claim 5.3.1**  $\{t_{s'} : s' \in S'\}$  is linearly ordered.

⊢ Suppose  $s, s' \in S'$ . Then  $F_{t_s}^{k^*} \cap F_{t_{s'}}^{k^*} \in D$ . Since elements of  $D$  are uncountable and by the pairwise disjointness of elements of  $S$ ,

$$(5.4) \quad \{\text{the } k^*\text{'th element of } u : u \in F_{t_s}^{k^*} \cap F_{t_{s'}}^{k^*}\}$$

is uncountable while  $T \downarrow t_s \cap T \downarrow t_{s'}$  is countable. Hence there is  $s'' \in F_{t_s}^{k^*} \cap F_{t_{s'}}^{k^*}$  such that  $t_s, t_{s'} \leq_T$  the  $k^*$ th element of  $s''$ . Since  $T$  is a tree, it follows that  $t_s$  and  $t_{s''}$  are compatible. ⊣ (Claim 5.3.1)

□ (Lemma 5.3)

**Lemma 5.4** Suppose that  $T$  is a tree without uncountable branches. Then the p.o.  $\langle \mathbb{P}_T, \leq_{\mathbb{P}_T} \rangle$  defined by

$$(5.5) \quad \mathbb{P}_T = \{p : p \text{ is a finite partial function from } T \text{ to } \omega \text{ such that,}$$

$$(5.5a) \quad \text{for any two distinct } t, t' \in \text{dom}(p),$$

$$p(t) = p(t') \text{ implies } t \text{ and } t' \text{ are incomparable}\}$$

with the ordering

$$(5.6) \quad p \leq_{\mathbb{P}_T} p' \Leftrightarrow p' \subseteq p$$

has the ccc.

**Proof.** Suppose that  $A \subseteq \mathbb{P}_T$  is uncountable. We have to show that there are two compatible elements in  $A$ . By  $\Delta$ -system Lemma and countability of  $\omega$ , we may assume without loss of generality, that  $\{\text{dom}(p) : p \in A\}$  builds a  $\Delta$ -system  $\subseteq [T]^{<\aleph_0}$  with the root  $s_0$  and that

$$(5.7) \quad p \upharpoonright s_0 \text{ is the same for all } p \in S.$$

By Lemma 5.3, there are  $p, p' \in A$  such that any  $t \in \text{dom}(p) \setminus s_0$  and  $t' \in \text{dom}(p') \setminus s_0$  are incomparable. Let  $p'' = p \cup p'$ .

**Claim 5.4.1**  $p'' \in \mathbb{P}_T$ .

$\vdash$   $p''$  is a partial function by (5.7). Thus it is enough to show that  $p''$  satisfies (5.5a).

Suppose  $p''(t) = p''(t')$  for two distinct  $t, t' \in \text{dom}(p'')$ . If  $t, t' \in \text{dom}(p)$  or  $t, t' \in \text{dom}(p')$  then  $t$  and  $t'$  are incomparable since  $p, p' \in \mathbb{P}_T$ . If  $t \in \text{dom}(p) \setminus s_0$  and  $t' \in \text{dom}(p') \setminus s_0$  — or  $t \in \text{dom}(p') \setminus s_0$  and  $t' \in \text{dom}(p) \setminus s_0$  — then  $t$  and  $t'$  are incomparable anyway by the choice of  $p$  and  $p'$ .  $\dashv$  (Claim 5.4.1)

Since  $p'' \supseteq p, p'$ , it follows that  $p'' \leq_{\mathbb{P}_T} p, p'$ . Thus  $p$  and  $p'$  are compatible in  $\mathbb{P}_T$ .  $\square$  (Lemma 5.4)

**Proof of Theorem 5.2:** Suppose that  $T$  is a tree without uncountable branches and  $|T| < \mathfrak{ma}$ . For each  $t \in T$  let

$$(5.8) \quad D_t = \{p \in \mathbb{P}_T : t \in \text{dom}(p)\}.$$

Then each  $D_t, t \in T$  is dense in  $T$ . Let  $\mathcal{D} = \{D_t : t \in T\}$ . Since  $|\mathcal{D}| < \mathfrak{ma}$  there is a  $\mathcal{D}$ -generic  $G$  over  $\mathbb{P}_T$ .

$f_G = \bigcup G$  is a function from  $T$  to  $\omega$  and witnesses the specialness of  $T$ .

$\square$  (Theorem 5.2)

A prominent consequence of Theorem 5.2 is that, under  $\mathbf{MA}_{\aleph_1}$ , every Aronszajn tree is special.

The following Lemma together with Theorem 5.2 implies that  $\mathfrak{Rfl}_{\text{Rado}}$  is strictly larger than  $\mathfrak{ma}$ .

Let

$$(5.9) \quad T_{\mathbb{R}} = \{t : t \text{ is an increasing sequence of elements of } \mathbb{R} \\ \text{of order-type of countable successor ordinal}\}$$

be the tree with the ordering

$$(5.10) \quad t \leq_{T_{\mathbb{R}}} t' \Leftrightarrow t' \text{ is an end-extension of } t$$

for  $t, t' \in T_{\mathbb{R}}$ . Clearly  $T_{\mathbb{R}}$  has no uncountable branch.

**Lemma 5.5** (Todorćević [15])  $T_{\mathbb{R}}$  is not special.

**Proof.** This follows immediately from Theorem 1 in [15].

$\square$  (Lemma 5.5)

**Corollary 5.6**  $\mathfrak{ma} < \mathfrak{Rfl}_{\text{Rado}}$ .

**Proof.**  $T_{\mathbb{R}}$  does not have any uncountable branch. By Theorem 5.2, it follows that all subtrees of  $T_{\mathbb{R}}$  of size  $< \mathfrak{ma}$  are special. Since  $T_{\mathbb{R}}$  itself is not special by Lemma 5.5,  $T_{\mathbb{R}}$  witnesses the inequality  $\mathfrak{ma} < \mathfrak{Rfl}_{\text{Rado}}$ .

$\square$  (Corollary 5.6)

The non-freeness in any variety is preserved by a ccc p.o.:

**Proposition 5.7** (Fuchino [6]) *For an algebra in a variety  $\mathcal{V}$ , if there is a ccc p.o.  $\mathbb{P}$  such that  $\Vdash_{\mathbb{P}}$  “ $A$  is free” then  $A$  is really free.*  $\square$

**Theorem 5.8** *Suppose that  $\kappa$  is a supercompact cardinal. If  $\mathbb{P}$  is the standard p.o. forcing Martin’s Axiom and  $2^{\aleph_0} = \kappa$  then we have*

$$(5.11) \quad \Vdash_{\mathbb{P}} \text{“MA} + \mathfrak{Rfl}_{\text{fBa}} = 2^{\aleph_0} < \mathfrak{Rfl}_{\text{Rado}} \text{”}.$$

**Proof.**  $\Vdash_{\mathbb{P}}$  “ $\dots = \dots$ ” follows from Proposition 5.7 and an argument similar to the proof of Theorem 4.2.  $\Vdash_{\mathbb{P}}$  “ $\dots < \dots$ ” follows from Corollary 5.6.  $\square$  (Theorem 5.8)

If we have a strongly compact cardinal above the supercompact  $\kappa$  in Theorem 5.8, we can conclude that  $\mathfrak{Rfl}_{\text{Rado}} < \infty$  and manipulate further the value of  $\mathfrak{Rfl}_{\text{Rado}}$ .

## References

- [1] J. Baumgartner, J. Malitz and W. Reinhardt, Embedding trees in the rationals, Proceedings of the National Academy of Sciences Vol.67, No.4, (1970), 1748–1753.
- [2] A. Dow, F.D. Tall and W.A.R. Weiss, New proofs of the consistency of the normal Moore space conjecture I, II, Topology and its APPLICATIONS, 37 (1990) 33-51, 115-129.
- [3] Akihiro Kanamori, The Higher Infinite, Springer–Verlag (1994/2003).
- [4] Philipp Doebler, Rado’s Conjecture implies that all stationary set preserving forcing are semiproper, Journal of Mathematical Logic Vol.13, 1 (2013).
- [5] Paul Erdős and András Hajnal, On chromatic number of graphs and set-systems, Acta Mathematica Hungarica, 17 (1–2), (1966), 61–99.
- [6] Sakaé Fuchino, On potential embedding and versions of Martin’s axiom, Notre Dame Journal of Logic, Vol.33, No.4 (1992), 481–492.
- [7] Sakaé Fuchino, István Juhász, Lajos Soukup, Zoltán Szentmiklóssy and Toshimichi Usuba, Fodor-type Reflection Principle and reflection of metrizable and meta-Lindelöfness, Topology and its Applications Vol.157, 8 (2010), 1415–1429.
- [8] Sakaé Fuchino, Sabine Koppelberg and Makoto Takahashi, On  $L_{\infty\kappa}$ -free Boolean algebras, Annals of Pure and Applied Logic 55, (1992), 265–284.
- [9] Sakaé Fuchino, Lajos Soukup, Hiroshi Sakai and Toshimichi Usuba, More about Fodor-type Reflection Principle, submitted.
- [10] Sakaé Fuchino and Assaf Rinot, Openly generated Boolean algebras and the Fodor-type Reflection Principle, Fundamenta Mathematicae 212, (2011), 261-283.
- [11] Sakaé Fuchino and Hiroshi Sakai, On reflection and non-reflection of countable list-chromatic number of graphs, Kôkyûroku, No.1790, (2012), 31–44.
- [12] Sakaé Fuchino, Hiroshi Sakai, Victor Torres-Perez and Toshimichi Usuba, Rado’s Conjecture and the Fodor-type Reflection Principle, in preparation.

- [13] A. Hajnal and I. Juhász, On spaces in which every small subspace is metrizable, *Bulletin de l'académie polonaise des sciences, Série des sciences math, astr. et phys.*, Vol.24, No.9, (1976), 728–731.
- [14] Hiroshi Sakai, Semistationary and stationary reflection, *Journal of Symbolic Logic* 73(1), (2008), 181–192.
- [15] Stevo Todorčević, On a conjecture of Rado, *Journal London Mathematical Society* Vol.s2-27, (1) (1983), 1–8.
- [16] Stevo Todorčević, Real functions on the family of all well-ordered subsets of a partially ordered set, the *Journal of Symbolic Logic*, Vol.48, No.1 (1983), 91–96.
- [17] Stevo Todorčević, Conjectures of Rado and Chang and Cardinal Arithmetic, in: *Finite and Infinite Sets in Combinatorics* (N. W. Sauer et al., eds), Kluwer Acad. Publ. (1993) 385-398.
- [18] Stevo Todorčević, Combinatorial dichotomies in set theory, the *Bulletin of Symbolic Logic*, Vol.17, No.1 (2011), 1–72.