

RMD  
OpenRheumatic &  
Musculoskeletal  
Diseases

## ORIGINAL ARTICLE

## Nardilysin is involved in autoimmune arthritis via the regulation of tumour necrosis factor alpha secretion

Takayuki Fujii,<sup>1</sup> Eiichiro Nishi,<sup>2,3</sup> Hiromu Ito,<sup>1</sup> Hiroyuki Yoshitomi,<sup>4</sup> Moritoshi Furu,<sup>5</sup> Namiko Okabe,<sup>6</sup> Mikiko Ohno,<sup>2</sup> Kiyoto Nishi,<sup>2</sup> Yusuke Morita,<sup>2</sup> Yugo Morita,<sup>1</sup> Masayuki Azukizawa,<sup>1</sup> Akinori Okahata,<sup>1</sup> Takuya Tomizawa,<sup>1</sup> Takeshi Kimura,<sup>2</sup> Shuichi Matsuda<sup>1</sup>

**To cite:** Fujii T, Nishi E, Ito H, *et al.* Nardilysin is involved in autoimmune arthritis via the regulation of tumour necrosis factor alpha secretion. *RMD Open* 2017;**3**:e000436. doi:10.1136/rmdopen-2017-000436

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/rmdopen-2017-000436>).

Received 16 January 2017

Revised 23 March 2017

Accepted 19 April 2017

## ABSTRACT

**Objective** Tumour necrosis factor alpha (TNF- $\alpha$ ) plays an important role in rheumatoid arthritis (RA). TNF- $\alpha$  is synthesised as a membrane-anchored precursor and is fully activated by a disintegrin and metalloproteinase 17 (ADAM17)-mediated ectodomain shedding. Nardilysin (NRDC) facilitates ectodomain shedding via activation of ADAM17. This study was undertaken to elucidate the role of NRDC in RA.

**Methods** NRDC-deficient (*Nrdc*<sup>-/-</sup>) mice and macrophage-specific NRDC-deficient (*Nrdc*<sup>delM</sup>) mice were examined in murine RA models, collagen antibody-induced arthritis (CAIA) and K/BxN serum transfer arthritis (K/BxN STA). We evaluated the effect of gene deletion or silencing of *Nrdc* on ectodomain shedding of TNF- $\alpha$  in macrophages or monocytes. NRDC concentration in synovial fluid from patients with RA and osteoarthritis (OA) were measured. We also examined whether local gene silencing of *Nrdc* ameliorated CAIA.

**Results** CAIA and K/BxN STA were significantly attenuated in *Nrdc*<sup>-/-</sup> mice and *Nrdc*<sup>delM</sup> mice. Gene deletion or silencing of *Nrdc* in macrophages or THP-1 cells resulted in the reduction of TNF- $\alpha$  shedding. The level of NRDC is higher in synovial fluid from RA patients compared with that from OA patients. Intra-articular injection of anti-*Nrdc*small interfering RNA ameliorated CAIA.

**Conclusion** These data indicate that NRDC plays crucial roles in the pathogenesis of autoimmune arthritis and could be a new therapeutic target for RA treatment.

## INTRODUCTION

Rheumatoid arthritis (RA) is a chronic inflammatory disease that causes joint inflammation and destruction affecting 0.5%–1.0% of the population.<sup>1</sup> Anti-tumour necrosis factor alpha (TNF- $\alpha$ ) biologic agents have led to a revolutionary improvement in the management of RA patients, indicating that this inflammatory cytokine plays a key role in RA pathogenesis. The importance of TNF- $\alpha$  has been also demonstrated in murine models of RA; for example, human TNF- $\alpha$  transgenic

## Key messages

## What is already known about this subject?

- Tumour necrosis factor alpha (TNF- $\alpha$ ), which is activated by ectodomain shedding, plays an important role in rheumatoid arthritis (RA).
- Nardilysin facilitates TNF- $\alpha$  shedding via enhancing a disintegrin and metalloproteinase 17 activity.

## What does this study add?

- Deletion or inhibition of nardilysin prevents autoantibody-induced arthritis, murine models of RA.
- Concentration of nardilysin in synovial fluid is markedly increased in patients with RA compared with that in patients with osteoarthritis, suggesting a crucial role of nardilysin in the pathogenesis of RA.

## How might this impact on clinical practice?

- Nardilysin could be a novel biomarker and therapeutic target for RA.

mice develop arthritis,<sup>2</sup> TNF- $\alpha$  is produced in synovial tissues of murine experimental models of RA,<sup>3</sup> murine experimental arthritis is attenuated in TNF- $\alpha$ -deficient mice<sup>4</sup> and neutralisation of TNF- $\alpha$  ameliorates murine collagen-induced arthritis.<sup>5</sup> TNF- $\alpha$  is produced mostly by macrophages in RA synovial tissues<sup>6</sup> and, alone or in combination with other cytokines or chemokines, recruits inflammatory cells such as B cells, T cells and neutrophils, and induces production of cytokines, chemokines, matrix metalloproteinases or receptor activator of nuclear factor  $\kappa$ B ligand, resulting in joint inflammation and bone resorption.<sup>7–11</sup>

One of the earliest subclinical features of RA is a breakdown of self-tolerance. Autoantibodies such as rheumatoid factor (RF) and anti-citrullinated peptide antibody (ACPA) are found in



CrossMark

For numbered affiliations see end of article.

## Correspondence to

Dr Eiichiro Nishi; [nishi@kuhp.kyoto-u.ac.jp](mailto:nishi@kuhp.kyoto-u.ac.jp) and Dr Hiromu Ito; [hiromu@kuhp.kyoto-u.ac.jp](mailto:hiromu@kuhp.kyoto-u.ac.jp)

the patients' serum long before arthritis occurs.<sup>12</sup> While the autoantigens have not been fully identified, autoantibodies may contribute to the pathogenesis of RA via immunocomplex (IC) formation and deposition in synovial tissues.<sup>13</sup> The importance of autoantibodies has been demonstrated in mouse autoantibody-induced arthritis (AAIA) models of RA, such as collagen antibody-induced arthritis (CAIA) and K/BxN serum transfer arthritis (K/BxN STA). Antibodies against intra-articular antigens, type II collagen in CAIA<sup>14 15</sup> and glucose-6-phosphoisomerase in K/BxN STA<sup>16 17</sup> are arthrogenic in mice. TNF- $\alpha$  has been demonstrated to be important in both models: anti-TNF- $\alpha$  antibody attenuates CAIA,<sup>18</sup> while TNF- $\alpha$ -deficient mice are resistant to K/BxN STA.<sup>4</sup>

Importantly, TNF- $\alpha$  is synthesised as a membrane-anchored precursor and is fully activated by a disintegrin and metalloproteinase 17 (ADAM17)-mediated ectodomain shedding,<sup>19</sup> indicating that ADAM17 is an attractive alternative target for treatment of RA. However, the multiple disease phenotypes of ADAM17-deficient patients and mice have raised concerns about the potential adverse effects of anti-ADAM17 therapy.<sup>20 21</sup> We have previously shown that ADAM17 activity and TNF- $\alpha$  shedding are enhanced by nardilysin, N-arginine dibasic convertase (NRDC), via its direct binding to ADAM17.<sup>22-24</sup>

NRDC is a zinc peptidase of the M16 family that selectively cleaves dibasic sites.<sup>25 26</sup> We identified NRDC on the cell surface as a specific binding partner of heparin-binding epidermal growth factor (EGF)-like growth factor (HB-EGF) and demonstrated that NRDC enhances ectodomain shedding of HB-EGF.<sup>27</sup> Our subsequent studies revealed that NRDC enhances ectodomain shedding of not only HB-EGF but also a wide range of other membrane proteins including TNF- $\alpha$ ,<sup>23 28</sup> neuregulin 1<sup>29</sup> and amyloid precursor protein.<sup>22</sup> We have also demonstrated that NRDC-induced ectodomain shedding plays critical roles in myelination, gastric cancer and Alzheimer's disease.<sup>22-24 28-31</sup>

We demonstrate here that NRDC-deficient (*Nrdc*<sup>-/-</sup>) mice and macrophage-specific NRDC-deficient (*Nrdc*<sup>delM</sup>) mice show attenuation of AAIA. We also demonstrate that gene deletion or silencing of *Nrdc* in mouse macrophages or human monocytic cells results in reduction of TNF- $\alpha$  shedding. Importantly, the level of NRDC is increased in synovial fluid from RA patients compared with that from osteoarthritis (OA) patients. Moreover, intra-articular injection of anti-*Nrdc* small interfering RNA (siRNA) ameliorates CAIA. Together, these data suggest that NRDC controls AAIA via the regulation of TNF- $\alpha$  shedding and could be a new therapeutic target for RA treatment.

## MATERIALS AND METHODS

The full methods are provided in the online supplementary material.

### Blood and synovial fluid collection

Synovial fluid was obtained during joint surgery from 20 patients with RA and 17 patients with OA, who had

been admitted to Kyoto University Hospital. All of the RA patients met 2010 American College of Rheumatology and European League against Rheumatism (ACR/EULAR) classification criteria for RA. Peripheral blood samples were obtained and the level of C-reactive protein (CRP), erythrocyte sedimentation rate (ESR), RF, and ACPA was examined before surgery. Disease activity score 28 (DAS28) was calculated according to the results of physical examinations and blood tests.<sup>32</sup> The full methods are provided in the online supplementary material.

## RESULTS

### *Nrdc*-deficient mice are resistant to CAIA


To explore the role of NRDC in RA, we compared wild-type (WT) and *Nrdc*<sup>-/-</sup> mice with a BALB/c background using the CAIA model. After the intraperitoneal administration of a five-clone cocktail of monoclonal anti-type II collagen antibodies, we assessed the severity of arthritis daily for 2 weeks. The onset of arthritis was delayed (on day 3 in WT mice and on day 6 in *Nrdc*<sup>-/-</sup> mice, respectively) and the average of arthritis score was significantly and markedly lower in *Nrdc*<sup>-/-</sup> mice (figure 1A,B). Ankle thickness was also significantly less in *Nrdc*<sup>-/-</sup> mice from days 4 to 9 compared with that in WT mice. Histological examination of the ankle joints demonstrated that synovial hypertrophy with infiltration of inflammatory cells and bone and cartilage erosion were reduced or absent in *Nrdc*<sup>-/-</sup> mice (figure 1C). These results suggest that NRDC is crucial for AAIA development and progression.

### Macrophage-specific *Nrdc*-deficient mice are resistant to CAIA

Because macrophages play an important role in the pathogenesis of RA, we next generated *Nrdc*<sup>delM</sup> mice by crossing *Nrdc*<sup>fllox/fllox</sup> mice with *Lyz2*-Cre mice, in which Cre recombinase is expressed under the control of the lysozyme 2 gene (*Lyz2*) promoter. Because the *Nrdc*<sup>fllox/fllox</sup> and *Nrdc*<sup>delM</sup> mice have a C57BL/6 genetic background, which is more resistant to CAIA than BALB/c, we administered 2.5 times more antibody to *Nrdc*<sup>fllox/fllox</sup> and *Nrdc*<sup>delM</sup> mice for the CAIA experiments compared with the amount that was used for the *Nrdc*<sup>-/-</sup> mice. The results showed that the onset of arthritis was delayed and the severity of arthritis was markedly reduced in *Nrdc*<sup>delM</sup> mice compared with controls (figure 2A, B). As was the case for *Nrdc*<sup>-/-</sup> mice, synovial hypertrophy and bone erosion were attenuated in *Nrdc*<sup>delM</sup> mice (figure 2C). These findings demonstrated that macrophage-specific deletion of NRDC almost completely recapitulated the resistance of *Nrdc*<sup>-/-</sup> mice to CAIA.

### Arthritis in the K/BxN STA model is attenuated in *Nrdc*<sup>delM</sup> mice

To confirm the role of NRDC in autoimmune arthritis, we examined *Nrdc*<sup>fllox/fllox</sup> and *Nrdc*<sup>delM</sup> mice in another murine model, the K/BxN STA model. K/BxN mouse serum has been reported to strongly induce arthritis in a wide range of mouse strains.<sup>33</sup> As shown in figure 2D,E,



**Figure 1** CAIA is attenuated in *Nrdc*<sup>-/-</sup> mice. (A) Photographs of the hind paws 4 days and 7 days after the induction of CAIA. Upper panels, wild-type (WT); lower panels, *Nrdc*<sup>-/-</sup> mice. (B) Arthritis score (upper) and ankle thickness (lower) were measured daily after the induction of CAIA. n=5 per genotype (mean ± SEM). \*p<0.05; \*\*p<0.01, Welch's t-test. (C) H&E staining of the ankle joints 14 days after the induction of CAIA in WT (upper panels) and *Nrdc*<sup>-/-</sup> (lower panels) mice. Scale bars: 500 μm in the 40× magnification and 200 μm in the 100× magnification. Symbols in the 100× magnification images: arrow, bone erosion; arrow head, cartilage erosion; \*, hypertrophic synovium.


K/BxN STA was significantly attenuated in *Nrdc*<sup>delM</sup> mice up to day 8 after injection. In contrast, there were no significant differences between *Nrdc*<sup>flox/flox</sup> and *Nrdc*<sup>delM</sup> mice in the arthritis score after day 9 or in ankle thickness, although both values tended to be lower in *Nrdc*<sup>delM</sup> mice (figure 2D,E). The histology of the hind paws at day 14 demonstrated that the synovial hypertrophy with infiltration of inflammatory cells, bone erosion and cartilage erosion were equivalently observed in both *Nrdc*<sup>flox/flox</sup> mice and *Nrdc*<sup>delM</sup> mice (figure 2F), which is consistent with the results of arthritis score and ankle thickness. Collectively, these data demonstrated that NRDC deficiency in macrophages attenuated the

severity of arthritis in two different experimental models of AAIA.

### Secretion of TNF-α from macrophages and monocytic cells is reduced by the deletion of *Nrdc*

Because macrophages are a major source of TNF-α in AAIA<sup>34</sup> and NRDC facilitates ectodomain shedding of TNF-α,<sup>23</sup> we hypothesised that the attenuation of AAIA in *Nrdc*<sup>-/-</sup> mice and *Nrdc*<sup>delM</sup> mice was due to the impairment of ectodomain shedding of TNF-α in macrophages. To test this hypothesis, we first examined TNF-α secretion from NRDC-deficient macrophages. We harvested bone marrow derived macrophages (BMDMs) from






**Figure 2** Macrophage-specific *Nrdc*-deficient (*Nrdc*<sup>delM</sup>) mice are resistant to CAIA and K/BxN STA. (A) Photographs of the hind paws 4 days and 7 days after the induction of CAIA. Upper panels, *Nrdc*<sup>flox/flox</sup>; lower panels, *Nrdc*<sup>delM</sup> mice. (B) Arthritis score (upper) and ankle thickness (lower) were measured daily after the induction of CAIA. n=5 per genotype (mean ± SEM). \*p<0.05; \*\*p<0.01, Welch's t-test. (C) H&E staining of the ankle joints 14 days after the induction of CAIA in *Nrdc*<sup>flox/flox</sup> (upper panels) and *Nrdc*<sup>delM</sup> (lower panels) mice. Scale bars: 500 μm in the 40× magnification and 200 μm in the 100× magnification. Symbols in the 100× magnification images: arrow, bone erosion; \*, synovial hypertrophy. (D) Photographs of the hind paws 3 days and 5 days after the induction of K/BxN STA. Upper panels, *Nrdc*<sup>flox/flox</sup>; lower panels, *Nrdc*<sup>delM</sup> mice. (E) Arthritis score (upper) and ankle thickness (lower) were measured daily after the induction of K/BxN STA. n=4 per genotype (mean ± SEM). \*p<0.05; \*\*p<0.01, Welch's t-test. (F) H&E staining of the ankle joints 14 days after the induction of K/BxN STA in *Nrdc*<sup>flox/flox</sup> (upper panels) and *Nrdc*<sup>delM</sup> (lower panels) mice. Scale bars: 500 μm in the 40× magnification and 200 μm in the 100× magnification. Symbols in the 100× magnification images: arrow, bone erosion; arrow head, cartilage erosion; \*, synovial hypertrophy.

*Nrdc*<sup>flox/flox</sup> and *Nrdc*<sup>delM</sup> mice and confirmed that NRDC protein expression was remarkably reduced in BMDMs from *Nrdc*<sup>delM</sup> mice (figure 3A). BMDMs were stimulated with or without a low concentration (10 ng/mL)

of lipopolysaccharide (LPS), then the concentration of secreted TNF-α in the supernatants and the levels of TNF-α messenger RNA (mRNA) in BMDMs were examined. The results showed that there was no significant



**Figure 3** Secretion of TNF- $\alpha$  from macrophages and monocytic cells were reduced by the deletion of *Nrdc*. (A) Western blot analysis of NRDC in BMDM from  $Nrdc^{lox/lox}$  (left) and  $Nrdc^{delM}$  mice (right).  $\beta$ -Actin was used as a loading control. (B) TNF- $\alpha$  mRNA levels in unstimulated or LPS-stimulated BMDMs were analysed by reverse transcription and quantitative PCR (qRT-PCR).  $\beta$ -Actin mRNA levels were used as internal control. n=6 (mean  $\pm$  SEM). (C) TNF- $\alpha$  secreted from BMDMs in conditioned medium was measured by ELISA. n=6 (mean  $\pm$  SEM). (D) Membrane-bound TNF- $\alpha$  in BMDMs was stained 3 hours after LPS stimulation and analysed by flow cytometry. Filled histogram, unstimulated control; blue open histogram,  $Nrdc^{lox/lox}$ ; red open histogram,  $Nrdc^{delM}$ . Representative histograms out of four independent experiments are shown. (E) ADAM17 mRNA levels in unstimulated or LPS-stimulated BMDMs were analysed by qRT-PCR.  $\beta$ -Actin mRNA levels were used as internal control. n=6 (mean  $\pm$  SEM). (F) NRDC mRNA levels in THP-1 cells, transfected with lentiviral vectors for negative control RNA interference (RNAi) (control microRNA (miR)) or RNAi targeting *Nrdc* (anti-NRDC miR), were analysed using qRT-PCR.  $\beta$ -Actin mRNA was used as an internal control. n=6 (mean  $\pm$  SEM). (G) NRDC and ADAM17 protein expressions in THP-1 cells transfected with lentiviral vectors for control miR (left) or anti-NRDC miR 1 (right) were analysed by Western blot.  $\beta$ -Actin was used as a loading control. (H) TNF- $\alpha$  mRNA levels in THP-1 cells, transfected with lentiviral vectors as described in (G), were analysed using qRT-PCR. n=8 (mean  $\pm$  SEM). (I) Secreted TNF- $\alpha$  (conditioned medium) and cell-contained TNF- $\alpha$  (cell extracts) in THP-1 cells described above were measured by ELISA. n=8 (mean  $\pm$  SEM). (J) F4/80 expression in peritoneal macrophage (PMs). Filled histogram, isotype control; open histogram, F4/80. Representative histograms out of three independent experiments are shown. (K) NRDC expression in PMs, isolated from  $Nrdc^{lox/lox}$  (left) and  $Nrdc^{delM}$  mice (right) after the reverse passive Arthus reaction, was examined by Western blot.  $\beta$ -Actin was used as a loading control. (L) Membrane-bound TNF- $\alpha$  was stained and analysed by flow cytometry. Filled histogram, unstimulated control; blue open histogram,  $Nrdc^{lox/lox}$ ; red open histogram,  $Nrdc^{delM}$ . Representative histograms out of three independent experiments are shown. (M) Measurement of secreted TNF- $\alpha$  concentration in the peritoneal lavage of  $Nrdc^{lox/lox}$  (n=5),  $Nrdc^{delM}$  (n=5) and control (n=4) mice. \*p<0.05; \*\*p<0.01, Welch's t-test.

difference in the mRNA level, but that there was a significant decrease in TNF- $\alpha$  secretion from NRDC-deficient BMDMs at 3 hours after stimulation (figure 3B,C). Because impairment of TNF- $\alpha$  shedding leads to accumulation of membrane-bound TNF- $\alpha$ ,<sup>35</sup> we analysed the cell surface expression of TNF- $\alpha$  on BMDMs by flow cytometry and found that membrane-bound TNF- $\alpha$  was increased in BMDMs from *Nrdc*<sup>delM</sup> mice (figure 3D). ADAM17 is essential for TNF- $\alpha$  shedding, and we previously reported that NRDC enhances the sheddase activity of ADAM17 via a direct interaction. Consistent with this, there were no significant differences in ADAM17 mRNA levels between BMDMs from *Nrdc*<sup>fllox/fllox</sup> and *Nrdc*<sup>delM</sup> mice (figure 3E).

We next assessed the effect of *NRDC* gene knockdown in a human monocytic cell line, THP-1. Efficient inhibition of *NRDC* mRNA and protein levels was confirmed in THP-1 cells infected with lentiviral vectors expressing RNAi targeting *NRDC* (figure 3F,G). LPS-induced TNF- $\alpha$  secretion in the supernatants was significantly decreased by gene knockdown of *NRDC*, whereas TNF- $\alpha$  mRNA levels and TNF- $\alpha$  protein concentrations in the cell lysates did not differ between control and *NRDC*-knocked down cells (figure 3H,I). Together, our results in mouse macrophages and human monocytic cells suggested that TNF- $\alpha$  activation via ectodomain shedding was inhibited by the deletion of *NRDC*.

To assess the *in vivo* role of *NRDC* on TNF- $\alpha$  secretion in a way more relevant to disease, we examined the effect of immunocomplexes (ICs) on TNF- $\alpha$  secretion. In the mechanism of AAIA, inflammatory cells are activated by ICs that are deposited in the joints. To mimic this response, we applied a reverse passive Arthus reaction peritonitis<sup>36</sup> model to *Nrdc*<sup>fllox/fllox</sup> and *Nrdc*<sup>delM</sup> mice. In this model, ovalbumin (OVA) and anti-OVA antibody are administered separately, and then combine to make ICs and trigger complement pathways. We confirmed that cells collected from peritoneal lavage 3 hours after the administration were mostly F4/80 positive, and therefore were peritoneal macrophages (PMs) (figure 3J). We also confirmed that *NRDC* expression was markedly reduced in PMs from *Nrdc*<sup>delM</sup> mice (figure 3K), and analysis by flow cytometry demonstrated that membrane-bound TNF- $\alpha$  was increased in PMs from *Nrdc*<sup>delM</sup> mice (figure 3L). In contrast, secreted TNF- $\alpha$  in the peritoneal lavage was lower in *Nrdc*<sup>delM</sup> mice than in *Nrdc*<sup>fllox/fllox</sup> mice (figure 3M). These results indicated that ectodomain shedding of TNF- $\alpha$  was impaired in *NRDC*-deficient macrophages.

### NRDC increased in synovial fluid of patients with RA

To assess the pathophysiological role of *NRDC* in patients, we collected synovial fluid from patients with RA or OA during joint surgery. Intriguingly, we found a remarkable and significant elevation of the level of *NRDC* and TNF- $\alpha$  in synovial fluid from RA patients compared with that from OA patients (table 1, figure 4A,B).

**Table 1** Patients' characteristics

	OA	RA	p Value
	(n=17)	(n=20)	
Age (years)	78±4.94	65.8±10.6	<0.01‡
Female (%)	14 (82.4)	16 (80)	0.999§
CRP (mg/dL)	0.0.69±0.079¶	1.845±1.88	<0.01‡
ESR (mm/hour)	24.4±14.6#	47.3±35.0	0.014‡
DAS28-ESR		4.25±1.12	
RF (U/mL)		73.5 (18.6 to 189.1)	
ACPA positive (%)		19 (95)	
OA grading*			
Grade I or II (%)	1 (5.9)		
Grade III or IV (%)	16 (94.1)		
RA stage†			
Stage I or II (%)		7 (35)	
Stage III or IV (%)		13 (65)	

Data are presented as mean±SD, n (%) or median (IQR).

\*The Kellgren-Lawrence Grading Scale.

†The Steinbrocker Staging system.

‡Data was not available in one patient.

§Data was not available in two patients.

¶Fisher's exact test.


#Welch's t-test.

Notably, the level of TNF- $\alpha$  in RA synovial fluid had a tendency of positive correlation with the level of *NRDC* ( $p=0.054$ ), suggesting the causative role of *NRDC* in TNF- $\alpha$  elevation (figure 4C). We next analysed the association of *NRDC* levels in RA synovial fluid with the patients' clinical condition. Interestingly, *NRDC* levels showed a significant but modest correlation with CRP levels, but no correlations with other factors including ESR, disease activity (disease activity score 28 (DAS28)-ESR), RF or ACPA (figure 4D and data not shown). These data suggested that *NRDC* could be a unique synovial fluid marker for the differential diagnosis of late RA and late OA.


### Gene silencing of *Nrdc* in the ankle joints ameliorated CAIA in WT mice

To confirm the pathological role *in vivo* of *NRDC* in autoimmune arthritis, we examined whether local gene silencing of *Nrdc* ameliorated CAIA. We first tested three siRNAs targeting *Nrdc* in three different cell lines (RAW264.7, WEHI 13 VAR and mouse embryonic fibroblasts) and selected the siRNA with the best silencing efficacy (siRNA3) (figure 5A). This anti-*Nrdc* siRNA was injected into the left ankle joint of mice 2 days after CAIA induction, while negative control siRNA was injected into the right ankle joint. The results showed that the severity of arthritis was clearly reduced in the left joint (anti-*NRDC* siRNA) compared with the right joint (control siRNA) (figure 5C). The measurement of ankle thickness revealed that anti-*Nrdc* siRNA significantly ameliorated CAIA from days 3 to 8 and days 12 to 14 (figure 5D).





**Figure 4** NRDC and TNF- $\alpha$  increase in synovial fluid from patients with RA. (A) NRDC concentration in synovial fluid from OA (n=17) and RA (n=20) patients. \*\* $p<0.01$ , Welch's t-test. (B) TNF- $\alpha$  concentration in synovial fluid from OA and RA patients. \*\* $p<0.01$ , Welch's t-test. (C) NRDC in synovial fluid has a tendency of positive correlation with TNF- $\alpha$  in RA patients. Solid line, approximate line; broken line, 95% CI.  $r=0.44$ ,  $p=0.054$ , Spearman's rank correlation test. (D) NRDC in synovial fluid is significantly correlated with CRP in RA patients. Solid line, approximate line; broken line, 95% CI.  $r=0.50$ ,  $p<0.05$ , Spearman's rank correlation test.



**Figure 5** Intra-articular *Nrdc* gene silencing with siRNA attenuates CAIA. (A) Screening of siRNAs against NRDC in RAW264.7, WEHI 13 VAR and mouse embryonic fibroblasts (MEF). (B) Schedule of the induction of CAIA and intra-articular injection of siRNA. (C) Photographs of the hind paws 4 days after the induction of CAIA. Anti-NRDC siRNA was injected in the left ankle, while control siRNA was injected in the right ankle joints of BALB/c WT mice. (D) Ankle thickness was measured daily after the induction of CAIA (n=5 per treatment). \* $p<0.05$ , Welch's t-test.

## DISCUSSION

In this study, we demonstrate for the first time in vivo that NRDC plays important roles in autoimmune arthritis. First, *Nrdc*<sup>-/-</sup> mice were resistant to CAIA, and resistance to two models of AAIA, CAIA and K/BxN STA, was confirmed in *Nrdc*<sup>delM</sup> mice, indicating that NRDC in macrophages is crucial for the pathogenesis of autoimmune arthritis. In both CAIA and K/BxN STA models, inhibition of TNF- $\alpha$  has been shown to attenuate arthritis.<sup>4,18</sup> Given that macrophages are a major source of TNF- $\alpha$  in the joints, reduced release of TNF- $\alpha$  from NRDC-deficient macrophages may strongly contribute to the attenuation of arthritis.<sup>6</sup>

TNF- $\alpha$  is synthesised as a type 2 membrane protein precursor, which is cleaved by ADAM17 at a juxtamembrane domain to be released from the cell surface. A comparison of the phenotypes of TNF- $\alpha$ -null mice and knockin mice expressing an uncleavable form of TNF- $\alpha$  (mice without soluble TNF- $\alpha$ ) revealed that soluble TNF- $\alpha$  is dispensable for the organogenesis of most lymphoid tissues, but is indispensable for optimal development of inflammatory lesions.<sup>19</sup> This finding indicated that TNF- $\alpha$  activation by ectodomain shedding plays important roles in inflammatory diseases. Indeed, myeloid cell-specific ADAM17-deficient mice were protected from K/BxN STA.<sup>34</sup> Moreover, the deletion of *iRHOM2*, a critical regulator of protein maturation and of the cell surface localisation of ADAM17, also protected against K/BxN STA,<sup>34</sup> further indicating that ADAM17-mediated TNF- $\alpha$  shedding is involved in AAIA. Our previous studies have demonstrated that NRDC enhances ectodomain shedding of TNF- $\alpha$  via activation of ADAM17.<sup>23,28,30</sup> NRDC is one of the rare proteins that have been shown to bind directly to ADAM17 and enhance its enzymatic activity. Interestingly, the peptidase activity of NRDC is not required for the enhancement of ADAM17 activity.<sup>22-24</sup> These previous findings prompted us to examine the role of NRDC in autoimmune arthritis, and we found that inhibition of NRDC had a protective effect against the disease. We have demonstrated here that macrophage-specific deletion of NRDC closely recapitulated the resistant phenotype of whole-body knockout mice in AAIA models. Using BMDMs, PMs and THP-1 cells, we showed that NRDC regulates ectodomain shedding of TNF- $\alpha$  in macrophages and monocytic cells. A deficiency of NRDC in all cell types resulted in a decrease in cleaved TNF- $\alpha$ , whereas TNF- $\alpha$  mRNA was not significantly affected. These results strongly suggest that NRDC in macrophages is involved in the pathogenesis of AAIA via the regulation of TNF- $\alpha$  shedding.

It was previously reported that the level of TNF- $\alpha$  is increased in RA synovial fluid.<sup>37</sup> We demonstrate here that the level of NRDC and TNF- $\alpha$  in synovial fluid is remarkably elevated in RA patients compared with that in OA patients. NRDC levels in RA synovial fluid correlated with synovial fluid TNF- $\alpha$  and serum CRP, suggesting that a local increase of NRDC in the RA joint may enhance TNF- $\alpha$  secretion, resulting in aggravation of joint

inflammation. These results suggest that NRDC plays important roles not only in mouse experimental models of RA but also in RA patients, and therefore that gene-targeted therapy with RNAi has potential in the treatment of RA.<sup>38</sup> Intra-articular injection of anti-TNF- $\alpha$  siRNA was shown to inhibit collagen-induced arthritis in mice,<sup>39</sup> and intravenous administration of double-stranded miR-451 reduced arthritis in SKG mice via downregulation of neutrophil chemotaxis.<sup>40</sup> We show here that intra-articular injection of anti-*Nrdc* siRNA ameliorated CAIA. In this experiment, arthritis was induced more rapidly than without intra-articular injection, probably because of the inflammatory response against injected reagents or trauma associated with the injection. This significant attenuation of arthritis by anti-*Nrdc* siRNA provided an indication of a novel treatment for RA. Although biologic TNF- $\alpha$  blockers are widely used for the management of RA patients, complete inhibition of TNF- $\alpha$  might cause adverse effects, mainly infections, because a physiological level of TNF- $\alpha$  secretion is necessary for the normal inflammatory response. As we show here, gene deletion or silencing of *NRDC* in macrophages and monocytic cells did not affect their baseline level of constitutive TNF- $\alpha$  production and secretion. Thus, anti-NRDC therapy could have the potential to reduce only excessive TNF- $\alpha$  secretion. In a similar sense, ADAM17 could be an alternative target for controlling TNF- $\alpha$  secretion. However, ADAM17 cleaves not only TNF- $\alpha$  but also a wide range of membrane proteins including EGF receptor ligands and Notch,<sup>41</sup> which may cause side effects of ADAM17 inhibitor treatment in the skin, intestine and liver.<sup>34,42</sup> In terms of substrates, NRDC also enhances ectodomain shedding of various membrane proteins including HB-EGF, neuregulin1 and amyloid precursor protein.<sup>22,24,29</sup> Other than TNF- $\alpha$ , several membrane proteins considered to be important for RA pathogenesis, such as TNF receptors and IL-6 receptors, are also functionally regulated by ectodomain shedding. NRDC-mediated regulation of the shedding of these molecules might contribute to the attenuation of arthritis by the deletion of NRDC.

Apart from its extracellular function as an enhancer of ectodomain shedding, a nuclear function of NRDC as a transcriptional coregulator has been recently clarified. The roles of NRDC in transcriptional regulation were illustrated by the interaction of NRDC with various nuclear proteins such as histone deacetylase 3, peroxisome proliferator-activated receptor gamma coactivator 1- $\alpha$  and *islet1*.<sup>43-45</sup> Intriguingly, the nuclear function of NRDC is crucially involved in adaptive thermogenesis<sup>44</sup> and insulin secretion in vivo.<sup>43</sup> Although we cannot exclude a possible involvement of nuclear NRDC in AAIA, we demonstrate here that mRNA levels of TNF- $\alpha$  and ADAM17 in macrophages are not affected by the inhibition of NRDC, suggesting that it is extracellular NRDC that mainly regulates AAIA.

Another limitation of this study is that we have not clarified NRDC function in neutrophils, which have a role in the pathogenesis of AAIA.<sup>33</sup> Because the *Ly2z*



gene is expressed not only in macrophages but also in neutrophils,<sup>46</sup> NRDC in neutrophils should be deleted in *Nrdc<sup>delM</sup>* mice.

In summary, our results demonstrated for the first time that NRDC is involved in AAIA at least partially via enhancement of TNF- $\alpha$  shedding in macrophages. NRDC levels in synovial fluid are increased in RA patients and correlate with systemic inflammation, providing a new insight into the pathogenesis of RA. In addition, we showed that *Nrdc* gene silencing ameliorates arthritis in mouse experimental models of RA, suggesting that NRDC could be a new therapeutic target for RA.

#### Author affiliations

<sup>1</sup>Department of Orthopaedic Surgery, Graduate School of Medicine, Kyoto University, Kyoto, Japan

<sup>2</sup>Department of Cardiovascular Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan

<sup>3</sup>Department of Pharmacology, Shiga University of Medical Science, Shiga, Japan

<sup>4</sup>Department of Tissue Regeneration, Institute for Frontier Medical Sciences, Kyoto University, Kyoto, Japan

<sup>5</sup>Department of Control for Rheumatic Diseases, Graduate School of Medicine, Kyoto University, Kyoto, Japan

<sup>6</sup>Department of Rheumatology and Clinical Immunology, Graduate School of Medicine, Kyoto University, Kyoto, Japan

**Acknowledgements** We gratefully acknowledge the patients who consented to synovial fluid and blood collection for this study. Detection of human NRDC in synovial fluid was supported by Rina Yamaguchi, Yuto Kurokawa and Yoshiyuki Amano (Sanyo Chemical Industries). We thank Sayaka Saijo, Shintaro Matsuda, Hideyuki Shibuya, Hiromi Iwai, Ryoko Nakanishi and Yoshiko Nakajima (Kyoto University) for their excellent help. We are also grateful to Diane Mathis and Christophe Benoist (Harvard Medical School) for providing us with KRN mice.

**Contributors** TF, HI and EN designed the studies. TF and EN wrote the manuscript. TF, HY, MO, KN, Yus M, Yug M, AO and TT performed the experiments and analysed the data. NO harvested K/BxN mice serum. TF, HI, MF and MA were involved in the analysis of human subjects. TK and SM supervised the works. All authors read and approved the final manuscript.

**Funding** This work was supported by Grants-in-Aid for Scientific Research (KAKENHI) (26293068, 25462368, 15K19513, 15K19376, 15H01557, 16K15216, 16K10890004). It was also supported by the Takeda Science Foundation and the Uehara Memorial Foundation.

**Competing interests** None declared.

**Patient consent** Obtained.

**Ethics approval** The ethics committee of Kyoto University Graduate School of Medicine.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** All available data can be obtained by contacting the corresponding authors.

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

#### REFERENCES

1. Terao C, Ikari K, Nakayamada S, *et al.* A twin study of rheumatoid arthritis in the Japanese population. *Mod Rheumatol* 2016;26:685–9.
2. Keffer J, Probert L, Cazlaris H, *et al.* Transgenic mice expressing human tumour necrosis factor: a predictive genetic model of arthritis. *Embo J* 1991;10:4025–31.
3. Müssener A, Litton MJ, Lindroos E, *et al.* Cytokine production in synovial tissue of mice with collagen-induced arthritis (CIA). *Clin Exp Immunol* 1997;107:485–93.
4. Ji H, Pettit A, Ohmura K, *et al.* Critical roles for interleukin 1 and tumor necrosis factor alpha in antibody-induced arthritis. *J Exp Med* 2002;196:77–85.
5. Williams RO, Feldmann M, Maini RN. Anti-tumor necrosis factor ameliorates joint disease in murine collagen-induced arthritis. *Proc Natl Acad Sci U S A* 1992;89:9784–8.
6. Kinne RW, Stuhl Müller B, Burmester GR. Cells of the synovium in rheumatoid arthritis. Macrophages. *Arthritis Res Ther* 2007;9:224.
7. Crotti TN, Dharmapatri AA, Alias E, *et al.* Osteoimmunology: major and costimulatory pathway expression associated with chronic inflammatory induced bone loss. *J Immunol Res* 2015;2015:1–13.
8. Iwata T, Ito H, Furu M, *et al.* Periarticular osteoporosis of the forearm correlated with joint destruction and functional impairment in patients with rheumatoid arthritis. *Osteoporos Int* 2016;27:691–701.
9. Kobayashi S, Murata K, Shibuya H, *et al.* A distinct human CD4+ T cell subset that secretes CXCL13 in rheumatoid synovium. *Arthritis Rheum* 2013;65:3063–72.
10. Takayanagi H. New developments in osteoimmunology. *Nat Rev Rheumatol* 2012;8:684–9.
11. Tanida S, Yoshitomi H, Nishitani K, *et al.* CCL20 produced in the cytokine network of rheumatoid arthritis recruits CCR6+ mononuclear cells and enhances the production of IL-6. *Cytokine* 2009;47:112–8.
12. Nielen MM, van Schaardenburg D, Reesink HW, *et al.* Specific autoantibodies precede the symptoms of rheumatoid arthritis: a study of serial measurements in blood donors. *Arthritis Rheum* 2004;50:380–6.
13. Song YW, Kang EH. Autoantibodies in rheumatoid arthritis: rheumatoid factors and anticitrullinated protein antibodies. *QJM* 2010;103:139–46.
14. Nandakumar KS, Svensson L, Holmdahl R. Collagen type II-specific monoclonal antibody-induced arthritis in mice: description of the disease and the influence of age, sex, and genes. *Am J Pathol* 2003;163:1827–37.
15. Terato K, Hasty KA, Reife RA, *et al.* Induction of arthritis with monoclonal antibodies to collagen. *J Immunol* 1992;148:2103–8.
16. Maccioni M, Zeder-Lutz G, Huang H, *et al.* Arthritogenic monoclonal antibodies from K/BxN mice. *J Exp Med* 2002;195:1071–7.
17. Matsumoto I, Staub A, Benoist C, *et al.* Arthritis provoked by linked T and B cell recognition of a glycolytic enzyme. *Science* 1999;286:1732–5.
18. Kagari T, Doi H, Shimozato T. The importance of IL-1 beta and TNF-alpha, and the noninvolvement of IL-6, in the development of monoclonal antibody-induced arthritis. *J Immunol* 2002;169:1459–66.
19. Ruuls SR, Hoek RM, Ngo VN, *et al.* Membrane-bound TNF supports secondary lymphoid organ structure but is subservient to secreted TNF in driving autoimmune inflammation. *Immunity* 2001;15:533–43.
20. Blaydon DC, Biancheri P, Di WL, WI D, *et al.* Inflammatory skin and bowel disease linked to ADAM17 deletion. *N Engl J Med* 2011;365:1502–8.
21. Chalaris A, Adam N, Sina C, *et al.* Critical role of the disintegrin metalloprotease ADAM17 for intestinal inflammation and regeneration in mice. *J Exp Med* 2010;207:1617–24.
22. Hiraoka Y, Ohno M, Yoshida K, *et al.* Enhancement of alpha-secretase cleavage of amyloid precursor protein by a metalloendopeptidase nardilysin. *J Neurochem* 2007;102:1595–605.
23. Hiraoka Y, Yoshida K, Ohno M, *et al.* Ectodomain shedding of TNF-alpha is enhanced by nardilysin via activation of ADAM proteases. *Biochem Biophys Res Commun* 2008;370:154–8.
24. Nishi E, Hiraoka Y, Yoshida K, *et al.* Nardilysin enhances ectodomain shedding of heparin-binding epidermal growth factor-like growth factor through activation of tumor necrosis factor-alpha-converting enzyme. *J Biol Chem* 2006;281:31164–72.
25. Chesneau V, Pierotti AR, Barré N, *et al.* Isolation and characterization of a dibasic selective metalloendopeptidase from rat testes that cleaves at the amino terminus of arginine residues. *J Biol Chem* 1994;269:2056–61.
26. Pierotti AR, Prat A, Chesneau V, *et al.* N-arginine dibasic convertase, a metalloendopeptidase as a prototype of a class of processing enzymes. *Proc Natl Acad Sci U S A* 1994;91:6078–82.
27. Nishi E, Prat A, Hospital V, *et al.* N-arginine dibasic convertase is a specific receptor for heparin-binding EGF-like growth factor that mediates cell migration. *EMBO J* 2001;20:3342–50.
28. Ishizu-Higashi S, Seno H, Nishi E, *et al.* Deletion of nardilysin prevents the development of steatohepatitis and liver fibrotic changes. *PLoS One* 2014;9:e98017.

29. Ohno M, Hiraoka Y, Matsuoka T, *et al.* Nardilysin regulates axonal maturation and myelination in the central and peripheral nervous system. *Nat Neurosci* 2009;12:1506–13.
30. Kanda K, Komekado H, Sawabu T, *et al.* Nardilysin and ADAM proteases promote gastric cancer cell growth by activating intrinsic cytokine signalling via enhanced ectodomain shedding of TNF- $\alpha$ . *EMBO Mol Med* 2012;4:396–411.
31. Ohno M, Hiraoka Y, Lichtenthaler SF, *et al.* Nardilysin prevents amyloid plaque formation by enhancing  $\alpha$ -secretase activity in an Alzheimer's disease mouse model. *Neurobiol Aging* 2014;35:213–22.
32. Prevoe ML, van 't Hof MA, Kuper HH, *et al.* Modified disease activity scores that include twenty-eight-joint counts. Development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. *Arthritis Rheum* 1995;38:44–8.
33. Christensen AD, Haase C, Cook AD, *et al.* K/BxN serum-transfer arthritis as a model for human inflammatory arthritis. *Front Immunol* 2016;7:213.
34. Issuree PD, Maretzky T, McIlwain DR, *et al.* iRHOM2 is a critical pathogenic mediator of inflammatory arthritis. *J Clin Invest* 2013;123:928–32.
35. McIlwain DR, Lang PA, Maretzky T, *et al.* iRhom2 regulation of TACE controls TNF-mediated protection against Listeria and responses to LPS. *Science* 2012;335:229–32.
36. Baumann U, Köhl J, Tschernig T, *et al.* A codominant role of Fc gamma RI/III and C5aR in the reverse Arthus reaction. *J Immunol* 2000;164:1065–70.
37. Tetta C, Camussi G, Modena V, *et al.* Tumour necrosis factor in serum and synovial fluid of patients with active and severe rheumatoid arthritis. *Ann Rheum Dis* 1990;49:665–7.
38. Pauley KM, Cha S. RNAi therapeutics in autoimmune disease. *Pharmaceuticals* 2013;6:287–94.
39. Khoury M, Courties G, Fabre S, *et al.* Adeno-associated virus type 5-mediated intraarticular administration of tumor necrosis factor small interfering RNA improves collagen-induced arthritis. *Arthritis Rheum* 2010;62:765–70.
40. Murata K, Yoshitomi H, Furu M, *et al.* MicroRNA-451 down-regulates neutrophil chemotaxis via p38 MAPK. *Arthritis Rheumatol* 2014;66:549–59.
41. Arribas J, Esselens C. ADAM17 as a therapeutic target in multiple diseases. *Curr Pharm Des* 2009;15:2319–35.
42. DasGupta S, Murumkar PR, Giridhar R, *et al.* Current perspective of TACE inhibitors: a review. *Bioorg Med Chem* 2009;17:444–59.
43. Nishi K, Sato Y, Ohno M, *et al.* Nardilysin is required for maintaining pancreatic  $\beta$ -cell function. *Diabetes* 2016;65:3015–27.
44. Hiraoka Y, Matsuoka T, Ohno M, *et al.* Critical roles of nardilysin in the maintenance of body temperature homeostasis. *Nat Commun* 2014;5:3224.
45. Li J, Chu M, Wang S, *et al.* Identification and characterization of nardilysin as a novel dimethyl H3K4-binding protein involved in transcriptional regulation. *J Biol Chem* 2012;287:10089–98.
46. Clausen BE, Burkhardt C, Reith W, *et al.* Conditional gene targeting in macrophages and granulocytes using LysMcre mice. *Transgenic Res* 1999;8:265–77.



## Nardilysin is involved in autoimmune arthritis via the regulation of tumour necrosis factor alpha secretion

Takayuki Fujii, Eiichiro Nishi, Hiromu Ito, Hiroyuki Yoshitomi, Moritoshi Furu, Namiko Okabe, Mikiko Ohno, Kiyoto Nishi, Yusuke Morita, Yugo Morita, Masayuki Azukizawa, Akinori Okahata, Takuya Tomizawa, Takeshi Kimura and Shuichi Matsuda

*RMD Open* 2017 3:

doi: 10.1136/rmdopen-2017-000436

---

Updated information and services can be found at:  
<http://rmdopen.bmj.com/content/3/1/e000436>

---

*These include:*

### References

This article cites 46 articles, 17 of which you can access for free at:  
<http://rmdopen.bmj.com/content/3/1/e000436#BIBL>

### Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

### Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Notes

---

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>