TITLE:

Thenar Dysplasia in Radial Polydactyly Depends on the Level of Bifurcation

AUTHOR(S):

Saito, Susumu; Ueda, Maho; Murata, Mai; Suzuki, Shigehiko

CITATION:

Saito, Susumu ...[et al]. Thenar Dysplasia in Radial Polydactyly Depends on the Level of Bifurcation. Plastic and Reconstructive Surgery 2018, 141(1): 85e-90e

ISSUE DATE:

2018-01-01

URL:

<http://hdl.handle.net/2433/228917>

RIGHT:

This is a non-final version of an article published in final form in (provide complete journal citation); The full-text file will be made open to the public on 01 January 2019 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

“Thenar Dysplasia in Radial Polydactyly Depends on the Level of Bifurcation.”

Author list: Susumu **Saito**, MD, PhD¹; Maho **Ueda**, MD¹; Mai **Murata**, MD¹; Shigehiko **Suzuki**, MD, PhD¹.

1. Department of Plastic and Reconstructive Surgery, Graduate School of Medicine and Faculty of Medicine, Kyoto University.

Corresponding author:

Susumu Saito MD, PhD.

Yoshida-Konoe-cho, Sakyo-ku

Kyoto, Japan

Postal code: 606-8501

Telephone number: +81-75-751-3613

Fax number: +81-75-751-4340

e-mail address: susumus@kuhp.kyoto-u.ac.jp

Financial Disclosure Statement: No funding was received for this article.

This study is registered in UMIN Clinical Trials Registry

Trial registry name: “カスタムメイド超音波 3D 画像による体表解剖に関する研究および臨床診断における有効性に関する研究” (Studies on superficial anatomies of human body by 3D ultrasound)

Registration identification number: UMIN000014396

URL: https://upload.umin.ac.jp/cgi-open-bin/ctr/ctr_view.cgi?recptno=R000016753

This study was accepted to be orally presented at 4th European Symposium on Pediatric Hand Surgery and Rehabilitation, which will be held in Paris, France in June 15-16, 2017.

Short Running Head

Thumb Dysplasia in Radial Polydactyly

1 Thenar Dysplasia in Radial Polydactyly Depends on the Level of Bifurcation

2

3 Abstract

4

5 **Background**

6 Little is known about thenar dysplasia in radial polydactyly, other than that thenar hypoplasia occasionally
7 occurs in radial polydactyly with triphalangism. In particular, the phenotype and level of duplication
8 associated with thenar dysplasia remains unclear.

9 **Methods**

10 The abductor pollicis brevis (APB) and flexor pollicis brevis (FPB) muscles were visualized using
11 three-dimensional ultrasound and their horizontal geometry was assessed using a biaxial level
12 classification system. Subjects were categorized into three phenotypes according to the developmental
13 condition of the radial thumb. The relationship between the level of distribution of the muscles and the level
14 of the bifurcation of the radial thumb was investigated.

15 **Results**

16 Nineteen patients with radial polydactyly without triphalangism were included. There were ten patients with
17 the non-floating type, three with the floating type, and six with the rudimentary type. All patients with
18 bifurcation at or more distal to the metacarpophalangeal (MCP) joint had normal thenar muscle distribution,

19 but the muscles in patients with bifurcation at or more proximal to the level of the metacarpals were
20 confined, regardless of phenotype. The level of muscle distribution was strongly correlated with the level of
21 the bifurcation of the radial thumb.

22 **Conclusions**

23 These findings suggest that formation of the thenar muscles in the longitudinal direction in radial polydactyly
24 might depend on the level of bifurcation of the radial thumb. The presence of thenar dysplasia even in
25 floating-type or rudimentary-type duplications is of clinical and etiological importance.

26

27 **Key Words**

28 Radial polydactyly, thenar muscle, dysplasia, ultrasound

29

30 **Introduction**

31 For radial polydactyly, reconstructive surgery has been widely accepted as a result of increased knowledge
32 about associated skeletal and soft tissue abnormalities¹⁻⁴. However, even though all these anomalies are
33 addressed intraoperatively, complete functional restoration of the thumb remains challenging^{5, 6}. In 1982,
34 Miura stated issues remaining after reconstruction for radial polydactyly are adduction contracture of the
35 thumb and thenar dysplasia⁷. Although more than 30 years have passed since Miura's statement, little is
36 known about thenar dysplasia in radial polydactyly apart from the fact that thenar hypoplasia occurs in
37 special cases of radial polydactyly with triphalangism⁸.

38 The objective of this study was to investigate developmental conditions of the thenar muscles in radial
39 polydactyly without triphalangism using three-dimensional (3D) ultrasonography. In particular, the
40 relationship between thenar dysplasia and the level of bifurcation of the radial thumb was evaluated.

41 **Materials and methods**

42 This study began in 2014 after it was approved by the ethics committee of our institution. The patients'
43 families received an explanation about the 3D ultrasound examination and surgery, and written informed
44 consent was obtained. Triphalangeal-thumb polydactyly or polysyndactyly, and a six-fingered hand were
45 not included because these types were known to have distinct pathogenic backgrounds^{9, 10}.

46 The 3D ultrasound method was described previously¹¹. Briefly, the thenar region was scanned using a
47 linear array transducer (Preirus; Hitachi Aloka Medical Co., Ltd., Tokyo, Japan) in the transverse direction

under precise speed regulation at 3 mm/s. During scan, the thumb was placed in the 45-degree radially-abducted position. Consecutive images were acquired at 0.2 mm intervals and processed to reconstruct 3D data using imaging software (ImageJ, U. S. National Institutes of Health (NIH), Bethesda, Maryland, USA). Anatomical identifications were accomplished on the basis of multi-planar observation, initially using coronal sections, followed by sagittal and axial sections. Especially for coronal sections, the outlines of the first metacarpal were memorized to be shown in more superficial sections as anatomical references. (Supplementary figure. 1).

To systematically investigate developmental conditions of the thenar muscles in patients with radial polydactyly of various degrees of severity and phenotypes, subjects were categorized according to the level of bifurcation and the developmental condition of the radial thumb. For bifurcation, nine levels were defined along the phalanges and metacarpal (Fig. 1). For developmental condition of the radial thumb, three phenotypes were defined. The non-floating type had a true articulation or an osseous continuity between the radial and ulnar thumbs. The floating type had a radial thumb connected to the ulnar thumb by fibrous tissues and at least one pair of articulated phalanges in the radial thumb. The rudimentary type had a more underdeveloped radial thumb that is comprised of soft tissues only or contains no more than one osseous component. To analyze horizontal geometry of the abductor pollicis brevis (APB) and flexor pollicis brevis (FPB), a biaxial level classification system was further defined (Fig. 1). Distributions of the insertion and origin of the FPB were classified by levels along the metacarpal axis and levels along the longitudinal axis,

respectively. The APB distribution was assessed only on the insertion side by levels along the metacarpal axis. The relationship between the level of muscle distribution and the level of the bifurcation of the radial thumb was analyzed using Pearson's correlation coefficients for samples bifurcating at Levels 1–7 because the levels from Level 1 to Level 7 were aligned regularly.

Results

Consecutive nineteen patients were included in this study. There were no bilateral cases. Additional examinations using nine unaffected hands of infants verified that the APB on the insertion side, FPB on the insertion side, and FPB on the origin side located at Levels 6, 6, and 5, respectively. Scatter plots of the distribution of the thenar muscles are shown in Fig. 2. In all hands with Level 6 or more distal bifurcation, muscle distribution was within the normal range for both APB and FPB regardless of phenotypic differences in the radial thumb (Supplementary figure 2). In contrast, in hands with Level 5 or more proximal bifurcation, the level of muscle distribution was strongly related to the level of the bifurcation, especially for APB and FPB on the insertion side. Notably, even the origin side of the FPB was distributed proximally according to the proximity of the bifurcation of the radial thumb, especially in cases with Level 3 or more proximal bifurcation. Representative cases were presented in Supplementary figure 3-6.

Discussion

In this study, a strong correlation between the level of APB and FPB distribution and the level of bifurcation of the radial thumb was found. Herein, we propose that etiologically, muscle formation in the

84 longitudinal direction is limited by the level of bifurcation. Surprisingly, proximal confinement of the thenar
85 muscles was found even in patients with a metacarpal duplication of the floating or rudimentary type,
86 suggesting that the mechanism we proposed may apply to different phenotypes of radial polydactyly,
87 although the presence of pathogenic factors in the transverse direction is speculated (Fig. 3). These results
88 provide two-fold clinical implications. One is that aberrant APB and FPB insertion can exist even in floating
89 and rudimentary types. The other is that longitudinal underdevelopment of the thenar muscles can influence
90 surgical outcomes. Lourie et al. reported unusual cases showing a zigzag deformity after ligation treatment
91 for an extra pedunculated thumb with at the radial aspect of the metacarpal¹². Similarly, we experienced
92 APB underdevelopment in cases of a rudimentary extra thumb with metacarpal bifurcation (Supplementary
93 figure 5 and 6). Although APB atrophy observed in the postoperative case (Supplementary figure 5)
94 indicates the possibility that detachment of the APB insertion might have occurred during surgery, APB
95 underdevelopment shown in the non-treated case (Supplementary figure 6) rather suggests that APB
96 hypoplasia originally existed. It is known that APB insufficiency causes joint deformities, thus potential APB
97 hypoplasia might be associated with secondary joint deformities in Wassel type V and VI^{13, 14}. Further
98 studies on surgical outcomes for radial polydactyly using assessment systems involving palmar abduction¹⁵,
99 ¹⁶ would clarify the clinical importance of the thenar dysplasia observed in this study.

101

102 Authors' role/participation in the authorship of the manuscript

103 SS conducted the project of this study and performed all examinations. SS, MM and MU analyzed the data.

104 All the authors contributed to the writing of the paper.

105

106

107 References

- 108 1. Marks TW and Bayne LG. Polydactyly of the thumb: Abnormal anatomy and treatment. *J Hand Surg.*
109 1978; 3:107-116.
- 110 2. Tada K, Yonenobu K, Tsuyuguchi Y, Kawai H, Egawa T. Duplication of the thumb. A retrospective review
111 of two hundred and thirty-seven cases. *J Bone Joint Surg Am.* 1983; 65:584-98.
- 112 3. Miura T. An appropriate treatment for postoperative Z-formed deformity of the duplicated thumb. *J Hand*
113 *Surg.* 1977; 2:380-386.
- 114 4. Salama R, Weissman SL. Congenital bilateral anomalous band between flexor and extensor pollicis
115 longus tendons. Report of a case. *Hand.* 1975; 7:25-26.
- 116 5. Kawabata H, Tada K, Masada K, Kawai H, Ono K. Revision of residual deformities after operations for
117 duplication of the thumb. *J Bone Joint Surg Am.* 1990; 72:988-998.
- 118 6. Naasan A, Page RE. Duplication of the thumb. A 20-year retrospective review. *J Hand Surg Br.* 1994;
119 19:355-360.
- 120 7. Miura T. Duplicated thumb. *Plast Reconstr Surg.* 1982; 69:470-481.
- 121 8. Graham JM, Jr., Brown FE and Hall BD. Thumb Polydactyly as a Part of the Range of Genetic
122 Expression for Thenar Hypoplasia. *Clin Pediatr (Phila).* 1987; 26:142-148.
- 123 9. Wieczorek D, Pawlik B, Li Y, Akarsu NA, Caliebe A, May KJ, et al. A specific mutation in the distant sonic

- 124 hedgehog (SHH) cis-regulator (ZRS) causes Werner mesomelic syndrome (WMS) while complete ZRS
- 125 duplications underlie Haas type polysyndactyly and preaxial polydactyly (PPD) with or without triphalangeal
- 126 thumb. *Hum Mutat.* 2010; 31:81-89.
- 127 10. Sun M, Ma F, Zeng X, Liu Q, Zhao XL, Wu FX, et al. Triphalangeal thumb-polysyndactyly syndrome and
- 128 syndactyly type IV are caused by genomic duplications involving the long range, limb-specific SHH
- 129 enhancer. *J Med Genet.* 2008; 45:589-595.
- 130 11. Saito S, Ueda M, Takahashi N, Kawakatsu M, Suzuki S. Three-dimensional Ultrasonography for
- 131 Visualization of Muscular Anomalies in Type VI and VII Radial Polydactyly. *Skel Radiol.* 2016; 45:541-547.
- 132 12. Lourie GM, Costas BL, Bayne LG. The zig-zag deformity in pre-axial polydactyly. A new cause and its
- 133 treatment. *J Hand Surg Br.* 1995; 20:561-564.
- 134 13. CH Yen, WL Chan, HB Leung, KH Mak. Thumb polydactyly: clinical outcome after reconstruction. *J*
- 135 *Orthop Surg.* 2006;14 :295-302.
- 136 14. Ogino T, Ishii S, Takahata S, Kato H. Long-term results of surgical treatment of thumb polydactyly. *J*
- 137 *Hand Surg Am.* 1996; 21:478-486.
- 138 15. Iba K, Horii E, Ogino T, Kazuki K, Kashiwa K. The Classification of Swanson for Congenital Anomalies
- 139 of Upper Limb Modified by the Japanese Society for Surgery of the Hand (JSSH). *Hand Surg.* 2015;
- 140 20:237-250.
- 141 16. Dijkman RR, van Nieuwenhoven CA, Selles RW, Habenicht R, Hovius SE. A multicenter comparative

142 study of two classification systems for radial polydactyly. *Plast Reconstr Surg.* 2014; 134:991-1001.

143

144 Figure legends

145

146 Fig. 1

147 The biaxial level classification system and its application

148 Seven levels are defined for the thenar region along two anatomical axes. In addition to the seven levels,

149 two levels are defined distally in order to classify the level of bifurcation of the radial thumb (top). An

150 ultrasound coronal image of a non-floating-type duplication with Level 5 bifurcation (bottom). In this case,

151 the APB on the insertion side, FPB on the insertion side, and FPB on the origin side all distribute at Levels 5.

152 APB, abductor pollicis brevis; FPB, flexor pollicis brevis; CMC, carpometacarpal; MCP,

153 metacarpophalangeal; IP, interphalangeal.

154

155 Fig. 2

156 Scatter plots and lines of best fit showing relationships between distributions of APB and FPB versus the

157 level of bifurcation. Pearson's correlation coefficients for patients with bifurcation at the level of the

158 metacarpals are also shown. Yellow bars indicate normal muscle distributions. APB, abductor pollicis

159 brevis; FPB, flexor pollicis brevis.

160

161 Fig. 3

162 Schematic illustration showing a proposed mechanism of thenar dysplasia associated with radial
163 polydactyly.

164

165 Supplementary figure 1

166 Ultrasound images showing normal thenar anatomy in a 1-year-old infant. **A**, coronal; **B**, sagittal; **C**, axial
167 sections. Lines with numbers indicate the location of the images. Green lines indicate the outline of the first
168 metacarpal. Yellow lines show the outline of the thenar muscles. FPL is used as a landmark that separates
169 the superficial and deep part of the muscles (Red circles). APB, abductor pollicis brevis; AD, adductor
170 pollicis; FPB, flexor pollicis brevis; FPL, flexor pollicis longus; OP, opponens pollicis; CT, carpal tunnel; CMC,
171 carpometacarpal joint; MCP, metacarpophalangeal joint; 1st MC, first metacarpal; d, depth from the skin.

172

173 Supplementary figure 2

174 Photos, radiographs, and coronal ultrasound images of patients with radial polydactyly bifurcating at Level 7
175 or more distally. Yellow lines show the outline of the superficial thenar muscles. Note that all the muscles
176 distribute distally to the level of the metacarpophalangeal joint (red arrowheads). Abbreviations are as
177 indicated in Fig. 2.

178

179 Supplementary figure 3

180 A 6-month-old patient with a non-floating-type duplication with Level 1 bifurcation. **A**, The thenar eminence
181 seems hypoplastic (right). Note that the proximal thenar crease is lacking (arrow). **B**, Radiograph. **C**,
182 Coronal; **D**, axial; and **E**, sagittal ultrasound sections. The thenar muscles are confined proximally to Level
183 2 (red arrowheads). The distal portion of the muscles is located in the direction of the MCP joint of the radial
184 thumb (yellow arrowheads). Yellow lines show the outline of the thenar muscles. Red circles indicate the
185 FPL in a cross-sectional view. Abbreviations are as indicated in Fig. 2. L and R indicate left and right,
186 respectively.

187

188 Supplementary figure 4

189 A 1-year-old patient with a floating-type extra thumb bifurcating at Level 3. **A**, The thenar eminence seems
190 slightly hypoplastic (right, arrow). **B**, Radiograph. **C**, Coronal; **D**, axial; and **E**, sagittal ultrasound sections.
191 The thenar muscles are confined proximally to Level 3 (red arrowheads). Note that there is no muscle
192 distribution in the direction of the MCP joint of the ulnar thumb (yellow arrowheads). Yellow lines show the
193 outline the thenar muscles. Red circles indicate the FPL in a cross-sectional view. Abbreviations used are
194 indicated in Fig. 2. L and R indicate left and right, respectively.

195

196 Supplementary figure 5

197 A 4-year-old patient who underwent ablative surgery for rudimentary-type duplication with Level 4

198 bifurcation four years previously. **A**, The thenar eminence seems underdeveloped radially (arrows). Palmar
199 abduction of the affected thumb is insufficient (bottom, right). **B**, Preoperative appearance (left) and
200 radiograph (right). **C**, Coronal; **D**, sagittal; and **E**, axial ultrasound sections. Axial sections at the level of the
201 epiphyseal line demonstrate APB atrophy (red arrowheads) although FPB is shown is less underdeveloped
202 (yellow arrowheads). Red circles indicate the FPL in a cross-sectional view. Abbreviations used are
203 indicated in Fig. 2. L and R indicate left and right, respectively.

204

205 Supplementary figure 6

206 A 14-year-old patient with untreated rudimentary-type duplication with Level 2 bifurcation. **A**, The thenar
207 eminence seems underdeveloped (top, black arrow). The involved thumb shows a deformity in the flexion
208 position (bottom, white arrow). **B**, Radiograph. **C**, Coronal; **D**, sagittal; and **E**, axial ultrasound sections. The
209 thenar muscles are confined proximally to Level 2 (red arrowheads). Axial sections at the level of the
210 epiphysis demonstrate muscle atrophy (yellow arrowheads). Yellow lines show the outline of the thenar
211 muscles. Red circles indicate the FPL in a cross-sectional view. Abbreviations used are indicated in Fig. 2. L
212 and R indicate left and right, respectively.

Fig. 1

Fig. 2

Fig. 3

Supplementary figure 1

Supplementary figure 2

Supplementary figure 3

Supplementary figure 4

Supplementary figure 5

Supplementary figure 6